



Temporal Trends and Geographic Comparisons



Hood Canal - 1999




1999 Hood Canal Summary


- Three *possibly impacted* samples, one in Port Gamble and two in Dabob Bay.
- The Benthic Index showed the poorest quality in over half of the samples.
- Weighted mean Triad Index (wmTriad) decreased from 71.3 in 1999 to 62.9 in 2004.

Triad Index	
Category	Score
Unimpacted	>81-100
Likely unimpacted	>57-81
Possibly impacted	>35-57
Likely impacted	>5-35
Clearly impacted	0-5
Inconclusive	


Chemistry Index




Toxicity Index




Benthic Index






Temporal Trends and Geographic Comparisons



Hood Canal - 2004




2004 Hood Canal Summary


- Five *possibly to likely impacted* samples, three in Dabob Bay, one in central Hood canal, and one near Liliwaup.
- The Benthic Index showed the poorest quality in over half of the samples.
- Weighted mean Triad Index (wmTriad) decreased from 71.3 in 1999 to 62.9 in 2004.

Triad Index	
Category	Score
Unimpacted	>81-100
Likely unimpacted	>57-81
Possibly impacted	>35-57
Likely impacted	>5-35
Clearly impacted	0-5
Inconclusive	

Chemistry Index



Toxicity Index



Benthic Index

