ISSUED BY
RPP-WTP PDC
6·6·00
INIT DATE

BNFL Inc.

Document title:

Final Work Plan for Screening Level Risk Assessment for the RPP-WTP

Contract deliverable:	B-1-35 / B-1-40
Department:	Environmental, Safety and Health Programs
Contract title:	RPP-WTP
Contract number:	DE-AC06-96RL13308
Author(s):	Various
Principal Author Signature:	
Document Number:	RPT-W375-EN00001, Rev. 1
Checked by:	J. R. Cook
Principal Checker Signature:	
Date of issue:	28 April 2000
Issue status:	Approved
Approved by:	O. Lee Bostic
Approver's position:	Environmental Safety Manager

Approver's signature

BNFL Inc.
RPP-WTP
3000 George Washington Way
Richland, WA 99352
United States of America
Tel: 509 371 3000
Fax: 509 371 3001

DM

BNFL Inc.

Circulation List

Purpose of issue: 1 For comment

2 For action
3 For information

Name	Location	No of copies	Remarks	Purpose of issue
Ian Wheeler	B101	1		3
Lee Bostic	P123	1		3

Please note that source, special nuclear, and byproduct materials, as defined in the Atomic Energy Act of 1954 (AEA) are regulated at the U. S. Department of Energy (DOE) facilities exclusively by DOE acting pursuant to its AEA authority. DOE asserts that pursuant to AEA, it has sole and exclusive responsibility and authority to regulate source, special nuclear, and byproduct materials at DOE-owned nuclear facilities. Information contained herein on radionuclides is provided for process description purposes only.

History Sheet

Rev	Date	Reason for revision	Revised by
A	12/98	Draft	Donald W. Edwards
B	3/24/99	Draft	Donald W. Edwards
C	5/07/99	Draft	Donald W. Edwards
D	11/15/99	Draft	Donald W. Edwards
E	3/1/00	Draft	O. Lee Bostic
0	3/31/00	Issue	O. Lee Bostic

Contents

Item — Page Number

Acronyms and Abbreviations viii

1. Introduction 1-1
2. Risk Assessment Approach 2-1
3. Engineering Description 3-1
   - 3.1. RPP-WTP Overview 3-1
   - 3.2. Pretreatment 3-2
   - 3.3. LAW Vitrification 3-6
   - 3.4. HLW Vitrification 3-9
   - 3.5. Cascade Ventilation System 3-13
   - 3.6. Fugitive Emissions 3-13
   - 3.7. Upset Conditions 3-14
   - 3.8. Facility Control Philosophy 3-15
4. Screening Level Risk Assessment and Constituents of Potential Concern 4-1
   - 4.1. Preliminary Chemicals of Potential Concern (COPCs) 4-2
   - 4.2. Preliminary Radionuclides of Potential Concern (ROPCs) 4-5
   - 4.3. Selection of Final COPCs/ROPCs for the Quantitative Preliminary Risk Assessment 4-5
   - 4.4. Summary for Identification of COPCs and ROPCs 4-9
5. Quantification of Emissions 5-1
   - 5.1. Emission Sources 5-1
   - 5.2. Emission Assumptions for the Preliminary Risk Assessment 5-1
   - 5.3. Measured Emission Rates for the Final Risk Assessment 5-10
   - 5.4. Uncertainties in Estimating Emission Rates 5-13
   - 5.5. Summary for Quantification of Emissions 5-16
6. Environmental Modeling 6-1
   - 6.1. Air Dispersion Modeling 6-1
   - 6.2. Soil Accumulation Modeling 6-10
   - 6.3. Terrestrial Plant and Animal Modeling 6-12
   - 6.4. Surface Water and Fish Bioaccumulation Modeling 6-16
   - 6.5. Uncertainties 6-21

Contents

Item — Page Number

6.6. Summary for Environmental Modeling 6-23

7. Screening Human Health Risk Assessment 7-1

7.1. Exposure Assessment 7-1

7.2. Toxicity Assessment 7-16

7.3. Risk Characterization 7-28

7.4. Uncertainty Assessment 7-30

7.5. Summary for Screening Human Health Risk Assessment 7-31

8. Screening Ecological Risk Assessment 8-1

8.1. Problem Formulation 8-1

8.2. Exposure Assessment 8-23

8.3. Effects Assessment Calculations 8-48

8.4. Risk Equations 8-53

8.5. Reporting of Major Ecological Risk Findings 8-54

8.6. Uncertainties in Ecological Risk Assessment 8-55

8.7. Summary for Screening Ecological Risk Assessment 8-58

9. Risk-Based Emissions Criteria 9-1

9.1. Human Health Risk-Based Emission Levels 9-1

9.2. Ecological Receptor-based Emissions Levels 9-1

10. Summary 10-1

11. References 11-1

Appendix A Constituents of Potential Concern Lists ..... A-i

Appendix B Chemical-Specific Parameters and Supporting Equations for Human Health Risk Assessment ..... B-i

Appendix C Chemical-Specific Parameters for Ecological Risk Assessment ..... C-i

Appendix D Selection of COPCs and ROPCs for Human Health and Ecological Risk Assessment ..... D-i

Appendix E COPCs and ROPCs for Which There are Emission Rates ..... E-i

Contents

Item | Page Number

FIGURES

Figure 2-1. Overview of Screening Level Risk Assessment Process for RPP-WTP Air Emissions 2-3
Figure 2-2. Overview of Screening Level Risk Assessment Process During the Preliminary Risk Assessment and the Final Risk Assessment for RPP-WTP Air Emissions 2-4
Figure 3-1. Location of the RPP-WTP on the Hanford Site 3-17
Figure 3-2. RPP-WTP Layout 3-18
Figure 3-3. Process Flow Diagram 3-19
Figure 4-1. Selection of Constituents of Potential Concern 4-10
Figure 6-1. Exposure Grids in Each concentric Zone 6-24
Figure 6-2. Sources of COPC and ROPC Concentrations in Soil 6-25
Figure 6-3. Sources of Constituents of Potential Concern Concentration in Plants 6-26
Figure 6-4. Generic Sources of Constituents of Potential Concern Concentrations in Animal Tissue, Eggs, and Dairy Products 6-27
Figure 6-5. Sources of Constituents of Potential Concern Concentrations in Tissue, Eggs, and Dairy Products of Domestic Farm Animals 6-28
Figure 6-6. Sources of Constituents of Potential Concern Concentrations in Tissue of Game Animals 6-29
Figure 6-7. Sources of Constituents of Potential Concern Concentrations in Fish 6-30
Figure 7-1. Human Health Conceptual Exposure Model 7-32
Figure 7-2. Exposure Distances (Far-Field) from the RPP-WTP at the Hanford Site 7-33
Figure 7-3. Exposure Distances (Near-Field) from the RPP-WTP at the Hanford Site 7-34
Figure 7-4 Hanford Site Existing Land Use Map – 1996 7-35
Figure 7-5 Hanford Site Projected Land Use Map – 2046 7-36
Figure 8-1. Ecological Conceptual Exposure Model 8-59
Figure 8-2. Recreation and Wildlife Areas and the Hanford Reach 8-60
Figure 8-3. Vegetation Types of the Hanford Site 8-61
Figure 8-4. Selected Raptor Nesting and Perching Locations on the Hanford Site 8-62
Figure 8-5. RPP-WTP Areas Vegetation Types (Simplified) 8-63
Figure 8-6. Fall Chinook Salmon Spawning Areas Along the Columbia River 8-64
Figure 8-7. Trophic Levels and Receptor Species Considered in Development of the 200 Area and Vicinity Terrestrial Conceptual Exposure Model 8-65
Figure 8-8. Trophic Levels and Receptor Species at the Columbia River Considered in Development of the 200 Area and Vicinity Aquatic Conceptual Exposure Model 8-66
Figure 8-9. Trophic Levels and Selected Receptor Species (names in bold) from the 200 Area and Vicinity Terrestrial Conceptual Exposure Model 8-67
Figure 8-10. Trophic Levels and Selected Receptor Species (names in bold) at the Columbia River Considered in Development of the 200 Area and Vicinity Aquatic Conceptual Exposure Model 8-68
Figure 8-11. Relationship Between Sources and Biotransfer Factors for Calculating Terrestrial and Aquatic Exposures 8-69

Contents

Item | Page Number

TABLES

Table 4-1. Human Health Chemical Toxicity Values for Organic Constituents of Potential Concern 4-11
Table 4-2. Human Health Toxicity Values for Inorganic Constituents of Potential Concern 4-23
Table 4-3. Human Health Toxicity Values for Radionuclides of Potential Concern 4-25
Table 4-4. Acute Inhalation Exposure Criteria for COPCs 4-27
Table 6-1. Specific Variables for Plant Uptake Modeling[1] 6-31
Table 6-2. Specific Variables of Uptake into Animal Products[a] 6-32
Table 7-1 Exposure Scenarios 7-37
Table 7-2 Exposure Parameters for Residential Scenarios 7-40
Table 7-3 Summary of Exposure Parameters for Native American Subsistence Scenarios[a] 7-46
Table 7-4 Summary of Exposure Parameters for Hanford Site Worker Scenario 7-53
Table 8-1. Policy Goals, Ecological Assessment Endpoints, Measures, and Decision Rules for 200 Areas and Surroundings 8-70
Table 8-2. Soil Ingestion Rates For Hanford Receptors 8-74
Table 8-3. Derivation of Soil-to-Body Weight Ratios for Prey of Hanford Receptors 8-75
Table 8-4. Derivation of Indirect Soil Ingestion Rate for Hanford Receptors 8-76
Table 8-5 Toxicity Equivalence Factors for Chlorinated Dioxins, Chlorinated Dibenzofurans, and PCBs[a] 8-77

Acronyms and Abbreviations

ADD	average daily dose
AE	absorption efficiency
AIEC	acute inhalation exposure criteria
ALI	annual limit on intake
APCS	air pollution control system
ASIL	acceptance source impact levels
AT	averaging time
ATSDR	Agency for Toxic Substances and Disease Registry
AUF	area-use factor
BAF	bioaccumulation factor
BAF-S	soil-to-tissue transfer factor
BAF-T	tissue-to-tissue transfer factor
BAS-F	sediment to tissue uptake factor
BCF	bioconcentration factors
BPIP	Building Profile Input Program
BSF	sediment-to-tissue uptake factor
BTU	British Thermal Unit
CAA	*Clean Air Act*
CAS	Chemical Abstract Service
CETRED	Combustion Emissions Technical Resource Document
COPC	constituent of potential concern and chemicals of potential concern
DEHP	bis(2-ethylhexyl)phthalate
DF	decontamination factors
DNA	deoxyribonucleic acid
DOE	U.S. Department of Energy
DOE-RL	U.S. Department of Energy, Richland Operations Office
DQO	Data Quality Objective
DRE	destruction and removal efficiency
DSCM	dry standard cubic meter
DST	double-shell tank
Ecology	Washington State Department of Ecology
ED	duration of exposure
EDL	estimated detection limit
EF	exposure frequency
EPA	U.S. Environmental Protection Agency
ERA	ecological risk assessment
ESU	evolutionarily significant units
FEALE	Fitzner-Eberhardt Arid Lands Ecology Reserve
FRA	final risk assessment
HAMMER	Hazardous Materials Management and Emergency Response
HAP	hazardous air pollutants
HAZWRAP	Hazardous Waste Remedial Actions Program
HEAST	Health Effects Assessment Summary Tables
HEME	high efficiency mist eliminator
HEPA	high efficiency particulate air
HHRA	human health risk assessment
HI	hazard index

Acronyms and Abbreviations

HLW	high-level waste
HQ	hazard quotient
IAEA	International Atomic Energy Agency
IHLW	immobilized high-level waste
ILAW	immobilized low-activity waste
IRIS	Integrated Risk Information System
ISCST3	Industrial Source Complex Short Term Model, Version 3
K_{ow}	octanol/water partition coefficient
LADD	lifetime average daily dose
LAW	low-activity waste
LIGO	Laser Interferometer Gravitational Wave Observatory
LOAEL	lowest-observed adverse effect level
MACT	Maximum Achievable Control Technology
MDL	method detection limit
MF	metabolism factor
MPRM	Meteorological Processor for Regulatory Models
MSL	mean sea level
MW	molecular weight
NAAQS	National Ambient Air Quality Standard
NaOH	sodium hyrdoxide
NAWQC	National Ambient Water Quality Criteria
NCEA	National Center for Environmental Assessment
NOAEL	no-observed adverse effect level
NWS	National Weather Surface
PAH	polycyclic aromatic hydrocarbon
PCB	polychlorinated biphenyl
PCDD	polychlorinated dibenzo-p-dioxin
PCDF	polychlorinated dibenzofuran
PIC	product of incomplete combustion
PM_{10}	particulates less than 10 microns in diameter
PODC	principal organic dangerous constituent
PRA	preliminary risk assessment
RDL	reported detection limit
RfC	reference concentration
RfD	reference dose
RFD	reverse flow diverters
RME	reasonable maximum exposure
RMP	risk management program
RPF	relative potency factor
RPP-WTP	River Protection Project - Waste Treatment Plant
SBS	submerged bed scrubber
SCR	selective catalytic reduction
SLRA	screening level risk assessment
SO_x	sulfur oxide
SP	slope factor
SP	soil-to-plant transfer factor
SRE	system removal efficiency

Acronyms and Abbreviations

SVOC	semi-volatile organic compound
T&E	threatened and endangered
TEF	toxicity equivalency factor
TEQ	toxicity equivalent
TIC	tentatively identified compound
TNC	The Nature Conservancy
TOC	total organic carbon
TOE	total organic emission
TRU	transuranic
TRV	toxicity reference value
TSD	treatment, storage, or disposal
TUF	temporal-use factor
TWINS	Tank Waste Information Network System
TWRS	Tank Waste Remediation System
UHC	Underlying Hazardous Constituents
UR	unit risk
USFWS	U.S. Fish and Wildlife Service
VOC	volatile organic compound
VSL	Vitreous State Laboratory
WDFW	Washington Department of Fish and Wildlife
WESP	wet electrostatic precipitator
Work Plan	*Final Work Plan for Screening Level Risk Assessment for the RPP-WTP*
WPPSS	Washington Public Power Supply System

1. Introduction

This document presents the risk assessment work plan (hereafter referred to as the Work Plan). The Work Plan addresses the risk assessment protocol for assessing risk to human health and the environment for the River Protection Project - Waste Treatment Plant (RPP-WTP) at the Hanford Site. The Hanford Site is located in southeastern Washington state, owned by the US Government, and operated by the US Department of Energy (DOE). The proposed facility will include two waste melter and vitrification facilities as well as pretreatment and offgas systems, and will be built in the 200 East Area of the Hanford Site.

The Work Plan establishes the criteria and methodology for conducting a risk assessment to identify potential human health and ecological risks associated with processing Hanford tank waste into a stable glassified form. This risk assessment will be used to establish waste feed acceptance limits as necessary to meet defined risk limits of operation of the RPP-WTP goals.

The Work Plan provides a description of key components that will be used in the screening level risk assessment (SLRA). The emphasis of the work is aerial dispersion of emissions and the potential for risk to human and ecological receptors.

The Work Plan contains a brief statement of the risk assessment approach (section 2) and an engineering description of the RPP-WTP (section 3). Sections 4-8 present the steps involved in both the human health and ecological screening risk assessments including the following:

- Identification of constituents of potential concern (COPCs) – section 4
- Quantification of airborne emissions – section 5
- Environmental modeling of the dispersed airborne emissions – section 6
- Screening human health risk assessment – section 7
- Screening ecological risk assessment –section 8.

Section 9 presents the methodology for establishing risk-based emissions criteria. Section 10 summarizes each area of the screening level risk assessment (SLRA). References are included in section 11 and are followed by data appendices A, B, C, D, and E.

2. Risk Assessment Approach

This section describes the overall programmatic screening level risk assessment (SLRA) approach (Figure 2-1) that will be used to support the *RPP-WTP Dangerous Waste Permit Application* (BNFL 1999a). The primary regulatory guidance followed for this assessment is *Human Health Risk Assessment Protocol for Hazardous Waste Combustion Facilities* (EPA 1998a). The major components of the SLRA process for airborne emissions will include the following:

- Work Plan for the SLRA – This Work Plan is submitted with the Dangerous Waste Permit Application and establishes the methodology for the future implementation of the SLRA. The preliminary risk assessment (PRA) and final risk assessment (FRA) are subparts of the SLRA, as described in the Work Plan.

- Preliminary risk assessment – The PRA will be performed while the Dangerous Waste Permit Application is being reviewed and before final approval of the permit.

- Final risk assessment – The scope of the FRA will be similar to the PRA. However, modeled emission rates will be replaced with the results of the environmental performance demonstration.

The PRA and FRA will address the potential operating life of the facility. The current Dangerous Waste Permit Application covers operations through the year 2018. However, the SLRA will initially assume the facility will operate at maximum capacity for the entire design life of the facility (40 years). This assumption will result in the SLRA addressing more than the portion of the total radioactivity (25 percent) and volume (10 percent) of Hanford tank waste that will be treated during the first stage of the RPP-WTP pretreatment and vitrification process. Risk from the remaining waste as well as cumulative risk from the Hanford Site, are outside the scope of the SLRA. Only 25 percent and 10 percent volumes are to be treated first. The general technical procedure for the SLRA is provided in Figure 2-2.

The Work Plan focuses on the PRA because it must be completed before issuance of the Dangerous Waste Permit (BNFL 1999a). The FRA will be performed when the results of the environmental performance demonstration become available. Based on the results of the environmental performance demonstration, the FRA may involve rerunning models, modeling additional chemicals, or changing model parameters. Information that may be updated in the FRA includes:

- toxicity data based on reevaluation of toxicity data sources identified in the work plan
- addition of compounds newly identified during the environmental performance demonstration
- emission rates updated based on environmental performance demonstration results
- stack gas parameters (i.e., flow rates, temperatures) updated based on environmental performance demonstration
- receptor locations based on land use changes, if any
- and address any errors in the PRA

The FRA reruns may also be dictated by the findings of the PRA.

As described in section 7, Screening Human Health Risk Assessment, and Section 4, Constituents of Potential Concern, there will be two exposure scenarios: worst-case exposures and plausible exposures.

The primary difference between these scenarios is the assumed location of the receptor populations. For the worst-case scenarios, receptor populations are assumed to be present at the location where emission concentrations are estimated to be greatest regardless of the likelihood of their presence. For the plausible scenarios, receptors are assumed to be located in areas where similar receptors are currently present or may be expected in the future. The scenarios are intended to provide a better understanding of the range of potential risks to human and ecological receptors under any land use condition at various points in time.

This approach is consistent with the *Environmental Protection Agency Risk Characterization Policy* (EPA 1995c) which directs the "use of several descriptors, rather than a single description, to enable the Environmental Protection Agency (EPA) to present a fuller picture of risk that corresponds to the range of different exposure conditions encountered by various individuals and populations."

Participants in the SLRA process are:

- BNFL Inc.
- US Department of Energy
- Washington State Department of Ecology (Ecology)
- EPA Region X
- Yakama Nation
- Confederated Tribes of the Umatilla Reservation
- Nez Perce Tribe
- The general public

Figure 2-1. Overview of Screening Level Risk Assessment Process for RPP-WTP Air Emissions



Hanford ERA - 001

Figure 2-2. Overview of Screening Level Risk Assessment Process During the Preliminary Risk Assessment and the Final Risk Assessment for RPP-WTP Air Emissions



Hanford ERA - 002

3. Engineering Description

The River Protection Project Waste Treatment Plant (RPP-WTP) is being developed to treat mixed waste stored at the US Department of Energy, Richland Operations Office (DOE-RL) Hanford Site. Waste treatment will be performed using a pretreatment system and two independent vitrification systems. Offgas generated by the pretreatment and vitrification processes will be treated in independent offgas treatment systems. This section provides an overview of the mixed waste treatment processes that will be used in the RPP-WTP. More detailed process information can be found in Chapter 4, Section 4.2, of the RPP-WTP Dangerous Waste Permit Application (BNFL Inc. 2000a).

The mixed waste to be treated in the RPP-WTP is consistent with the definition of high level waste found in 10 CFR 72.3. The mixed waste is further classified into a low activity waste (LAW) feed and a high level waste (HLW) feed. The LAW feed consists primarily of the aqueous phase supernatant containing soluble solids presently stored in the Hanford tank system. The HLW feed is primarily an aqueous slurry with a higher solids content than the LAW feed. The composition of the liquid fraction of the HLW feed is similar to LAW feed, whereas the HLW feed solids consist of precipitated metals, and radioisotopes of uranium, plutonium, and strontium.

Two vitrification systems are being designed to immobilize the radioactive waste: one for LAW feed and the second for HLW feed. Waste will be immobilized in the form of glass monoliths contained in metal containers. These IHLW and ILAW containers, and other containers of various secondary wastes generated during treatment operations, will be temporarily stored at the RPP-WTP and then transferred to other permitted treatment, storage, or disposal (TSD) facilities.

3.1. RPP-WTP Overview

BNFL Inc. has entered into a contract with DOE-ORP for pretreatment and vitrification of radioactive waste stored in underground tanks at the Hanford Site. Underground storage tanks located at the Hanford Site consist of single shell and double shell tanks operated by the DOE Office of River Protection. Waste contained in the single shell tanks will be transferred to double shell tanks prior to being transferred to the RPP-WTP. The LAW feed and HLW feed will be retrieved and transferred, by DOE or its agent, from the double shell tanks to facilities designed, built, and operated by BNFL Inc. for pretreatment and vitrification. Vitrified immobilized waste and secondary waste generated by the processes will be transferred to permitted storage, and/or disposal units for disposition.

DOE is leasing property at the Hanford Site to BNFL Inc. for construction and operation of the RPP-WTP. The leased land lies at the eastern end of the 200 East Area of the Hanford Site, near the former Grout Treatment Facility, 241-AP Tank Farm Complex, and Plutonium Uranium Extraction Plant (PUREX). The location of the RPP-WTP on the Hanford Site is shown in Figure 3-1.

As shown in Figure 3-2, the RPP-WTP will include several primary buildings. Four main process buildings will contain most of the dangerous waste management operations, and include major areas for pretreating HLW and LAW feed, and immobilizing tank waste in the LAW vitrification plant and HLW vitrification plant. Other smaller support buildings such as the wet chemical storage building, the glass-former storage building, the central waste storage area, and two melter storage buildings will provide for storage or transfer of materials used in the treatment process, and for storage of wastes.

The waste treatment plant will store and treat HLW feed and LAW feed from the Hanford Site DST system unit. The treatment processes are designed to pretreat both LAW feed and HLW feed, immobilize waste in a glass matrix through vitrification, and treat the offgas. Two separate melter or vitrification

processes will be used: one melter process will produce ILAW, and the other will produce IHLW. Figure 3-3 provides simplified process flow diagrams of the treatment processes.

3.2. Pretreatment

The RPP-WTP will contain processes for pretreating and immobilizing both LAW feed and HLW feed, as indicated in the simplified process flow diagram shown in Figure 3-3. Characterized LAW and HLW feeds will be transferred from the Hanford tank system to the pretreatment plant. All waste transfers will be made through double-contained transfer pipes with leak-detection systems.

The following four waste types, or envelopes will be treated in the RPP-WTP.

- Envelope A. This LAW feed envelope will contain cesium and technetium at concentrations high enough to warrant removal during pretreatment in order to ensure the immobilized LAW glass meets applicable requirements.

- Envelope B. This LAW feed envelope will contain higher concentrations of cesium than Envelope A. Both cesium and technetium must be removed to comply with the contract specifications for ILAW. This envelope will also contain concentrations of chlorine, chromium, fluorine, phosphates, and possibly sulfates, than those found in Envelope A, and which may limit the rate of waste incorporation into the glass.

- Envelope C. This LAW feed envelope will contain organically complexed strontium and transuranic (Sr/TRU) compounds that will require removal in a processing step unique to this waste envelope. As with Envelopes A and B, cesium contract ILAW specifications can be met.

- Envelope D. HLW feed envelope will be in the form of a slurry containing 10-200 grams of solids per liter of slurry. The liquid fraction of the slurry will be comprised of Envelope A, B, or C waste, and the solid fraction will be Envelope D waste. Envelope D waste will be delivered via pipeline into a receipt vessel or vessels located in the pretreatment process building.

LAW feed and HLW feed will first be treated in the pretreatment system. As shown in Figure 3-3, the pretreatment system will be divided into LAW and HLW components.

3.2.1. LAW Pretreatment

Pretreatment of the LAW feed will occur in two plants: the pretreatment plant and the LAW pretreatment plant. Process steps include reducing radionuclide concentrations and reducing the volume of waste being fed to the LAW vitrification system. In addition, in order to ensure that the limits for the concentrations of the following constituents in the ILAW glass will be met, the LAW pretreatment process will remove entrained solids, Sr and TRU compounds (Envelope C only), Cs and Tc. The removed constituents, with the exception of entrained solids, will be incorporated into the HLW melter feed. Entrained solids will either be returned to another permitted unit located at the Hanford Site or will be incorporated in the HLW or LAW melter feed.

3.2.1.1. LAW Feed Receipt

LAW feed will be pumped from the Hanford tank system via a double-walled pipeline into the LAW feed receipt vessels. These six receipt vessels are identical in design and are located in one cell. Two identical pipelines, one in operation and the other on standby, will be used for the transfer. Each pipeline will be

equipped with a leak detection system. Both rejected LAW feed and entrained solids from LAW ultrafiltration may be returned to the Hanford tank system via the same pipelines.

Each feed receipt pipeline can be routed to any of the six feed receipt vessels. Prior to a transfer, four of the vessels will be designated as the "active" receptors of the LAW feed. The remaining two vessels will be available to provide contingency space to avoid emergency transfers back to the Hanford tank system, in the event of a leak, overflow, or spill from the other vessels. During the transfer, no operations such as mixing may occur in the active vessels, nor may any other transfers be made to or from the active vessels. After delivery completion, the transferred feed volume will be confirmed, and the feed will be sampled and analyzed. Depending on the results of the analysis, the feed will be either transferred to LAW feed evaporation or transferred back to the Hanford tank system. Rejection of a completed feed transfer should never occur, but the option is left open, if confirmation sampling and analyses show that the waste is significantly different than indicated by previous data.

3.2.1.2. LAW Feed Evaporator

The LAW feed stream will be evaporated as needed in the feed evaporator to provide a consistent feed composition for ultrafiltration, in order to minimize the amount of reagents used for strontium and TRU removal (in envelope C) and to optimize ion-exchange performance. The LAW feed evaporator will be a continuous forced circulation vacuum evaporator that will concentrate the feed. The LAW feed will be recirculated through the evaporator reboiler until the sodium content of the stream reaches the desired concentration. The concentrated LAW feed will then be pumped to buffer tanks. The vapor stream from the evaporator will be condensed and routed to the process condensate tanks. If the condensate exceeds radioactivity limits, it will be routed to a contaminated condensate vessel and then back to the LAW feed evaporator. If it does not exceed radioactivity limits, it will be routed to a process condensate vessel, from which it may be routed to the Hanford Liquid Effluent Retention Facility/Effluent Treatment Facility (LERF/ETF), recycled as process water or routed to the contaminated condensate vessel.

3.2.1.3. Ultrafiltration

Following evaporation, the concentrated LAW feed will be sent to the ultrafiltration process to separate entrained solids and, in the case of envelope C, to remove strontium and TRU.

The entrained solids for all envelopes will be concentrated, washed to remove soluble components, and then sampled and analyzed to determine their final destination. For envelopes A and B, the permeate from separation of the entrained solids will be transferred to the cesium ion-exchange process.

After the entrained solids have been removed from envelope C wastes, strontium and TRU elements will be removed by precipitation. The resulting precipitate will be concentrated and washed in the ultrafiltration process before being routed to HLW feed receipt and storage. The separated Sr and TRU precipitate will be mixed with HLW feed and with pretreated cesium and technetium eluates (see section 3.2.1.5 for technetium eluate) from the ion-exchange process. The waste feed will then be incorporated into the HLW melter feed.

Ultimately, the concentrated entrained solids from envelopes A, B and C will either be transferred to a Hanford Site TSD unit or incorporated into the HLW or LAW melter feed.

3.2.1.4. Ion Exchange

To meet ILAW specifications, the radioactive cesium content of the LAW feed must be reduced. Permeate from ultrafiltration will enter the ion exchange system where the cesium concentration will be reduced by passing the feed through successive ion exchange systems.

The system will have two sets of ion exchange columns arranged in parallel and one final ion exchange column. Each set will consist of two columns arranged in series, one in the lead position and the second in the lag position. One set from each system will process feed, while elution and regeneration take place on the lead column of the other set. When cesium loading on the lead column has reached approximately 50 % cesium breakthrough, or when any significant breakthrough is seen after the lag column, the flow to that set of columns will be suspended and the LAW feed diverted to the other set of columns. The final, or trailing, column will function as a third column in series for either pair of main columns during loading.

At this stage, cesium will be removed from the lead column using a nitric acid solution and the resins will be regenerated for reuse. The resin will have a finite useful life, after which it will be removed from the columns and replaced. Spent resin will ultimately be transferred to a Hanford Site TSD facility for disposal. Ion exchange media or resins are discussed further in Chapter 4 of the RPP-WTP DWPA.

Using evaporation, nitric acid will be recovered from the cesium eluate taken from the ion-exchange column. This recovery will allow reuse of recovered nitric acid and will also reduce the storage volume of the intermediate waste. An evaporation system will be used to concentrate the cesium eluate. The resultant concentrates will be stored prior to blending with the pretreated HLW feed, the separated strontium and TRU, and the technetium concentrate (as discussed below).

3.2.1.5. LAW Pretreatment Plant

Following the removal of cesium, the LAW feed will be transferred to the LAW pretreatment plant for removal of technetium in an ion exchange system. The technetium removal system will consist of four ion exchange columns, arranged as two sets of two columns in parallel. One set will operate in the loading cycle while the other set is being regenerated. One set will process feed, while elution and regeneration take place on the lead column of the other set.

When technetium loading on the lead column has reached approximately 50 % breakthrough or a significant proportion of the constituent has been detected in the effluent from its respective lag column, the two columns will be taken offline and the resin in the lead column will be regenerated. At this stage technetium will be eluted with water and the resins will be regenerated for reuse. The resultant technetium concentrates will routed back to the pretreatment plant for where it will be stored prior to blending with cesium concentrates, the pretreated HLW feed, and the separated strontium and TRU.

Resins used in the ion exchange columns cannot be regenerated indefinitely. Therefore, the resin must periodically be removed from the ion exchange columns and replaced. The spent resin collection and dewatering system fluidizes the ion exchange columns using a flush liquor that hydraulically discharges the contents into spent resin collection vessels. Here, the resin slurry will be circulated, monitored for technetium content, and delivered to a sampling system to determine if the resin meets Hanford Site acceptance criteria.

Spent resin that does not meet the criteria will be routed back to the ion exchange columns for re-elution. Spent resin that meets the criteria will be transferred to dewatering containers where water will be

removed and will ultimately be transferred to a Hanford Site TSD facility for disposal. Liquor from the dewatering process will be reused as flush liquor for the ion exchange columns.

Following the ion exchange step, the LAW vitrification feed preparation system will concentrate the waste by evaporation, as needed, to reduce the volume that must be processed through the LAW melters. Concentration of the product will be controlled to avoid precipitation of soluble compounds.

The concentrated waste feed will be discharged to the concentrated LAW melter feed holding tank. The solution will then be transferred to a receipt tank or tanks in the LAW vitrification building. The condensate from the LAW melter feed evaporator will be combined with condensate from the LAW feed evaporator, characterized, and either recycled within the process or transferred to the Hanford Liquid Effluent Retention Facility/Effluent Treatment Facility (LERF/ETF).

3.2.2. HLW Pretreatment

HLW feed will be transferred from the Hanford tank system to the HLW feed receipt tanks. Following receipt, the HLW feed will be sent to the ultrafiltration vessels to separate HLW solids. In the ultrafiltration vessels, solids will be concentrated, and washed (using either process water or caustic solution) to remove soluble components (mainly Na salts). The ultrafiltration permeate will be transferred to the LAW pretreatment system and treated in the same manner as the LAW feed. The solids concentrate from the ultrafilter will be stored in the HLW feed lag storage vessels. The washed sludge will be combined with Cs, Tc, and Sr and TRU removed in the LAW pretreatment system, and will be transferred to the HLW vitrification system.

3.2.3. Pretreatment Emissions Control

Gaseous emissions will be produced within the pretreatment plant and the LAW pretreatment plant, from fluid transfer devices and agitators, exhausters, evaporator overheads, ejector transfers, and filling or emptying of vessels. Pretreatment emissions will be passed into the vessel ventilation system and treated in an emissions treatment system, which will consist of:

- High efficiency mist eliminators (HEMEs), to remove entrained droplets and particulates.
- Counter-current scrubber, to remove acid gases. Excess scrubber solution will be transferred to the LAW pretreatment system.
- High efficiency particulate air (HEPA) filtration with preheater.

The LAW pretreatment plant vessel ventilation system will include a VOC oxidation unit, HEPA filters, and carbon bed adsorbers. Primary functions of this system include:

- Collect offgas from LAW pretreatment vessel vents
- Remove particulates from vent system offgas
- Prevent condensation in the HEPA filters
- Adsorb and/or oxidize organic gases
- Collect exhaust streams from LAW pretreatment system RFD and PJM jet pumps

The combined LAW pretreatment plant treated vessel ventilation offgas and treated RFD and PJM exhaust streams will be routed to the Pretreatment plant. Pretreatment and LAW pretreatment cleaned

emissions streams will be monitored for radionuclides and other constituents, and released through the pretreatment building stack.

3.3. LAW Vitrification

The pretreated LAW feed will be processed in three melters. Offgas produced as a result of waste treatment will be treated in an offgas treatment system. The following sections address the LAW vitrification system, offgas treatment systems, waste feed cutoff system, and control devices. Figure 3-3 depicts the vitrification and offgas treatment systems.

3.3.1. LAW Vitrification System

The primary functions of the LAW vitrification system are to:

- Convert blended waste slurry and glass-forming chemical additives into molten glass
- Discharge molten glass to metal containers

A blend of glass-forming chemical additives will be used in the LAW glass recipe. For example, silica, alumina, boric acid, calcium silicate (wollastonite), ferric oxide, lithium carbonate, magnesium silicate (olivine), zircon sand, and zinc oxide will be used. Each glass recipe will be prepared from formulations, developed by Vitreous State Laboratory (VSL), that were found to produce durable glass (Matlock and Pegg 1999). These chemical additives will be stored in silos prior to use. Reductant (sucrose) will be added with the glass-forming chemicals to enhance melter performance.

From the storage silos, dry chemicals will be weighed and transferred into a pneumatic blending silo, which will use compressed air to blend a batch of dry chemicals. Each dry chemical will be weighed, then transferred to the blend hopper. After blending, the glass-formers will be transferred to a feed hopper within the main facility and held until they are required for use. Weighing will be used to confirm dry chemical transfers. Each melter will have a glass-former feed hopper sized for preparation of a melter feed batch. The entire contents of the feed hopper are used in the preparation of a melter feed batch.

Following process related sampling and analysis, the concentrated waste will be combined with glass forming chemicals and then mixed. This material will be transferred as slurry to the LAW melter feed tanks in order to provide continuous feed to each of the three LAW melters. Three electric powered, joule heated LAW melters will operate in parallel. Each melter will have a nominal throughput capacity of 10 metric tons and a maximum of 16.6 tons of glass per day. The operating temperature of the melter will be between 950 °C and 1,250 °C. As the LAW feed enters the melter, it will form a cold cap of dried solids on the surface of the molten glass. An air agitation system will be used to incorporate the cold cap into the melt and mix the melt to enhance melter performance. The outer shell of the melter containing the molten glass bath will be water cooled in order to minimize migration of molten glass within the melter refractory and to reduce the heat load to the melter cave.

Feed components will be converted to their respective oxides, and form the glass melt. As these materials are heated, gases will be released into the melter offgas system. Molten glass will be discharged into stainless steel containers. The containers will be cooled, welded closed, decontaminated, stored, and finally transferred to a permitted disposal unit. The process will yield a durable glass containing immobilized LAW.

3.3.2. LAW Offgas Treatment System

The principal components of the offgas will be nitrogen and oxygen, which will result from the controlled introduction of air into the melter, the operation of the air agitation system, and the chemical decomposition reactions occurring in the melter. Water in the feed will be evaporated or flashed to steam in the melter and condensed in the offgas system. Also contained within the offgas will be radionuclides, primarily as aerosols, nitrogen oxides (NO_x) generated from the decomposition of nitrates in the waste feed, organic compounds, chloride, fluoride, and sulfur oxides.

The LAW offgas treatment system will treat normal offgas flows and intermittent surges of up to seven times the normal flow rate for condensable gases, such as steam, and up to three times the flow rate for noncondensable gases.

The LAW offgas treatment system will be composed of four components: a primary melter offgas treatment system, a secondary offgas treatment system, a vessel ventilation treatment system, and a standby offgas system.

3.3.2.1. Primary LAW Melter Offgas Treatment System

The purpose of the primary melter offgas system is to cool the offgas and remove the aerosols generated by each melter. It will consist of the following components:

- Film cooler
- Submerged bed scrubber (SBS)
- Wet electrostatic precipitator (WESP)

The following paragraphs describe these components in greater detail.

Film cooler.

Offgas will exit the melt chamber and be mixed with compressed air in the LAW melter offgas film cooler. The film cooler will be an integral part of the melter, and will consist of a double-walled pipe designed to introduce steam and/or air (or a mixture of steam and air) along the walls of the offgas pipe through a series of holes or slots in the inner wall. This process will provide a film of air (or mixture of steam and air) flow along the pipe wall to cool the offgas. It will also ensure the offgas maintains a high velocity, to minimize the deposition of aerosols. A water spray ring and a mechanical reamer inserted into the offgas discharge line from the melter will be provided to remove buildup, if necessary. Dislodged particulates will either fall back into the melt pool, or be carried into the offgas stream and removed by the SBS.

Submerged bed scrubber.

The submerged bed scrubber (SBS) will remove entrained gross particulate, soluble material and organic components from the offgas, and will also cool the offgas prior to subsequent treatment. Offgas will be scrubbed by drawing it into a packed bed covered with liquid. It will enter the SBS through an inlet pipe that will run down through the center of the packed bed. Gas bubbles formed underneath the packing support plate will be broken up and scrubbed as the gas rises through the packing. Steam generated in the melter will condense in the submerged bed scrubber, and the resulting water overflow will be directed to the LAW offgas condensate collection vessel. The offgas condensate will be recycled to the pretreatment system.

Wet electrostatic precipitator.

After larger aerosols have been removed in the submerged bed scrubber, the cooled offgas will be routed to a wet electrostatic precipitator (WESP). Ionizing electrodes located inside tubes within the WESP will create strong electrical fields that will charge aerosols entrained in the offgas stream. The charged aerosols will be removed from the offgas as they adhere to the oppositely-charged tube walls. The aerosols will be washed from the tube walls and collected in the sump, which will drain into the WESP drain.

3.3.2.2. Secondary LAW Offgas Treatment System

The secondary LAW offgas treatment system will consist of the following components:

- HEPA filtration
- Thermal catalytic oxidation unit
- Selective catalytic reduction (SCR) unit
- Caustic Scrubber

The following paragraphs describe these components in greater detail.

High-efficiency particulate air filtration.

After leaving the WESP, the offgas will pass through the HEPA preheater to raise its temperature above the dewpoint in order to prevent condensation as it passes through HEPA filtration. HEPA filters are capable of removing particulates 0.3 micrometer in diameter and larger, with an efficiency of 99.97%.

Thermal catalytic oxidation unit.

The offgas will pass through a heater to raise its temperature to the desired operating range for removal of volatile organics in a catalytic oxidation unit. This unit will use a precious metal catalyst to facilitate the decomposition of organics to carbon dioxide (CO_2) and water vapor. The treated offgas will be discharged to the SCR unit.

Selective catalytic reduction unit.

Gaseous NO_x will be removed in a SCR unit and converted to nitrogen and water, using ammonia as the reducing agent. The reaction will occur in catalyst beds within the SCR column. Depending on inlet gas concentration, it will be possible to achieve high NO_x removal efficiencies. The catalyst will be replaced periodically to ensure that offgas treatment adheres to design specifications. Careful control of the ammonia addition will be necessary to minimize residual ammonia in the offgas stream.

Caustic scrubber

Offgas exiting the SCR unit will enter a caustic scrubber. Contaminants in the offgas stream will be absorbed into a caustic liquid stream flowing counter-current to the offgas. The caustic scrubbing liquid will remove acid gases and carbon-14 (as carbon dioxide) from the offgas. These contaminants will be absorbed into the liquid stream through interaction of the gas, liquid, and scrubber column packing media. To neutralize the collected acid gases, sodium hydroxide solution will be added to the top of the scrubbing column and will serve as the scrubbing liquid. The scrubbing liquid will be recirculated

through the column. After scrubbing, the treated offgas will be monitored and released through the LAW building stack.

3.3.2.3. LAW Vessel Ventilation System

Within the LAW vitrification plant, gaseous emissions will be produced from vessel vents and various equipment related to the LAW melter system. All vessel ventilation lines will be connected to the main vessel ventilation header. This main header will join the primary offgas treatment system lines downstream from the WESP and immediately upstream from the preheater for the secondary offgas system HEPA filters. The combined melter offgas streams and the vessel ventilation offgas stream will then go into the LAW secondary offgas treatment system described above.

3.3.2.4. Alternate Route

Infrequently, each melter may experience a reduction in vacuum, which will actuate an alternate route into the offgas treatment system. The following are abnormal situations that would require diversion:

- Plugging of the air film cooler line from the melter to the submerged bed scrubber
- Melter surge in excess of the design basis of seven times the volume of condensable gases and three times noncondensable gases

Automated gas pressure monitoring, alarm and control devices will detect the above conditions. Each LAW melter will have an alternate route into the offgas treatment system. The alternate route will consist of a fast-actuating butterfly valve and a redundant bypass film cooler line from the melter to the submerged bed scrubber.

In the event of a plug developing in the line from the melter to the submerged bed scrubber, or a surge in offgas resulting in reduction of vacuum below the minimum normal set point, the butterfly valve will open the alternate route to the submerged bed scrubber. On the alternate route, air and steam will flow through the bypass film cooler to cool the offgas and minimize the deposition of solids on the pipe wall.

A general power failure is estimated to occur on average once in 10 years. In such a situation, the standby generators may take from 15 to 30 seconds to come online. During this transition to standby power, the offgas exhaust fans will be without power, and offgas would leak from the melter to the cave. Ventilation air for the melter cell will be subject to all C5 ventilation maximum contamination controls, and will be HEPA filtered to prevent or control the release of emissions during or following such an occurrence. (See 3.5 Cascade Ventilation System.)

3.3.3. Automatic Waste Feed Cutoff System

The automatic waste feed cutoff system will prevent the introduction of LAW feed into the melter when key process conditions fall outside of the predetermined range. As a minimum, the automatic waste feed cutoff system will automatically lock out operation of the waste feed system until proper operating conditions are restored.

3.4. HLW Vitrification

The pretreated HLW feed will be vitrified in one melter. The HLW vitrification plant is being designed with a second HLW melter cave for the installation of a second melter if increased waste processing is

warranted. Offgas produced as a result of HLW vitrification will be cleaned in an offgas treatment system. The following sections address the HLW vitrification system, offgas treatment system, waste feed cutoff system, and control devices. Figure 3-3 depicts the HLW vitrification and offgas treatment systems.

3.4.1. HLW Vitrification System

The primary functions of the HLW vitrification system are to:

- Convert blended waste slurry and glass-forming chemical additives into molten glass
- Discharge molten glass to metal containers

A blend of glass-forming chemical additives will be used in the HLW glass recipe. For example, silica, boric acid, calcium silicate (wollastonite), ferric oxide, and lithium carbonate will be used. Not all of these additives will be used in each glass recipe. Each glass recipe will be prepared from formulations, developed by Vitreous State Laboratory (VSL), that were found to produce durable glass (Matlock and Pegg 1999).

The glass-forming chemicals will be stocked in storage silos. Each dry chemical will be conveyed from its storage silo, weighed, and then transferred into a pneumatic blending silo where compressed air will be used to blend dry chemicals for the HLW melter. After blending, the glass-forming chemicals will be transferred to the feed hopper located within the main facility until they are required for use. A reductant (sucrose) will be added with the glass-forming chemicals to enhance melter performance. Weighing will be used to confirm dry chemical transfers, once the chemicals have been placed in the feed hopper. The entire contents of the feed hopper will be used in the preparation of a melter feed batch.

Waste feed will be transferred to the HLW melter feed vessel, from which it will be fed directly to the HLW melter. The HLW melter will have a nominal throughput capacity of 1.5 metric tons per day and a maximum capacity of 3 metric tons per day. The operating temperature of the melter will be between 950 °C and 1,250 °C. As the HLW feed enters the melter, it will form a cold cap of dried solids on the surface of the molten glass. An air bubbler agitation system will be used to incorporate the cold cap into the melt and mix the melt to enhance melter performance. The steel outer shell of the melter will be water cooled to minimize migration of molten glass within the melter refractory and to reduce the heat load to the melter cave.

The feed components will undergo chemical reaction, be converted to their respective oxides, and form the glass melt. As these materials are heated, gases will be released into the melter offgas system. Molten glass will be discharged to stainless steel containers for cooling, solidification, and storage. The process will yield a durable glass containing immobilized HLW.

3.4.2. HLW Offgas Treatment System

The HLW melter offgas will consist of steam, products of chemical decomposition reactions that occur in the melter, organic compounds, and radionuclides. Its volume will be smaller than that of the LAW melter offgas and will contain little NO_x; however, it will have a considerably higher radioactivity level.

The HLW offgas treatment system will treat normal offgas flows and intermittent surges of up to seven times the normal flow rate for condensable gases, such as steam, and up to three times the normal flow rate for noncondensable gases. The system will be composed of four components: a primary melter

offgas treatment system, a secondary offgas treatment system, a vessel ventilation treatment system, and a standby offgas system.

3.4.2.1. Primary HLW Melter Offgas Treatment System

The primary HLW melter offgas treatment system will consist of the following components:

- Film cooler
- SBS
- WESP
- HEME
- HEPA filtration

The following paragraphs describe these components in greater detail.

Film cooler.

Offgas will exit the melt chamber and be mixed with compressed air in the HLW melter offgas film cooler. The film cooler will be an integral part of the melter, and will consist of a double-walled pipe designed to introduce steam and/or air (or a mixture of steam and air) along the walls of the offgas pipe through a series of holes or slots in the inner wall. This process will provide a film of air (or mixture of steam and air)flow along the pipe wall to cool the offgas. It will also ensure the offgas maintains a high velocity, to minimize the deposition of aerosols. A water spray ring and a mechanical reamer inserted into the offgas discharge line from the melter will be provided to remove buildup, if necessary. Dislodged particulates will either fall back into the melt pool, or be carried into the offgas stream and removed by the SBS.

Submerged bed scrubber.

The SBS will remove entrained gross particulate, soluble material and organic components from the offgas, and will also cool the offgas prior to subsequent treatment. Offgas will be scrubbed by drawing it into a packed bed covered with liquid. It will enter the SBS through an inlet pipe that will run down through the center of the packed bed. Gas bubbles formed underneath the packing support plate will be broken up and scrubbed as the gas rises through the packing. Steam generated in the melter will condense in the submerged bed scrubber, and the resulting water overflow will be directed to the HLW offgas condensate collection vessel. The offgas condensate will be recycled to the pretreatment system.

Wet electrostatic precipitator.

After the larger aerosols have been removed in the SBS, the cooled offgas will be routed to a WESP. Ionizing electrodes located inside tubes within the WESP will create strong electrical fields that will charge aerosols entrained in the offgas stream. The charged aerosols will be removed from the offgas as they adhere to the oppositely charged tube walls. The aerosols will be washed from the tube walls and collected in the sump, which will drain into the WESP drain collection tank.

High efficiency mist eliminator.

The HEME, a high efficiency demister with a removal efficiency of about 99% for aerosols down to submicron size, will further remove aerosols from the melter offgas stream. Aerosols adhere to fiber-packing filaments within the HEME by surface tension. As the droplets grow they acquire enough mass

to fall by gravity to the bottom of the unit, where they will be collected and drain into the HEME collection tank.

High efficiency particulate air filtration.

From the HEME, the offgas will pass through the HEPA preheater, where it will be heated to above the dewpoint in order to prevent condensation within the HEPA filters. The HEPA filters are capable of removing particulates of 0.3 micron and greater, with an efficiency of 99.97%. The cleaned offgas will be discharged to the secondary HLW offgas treatment system.

3.4.2.2. Secondary HLW Offgas Treatment System

The secondary HLW offgas treatment system will consist of the following components:

- Caustic scrubber
- Thermal catalytic oxidation unit

The following paragraphs describe these components in greater detail.

- Caustic scrubber. The offgas flow from the HEPA filters will be treated by the HLW offgas caustic scrubber. The scrubber will use a solution of sodium hydroxide (NaOH) to remove soluble organics, acid gases, NO_x, and CO_2 (including radioactive $^{14}CO_2$) from the offgas. The scrubber liquor will be combined with condensate or purge from the HLW melter submerged bed scrubber, and then will be either returned to the LAW feed evaporator, or sampled and transferred to the permitted Hanford Liquid Effluent Retention Facility/Effluent Treatment Facility (LERF/ETF).

- Thermal catalytic oxidation unit. After exiting the wet scrubber, the HLW offgas will pass through an electric preheater to raise its temperature to the desired operating range for removal of residual volatile organics in a catalytic oxidation unit. This unit will use a precious metal to facilitate the decomposition of organics to carbon dioxide (CO_2) and water vapor. The treated offgas will be monitored and discharged through the HLW plant stack.

3.4.2.3. HLW Vessel Ventilation Treatment System

Within the HLW vitrification plant, gaseous emissions will be produced from vessel vents and various equipment related to the HLW melter system. All vessel ventilation lines will be connected to the main vessel ventilation header. The main header will join the HLW offgas treatment system lines downstream from the WESP and upstream from the HEME. The combined melter offgas stream and the vessel ventilation offgas stream will be treated in the remaining portion of the HLW primary offgas treatment system and the HLW secondary offgas treatment system.

3.4.2.4. Alternate Route

In the event of a plug developing in the film cooler line from the melter to the submerged bed scrubber, or a surge in offgas that reduces the melter vacuum below the minimum normal set point, a butterfly valve will open the alternate route to the submerged bed scrubber. On the alternate route, air and steam will flow through the bypass film cooler line to cool the offgas and minimize the deposition of solids on the pipe wall.

A general power failure is estimated to occur on average once in 10 years. In such a situation, the standby generators may take from 15 to 30 seconds to come online. During this transition to standby power, the offgas exhaust fans will be without power, and offgas would leak from the melter to the cell. Ventilation air for the melter cell will be subject to all C5 ventilation maximum contamination controls, and will be HEPA filtered after power is restored to prevent or control the release of emissions during such an occurrence. Much of the offgas leaked from a melter during such a short power outage would actually be recaptured into the melter offgas treatment system after power is restored, because a large portion of the offgas is normally drawn from the ambient air in the immediate vicinity of the melter. (See 3.5 Cascade Ventilation System.)

3.4.3. Automatic Waste Feed Cutoff System

The automatic waste feed cutoff system will prevent the introduction of HLW feed when key process conditions fall outside of a predetermined range. As a minimum, the automatic waste feed cutoff system will automatically lock out operation of the waste feed system until proper operating conditions have been restored.

Parameters that will initiate the automatic waste feed cutoff system are being designed. When these parameters have been finalized, a list of automatic waste feed cutoff initiating parameters will be provided in this section. See Appendix A for the proposed information completion schedule.

3.5. Cascade Ventilation System

A primary factor in the design of the ventilation system for the RPP-WTP is the need to isolate the sources of radiation, and radiological and dangerous waste contamination, to protect human health and the environment during normal and abnormal operating conditions. Barriers or barrier systems, including ventilation systems, will contain and minimize the release of radionuclides and contaminants. The ventilation systems are designed to conform to stringent nuclear facility ventilation standards, and fugitive emissions from the pretreatment and vitrification facilities will be minimized.

The pretreatment plant, LAW pretreatment plant, LAW vitrification plant and HLW vitrification plant will be divided into four numbered zones, with the higher number indicating greater contamination potential and therefore a requirement for a greater degree of control or restriction. A separate zoning system for the ventilation systems will be based on the system for classifying building areas for potential contamination. Zones classified as C5 will have the potential for the greatest contamination and will include the pretreatment cells, melter cells, and glass pouring and cooling cells. All C5 zones will be operated remotely. Zones classified as C1 will be those areas that have no risk of contamination, such as equipment rooms and offices.

Confinement will be achieved by maintaining C5 areas at the greatest negative pressure, with airflows cascaded through engineered routes from C2 areas via C3 areas to the C5 areas. The principle of a cascade system, in which air passes through more than one area, effectively reduces the number of separate ventilation streams and hence the amount of air requiring treatment. Adherence to these principles in the design and operation of the RPP-WTP will ensure that the plant will not become a significant source of radiological or dangerous waste exposure to operators, or emissions to the environment.

3.6. Fugitive Emissions

The following sections address fugitive emissions. Because of the stringent design of the processing facilities, it is not anticipated that fugitive emissions will be emitted from the pretreatment, LAW

vitrification, or HLW vitrification processes. Insignificant amounts of fugitive emissions are anticipated from the materials storage areas that will support the processing plants.

3.6.1. Waste Treatment Processing Facilities

The RPP-WTP ventilation system will be designed to confine all sources of contamination and, thus, fugitive releases of COPCs to the environment are not anticipated. Confinement will be achieved through a combination of barriers or barrier systems, coupled with a cascade ventilation system that will be designed to conform to stringent nuclear facility ventilation standards.

The key to prevention of fugitive emissions will be that the more contaminated areas will be maintained at a negative pressure with respect to the less contaminated areas. That is, C5 areas will be maintained at the greatest negative pressure, with air flows cascaded through engineered routes from C2 areas via C3 areas to C5 areas. The pressure gradient will ensure that no fugitive emissions are released from the processing plants (pretreatment, LAW vitrification, and HLW vitrification). The air flow that reaches the C5 area will then be HEPA filtered and routed to the building stack.

3.6.2. Material Storage Systems

Two storage facilities not directly associated with the main processing plant will have the potential to be minor sources of fugitive emissions: the glass-former storage system, and the nitric acid storage system.

The glass-former storage bins will provide bulk storage for the alumina, boric acid, iron oxide, and similar materials that will be used as part of a recipe to support glass formation in the melters. These materials will have no radiological contamination and will not contain COPCs, but very small levels of particulate emissions may occur during periodic refilling of the bins by outside vendors. The storage systems will have filters installed to minimize emissions during filling and normal operations.

The wet chemical storage system will be maintained to provide the nitric acid and other bulk chemicals used in operating the process plant. Emissions from this facility are not expected to be significant.

3.7. Upset Conditions

Process upset conditions include periods of startup, shutdown, or process malfunction; that is, the unit is operating outside the permitted operating conditions. Process upset conditions are generally assumed to result in greater than normal stack emissions during the short period of the upset. The potential for increased emissions during upset conditions will be addressed in this section through an analysis of the processing system of each of the three main components of the facility: pretreatment system, LAW vitrification, and HLW vitrification.

3.7.1. Upset Conditions in the Pretreatment Systems

The pretreatment plant and LAW pretreatment plant ventilation systems will be designed to accommodate several inputs from various unit operations and components. Inputs from these unit operations will be composed, in part, of streams originating from evaporator operations and acid/base reactions. It is expected that normal operations will produce volatilization of semivolatile components and that these components will enter the vessel ventilation system, as designed. The ventilation system will capture these components at prescribed decontamination factors through a series of condensers, scrubbers, HEMEs, filters and other ventilation components prior to entering the discharge stream.

Since normal design operating temperatures and pressures are at the boiling point of the process stream, volatilization of semivolatile components will occur under normal operation. Consequently, an upset condition that impacts temperature or pressure will not produce emissions of semivolatile components in excess of those found during normal operations.

The failure of a ventilation component, such as a condenser, may result in the localized elevation of a semivolatile component concentration. However, the safeguards and control system interlocks for the ventilation system will be in place to notify the operators and initiate a safe shut down of the affected systems. The safe shut down of the ventilation system will reduce the volumetric throughput of the ventilation system and reduce the likelihood that decontamination factors would be exceeded.

In summary, the design of the ventilation system associated with the pretreatment process and LAW pretreatment process will adequately handle the emissions from upset conditions. There are no other anticipated upset conditions associated with the pretreatment systems that would result in additional emissions to the environment.

3.7.2. Upset Conditions in the LAW and HLW Melters

The most likely process upset in the LAW and HLW melters is a loss of vacuum in the melters. The melter offgas system is designed to treat normal offgas flow rates and intermittent surges of 7 times the normal flow rate for condensable gas flow and 3 times the noncondensable gas flow. Should a loss or significant reduction of vacuum occur due to a gas surge or clogging of the film cooler pipe, the upset will be controlled by automatic opening of the fast acting butterfly valve, allowing flow through the bypass film cooler line to the submerged bed scrubber.

If the "bypass" to the SBS is activated, waste feed to the melter will be stopped, the cold cap on the melter allowed to be consumed, and the melter brought to idle condition until the cause for the activation is investigated and necessary repairs made. In a bypass condition, the offgas will be treated by the backup film cooler, SBS and all the downstream offgas treatment units, therefore, no pollution control equipment is bypassed.

In the event of a complete power failure, a loss of vacuum could occur for a brief time, one minute or less, until backup power is available. The melter offgas would leak into the melter cave during that time, because the offgas exhaust fans would be without power. Within the building C5 ventilation system, the offgas will be passed through HEPA filters after power is restored, monitored for compliance with emission standards, and released through the building stack.

3.8. Facility Control Philosophy

This section presents an overall control philosophy for the RPP-WTP. The facility control philosophy goal is to satisfy the following criteria:

- Preservation of worker and public safety
- Protection of the environment
- Preservation of equipment integrity
- Assurance of product quality
- Minimization of plant lifetime costs

The measures proposed to achieve these goals are as follows:

The possibility of human error in facility operation will be addressed in the design hazards analysis. A simple and consistent operator interface will be developed for use throughout the facility.

- Automation will be used to optimize throughput and reduce operator radiological exposure. Where necessary, the system will be designed to allow human decisions required to maximize throughput of the RPP-WTP, meet product quality, or ensure environmental protection.

- Diagnostics will be provided to reduce downtime (for instance, overload and emergency stop signals will be monitored by the control system).

- Control modes will be simplified and the scope of fallback and recovery control provisions will be reduced to contain operating costs.

- A plant information computer with data entry and reporting capabilities will be provided to support product quality and tracking requirements, and to provide process information needed for facilitating plant optimization.

- Provisions will be made for overview and scheduling information.

The RPP-WTP requires a combination of batch and continuous monitoring and control. Process equipment will be contained within cells, due to the radioactivity levels of the waste.

The mechanical handling aspects of the RPP-WTP will be, for the most part, at the final stages of processing, and will require sequential control of mechanical equipment. The mechanical handling equipment will be contained within caves. Since direct access to the plant and equipment will not normally be achievable, viewing of the equipment will be provided by viewing windows, closed circuit television, or both.

Process services, effluent and offgas treatment, and ventilation services will require sequence control for startup and shutdown, and continuous monitoring and control.

Figure 3-1. Location of the RPP-WTP on the Hanford Site



Figure 3-2. RPP-WTP Layout



Figure 3-3. Process Flow Diagram



4. Screening Level Risk Assessment and Constituents of Potential Concern

The SLRA must serve several purposes including: (1) meeting EPA guidance specifications; (2) providing risk information for additional stakeholders, including Native American tribes and other public members; (3) identifying potential financial risk associated with building and operating the River Protection Project – Waste Treatment Plant (RPP-WTP); and (4) providing the information necessary to determine what, if any, additional permit conditions are necessary for the operation of the RPP-WTP to be protective of human health and the environment. For these reasons, the overall approach for the screening level risk assessment (SLRA) will be to identify potential risk associated with a worst-case exposure scenario as well as a plausible exposure scenario.

- The worst-case exposure scenario represents worst-case assumptions regarding the location of receptors, exposure pathways, and activity patterns (e.g., subsistence fishing). The receptor locations used in the worst-case scenario are considered hypothetical since the assumed activities (e.g., resident, subsistence farmer) do not currently occur in these locations nor are they expected to occur in the future.

- The plausible exposure scenario represents more realistic assumptions regarding the location of receptors. It reflects current uses of the surrounding land and habitats and reasonable assumptions about future land uses.

As with the worst-case exposure scenario, the plausible exposure scenario will incorporate conservative assumptions regarding human and ecological exposure patterns as the standard for the deterministic (as opposed to probabilistic) risk modeling.

The preliminary risk assessment (PRA) will be conducted prior to construction of the RPP-WTP. This PRA will rely on assumptions of potential emissions and operating conditions. The final risk assessment (FRA) will be conducted after construction of the RPP-WTP and completion of the environmental performance demonstration. The FRA will utilize real emissions data from the environmental performance demonstrations and focus on characterizing any potentially significant risks identified in the PRA.

The PRA will be performed first and is discussed in more detail in this Work Plan. Requirements for the FRA will be partially dictated by the results of the PRA and environmental performance demonstrations. The FRA will include estimated emissions based on engineering calculations (pretreatment system emissions and abated fugitive emissions) and environmental performance demonstration for the LAW and HLW vitrification systems.

Human Health Risk Assessment Protocol for Hazardous Waste Combustion Facilities (EPA 1998a) recommends the selection of Constituents of Potential Concern (COPCs) to focus on compounds that (1) are likely to be emitted, due to the presence of the compound or its precursors in the waste feed; (2) are potential products of incomplete combustion, (3) are potentially toxic to humans; and/or (4) have a definite propensity for bioaccumulating or bioconcentrating in human and ecological food chains. The COPC selection process described in EPA 1998a includes six steps:

1. Start with a list of all compounds analyzed for in the environmental performance demonstrations and note which compounds were detected according to analytical tests listed in the *Quality Assurance Project Plan for the Environmental Performance Demonstration* (BNFL 1999b).

2. Evaluate the type of waste in Envelopes A through D to determine whether any of the compounds that were not detected should be retained as COPCs because they are present in the waste.

3. Exclude compounds that are not detected, are not components of the waste, and do not have toxicological data.

4. Exclude compounds that are not detected, are not components of the waste, and do not have a high potential to be products of incomplete combustion (PICs).

5. Evaluate the 30 largest tentatively identified compounds (TICs) to determine whether any of these compounds have toxicities similar to the detected compounds. If they do not, consider surrogate toxicity data.

6. Evaluate compounds that may be of concern due to other site-specific factors. Include as COPCs any compounds that are a concern due to site-specific factors and may be emitted by the combustion unit.

This process will utilize data collected during the environmental performance demonstration. Because the PRA will be performed before the environmental performance demonstration results are available, it will be necessary to develop a list of COPCs and a list of ROPCs based on information and assumptions rather than emissions measurements. The following sections describe the strategy for identifying PRA chemicals of potential concern (COPCs) and radionuclides of potential concern (ROPCs). The COPC/ROPC selection process is summarized in Figure 4-1. The term COPC is used to represent chemicals associated with the tank waste and melting process, while the term ROPC refers to radioactive constituents of potential concern.

Vapor phase emissions are those emitted as volatile gases. Particle phase emissions are particulate constituents (generally metals or other inorganics). Particle-bound emissions are gaseous constituents that are bound to other particles.

4.1. Preliminary Chemicals of Potential Concern (COPCs)

The preliminary list of COPCs was compiled using input from *the Regulatory Data Quality Objectives Supporting Tank Waste Remediation System Privatization Project* (Wiemers et al. 1998), the list of PICs identified in Table A.1 of *Human Health Risk Assessment Protocol for Hazardous Waste Combustion Facilities* (EPA 1998a) and the criteria pollutants regulated by the National Ambient Air Quality Standards. Tables A-1 through A-5 in Appendix A of this Work Plan list 424 nonradioactive Preliminary COPCs evaluated for the SLRA. Final COPCs carried through the quantitative risk assessment will be all COPCs for which necessary toxicity data, physical/chemical parameters, and analytical methods are available. COPCs not included in the quantitative risk assessment will be qualitatively addressed in the uncertainty assessment. The COPC list will be re-evaluated for the FRA following the environmental performance demonstration. The derivation of Tables A-1 through A-6 is discussed below and summarized in Figure 4-1.

4.1.1. Regulatory Data Quality Objectives

Analytes considered during the *Regulatory Data Quality Objectives* process (Wiemers et al 1998) were selected from a large universe of regulated constituents using technically defensible decision logic. The

decision logic selected compounds that could plausibly be in the waste feed and of concern relative to the risk assessment and permitting activities. A consolidated list of 850 chemical compounds (Wiemers et al. 1998) was used as the input for the regulatory data quality objectives (DQOs) process. This list of compounds included:

- Toxic Air Pollutant lists Class A (WAC 173-460-150, toxic air pollutants: known, probable, and potential human carcinogens and acceptable source impact levels) and Class B (WAC 173-460-160, toxic air pollutants, and acceptable source impact levels).
- Underlying Hazardous Constituents (UHC) list (40 *CFR* 268.48).
- Universal Treatment Standards list (40 *CFR* 268.48).
- *Double-Shell Tank (DST) System Dangerous Waste Permit Application* (DOE-RL 1991) constituents, except for waste code F039. To date, no landfill leachate has been added to the tanks. Therefore, these compounds were not included in the regulatory DQO database used to select the COPCs.
- DST Waste Stream Profile Sheet constituents.

The list of 850 compounds was screened to arrive at a final list consisting of 139 organic and 49 inorganic compounds. The initial list of 125 organic compounds includes polychlorinated biphenyls (PCBs). This class of compounds includes 209 separate congeners. Fourteen of these congeners (the coplanar PCBs) are considered to be "dioxin-like" and are evaluated individually. These 14 coplanar PCBs were added to the regulatory DQO list of 125 resulting in a total of 139 organic compounds. Tables A-1 and A-2 in Appendix A identify these organic and inorganic COPCs, respectively. A brief discussion of the methodology and criteria used in the regulatory DQOs to narrow the initial input list is included below. Additional details regarding this process are provided in Wiemers et al. 1998.

The organics were screened based upon the following:

- Detectability in the single-shell/double-shell waste
- Stability in the DST environment
- Toxicity and carcinogenicity
- Availability of SW-846 (EPA 1986) analytical methods
- Association with the operations at the Hanford Site

The inorganics were established by the following:

- Listing the inorganic compounds and metals in the input of the starting lists
- Consolidating the list of metals and ions
- Comparing the resulting list to the Hanford Site waste inventories
- Considering the applicability of SW-846 (EPA 1986) analytical methods
- Assessing alternative sources of information

Toxicity criteria were not used to screen inorganic chemicals because the starting list of inorganics was so much shorter than the list of organic chemicals and there was not as large an unknown component to the inorganics (i.e., not as many compounds that were not detected).

4.1.2. Environmental Protection Agency Products of Incomplete Combustion

The organics and inorganics retained by the DQO process were compared to the EPA list of recommended and potential PICs contained in Table A.1 from EPA 1998a. The 139 organic compounds listed in Table A-3 in this Work Plan are the "Chemicals Recommended for Identification" and the "Chemicals for Potential Identification" that were originally identified in Tables 1 and 2 of the *Draft Exposure Assessment Guidance for Resource Conservation and Recovery Act Hazardous Waste Combustion Facilities* (EPA 1994a). In addition, compounds identified in combustion unit emissions and stack emissions originally identified in the *Review Draft Addendum to the Methodology for Assessing Health Risks Associated with Indirect Exposure to Combuster Emissions* (EPA 1993d) were included. All 148 PICs from Table A.1 of EPA 1998a not already included as part of the DQO COPCs were added. These additional 148 COPCs are listed in Table A-3 in Appendix A of this report. This was done to assure a conservative approach.

4.1.3. Site-Specific Products of Incomplete Combustion

A bench-scale test of the melter technology was conducted at The Catholic University of America Vitreous State Laboratory in December 1998 (Matlock and Pegg 1999). A surrogate waste feed was used for this test that likely does not represent the constituents in the actual tanks. This surrogate waste was designed to represent the most difficult-to-destroy chemicals potentially present in the tank waste and, thus, to provide a conservative estimate of potential PICs. This test identified 16 additional potential PICs. These chemicals are listed in Table A-4 of Appendix A.

4.1.4. Chemicals Screened in the DQO Process

EPA Region X (1999) does not agree with the screening of chemicals based on toxicity. Therefore, all chemicals not included in the DQO due to low toxicity, regardless of other factors (e.g., number of detects) are included in the list of preliminary COPCs.

The DQO process identified 146 regulated organic chemicals with positive detects in the liquid or vapor phase of the tank waste (PNNL, 1998 – Table B.5). These chemicals were carried through a toxicity ranking which identified 19 higher toxicity chemicals (PNNL, 1998 – Table B.6) and 127 lower toxicity chemicals. All 19 higher toxicity chemicals were retained for analysis (i.e., included with the 188 chemicals identified by the DQO in Figure 4-1 of this RAWP). Eighty-one of the lower toxicity chemicals were also retained for analysis (PNNL, 1998 – Table B.34). The remaining 46 chemicals were eliminated based on the following: 42 had $\leq$10 detects (PNNL, 1998 – Table B.8), 2 were common laboratory contaminants (PNNL, 1998 – Table B.30), and 2 required unique methods for analysis (PNNL, 1998 – Table B.32).

The 46 low toxicity regulated organic chemicals, with positive detects eliminated from the DQO, were compared to the PICs previously added to the list of preliminary COPCs (sections 4.1.2 and 4.1.3 of this RAWP). Based on this comparison, an additional 26 regulated organic chemicals with positive detects eliminated from the DQO due to low toxicity and having $\leq$10 detects or unique methods for analysis were added to the list of preliminary COPCs. These 26 additional chemicals are listed in Table A-5 under the heading "Detected Chemicals eliminated due to low toxicity".

The DQO process further identified 90 regulated organic chemicals that could plausibly be in the tanks but which were never detected (PNNL, 1998 – Table B.22). These chemicals were carried through a toxicity ranking which identified 25 higher toxicity chemicals (PNNL, 1998 – Table B.24) and 65 lower toxicity chemicals (PNNL, 1998 – Table B.23). All 25 higher toxicity chemicals were retained for

analysis (i.e., included with the 188 chemicals identified by the DQO in Figure 4-1 of this RAWP). The 65 lower toxicity chemicals were eliminated.

The 65 low toxicity regulated organic chemicals with no detects, previously eliminated from the DQO, were compared to the PICs previously added to the list of preliminary COPCs (sections 4.1.2 and 4.1.4 of this RAWP). Based on this comparison, an additional 41 regulated organic chemicals with no detects eliminated from the DQO due to low toxicity were added to the list of preliminary COPCs. These 41 additional chemicals were listed in Table A-5 under the heading "Non-detected chemicals eliminated due to low toxicity".

4.1.5. Criteria Pollutants

National Ambient Air Quality Standards (NAAQS) have been established for six criteria pollutants (sulfur dioxide, particulate matter, carbon monoxide, ozone, nitrogen dioxide, and lead). These criteria pollutants will be addressed in the Prevention of Significant Deterioration (PSD) permit application for this facility. In addition, five of these criteria pollutants are added to the list of preliminary COPCs. Lead was previously included in the COPCs identified by the regulatory DQO process.

4.2. Preliminary Radionuclides of Potential Concern (ROPCs)

The preliminary radionuclides of potential concern (ROPCs) were established based upon *Standard Inventories of Chemicals and Radionuclides in Hanford Site Tank Wastes* (Kupfer et al. 1997). Information used to establish the global inventories originated from key historical records, various chemical flowsheets used in reprocessing of irradiated Hanford Site reactor fuels, and from calculations of radionuclide isotope generation and decay. *Predominant Radionuclides in Hanford Site Waste Tanks* (Boothe 1996) provides the basis to consider 40 radionuclides that are estimated to present over 99% of the inherent radiological risks in each of the following five categories for human receptors:

- Long-term or short-term inhalation hazards
- Long-term groundwater hazards
- Long-term external radiation hazards
- Short-term shielding concerns
- Short-term volatile hazards

An additional six radionuclides (yttrium-90, barium-137m, europium-155, curium-242, curium-243, and plutonium-242) were added to the list of 40 because they are important in the source term. The resulting list of 46 radionuclides shown in Table A-6 in Appendix A of this Work Plan comprises the ROPCs. These 46 radionuclides represent around 99% of the activity associated with the 177 single-shell and double-shell waste tanks at the Hanford Site.

4.3. Selection of Final COPCs/ROPCs for the Quantitative Preliminary Risk Assessment

The preliminary COPCs and ROPCs identified in Tables A-1 through A-6 include an extensive list of chemicals and radionuclides (1) potentially present in the waste to be processed and (2) potentially produced as PICs during the processing of waste. Constituents are deleted from the final list of COPCs/ROPCs to be carried through the quantitative risk assessment because:

- Appropriate human health or ecological toxicity data are not available to quantitatively evaluate potential effects of the constituent;

analysis (i.e., included with the 188 chemicals identified by the DQO in Figure 4-1 of this RAWP). The 65 lower toxicity chemicals were eliminated.

The 65 low toxicity regulated organic chemicals with no detects, previously eliminated from the DQO, were compared to the PICs previously added to the list of preliminary COPCs (sections 4.1.2 and 4.1.4 of this RAWP). Based on this comparison, an additional 41 regulated organic chemicals with no detects eliminated from the DQO due to low toxicity were added to the list of preliminary COPCs. These 41 additional chemicals were listed in Table A-5 under the heading "Non-detected chemicals eliminated due to low toxicity".

4.1.5. Criteria Pollutants

National Ambient Air Quality Standards have been established for six criteria pollutants (sulfur dioxide, particulate matter, carbon monoxide, ozone, nitrogen dioxide, and lead). These criteria pollutants will be addressed in the Prevention of Significant Deterioration permit application for this facility. In addition, five of these criteria pollutants are added to the list of preliminary COPCs. Lead was previously included in the COPCs identified by the regulatory DQO process.

4.2. Preliminary Radionuclides of Potential Concern (ROPCs)

The preliminary radionuclides of potential concern (ROPCs) were established based upon *Standard Inventories of Chemicals and Radionuclides in Hanford Site Tank Wastes* (Kupfer et al. 1997). Information used to establish the global inventories originated from key historical records, various chemical flowsheets used in reprocessing of irradiated Hanford Site reactor fuels, and from calculations of radionuclide isotope generation and decay. *Predominant Radionuclides in Hanford Site Waste Tanks* (Boothe 1996) provides the basis to consider 40 radionuclides that are estimated to present over 99% of the inherent radiological risks in each of the following five categories for human receptors:

- Long-term or short-term inhalation hazards
- Long-term groundwater hazards
- Long-term external radiation hazards
- Short-term shielding concerns
- Short-term volatile hazards

An additional six radionuclides (yttrium-90, barium-137m, europium-155, curium-242, curium-243, and plutonium-242) were added to the list of 40 because they are important in the source term. The resulting list of 46 radionuclides shown in Table A-6 in Appendix A of this Work Plan comprises the ROPCs. These 46 radionuclides represent around 99% of the activity associated with the 177 single-shell and double-shell waste tanks at the Hanford Site.

4.3. Selection of Final COPCs/ROPCs for the Quantitative Preliminary Risk Assessment

The preliminary COPCs and ROPCs identified in Tables A-1 through A-6 include an extensive list of chemicals and radionuclides (1) potentially present in the waste to be processed and (2) potentially produced as PICs during the processing of waste. Constituents are deleted from the final list of COPCs/ROPCs to be carried through the quantitative risk assessment because:

- Appropriate human health or ecological toxicity data are not available to quantitatively evaluate potential effects of the constituent;

- Appropriate physical/chemical parameters are not available to quantitatively evaluate potential emissions or fate and transport behavior of the constituent through the environment; or

- Emission estimates are not available .

Constituents not included in the quantitative risk assessment will be discussed qualitatively as part of the uncertainty assessment.

4.3.1. Selection of Organic COPCs for Inclusion in the Quantitative PRA

The 370 organic preliminary COPCs identified as described in section 4.1 and Figure 4-1 are listed in Table 4-1 and have been grouped into the following classes:

Volatile Organics Compounds (VOCs)	Semi-volatile Organics Compounds (SVOCs)
Aromatic Halogenated Hydrocarbons	Dioxin and Furan Compounds
Aromatic Nonhalogenated Hydrocarbons	Polychlorinated Biphenyls (PCBs)
Non-aromatic Nonhalogenated Hydrocarbons	Phthalates
Non-aromatic Halogenated Hydrocarbons	Light Polynuclear Aromatic Hydrocarbons (PAHs) [molecular weight (MW)<200 g/mole]
	Heavy PAHs (MW>200 g/mole)
	Light Substituted Benzene Compounds (MW<200 g/mole)
	Other Light SVOCs (MW<200 g/mole)
	Other Heavy SVOCs (MW>200 g/mole)
	Herbicides and Organochlorinated Pesticides

EPA 1994a has identified several of these categories (dioxins/furans, PAHs, PCBs, phthalates, other chlorinated organics and nitroaromatics) as having the highest potential to cause increased risk to human health via indirect exposures.

Organic Chemicals with Adequate Toxicity Data

Toxicity data appropriate for the evaluation of potential impacts to human health due to chronic or acute exposures are available from the sources listed in section 7.2 of this workplan.

Toxicity data appropriate for evaluation of chronic exposures to organic COPCs are summarized in Appendix C. Of the 370 organic preliminary COPCs, about 200 have appropriate toxicity data available for the evaluation of potential chronic effects (cancer and/or non-cancer). More than another 30 organic COPCs can be evaluated for potential chronic effects using toxicity equivalency factors (see sections 7.2.3, 7.2.4, and 7.2.5).

Toxicity data appropriate for the evaluation of potential effects resulting from acute (i.e., 1 hour) exposure to organic COPCs are provided in Table 4-4. This table identifies over 260 of the organic preliminary COPCs with appropriate toxicity data available for evaluation of potential acute effects.

Sources of toxicity data for evaluation of potential impacts to ecological receptor populations are described in section 8.3, Effects Assessment Calculations, of this workplan. These data are summarized for organic preliminary COPCs in various tables in Appendix C. Different toxicity data are provided for each of the different types of receptor populations to be evaluated (e.g., aquatic invertebrates). Organic constituents having appropriate toxicity data available for at least one ecological receptor of interest are found in Appendix C.

Only estimated emission rates are needed to model COPC concentrations in air and soil. Additional physical/chemical data are required to model COPC transport through and uptake by other environmental media (e.g., plants, beef, fish, and other biota). Quantitative exposure modeling will be performed for all pathways and COPCs with appropriate physical/chemical data. If physical/chemical data are available for some but not all pathways (e.g., plant uptake factors are available but beef uptake factors are not) for a COPC, those pathways with available data will be included in the quantitative risk assessment.

Organic Chemicals with Emission Rate Estimates

Potential emission rates will be estimated as described in the Air Quality Permit Application and especially Appendix B of the three parts of the Application, and Section 3 of this Work Plan. The list of COPCs for which there are emission rates is provided in Appendix E. Estimated emission rates will be available for 269 organic COPCs.

The final list of organic COPCs that will be included in the quantitative PRA is provided in Appendix D. COPCs not carried through the quantitative PRA will be discussed qualitatively in the uncertainty evaluation.

4.3.2. Selection of Final Inorganic COPCs for Inclusion in the Quantitative PRA

Inorganic COPCs are summarized in Table 4-2.

Inorganic Chemicals with Adequate Toxicity Data

Toxicity data appropriate for the evaluation of potential impacts to human health due to chronic or acute exposures are available from the sources listed in section 7.2 of this workplan.

Toxicity data appropriate for evaluation of chronic exposures to inorganic COPCs are summarized in Appendix C. Of the 54 (the 49 and the 5 criteria pollutants) inorganic preliminary COPCs, about 20 have appropriate toxicity data available for the evaluation of potential chronic effects (cancer and non-cancer). Two other chemicals without toxicity values can be evaluated for media-specific health based effects; see sections 7.2.7 and 7.2.9.

Toxicity data appropriate for the evaluation of potential effects resulting from acute (i.e., 1 hour) exposure to inorganic COPCs are provided in Table 4-4. This table identifies about 50 of the inorganic preliminary COPCs with appropriate toxicity data available for evaluation of potential acute effects.

Sources of toxicity data for evaluation of potential impacts to ecological receptor populations are described in section 8.3 of this workplan. These data are summarized for inorganic preliminary COPCs in various tables in Appendix C. Different toxicity data are provided for each of the different types of receptor populations to be evaluated (e.g., aquatic invertebrates). Inorganic constituents having appropriate toxicity data available for at least one ecological receptor of interest are found in Appendix C.

Inorganic Chemicals with Adequate Physical/Chemical Data

As previously described for organic COPCs, only estimated emission rates are needed to model COPC concentrations in air and soil. Additional physical/chemical data are required to model COPC transport through and uptake by other environmental media (e.g., plants, beef, fish, and other biota). Quantitative exposure modeling will be performed for all pathways and COPCs with appropriate physical/chemical data. If physical/chemical data are available for some but not all pathways (e.g., plant uptake factors are

available but beef uptake factors are not) for a COPC, those pathways with available data will be included in the quantitative risk assessment.

Inorganic Chemicals Emission Rate Estimates

Estimated emission rates will be available for 53 of 54 inorganic COPCs. They will be developed in the Air Quality Permit Application and especially Appendix B of the three parts of the Application.

The final list of inorganic COPCs that will be included in the quantitative PRA is provided in Appendix D. COPCs not carried through the quantitative PRA will be discussed qualitatively in the uncertainty evaluation. In addition to the inorganic COPCs, ROPCs having potential health effects not associated with radioactivity will be evaluated as inorganic COPCs.

4.3.3. Selection of Final ROPCs for Inclusion in the Quantitative PRA

Preliminary ROPCs are summarized in Table 4-3.

Radionuclides with adequate Toxicity Data

Toxicity data appropriate for the evaluation of potential impacts to human health due to chronic or acute exposures are available from the sources listed in section 7.2 of this workplan.

Toxicity data appropriate for evaluation of chronic exposures to ROPCs are summarized in Table 4-3. Of the 46 preliminary ROPCs, 46 have appropriate toxicity data available for the evaluation of potential chronic effects. Toxicity data appropriate for the evaluation of potential effects resulting from acute (i.e., 1 hour) exposure to ROPCs are provided in Table 4-4.

Sources of toxicity data for evaluation of potential impacts to ecological receptor populations are described in section 8.3 of this workplan. These data are summarized for preliminary ROPCs in Appendix C. ROPCs having appropriate toxicity data available for ecological receptors are found in Appendix C.

Radionuclides Chemicals with Adequate Physical/Chemical Data

As previously described for COPCs, only estimated emission rates are needed to model ROPC concentrations in air and soil. Additional physical/chemical data are required to model ROPC transport through and uptake by other environmental media (e.g., plants, beef, fish, and other biota). Quantitative exposure modeling will be performed for all pathways and ROPCs with appropriate physical/chemical data. If physical/chemical data are available for some but not all pathways (e.g., plant uptake factors are available but beef uptake factors are not) for an ROPC, those pathways with available data will be included in the quantitative risk assessment.

Radionuclides with Emission Rate Estimates

Estimated emission rates will be available for 18 of 46 preliminary ROPCs. They will be developed in the Air Quality Permit Application and especially Appendix B of the three parts of the Application.

The final list of ROPCs that will be included in the quantitative PRA is provided in Appendix D. ROPCs not carried through the quantitative PRA will be discussed qualitatively in the uncertainty evaluation.

4.4. Summary for Identification of COPCs and ROPCs

Figure 4-1 provides a summary of the process used to identify preliminary COPCs and ROPCs for the PRA. The inorganic and organic preliminary COPCs and preliminary ROPCs are listed in Appendix A and summarized in Tables A-1 through A-6. They are also listed in Tables 4-1 (organics), 4-2 (inorganics), and 4-3 (radionuclides). Additional human health data are found in Appendix B and ecological data are found in Appendix C.

The list of 470 COPCs/ROPCs selected for the PRA includes many more compounds than are expected in actual facility emissions. The list is long because assumptions are used to compensate for the high level of uncertainty regarding the exact make-up of the waste and the lack of environmental performance demonstration data (i.e., it was assumed that all chemicals potentially present in the waste will be emitted along with all chemicals identified as PICs from any type of combustion unit. The list of preliminary organic COPCs includes numerous chemicals that have never been detected in the tank waste.

Appendix Tables D-1 through D-6 summarize the current availability of data to quantitatively evaluate the preliminary COPCs/ROPCs. These tables also provide a list of the COPCs/ROPCs that will be quantitatively evaluated in the PRA. Preliminary COPCs/ROPCs not included in the PRA will be addressed qualitatively in the uncertainty assessment.

Figure 4-1. Selection of Constituents of Potential Concern



Table 4-1. Human Health Chemical Toxicity Values for Organic Constituents of Potential Concern

Organic COPC	CAS Number	G.I. Absorption Factor [a] (unitless)	Chronic Noncarcinogenic Toxicity: Oral RfD [b] mg/(kg-d)	Dermal RfD [c] mg/(kg-d)	Inhalation RfC [b] mg/m³	Inhalation RfD [d] mg/(kg-d)	Carcinogenic Slope Factor: Oral [b] (kg-day/mg)	Dermal [c] (kg-day/mg)	Inhalation [e] (kg-day/mg)	EPA Class [b]
Aromatic Halogenated Hydrocarbons										
4-Chloro-3-methylphenol	59-50-7	-	-		-	-	-		-	-
2,3,4,6-Tetrachlorophenol	58-90-2	0.50	3.00E-02	1.5E-02	-	-	-	-	-	-
Aromatic Nonhalogenated Hydrocarbons										
2-Nitrotoluene	88-72-2	0.80	1.00E-02[f]	8.0E-03	-	-	-	-	-	-
4-Nitrobiphenyl	92-93-3	-	-	-	-	-	-	-	-	-
Benzaldehyde	100-52-7	0.80	1.00E-01	8.0E-02	-	-	-	-	-	-
Benzene	71-43-2	0.97	-	-	-	-	2.90E-02	2.99E-02	2.90E-02	A
Benzyl alcohol	100-51-6	0.66	3.00E-01[f]	1.98E-01	-	-	-	-	-	-
Ethyl benzene	100-41-4	0.97	1.00E-01	9.70E-02	1.00E+00	2.86E-01	-	-	-	D
m-Xylene	108-38-3	0.80	2.00E+00[f]	1.60E+00	-	-	-	-	-	-
o-Xylene	95-47-6	0.80	2.00E+00[f]	1.60E+00	-	-	-	-	-	-
p-Xylene	106-42-3	0.80	-	-	-	-	-	-	-	-
Styrene	100-42-5	0.80	2.00E-01	1.60E-01	1.00E+00	2.86E-01	-	-	-	-
Toluene	108-88-3	0.80	2.00E-01	1.60E-01	4.00E-01	1.14E-01	-	-	-	D
Non-aromatic Nonhalogenated Hydrocarbons										
1,2-Epoxybutane	106-88-7	0.50	-	-	2.00E-02	5.71E-03	-	-	-	-
1,3-Butadiene	106-99-0	0.80	-	-	-	-	-	-	1.80E+00	B2
1,4-Dioxane	123-91-1	0.80	-	-	-	-	1.10E-02	1.38E-02	-	B2
1-Methylpropyl alcohol	78-92-2	-	-	-	-	-	-	-	-	-
1-Nitropropane	108-03-2	-	-	-	-	-	-	-	-	-
2,2,4-Trimethylpentane	540-84-1	-	-	-	-	-	-	-	-	-
2-Butanone	78-93-3	0.80	6.00E-01	4.80E-01	1.00E+00	2.86E-01	-	-	-	D
2-Butenaldehyde (2-Butenal)	4170-30-3	-	-	-	-	-	-	-	-	-
2-Ethoxyethanol	110-80-5	0.50	4.00E-01[f]	2.00E-01	2.00E-01	5.71E-02	-	-	-	-
2-Heptanone	110-43-0	-	-	-	-	-	-	-	-	-
2-Hexanone	591-78-6	0.66	-	-	-	-	-	-	-	-
2-Methoxyethanol	109-86-4	0.50	1.00E-03[f]	5.0E-04	2.00E-02	5.71E-03	-	-	-	-
2-Methyl-2-propanol	75-65-0	-	-	-	-	-	-	-	-	-
2-Methyl-2-propenenitrile	126-98-7	0.80	1.00E-04	8.00E-05	7.00E-04[f]	2.00E-04	-	-	-	-
2-Methylaziridine	75-55-8	-	-	-	-	-	-	-	-	-
2-Methylpropyl alcohol	78-83-1	0.80	3.00E-01	2.40E-01	-	-	-	-	-	-
2-Pentanone	107-87-9	-	-	-	-	-	-	-	-	-
2-Propanone (Acetone)	67-64-1	0.83	1.00E-01	8.30E-02	-	-	-	-	-	D

Table 4-1. Human Health Chemical Toxicity Values for Organic Constituents of Potential Concern

Organic COPC	CAS Number	G.I. Absorption Factor [a] (unitless)	Chronic Noncarcinogenic Toxicity				Carcinogenic Slope Factor			EPA Class [b]
			Oral RfD [b] mg/(kg-d)	Dermal RfD [c] mg/(kg-d)	Inhalation RfC [b] mg/m³	Inhalation RfD [d] mg/(kg-d)	Oral [b] (kg-day/mg)	Dermal [c] (kg-day/mg)	Inhalation [e] (kg-day/mg)	
2-Propene-1-ol	107-18-6	0.50	5.00E-03	2.50E-03	-	-	-	-	-	-
2-Propyl alcohol	67-63-0	1.00	-	-	-	-	-	-	-	-
3-Heptanone	106-35-4	-	-	-	-	-	-	-	-	-
3-Methyl-1-butanol	123-51-3	-	-	-	-	-	-	-	-	-
3-Methyl-2-butanone	563-80-4	-	-	-	-	-	-	-	-	-
3-Pentanone	96-22-0	-	-	-	-	-	-	-	-	-
4-Heptanone	123-19-3	-	-	-	-	-	-	-	-	-
4-Methyl-2-pentanone	108-10-1	0.80	8.00E-02[f]	6.40E-02	8.00E-02[f]	2.29E-02	-	-	-	-
4-Methyl-3-penten-2-one	141-79-7	-	-	-	-	-	-	-	-	-
5-Methyl-2-hexanone	110-12-3	-	-	-	-	-	-	-	-	-
Acetaldehyde	75-07-0	0.80	-	-	9.00E-03	2.57E-03	-	-	7.70E-03[g]	B2
Acetamide	60-35-5	-	-	-	-	-	-	-	-	-
Acetic acid	64-19-7	-	-	-	-	-	-	-	-	-
Acetic acid ethyl ester	141-78-6	0.80	9.00E-01	7.20E-01	-	-	-	-	-	-
Acetic acid *n*-butyl ester	123-86-4	-	-	-	-	-	-	-	-	-
Acetonitrile	75-05-8	0.80	6.00E-03	4.80E-03	5.00E-02[f]	1.43E-02	-	-	-	-
Acrolein	107-02-8	0.80	2.00E-02[f]	1.60E-02	2.00E-05	5.71E-06	-	-	-	C
Acrylonitrile	107-13-1	0.80	1.00E-03[f]	8.00E-04	2.00E-03	5.71E-04	5.40E-01	6.75E-01	2.40E-01	B1
Bis(isopropyl)ether	108-20-3	-	-	-	-	-	-	-	-	-
Butane	106-97-8	-	-	-	-	-	-	-	-	-
Carbon disulfide	75-15-0	0.63	1.00E-01	6.30E-02	7.00E-01	2.00E-01	-	-	-	-
Cyanogen	460-19-5	0.80	4.00E-02	3.20E-02	-	-	-	-	-	-
Cyclohexane	110-82-7	-	-	-	-	-	-	-	-	-
Cyclohexanone	108-94-1	0.80	5.00E+00	4.00E+00	-	-	-	-	-	-
Cyclohexene	110-83-8	-	-	-	-	-	-	-	-	-
Cyclopentane	287-92-3	-	-	-	-	-	-	-	-	-
Ethyl alcohol	64-17-5	-	-	-	-	-	-	-	-	-
Ethyl ether	60-29-7	0.80	2.00E-01	1.60E-01	-	-	-	-	-	-
Ethyl methacrylate	97-63-2	0.80	9.00E-02[f]	7.20E-02	-	-	-	-	-	-
Formaldehyde	50-00-0	0.80	2.00E-01	1.60E-01	-	-	-	-	4.50E-02	B1
Formamide	75-12-7	-	-	-	-	-	-	-	-	-
Formic acid	64-18-6	0.50	2.00E+00[f]	1.0E+00	-	-	-	-	-	-
Formic acid, methyl ester	107-31-3	-	-	-	-	-	-	-	-	-
Glycidylaldehyde	765-34-4	0.50	4.00E-04	2.00E-04	1.00E-03[f]	2.86E-04	-	-	-	B2
Methyl acetate	79-20-9	0.50	1.00E+00[f]	5.00E-01	-	-	-	-	-	-

Table 4-1. Human Health Chemical Toxicity Values for Organic Constituents of Potential Concern

Organic COPC	CAS Number	G.I. Absorption Factor [a] (unitless)	Chronic Noncarcinogenic Toxicity				Carcinogenic Slope Factor			EPA Class [b]
			Oral RfD [b] mg/(kg-d)	Dermal RfD [c] mg/(kg-d)	Inhalation RfC [b] mg/m³	Inhalation RfD [d] mg/(kg-d)	Oral [b] (kg-day/mg)	Dermal [c] (kg-day/mg)	Inhalation [c] (kg-day/mg)	
Methyl alcohol	67-56-1	0.80	5.00E-01	4.00E-01	-	-	-	-	-	-
Methyl isocyanate	624-83-9	-	-	-	-	-	-	-	-	-
Methyl methacrylate	80-62-6	0.80	1.40E+00[f]	1.12E+00	7.00E-01	2.00E-01	-	-	-	E
Methyl tert-butyl ether	1634-04-4	0.80	-	-	3.00E+00	8.57E-01	-	-	-	-
Methylacetylene	74-99-7	-	-	-	-	-	-	-	-	-
Methylcyclohexane	108-87-2	0.80	-	-	3.00E+00[f]	8.57E-01	-	-	-	-
N,N-Dimethylacetamide	127-19-5	-	-	-	-	-	-	-	-	-
n-Butyl alcohol	71-36-3	0.50	1.00E-01	5.00E-02	-	-	-	-	-	D
n-Heptane	142-82-5	0.80	-	-	-	-	-	-	-	D
n-Hexane	110-54-3	0.80	6.00E-02[f]	4.80E-02	2.00E-01	5.71E-02	-	-	-	-
Nitromethane	75-52-5	-	-	-	-	-	-	-	-	-
n-Nonane	111-84-2	-	-	-	-	-	-	-	-	-
n-Octane	111-65-9	-	-	-	-	-	-	-	-	-
n-Pentane	109-66-0	-	-	-	-	-	-	-	-	-
n-Propionaldehyde	123-38-6	-	-	-	-	-	-	-	-	-
n-Propyl alcohol	71-23-8	-	-	-	-	-	-	-	-	-
n-Valeraldehyde	110-62-3	-	-	-	-	-	-	-	-	-
Oxirane	75-21-8	0.80	-	-	-	-	1.02E+00[f]	1.28E+00	3.50E-01[f]	B1[f]
p-Cymene	99-87-6	-	-	-	-	-	-	-	-	-
Phosgene	75-44-5	-	-	-	-	-	-	-	-	-
Propargyl alcohol	107-19-7	0.80	2.00E-03	1.60E-03	-	-	-	-	-	-
Propionic acid	79-09-4	-	-	-	-	-	-	-	-	-
Propionitrile	107-12-0	-	-	-	-	-	-	-	-	-
Propylene gylcol monomethyl ether	107-98-2	0.50	7.00E-01[f]	3.50E-01	2.00E+00	5.71E-01	-	-	-	-
p-tert-Butyltoluene	98-51-1	-	-	-	-	-	-	-	-	-
Triethylamine	121-44-8	0.80	-	-	7.00E-03	2.00E-03	-	-	-	-
Trimethylamine	75-50-3	-	-	-	-	-	-	-	-	-
Vinyl acetate	108-05-4	0.65	1.00E+00[f]	6.50E-01	2.00E-01	5.71E-02	-	-	-	-
Non-aromatic Halogenated Hydrocarbons										
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	-	-	-	-	-	-	-	-	-
1,1,1,2-Tetrachloroethane	630-20-6	0.80	3.00E-02	2.40E-01	-	-	2.60E-02	3.25E-02	2.60E-02	C
1,1,1-Trichloroethane	71-55-6	0.90	2.00E-01[h]	1.80E-01	1.00E+00[h]	2.86E-01	-	-	-	D
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	-	-	-	-	-	-	-	-	-
1,1,2,2-Tetrachloroethane	79-34-5	0.70	-	-	-	-	2.00E-01	2.86E-01	2.00E-01	C
1,1,2,2-Tetrachloroethene	127-18-4	1.00	1.00E-02	1.00E-02	6.00E-01[h]	1.71E-01	5.20E-02[h]	5.20E-02	2.00E-03[h]	-

Table 4-1. Human Health Chemical Toxicity Values for Organic Constituents of Potential Concern

Organic COPC	CAS Number	G.I. Absorption Factor [a] (unitless)	Chronic Noncarcinogenic Toxicity: Oral RfD [b] mg/(kg-d)	Dermal RfD [c] mg/(kg-d)	Inhalation RfC [b] mg/m³	Inhalation RfD [d] mg/(kg-d)	Carcinogenic Slope Factor: Oral [b] (kg-day/mg)	Dermal [c] (kg-day/mg)	Inhalation [e] (kg-day/mg)	EPA Class [b]
1,1,2-Trichloroethane	79-00-5	0.81	4.00E-03	3.24E-03	-	-	5.70E-02	7.04E-02	5.70E-02	C
1,1,2-Trichloroethylene	79-01-6	0.15	6.00E-03[h]	9.00E-04	-	-	1.10E-02[h]	7.33E-02	6.00E-03[h]	-
1,1-Dichloroethane	75-34-3	1.00	1.00E-01[f]	1.00E-01	5.00E-01[f]	1.43E-01	-	-	-	C
1,1-Dichloroethene	75-35-4	1.00	9.00E-03	9.00E-03	-	-	6.00E-01	6.00E-01	1.20E+00	C
1,2,2-Trichloro-1,1,2-trifluoroethane	76-13-1	0.80	3.00E+01	2.40E+00	3.00E+01[f]	8.57E+00	-	-	-	-
1,2,3-Trichloropropane	96-18-4	0.80	6.00E-03	4.80E-03	-	-	7.00E+00[f]	8.75E+00	-	B2[f]
1,2-Dibromo-3-chloropropane	96-12-8	0.80	-	-	2.00E-04	5.71E-05	1.40E+00[f]	1.75E+00	2.40E-03	B2[f]
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	-	-	-	-	-	-	-	-	-
1,2-Dichloroethane	107-06-2	1.00	-	-	-	-	9.10E-02	9.10E-02	-	B2
1,2-Dichloroethylene	540-59-0	0.80	9.00E-03[f]	7.20E-03	-	-	-	-	-	-
1,2-Dichloropropane	78-87-5	0.74	-	-	4.00E-03	1.14E-03	6.80E-02[f]	9.19E-02	-	B2[f]
1,3-Dichloropropene	542-75-6	0.55	3.00E-04	1.65E-04	2.00E-02	5.71E-03	1.80E-01[f]	3.27E-01	1.30E-01	B2
1,4-Dichloro-2-butene	764-41-0	0.80	-	-	-	-	-	-	9.30E+00	B2[f]
1-Chloroethene	75-01-4	1.00	-	-	-	-	1.90E+00[f]	1.90E+00	3.00E-01	A[f]
2,2-Dichloropropionic acid	75-99-0	0.50	3.00E-02	1.50E-02	-	-	-	-	-	-
2-Chloropropane	75-29-6	0.80	-	-	1.00E-01[f]	2.86E-02	-	-	-	-
3-Chloropropene (allyl chloride)	107-05-1	0.80	-	-	1.00E-03	2.86E-04	-	-	-	C
Bromochloromethane	74-97-5	0.80	-	-	-	-	-	-	-	D
Bromodichloromethane	75-27-4	0.98	2.00E-02	1.96E-02	-	-	6.20E-02	6.33E-02	-	B2
Bromoethene	593-60-2	0.80	-	-	3.00E-03	8.57E-04	-	-	-	B2[f]
Bromoform	75-25-2	0.60	2.00E-02	1.20E-02	-	-	7.90E-03	1.32E-02	3.90E-03	B2
Bromomethane	74-83-9	0.80	1.40E-03	1.12E-03	5.00E-03	1.43E-03	-	-	-	D
Carbon tetrachloride	56-23-5	0.65	7.00E-04	4.55E-04	-	-	1.30E-01	2.00E-01	5.30E-02	B2
Chlorodibromomethane	124-48-1	0.60	2.00E-02	1.20E-02	-	-	8.40E-02	1.40E-01	-	C
Chlorodifluoromethane	75-45-6	0.80	-	-	5.00E+01	1.43E+01	-	-	-	-
Chloroethane	75-00-3	0.80	-	-	1.00E+01	2.86E+00	-	-	-	-
Chloroform	67-66-3	0.20	1.00E-02	2.00E-03	-	-	6.10E-03	3.05E-02	8.10E-02	B2
Chloromethane	74-87-3	0.80	-	-	-	-	1.30E-02[f]	1.63E-02	6.30E-03	C[f]
Chloropentafluoroethane	76-15-3	-	-	-	-	-	-	-	-	-
cis-1,2-Dichloroethene	156-59-2	1.00	1.00E-02[f]	1.00E-02	-	-	-	-	-	D
cis-1,3-Dichloropropene	10061-01-5	-	-	-	-	-	-	-	-	-
Cyanogen bromide	506-68-3	0.80	9.00E-02	7.2E-02	-	-	-	-	-	-
Cyanogen chloride	506-77-4	0.80	5.00E-02	4.00E-02	-	-	-	-	-	-
Dichlorodifluoromethane	75-71-8	0.23	2.00E-01[f]	4.60E-02	2.00E-01[f]	5.71E-02	-	-	-	-

Table 4-1. Human Health Chemical Toxicity Values for Organic Constituents of Potential Concern

Organic COPC	CAS Number	G.I. Absorption Factor [a] (unitless)	Chronic Noncarcinogenic Toxicity				Carcinogenic Slope Factor			EPA Class [b]
			Oral RfD [b] mg/(kg-d)	Dermal RfD [c] mg/(kg-d)	Inhalation RfC [b] mg/m³	Inhalation RfD [d] mg/(kg-d)	Oral [b] (kg-day/mg)	Dermal [c] (kg-day/mg)	Inhalation [e] (kg-day/mg)	
Dichlorofluoromethane	75-43-4	-	-	-	-	-	-	-	-	-
Dichloromethane	75-09-2	0.95	6.00E-02	5.70E-02	3.00E+00[f]	8.57E-01	7.50E-03	7.89E-03	1.65E-03[g]	B2
Difluorodibromomethane	75-61-6	-	-	-	-	-	-	-	-	-
Hexafluoroacetone	684-16-2	-	-	-	-	-	-	-	-	-
Iodomethane	74-88-4	-	-	-	-	-	-	-	-	-
Methylene bromide	74-95-3	0.80	1.00E-02[f]	8.00E-03	-	-	-	-	-	-
Pentachloroethane	76-01-7	-	-	-	-	-	-	-	-	-
trans-1,2-Dichloroethene	156-60-5	1.00	2.00E-02	2.00E-02	-	-	-	-	-	-
trans-1,3-Dichloropropene	10061-02-6	-	-	-	-	-	-	-	-	-
Trichloroacetic acid	76-03-9	0.50	-	-	-	-	-	-	-	C
Trichlorofluoroethane	27154-33-2	-	-	-	-	-	-	-	-	-
Trichlorofluoromethane	75-69-4	0.23	3.00E-01	6.90E-02	7.00E-01[f]	2.00E-01	-	-	-	-
Trifluorobromomethane	75-63-8	-	-	-	-	-	-	-	-	-
Dioxin and Furan Compounds										
1,2,3,4,6,7,8-Heptachlorodibenzo(*p*)dioxin[i]	35822-46-9	-	-	-	-	-	-	-	-	-
1,2,3,4,6,7,8-Heptachlorodibenzofuran[i]	67562-39-4	-	-	-	-	-	-	-	-	-
1,2,3,4,7,8,9-Heptachlorodibenzofuran[i]	55673-89-7	-	-	-	-	-	-	-	-	-
1,2,3,4,7,8-Hexachlorodibenzo(*p*)dioxin[i]	39227-28-6	-	-	-	-	-	-	-	-	-
1,2,3,4,7,8-Hexachlorodibenzofuran[i]	70648-26-9	-	-	-	-	-	-	-	-	-
1,2,3,6,7,8-Hexachlorodibenzo(*p*)dioxin[i]	57653-85-7	-	-	-	-	-	-	-	-	-
1,2,3,6,7,8-Hexachlorodibenzofuran[i]	57117-44-9	-	-	-	-	-	-	-	-	-
1,2,3,7,8,9-Hexachlorodibenzo(*p*)dioxin[i]	19408-74-3	0.50	-	-	-	-	-	-	-	-
1,2,3,7,8,9-Hexachlorodibenzofuran[i]	72918-21-9	-	-	-	-	-	-	-	-	-
1,2,3,7,8-Pentachlorodibenzo(*p*)dioxin[i]	40321-76-4	-	-	-	-	-	-	-	-	-
1,2,3,7,8-Pentachlorodibenzofuran[i]	57117-41-6	0.50	-	-	-	-	-	-	-	-
2,3,4,6,7,8-Hexachlorodibenzofuran[i]	60851-34-5	-	-	-	-	-	-	-	-	-
2,3,4,7,8-Pentachlorodibenzofuran[i]	57117-31-4	0.50	-	-	-	-	-	-	-	-
2,3,7,8-Tetrachlorodibenzo(*p*)dioxin	1746-01-6	0.50	-	-	-	-	1.50E+05[f]	3.00E+05	1.50E+05	B2[f]
2,3,7,8-Tetrachlorodibenzofuran[i]	51207-31-9	0.50	-	-	-	-	-	-	-	-
Dibenzofuran	132-64-9	-	-	-	-	-	-	-	-	-
Octachlorodibenzo(*p*)dioxin[i]	3268-87-9	0.50	-	-	-	-	-	-	-	-
Octachlorodibenzofuran[i]	39001-02-0	0.50	-	-	-	-	-	-	-	-
PCBs [j]: Coplanar										
2,2',3,3',4,4',5- Heptachlorobiphenyl	35065-30-6	-	-	-	-	-	-	-	-	-

Table 4-1. Human Health Chemical Toxicity Values for Organic Constituents of Potential Concern

Organic COPC	CAS Number	G.I. Absorption Factor [a] (unitless)	Chronic Noncarcinogenic Toxicity: Oral RfD [b] mg/(kg-d)	Dermal RfD [c] mg/(kg-d)	Inhalation RfC [b] mg/m³	Inhalation RfD [d] mg/(kg-d)	Carcinogenic Slope Factor: Oral [b] (kg-day/mg)	Dermal [c] (kg-day/mg)	Inhalation [e] (kg-day/mg)	EPA Class [b]
2,2',3,4,4',5,5'- Heptachlorobiphenyl	35065-29-3	-	-	-	-	-	-	-	-	-
2,3,3',4,4',5'- Hexachlorobiphenyl	69782-90-7	-	-	-	-	-	-	-	-	-
2,3,3',4,4',5- Hexachlorobiphenyl	38380-08-4	-	-	-	-	-	-	-	-	-
2,3,3',4,4',5,5'- Heptachlorobiphenyl	no cas #	-	-	-	-	-	-	-	-	-
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	-	-	-	-	-	-	-	-	-
2,3,4,4',5- Pentachlorobiphenyl	74472-37-0	-	-	-	-	-	-	-	-	-
2',3,4,4',5- Pentachlorobiphenyl	no cas #	-	-	-	-	-	-	-	-	-
2,3',4,4',5- Pentachlorobiphenyl	31508-00-6	-	-	-	-	-	-	-	-	-
2,3',4,4',5,5'- Hexachlorobiphenyl	no cas #	-	-	-	-	-	-	-	-	-
3,3',4,4',5- Pentachlorobiphenyl	no cas #	-	-	-	-	-	-	-	-	-
3,3',4,4',5,5'- Hexachlorobiphenyl	32774-16-6	-	-	-	-	-	-	-	-	-
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	-	-	-	-	-	-	-	-	-
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	-	-	-	-	-	-	-	-	-
PCBs [j]: Non-coplanar										
Polychlorinated biphenyls	1336-36-3	0.90	2.00E-05	1.80E-05	-	-	2.00E+00 [k,l]	2.22E+00	2.00E+00 [k,l]	B2
Phthalates										
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	0.19	2.00E-02	3.80E-03	-	-	1.40E-02		-	B2
Butylbenzyl phthalate	85-68-7	0.61	2.00E-01	1.22E-01	-	-	-	-	-	C
Dibutyl phthalate	84-74-2	1.00	1.00E-01	1.00E-01	-	-	-	-	-	D
Diethyl phthalate	84-66-2	0.90	8.00E-01	7.20E-01	-	-	-	-	-	D
Dimethylphthalate	131-11-3	0.90	-	-	-	-	-	-	-	D
n-Dioctyl phthalate	117-84-0	0.90	2.00E-02 [f]	1.80E-02	-	-	-	-	-	-
Light Polycyclic Aromatic Hydrocarbons (molecular weight <200 g/mole)										
2-Chloronapthalene	91-58-7	0.50	8.00E-02	4.00E-02	-	-	-	-	-	-
2-Methyl naphthalene	91-57-6	0.80	-	-	-	-	-	-	-	-
5-Nitroacenaphthene	602-87-9	-	-	-	-	-	-	-	-	-
Acenaphthene	83-32-9	0.31	6.00E-02	1.86E-02	-	-	-	-	-	-
Acenaphthylene	208-96-8	0.31	-	-	-	-	-	-	-	-
Anthracene	120-12-7	0.76	3.00E-01	2.28E-01	-	-	-	-	-	D
Fluorene	86-73-7	0.50	4.00E-02	2.00E-02	-	-	-	-	-	D
Indene	95-13-6	-	-	-	-	-	-	-	-	-
Naphthalene	91-20-3	0.80	2.00E-02	1.60E-02	3.00E-03	8.57E-04	-	-	-	C
Phenanthrene	85-01-8	0.73	-	-	-	-	-	-	-	D
Pyrene	129-00-0	0.31	3.00E-02	9.30E-03	-	-	-	-	-	D

Table 4-1. Human Health Chemical Toxicity Values for Organic Constituents of Potential Concern

Organic COPC	CAS Number	G.I. Absorption Factor [a] (unitless)	Chronic Noncarcinogenic Toxicity: Oral RfD [b] mg/(kg-d)	Chronic Noncarcinogenic Toxicity: Dermal RfD [c] mg/(kg-d)	Chronic Noncarcinogenic Toxicity: Inhalation RfC [b] mg/m³	Chronic Noncarcinogenic Toxicity: Inhalation RfD [d] mg/(kg-d)	Carcinogenic Slope Factor: Oral [b] (kg-day/mg)	Carcinogenic Slope Factor: Dermal [c] (kg-day/mg)	Carcinogenic Slope Factor: Inhalation [e] (kg-day/mg)	EPA Class [b]
Heavy Polycyclic Aromatic Hydrocarbons (molecular weight >200 g/mole)										
3-Methylcholanthrene	56-49-5	-	-	-	-	-	-	-	-	-
5-Methylchrysene	3697-24-3	-	-	-	-	-	-	-	-	-
Benzo(*a*)anthracene[m]	56-55-3	0.31	-	-	-	-	-	-	-	-
Benzo(*a*)pyrene	50-32-8	0.31	-	-	-	-	7.30E+00	2.35E+01	3.10E+00[n]	B2
Benzo(*b*)fluoranthene[m]	205-99-2	0.31	-	-	-	-	-	-	-	-
Benzo(*e*)pyrene	192-97-2	-	-	-	-	-	-	-	-	-
Benzo(*g,h,i*)perylene	191-24-2	0.31	-	-	-	-	-	-	-	D
Benzo(*j*)fluoranthene	205-82-3	-	-	-	-	-	-	-	-	-
Benzo(*k*)fluoranthene[m]	207-08-9	0.31	-	-	-	-	-	-	-	-
Benzo[a,i]pyrene	191-30-0	-	-	-	-	-	-	-	-	-
Chrysene[m]	218-01-9	0.31	-	-	-	-	-	-	-	-
Dibenz(*a,h*)anthracene[m]	53-70-3	0.31	-	-	-	-	-	-	-	-
Dibenz[a,h]acridine	226-36-8	-	-	-	-	-	-	-	-	-
Dibenz[a,j]acridine	224-42-0	-	-	-	-	-	-	-	-	-
Dibenzo(*a,e*)fluoranthene	5385-75-1	-	-	-	-	-	-	-	-	-
Dibenzo(*a,h*)fluoranthene	no cas #	-	-	-	-	-	-	-	-	-
Dibenzo[a,e]pyrene	192-65-4	-	-	-	-	-	-	-	-	-
Dibenzo[a,h]pyrene	189-64-0	-	-	-	-	-	-	-	-	-
Dibenzo[a,i]pyrene	189-55-9	-	-	-	-	-	-	-	-	-
Fluoranthene	206-44-0	0.31	4.00E-02	1.24E-02	-	-	-	-	-	D
Hexachloronaphthalene	1335-87-1	-	-	-	-	-	-	-	-	-
Indeno(1,2,3-*cd*)pyrene[m]	193-39-5	0.31	-	-	-	-	-	-	-	-
Octachloronaphthalene	2234-13-1	-	-	-	-	-	-	-	-	-
Pentachloronaphthalene	1321-64-8	-	-	-	-	-	-	-	-	-
Tetrachloronaphthalene	1335-88-2	-	-	-	-	-	-	-	-	-
Trichloronaphthalene	1321-65-9	-	-	-	-	-	-	-	-	-
Light Substituted Benzene Compounds (MW <200 g/mole)										
1,2,3-Trichlorobenzene	87-61-6	-	-	-	-	-	-	-	-	-
1,2,4-Trichlorobenzene	120-82-1	0.97	1.00E-02	9.70E-03	2.00E-01[f]	5.71E-02	-	-	-	D
1,2,4-Trimethyl benzene	95-63-6	-	-	-	-	-	-	-	-	-
1,2-Dichlorobenzene	95-50-1	0.80	9.00E-02	7.20E-02	2.00E-01[f]	5.71E-02	-	-	-	D
1,3,5-Trimethyl benzene	108-67-8	-	-	-	-	-	-	-	-	-
1,3-Dichlorobenzene	541-73-1	0.80	-	-	-	-	-	-	-	D
1,3-Dinitrobenzene	99-65-0	0.65	1.00E-04	6.50E-05	-	-	-	-	-	D

Table 4-1. Human Health Chemical Toxicity Values for Organic Constituents of Potential Concern

Organic COPC	CAS Number	G.I. Absorption Factor [a] (unitless)	Chronic Noncarcinogenic Toxicity				Carcinogenic Slope Factor			EPA Class [b]
			Oral RfD [b] mg/(kg-d)	Dermal RfD [c] mg/(kg-d)	Inhalation RfC [b] mg/m³	Inhalation RfD [d] mg/(kg-d)	Oral [b] (kg-day/mg)	Dermal [c] (kg-day/mg)	Inhalation [e] (kg-day/mg)	
1,4-Dichlorobenzene	106-46-7	0.90	-	-	8.00E-01	2.29E-01	2.40E-02[f]	2.67E-02	-	C[f]
1,4-Dinitrobenzene	100-25-4	0.50	4.00E-04[f]	2.00E-04	-	-	-	-	-	-
2,4,5-Trichlorophenol	95-95-4	0.50	1.00E-01	5.00E-02	-	-	-	-	-	-
2,4,6-Trichlorophenol	88-06-2	0.50	-	-	-	-	1.10E-02	2.20E-02	1.00E-02	B2
2,4-Dichlorophenol	120-83-2	0.82	3.00E-03	2.46E-03	-	-	-	-	-	-
2,4-Dimethylphenol	105-67-9	0.50	2.00E-02	1.00E-02	-	-	-	-	-	-
2,4-Dinitrophenol	51-28-5	1.00	2.00E-03	2.00E-03	-	-	-	-	-	-
2,4-Dinitrotoluene	121-14-2	0.85	2.00E-03	1.70E-03	-	-	6.80E-01[e]	8.00E-01	-	B2
2,6-Dinitrotoluene	606-20-2	0.85	1.00E-03[f]	8.50E-04	-	-	6.80E-01[e]	8.00E-01	-	B2
2-Chlorophenol	95-57-8	0.50	5.00E-03	2.50E-03	-	-	-	-	-	-
2-Chlorotoluene	95-49-8	0.80	2.00E-02	1.60E-02	-	-	-	-	-	-
2-Nitrophenol	88-75-5	-	-	-	-	-	-	-	-	-
4,6-Dinitro-*o*-cresol	534-52-1	1.00	-	-	-	-	-	-	-	-
4-Chlorotoluene	106-43-4	-	-	-	-	-	-	-	-	-
4-Nitrophenol	100-02-7	1.00	-	-	-	-	-	-	-	-
alpha-Methylstyrene	98-83-9	0.80	7.00E-02[f]	5.60E-02	-	-	-	-	-	-
Aniline	62-53-3	0.50	-	-	1.00E-03	2.86E-04	5.70E-03	1.14E-01	-	B2
Benzotrichloride	98-07-7	0.50	-	-	-	-	1.30E+01	2.60E+01	-	B2
Benzyl chloride	100-44-7	0.80	-	-	-	-	1.70E-01	2.13E-01	-	B2
Bromobenzene	108-86-1	-	-	-	-	-	-	-	-	-
Chlorobenzene	108-90-7	0.31	2.00E-02	6.20E-03	2.00E-02[f]	5.71E-03	-	-	-	D
Cumene	98-82-8	0.80	1.00E-01	8.00E-02	4.00E-01	1.14E-01	-	-	-	D
m-Cresol	108-39-4	0.50	5.00E-02	2.50E-02	-	-	-	-	-	C
n-Butyl benzene	104-51-8	-	-	-	-	-	-	-	-	-
Nitrobenzene	98-95-3	0.97	5.00E-04	4.85E-04	2.00E-03[f]	5.71E-04	-	-	-	D
n-Propyl benzene	103-65-1	-	-	-	-	-	-	-	-	-
o-Cresol	95-48-7	0.50	5.00E-02	2.50E-02	-	-	-	-	-	C
o-Dinitrobenzene	528-29-0	0.93	4.00E-04[f]	3.72E-04	-	-	-	-	-	D
o-Nitroaniline	88-74-4	0.50	-	-	2.00E-04[f]	5.71E-05	-	-	-	-
o-Toluidine	95-53-4	0.50	-	-	-	-	2.40E-01[f]	4.80E-01	-	B2[f]
p-Chloroaniline	106-47-8	0.50	4.00E-03	2.00E-03	-	-	-	-	-	-
p-Cresol	106-44-5	0.65	5.00E-03[f]	3.25E-03	-	-	-	-	-	C
Phenol	108-95-2	0.90	6.00E-01	5.40E-01	-	-	-	-	-	D
p-Nitrochlorobenzene	100-00-5	-	-	-	-	-	-	-	-	-

Table 4-1. Human Health Chemical Toxicity Values for Organic Constituents of Potential Concern

Organic COPC	CAS Number	G.I. Absorption Factor [a] (unitless)	Chronic Noncarcinogenic Toxicity: Oral RfD [b] mg/(kg-d)	Chronic Noncarcinogenic Toxicity: Dermal RfD [c] mg/(kg-d)	Chronic Noncarcinogenic Toxicity: Inhalation RfC [b] mg/m³	Chronic Noncarcinogenic Toxicity: Inhalation RfD [d] mg/(kg-d)	Carcinogenic Slope Factor: Oral [b] (kg-day/mg)	Carcinogenic Slope Factor: Dermal [c] (kg-day/mg)	Carcinogenic Slope Factor: Inhalation [e] (kg-day/mg)	EPA Class [b]
p-Toluidine	106-49-0	0.50	-	-	-	-	1.90E-01[f]	3.80E-01	-	C[f]
sec-Butyl benzene	135-98-8	-	-	-	-	-	-	-	-	-
tert-Butyl benzene	98-06-6	-	-	-	-	-	-	-	-	-
Toluene-2,6-diamine	823-40-5	0.50	2.00E-01[f]	1.00E-01	-	-	-	-	-	-
Trimethyl benzene	25551-13-7	0.97	-	-	-	-	-	-	-	-
Other Light Semivolatile Compounds (molecular weight <200 g/mole)										
1,1`-Biphenyl	92-52-4	0.50	5.00E-02	2.50E-02	-	-	-	-	-	D
1,1-Dimethylhydrazine	57-14-7	0.50	-	-	-	-	-	-	-	-
1,2-Dimethylhydrazine	540-73-8	0.50	-	-	-	-	-	-	-	B2[f]
1,2-Diphenylhydrazine	122-66-7	0.50	-	-	-	-	8.00E-01	1.60E+00	8.00E-01	B2
1,3-Propane sultone	1120-71-4	-	-	-	-	-	-	-	-	-
2,4-Toluene diisocyante	584-84-9	-	-	-	-	-	-	-	-	-
2-Chloroacetophenone	532-27-4	0.50	-	-	3.00E-05	8.57E-06	-	-	-	-
2-Propenoic acid	79-10-7	0.50	5.00E-01	2.50E-01	1.00E-03	2.86E-04	-	-	-	-
4,4-Methylenedianiline	101-77-9	0.50	-	-	-	-	-	-	-	-
Acetophenone	98-86-2	0.80	1.00E-01	8.00E-02	-	-	-	-	-	D
Benzoic acid	65-85-0	1.00	4.00E+00	4.00E+00	-	-	-	-	-	D
bis(2-Chloroethoxy)methane	111-91-1	0.50	-	-	-	-	-	-	-	D
bis(2-Chloroethyl) ether	111-44-4	0.50	-	-	-	-	1.10E+00	2.20E+00	1.10E+00	B2
Chlorocyclopentadiene	41851-50-7	0.50	-	-	-	-	-	-	-	D
Cyclohexanol	108-93-0	-	-	-	-	-	-	-	-	-
Dichloroisopropyl ether	108-60-1	0.50	-	-	-	-	7.00E-02[f]	1.40E-01	3.50E-02	C[f]
Dichloromethyl ether	542-88-1	0.80	-	-	-	-	2.20E+02	2.75E+02	2.20E+02	A
Dichloropentadiene	no cas #	-	-	-	-	-	-	-	-	-
Dimethyl sulfate	77-78-1	0.50	-	-	-	-	-	-	-	B2
Dimethylaniline	121-69-7	0.50	2.00E-03	1.00E-03	-	-	-	-	-	-
Di-*n*-propylnitrosamine	621-64-7	0.25	-	-	-	-	7.00E+00	2.80E+01	-	B2
Diphenyl ether	101-84-8	-	-	-	-	-	-	-	-	-
Epichlorohydrin	106-89-8	0.80	2.00E-03[f]	1.60E-03	1.00E-03	2.86E-04	9.90E-03	1.24E-02	4.20E-03	B2
Ethyl carbamate (urethane)	51-79-6	-	-	-	-	-	-	-	-	-
Ethyl methanesulfonate	62-50-0	-	-	-	-	-	-	-	-	-
Ethylene dibromide	106-93-4	0.80	-	-	2.00E-04[f]	5.71E-05	8.50E+01	1.06E+02	7.60E-01	B2
Ethylene glycol	107-21-1	0.50	2.00E+00	1.00E+00	-	-	-	-	-	-
Ethylene glycol monobutyl ether	111-76-2	0.50	5.00E-01	2.50E-01	1.30E+01	3.71E+00	5.00E-01	-	-	-

Table 4-1. Human Health Chemical Toxicity Values for Organic Constituents of Potential Concern

Organic COPC	CAS Number	G.I. Absorption Factor [a] (unitless)	Chronic Noncarcinogenic Toxicity				Carcinogenic Slope Factor			EPA Class [b]
			Oral RfD [b] mg/(kg-d)	Dermal RfD [c] mg/(kg-d)	Inhalation RfC [b] mg/m^3	Inhalation RfD [d] mg/(kg-d)	Oral [b]	Dermal [c]	Inhalation [c]	
							(kg-day/mg)			
Ethylene glycol monoethyl ether acetate	111-15-9	0.50	3.00E-01[f]	1.50E-01	-	-	-	-	-	-
Ethylene thiourea	96-45-7	0.50	8.00E-05	4.00E-05	-	-	1.10E-01[f]	2.20E-01	-	B2[f]
Furfural	98-01-1	0.50	3.00E-03	1.50E-03	5.00E-02[f]	1.43E-02	-	-	-	-
Maleic hydrazide	123-33-1	0.50	5.00E-01	2.50E-01	-	-	-	-	-	-
Malononitrile	109-77-3	0.80	2.00E-05[f]	1.60E-05	-	-	-	-	-	-
Methyl styrene (mixed isomers)	25013-15-4	0.80	6.00E-03[f]	4.80E-03	4.00E-02[f]	1.14E-02	-	-	-	-
Methylhydrazine	60-34-4	0.50	-	-	-	-	-	-	-	-
N,N-Diphenylamine	122-39-4	0.50	2.50E-02	1.25E-02	-	-	-	-	-	-
Nitric acid, propyl ester	627-13-4	-	-	-	-	-	-	-	-	-
N-Nitrosodi-*n*-butylamine	924-16-3	0.50	-	-	-	-	5.40E+00	1.08E+01	5.40E+00	B2
N-Nitrosomorpholine	59-89-2	-	-	-	-	-	-	-	-	-
N-Nitroso-N,N-dimethylamine	62-75-9	0.50	-	-	-	-	5.10E+01	1.02E+02	5.10E+01	B2
o-Anisidine	90-04-0	-	-	-	-	-	-	-	-	-
Oxalic acid	144-62-7	-	-	-	-	-	-	-	-	-
Phthalic anhydride	85-44-9	0.50	2.00E+00	1.00E+00	1.20E-01[f]	3.43E-02	-	-	-	-
p-Phthalic acid	100-21-0	0.50	1.00E+00[f]	5.00E-01	-	-	-	-	-	-
Pyridine	110-86-1	0.50	1.00E-03	5.00E-04	-	-	-	-	-	-
Quinoline	91-22-5	0.50	-	-	-	-	1.20E+01[f]	2.40E+01	-	C[f]
Quinone	106-51-4	-	-	-	-	-	-	-	-	-
Safrole	94-59-7	-	-	-	-	-	-	-	-	-
Tetrahydrofuran	109-99-9	-	-	-	-	-	-	-	-	-
Other Heavy Semivolatile Compounds (molecular weight >200 g/mole)										
1,2,4,5-Tetrachlorobenzene	95-94-3	0.80	3.00E-04	2.40E-04	-	-	-	-	-	-
1,3,5-Trinitrobenzene	99-35-4	0.65	3.00E-02	1.95E-02	-	-	-	-	-	-
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	-	-	-	-	-	-	-	-	-
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	0.50	2.00E-03	1.00E-03	-	-	-	-	-	-
2-*sec*-Butyl-4,6-dinitrophenol	88-85-7	0.50	1.00E-03	5.00E-04	-	-	-	-	-	D
3,3'-Dimethoxybenzidine	119-90-4	0.50	-	-	-	-	1.40E-02[f]	2.80E-02	-	B2[f]
3,3-Dichlorobenzidine	91-94-1	0.50	-	-	-	-	4.50E-01	9.00E-01	-	B2
4-Bromophenylphenyl ether	101-55-3	0.50	-	-	-	-	-	-	-	D
Ammonium perfluorooctanoate	3825-26-1	-	-	-	-	-	-	-	-	-
Azobenzene	103-33-3	0.50	-	-	-	-	1.10E-01	2.22E-01	1.10E-01	B2
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	-	-	-	-	-	-	-	-	-
Captan	133-06-2	0.50	1.30E-01	6.50E-02	-	-	3.50E-03[f]	7.00E-03	-	B2[f]

Table 4-1. Human Health Chemical Toxicity Values for Organic Constituents of Potential Concern

Organic COPC	CAS Number	G.I. Absorption Factor [a] (unitless)	Chronic Noncarcinogenic Toxicity				Carcinogenic Slope Factor			EPA Class [b]
			Oral RfD [b] mg/(kg-d)	Dermal RfD [c] mg/(kg-d)	Inhalation RfC [b] mg/m³	Inhalation RfD [d] mg/(kg-d)	Oral [b]	Dermal [c]	Inhalation [e]	
							(kg-day/mg)			
Chlorobenzilate	510-15-6	0.50	2.00E-02	1.00E-02	-	-	2.70E-01[f]	5.40E-01	2.70E-01[f]	B2
Dibutylphosphate	107-66-4	-	-	-	-	-	-	-	-	-
Dimethyl aminoazobenzene	60-11-7	-	-	-	-	-	-	-	-	-
Hexachlorobenzene	118-74-1	0.50	8.00E-04	4.00E-04	-	-	1.60E+00	3.20E+00	1.60E+00	B2
Hexachlorobutadiene	87-68-3	0.50	2.00E-04[f]	1.00E-04	-	-	7.80E-02	1.56E-01	7.80E-02	C
Hexachlorocyclopentadiene	77-47-4	0.50	7.00E-03	3.50E-03	7.00E-05[f]	2.00E-05	-	-	-	D
Hexachloroethane	67-72-1	0.50	1.00E-03	5.00E-04	-	-	1.40E-02	2.80E-02	1.40E-02	C
Hexachlorophene	70-30-4	0.50	3.00E-04	1.50E-04	-	-	-	-	-	-
Hexamethylene-1,5-diisocyanate	822-06-0	0.50	-	-	1.00E-05	2.86E-06	-	-	-	-
Mirex	2385-85-5	0.50	2.00E-04	1.00E-04	-	-	-	-	-	B2[f]
Nitrofen	1836-75-5	-	-	-	-	-	-	-	-	-
Pentachlorobenzene	608-93-5	0.80	8.00E-04	6.40E-04	-	-	-	-	-	D
Pentachloronitrobenzene	82-68-8	0.80	3.00E-03	2.40E-03	-	-	2.60E-01[f]	3.25E-01	-	C[f]
Pentachlorophenol	87-86-5	1.00	3.00E-02	3.00E-02	-	-	1.20E-01		-	B2
Picric acid	88-89-1	-	-	-	-	-	-	-	-	-
Pronamide	23950-58-5	0.50	7.50E-02	3.75E-02	-	-	-	-	-	-
Strychnine	57-24-9	0.50	3.00E-04	1.50E-04	-	-	-	-	-	-
Terphenyls	26140-60-3	-	-	-	-	-	-	-	-	-
Tributyl phosphate	126-73-8	-	-	-	-	-	-	-	-	-
Trifluralin	1582-09-8	0.50	7.50E-03	3.75E-03	-	-	7.70E-03	1.54E-02	-	C
Triphenylamine	603-34-9	-	-	-	-	-	-	-	-	-
Herbicides and Organochlorinated Pesticides										
2,4,5-T	93-76-5	0.50	1.00E-02	5.00E-03	-	-	-	-	-	-
2,4-D and esters	94-75-7	0.80	1.00E-02	8.00E-03	-	-	-	-	-	-
4,4-DDD	72-54-8	0.70	-	-	-	-	2.40E-01	3.43E-01	-	B2
4,4-DDE	72-55-9	0.70	-	-	-	-	3.40E-01	4.86E-01	-	B2
4,4-DDT	50-29-3	0.70	5.00E-04	3.50E-04	-	-	3.40E-01	4.86E-01	3.40E-01	B2
Aldrin	309-00-2	0.50	3.00E-05	1.50E-05	-	-	1.70E+01	3.4E+01	1.70E+01	B2
alpha-BHC	319-84-6	0.97	-	-	-	-	6.30E+00	6.49E+00	6.30E+00	B2
beta-BHC	319-85-7	0.91	-	-	-	-	1.80E+00	1.98E+00	1.80E+00	C
Chlordane	57-74-9	0.50	5.00E-04	2.50E-04	7.00E-04	2.00E-04	3.50E-01	7.00E-01	1.30E+00	B2
Delta-BHC	319-86-8	0.50	-	-	-	-	-	-	-	D
Dieldrin	60-57-1	0.50	5.00E-05	2.50E-05	-	-	1.60E+01	3.2E+01	1.60E+01	B2
Endothall	145-73-3	0.50	2.00E-02	1.00E-02	-	-	-	-	-	-
Endrin	72-20-8	0.02	3.00E-04	6.00E-06	-	-	-	-	-	D

Table 4-1. Human Health Chemical Toxicity Values for Organic Constituents of Potential Concern

Organic COPC	CAS Number	G.I. Absorption Factor [a] (unitless)	Chronic Noncarcinogenic Toxicity				Carcinogenic Slope Factor			EPA Class [b]
			Oral RfD [b] mg/(kg-d)	Dermal RfD [c] mg/(kg-d)	Inhalation RfC [b] mg/m³	Inhalation RfD [d] mg/(kg-d)	Oral [b]	Dermal [c]	Inhalation [e]	
							(kg-day/mg)			
gamma-BHC (Lindane)	58-89-9	0.97	3.00E-04	2.91E-04	-	-	1.30E+00[f]	1.34E+00	-	B2[f]
Heptachlor	76-44-8	0.72	5.00E-04	3.60E-04	-	-	4.50E+00	6.25E+00	4.50E+00	B2
Isodrin	465-73-6	-	-	-	-	-	-	-	-	-
Methoxychlor	72-43-5	0.50	5.00E-03	2.50E-03	-	-	-	-	-	D
Silvex (2,4,5-TP)	93-72-1	0.50	8.00E-03	4.00E-03	-	-	-	-	-	D
Toxaphene	8001-35-2	0.50	-	-	-	-	1.10E+00	2.20E+00	1.10E+00	B2

- = not available
RfC = reference concentration
RfD = reference dose

[a] Gastrointestinal absorption factors are from the Risk Assessment Information System (RAIS) maintained by the Oak Ridge National Laboratory (ORNL); see the web site http://risk.lsd.ornl.gov/tox/tox_values.html. These values are used in the calculation of dermal toxicity values from oral toxicity values.

[b] Oral RfDs, inhalation RfCs, oral slope factors, and EPA classifications are from IRIS (EPA 1998c), unless otherwise noted.

[c] Dermal toxicity values are calculated from oral toxicity values as follows: Dermal RfD = (Oral RfD) x (G.I. Absorption Factor); Dermal Slope Factor = (Oral Slope Factor)/(G. I. Absorption Factor).

[d] Inhalation RfDs are calculated from inhalation RfCs; see section 7.2.1 (Noncarcinogenic Health Effects).

[e] Inhalation slope factors are from HEAST (EPA 1997a), unless otherwise noted.

[f] Toxicity value obtained from HEAST (EPA 1997a).

[g] The inhalation slope factor was calculated from the inhalation unit risk, which is provided by IRIS (EPA 1998c). See section 7.2.2.

[h] Provisional value obtained from Superfund Health Risk Technical Support Center, which can be contacted at (513) 569-7300.

[i] Toxicity Equivalency Factor (TEF) approach is used to evaluate these dioxins/furans, with exposure concentrations converted to equivalent concentrations of 2,3,7,8-TCDD by multiplying the concentration by the appropriate TEF. See section 7.2.3.

[j] Coplanar PCBs are evaluated with the TEF approach, using 2,3,7,8-TCDD as the basis; see section 7.2.5. Non-coplanar PCBs are evaluated with the toxicity of PCBs (including RfDs for Aroclor-1254).

[k] Toxicity value obtained from IRIS (EPA 1998c).

[l] The cancer potency of PCB mixtures is determined using a three tiered approach that depends on the information provided. Criteria for use of the High Risk and Persistence Tier include: food chain exposure; sediment or soil ingestion; dust or aerosol inhalation; dermal exposure if an absorption factor has been applied; any early-life exposure; and the presence of dioxin-like, tumor-promoting, or persistent congeners. The value presented, 2.00E+00 $(mg/kg\text{-}day)^{-1}$, is the upper bound slope factor for the High Risk and Persistence Tier. The central-estimate slope factor for the tier is 1.00E+00 $(mg/kg\text{-}day)^{-1}$.

[m] TEF approach is used to evaluate these carcinogenic PAHs, with exposure concentrations converted to equivalent concentrations of benzo(*a*)pyrene by multiplying the concentration by the appropriate TEF. See section 7.2.4.

[n] Provisional inhalation toxicity value developed by NCEA.

[o] Cancer Slope factor is from IRIS (EPA 1998c), but is for "Dinitrotoluene mixture, 2,4/2,6-." The value presented is based on a study using technical grade DNT.

Table 4-2. Human Health Toxicity Values for Inorganic Constituents of Potential Concern

Inorganic COPC	CAS Number	G.I. Absorption Factor [a] (unitless)	Chronic Noncarcinogenic Toxicity				Carcinogenic Slope Factor			EPA Class [b]
			Oral RfD [b] mg/(kg-d)	Dermal RfD [c] mg/(kg-d)	Inhalation RfC [b] mg/m³	Inhalation RfD [d] mg/(kg-d)	Oral [b] (kg-day/mg)	Dermal [c] (kg-day/mg)	Inhalation [e] (kg-day/mg)	
Metals										
Aluminum	7429-90-5	0.10	-	-	-	-	-	-	-	-
Antimony	7440-36-0	0.02	4.00E-04	8.00E-06	-	-	-	-	-	-
Arsenic	7440-38-2	0.41	3.00E-04	1.23E-04	-	-	1.50E+00	3.66E+00	-	A
Barium	7440-39-3	0.07	7.00E-02	4.90E-03	5.00E-04 [f]	1.43E-04	-	-	-	D
Beryllium	7440-41-7	0.01	2.00E-03	2.00E-05	2.00E-02	5.71E-03	-	-	8.40E+00	B1
Bismuth	7440-69-9	-	-	-	-	-	-	-	-	-
Boron	7440-42-8	0.90	9.00E-02	8.10E-02	2.00E-02 [f]	5.71E-03	-	-	-	-
Cadmium	7440-43-9	0.01	1.00E-03	1.00E-05	-	-	-	-	-	B1
Calcium	7440-70-2	-	-	-	-	-	-	-	-	-
Chromium (particulates) [g]	7440-47-3	0.02	3.00E-03	6.00E-05	1.00E-04	2.86E-05	-	-	4.10E+01	A
Chromium (chromic acid mist) [g]	7440-47-3	0.02	3.00E-03	6.00E-05	8.00E-06	2.29E-06	-	-	4.10E+01	A
Cobalt	7440-48-4	0.80	-	-	-	-	-	-	-	-
Copper	7440-50-8	0.30	-	-	-	-	-	-	-	D
Iron	7439-89-6	0.15	-	-	-	-	-	-	-	-
Lead	7439-92-1	0.15	-	-	-	-	-	-	-	B2
Lithium	7439-93-2	0.80	-	-	-	-	-	-	-	-
Magnesium	7439-95-4	0.20	-	-	-	-	-	-	-	-
Manganese	7439-96-5	0.04	4.60E-02 [h]	1.84E-03	5.00E-05	1.43E-05	-	-	-	D
Mercury	7439-97-6	0.07	-	-	-	-	-	-	-	C
Molybdenum	7439-98-7	0.38	5.00E-03	1.90E-03	-	-	-	-	-	-
Nickel [i]	7440-02-0	0.27	2.00E-02 [i]	5.40E-03	-	-	-	-	8.40E-01 [i]	A [i]
Potassium	7440-09-7	-	-	-	-	-	-	-	-	-
Rhodium	7440-16-6	-	-	-	-	-	-	-	-	-
Selenium	7782-49-2	0.44	5.00E-03	2.20E-03	-	-	-	-	-	D
Silicon	7440-21-3	-	-	-	-	-	-	-	-	-
Silver	7440-22-4	0.18	5.00E-03	9.00E-04	-	-	-	-	-	D
Sodium	7440-23-5	1.00	-	-	-	-	-	-	-	-
Strontium	7440-24-6	0.20	6.00E-01	1.20E-01	-	-	-	-	-	-
Tantalum	7440-25-7	-	-	-	-	-	-	-	-	-
Thallium	7440-28-0	0.15	-	-	-	-	-	-	-	-
Tin	7440-31-5	0.10	6.00E-01 [f]	6.00E-02	-	-	-	-	-	-
Tungsten	7440-33-7	-	-	-	-	-	-	-	-	-
Uranium	7440-61-1	0.85	3.00E-03	2.55E-03	-	-	-	-	-	-
Vanadium	7440-62-2	0.01	7.00E-03 [f]	7.00E-05	-	-	-	-	-	-
Yttrium	7440-65-5	-	-	-	-	-	-	-	-	-
Zinc	7440-66-6	0.20	3.00E-01	6.00E-02	-	-	-	-	-	D
Zirconium	7440-67-7	0.80	-	-	-	-	-	-	-	-
Non-metals and Anions										
Ammonia/ Ammonium	7664-41-7	0.20	-	-	1.00E-01	2.86E-02	-	-	-	-
Bromide	24959-67-9	-	-	-	-	-	-	-	-	-
Chloride	16887-00-6	1.00	-	-	-	-	-	-	-	-
Cyanide	57-12-5	0.17	2.00E-02	3.40E-03	-	-	-	-	-	D
Fluoride [j]	16984-48-8	0.97	6.00E-02	5.82E-02	-	-	-	-	-	-

Table 4-2. Human Health Toxicity Values for Inorganic Constituents of Potential Concern

Inorganic COPC	CAS Number	G.I. Absorption Factor [a] (unitless)	Chronic Noncarcinogenic Toxicity				Carcinogenic Slope Factor			EPA Class [b]
			Oral RfD [b] mg/(kg-d)	Dermal RfD [c] mg/(kg-d)	Inhalation RfC [b] mg/m³	Inhalation RfD [d] mg/(kg-d)	Oral [b] (kg-day/mg)	Dermal [c] (kg-day/mg)	Inhalation [e] (kg-day/mg)	
Hydroxide	14280-30-9	-	-	-	-	-	-	-	-	-
Iodine	7553-56-2	-	-	-	-	-	-	-	-	-
Nitrate	14797-55-8	0.50	1.60E+00	8.00E-01	-	-	-	-	-	-
Nitrite	14797-65-0	0.50	1.00E-01	5.00E-02	-	-	-	-	-	-
Phosphate	14265-44-2	-	-	-	-	-	-	-	-	-
Phosphorus	7723-14-0	0.20	-	-	-	-	-	-	-	-
Sulfate	14808-79-8	0.20	-	-	-	-	-	-	-	-
Total Sulfur	63705-05-5	-	-	-	-	-	-	-	-	-
Criteria Pollutants										
Carbon dioxide	124-38-9	-	-	-	-	-	-	-	-	-
Nitrogen dioxide	10102-44-0	0.20	-	-	-	-	-	-	-	-
Ozone	10028-15-6	-	-	-	-	-	-	-	-	-
Particulate matter	no cas #	-	-	-	-	-	-	-	-	-
Sulfur dioxide	7446-09-5	-	-	-	-	-	-	-	-	-

CAS = Chemical Abstract Service
- = not available
NOAEL = no observed adverse effect level
RfC = reference concentration
RfD = reference dose
TRV = toxicity reference value

[a] Gastrointestinal absorption factors are from the Risk Assessment Information System (RAIS) maintained by the Oak Ridge National Laboratory (ORNL); see the web site http://risk.lsd.ornl.gov/tox-values.html. These values are used in the calculation of dermal toxicity from oral toxicity values.

[b] Oral RfDs, inhalation RfCs, oral slope factors, and EPA classifications are from IRIS (EPA 1998c), unless otherwise noted.

[c] Dermal toxicity values are calculated from oral toxicity values as follows: Dermal RfD = (Oral RfD) × (G.I. Absorption Factor); Dermal Slope Factor = (Oral Slope Factor)/(G.I. Absorption Factor).

[d] Inhalation RfDs are calculated from inhalation RfCs; see section 7.2.1 (Noncarcinogenic Health Effects).

[e] Inhalation slope factors are from HEAST (EPA 1997a), unless otherwise noted.

[f] Toxicity value obtained from HEAST (EPA 1997a).

[g] Toxicity for Chromium VI (chromic acid mist) is used to evaluate chromium in water; otherwise, the toxicity for Chromium VI (particulates) is used to evaluate chromium.

[h] The chronic oral RfD for the total oral intake of manganese is 1.40E-01. However, when assessing exposure to manganese from drinking water or soil, IRIS (EPA 1998c) recommends using a modifying factor of 3, thereby lowering the RfD to 4.60E-02.

[i] Nickel will be evaluated as a potential carcinogen via the inhalation pathway using a slope factor calculated from the inhalation unit risk factor for nickel refinery dust. The oral RfD for nickel soluble salts will be used to evaluate noncarcinogenic effects for nickel. See section 7.2.8.

[j] Toxicity for Fluorine (Soluble Fluoride) is used to evaluate Fluoride.

Table 4-3. Human Health Toxicity Values for Radionuclides of Potential Concern

Radionuclide	CAS Number	Ingestion [a] (risk/pCi)	Inhalation [a] (risk/pCi)	External Radiation to Soil [a] (risk/year per pCi/g soil)	External Radiation for Air Submersion [b] (m^3/Bq-sec)	Isotopes used to calculate Slope Factor for External Radiation for Air Submersion [c]
Actinium-227+D [d]	14952-40-0	6.26E-10	7.87E-08	9.30E-07	1.34E-15	Ac-227, Th-227 (98.62%), Fr-223 (1.38%), Ra-223, Rn-219, Po-215, Pb-211, Bi-211, Tl-207 (99.727%), Po-211 (0.273%), Pb-207
Americium-241	14596-10-2	3.28E-10	3.85E-08	4.59E-09	5.00E-17	Am-241
Americium-243+D [d]	14993-75-0	3.31E-10	3.82E-08	2.66E-07	6.83E-16	Am-243, Np-239
Antimony-125+D [d]	14234-35-6	3.54E-12	5.85E-12	1.34E-06	1.52E-15	Sb-125, Te-125
Barium-137m [e]	13981-97-0	2.43E-15	1.57E-15	2.21E-06	2.16E-15	
Cadmium-113	no cas #	NA	NA	NA	3.29E-19	
Carbon-14	14762-75-5	1.03E-12	6.99E-15	0.00E+00	3.66E-20	
Cesium-134	13967-70-9	4.73E-11	2.89E-11	5.88E-06	5.68E-15	
Cesium-137+D [d]	10045-97-3	3.16E-11	1.91E-11	2.09E-06	1.37E-18	Cs-137
Cobalt-60	10198-40-0	1.89E-11	6.88E-11	9.76E-06	9.63E-15	
Curium-242	15510-73-3	3.83E-11	3.16E-09	2.34E-11	2.59E-19	Cm-242
Curium-243+D [d]	15757-87-6	2.51E-10	2.89E-08	1.71E-07	4.16E-16	Cm-243
Curium-244	13981-15-2	2.11E-10	2.43E-08	2.07E-11	2.15E-19	Cm-244
Europium-152	14683-23-9	5.73E-12	7.91E-11	4.08E-06	4.25E-15	Eu-152, Gd-152
Europium-154	15585-10-1	9.37E-12	9.15E-11	4.65E-06	4.63E-15	
Europium-155	14391-16-3	1.65E-12	9.60E-12	6.08E-08	1.64E-16	
Iodine-129	15046-84-1	1.84E-10	1.22E-10	2.69E-09	1.85E-17	
Neptunium-237+D [d]	13994-20-2	3.00E-10	3.45E-08	4.62E-07	7.43E-16	Np-237, Pa-233
Nickel-59	14336-70-0	1.85E-13	4.01E-13	0.00E+00	0.00E+00	
Nickel-63	13981-37-8	5.50E-13	1.01E-12	0.00E+00	0.00E+00	
Niobium-93m [e]	7440-03-1	6.64E-13	4.33E-12	3.64E-11	1.92E-19	
Plutonium-238	13981-16-3	2.95E-10	2.74E-08	1.94E-11	2.28E-19	Pu-238
Plutonium-239	15117-48-3	3.16E-10	2.78E-08	1.26E-11	2.56E-19	Pu-239
Plutonium-240	14119-33-6	3.15E-10	2.78E-08	1.87E-11	2.24E-19	Pu-240
Plutonium-241+D [d]	14119-32-5	5.20E-12	2.81E-10	0.00E+00	4.89E-21	Pu-241
Plutonium-242	13982-10-0	3.00E-10	2.64E-08	1.55E-11	1.91E-19	Pu-242
Protactinium-231	14331-85-2	1.49E-10	2.42E-08	2.71E-08	1.24E-16	Pa-231
Radium-226+D [d]	13982-63-3	2.96E-10	2.75E-09	6.74E-06	6.74E-15	Ra-226, Rn-222, Po-218, Pb-214, Bi-214, Po-214 (99.979%), Pb-210, Bi-210, Po-210, Pb-206
Radium-228+D [d]	15262-20-1	2.48E-10	9.94E-10	3.28E-06	9.77E-15	Ra-228, Ac-228, Th-228, Ra-224, Rn-220, Po-216, Pb-212, Bi-212, Po-212 (64.07%), Tl-208 (35.93%), Pb-208

Table 4-3. Human Health Toxicity Values for Radionuclides of Potential Concern

Radionuclide	CAS Number	Ingestion [a] (risk/pCi)	Inhalation [a] (risk/pCi)	External Radiation to Soil [a] (risk/year per pCi/g soil)	External Radiation for Air Submersion [b] (m^3/Bq-sec)	Isotopes used to calculate Slope Factor for External Radiation for Air Submersion [c]
Ruthenium-106+D [d]	13967-48-1	3.45E-11	1.15E-10	7.57E-07	7.85E-16	Ru-106, Rh-106, Pd-106
Samarium-151	15715-94-3	4.60E-13	4.63E-12	2.92E-13	1.52E-21	
Selenium-79	no cas #	NA	NA	NA	5.39E-20	
Strontium-90+D [d]	10098-97-2	5.59E-11	6.93E-11	1.40E-18	1.40E-18	Sr-90
Technetium-99	14133-76-7	1.40E-12	2.89E-12	6.19E-13	3.72E-19	
Thorium-229+D [d]	15594-54-4	3.56E-10	8.26E-08	7.68E-07	1.08E-15	Th-229, Ra-225, Ac-225, Fr-221, At-217, Bi-213, Po-213 (97.84%), Tl-209 (2.16%), Pb-209
Thorium-232	7440-29-1	3.28E-11	1.93E-08	1.97E-11	5.35E-19	Th-232
Tin-126	15832-50-5	2.12E-11	4.26E-11	3.32E-08	1.04E-14	Sn-126, Sb-126
Tritium	10028-17-8	7.15E-14	9.59E-14	0.00E+00	0.00E+00	
Uranium-232	14158-29-3	8.12E-11	5.29E-08	3.42E-11	8.67E-19	U-232
Uranium-233	13968-55-3	4.48E-11	1.41E-08	3.52E-11	1.09E-18	U-233
Uranium-234	13966-29-5	4.44E-11	1.40E-08	2.14E-11	1.56E-18	U-234, Th-230
Uranium-235+D [d]	15117-96-1	4.70E-11	1.30E-08	2.65E-07	5.43E-16	U-235, Th-231
Uranium-236	13982-70-2	4.21E-11	1.32E-08	1.72E-11	2.67E-19	U-236
Uranium-238+D [d]	7440-61-1	6.20E-11	1.24E-08	6.57E-08	7.04E-15	U-238, Th-234, Pa-234
Yttrium-90	10098-91-6	1.50E-11	9.90E-12	0.00E+00	1.66E-14	Y-90, Zr-90, Nb-90
Zirconium-93	15751-77-6	5.21E-13	5.26E-12	0.00E+00	0.00E+00	Zr-93

NA = slope factor is not available

[a] Cancer slope factors are from HEAST (EPA 1995d).

[b] External exposure slope factors for air submersion (for morbidity) are from Table 2.3 in EPA Federal Guidance Report No. 13 (EPA 1999b).

[c] Isotopes used to calculate the external exposure slope factors for air submersion are listed for parent isotopes only. All daughters in the decay chain are included in the calculation of the slope factor, until another isotope on the ROPC list is encountered. For example, Neptunium-237 decays into Protactinium-233, which decays into Uranium-233, which decays into other isotopes. Since Uranium-233 is on the ROPC list, its slope factor is not used to calculate the slope factor for Neptunium-237. The external exposure slope factor for air submersion for Neptunium-237 is obtained by adding the values in Table 2.3 (from EPA 1999b) for Neptunium-237 and Protactinium-233.

[d] +D slope factors from HEAST include contributions from short-lived daughter products.

[e] m designates isotopes in a metastable state.

Table 4-4. Acute Inhalation Exposure Criteria for COPCs

COPC	CAS Number	NCEA Provis. Value[a]		AEGL-1[b]		ERPG-1[c]		AREL[d]		TEEL-1[e]		Conv. Factor[f]	AIEC[g] (mg/m³)
ORGANICS	-	-	-	-	-	-	-	-	-	-	-	-	-
Aromatic Halogenated Hydrocarbons													
2,3,4,6-Tetrachlorophenol	58-90-2	-	-	-	-	-	-	-	-	-	-	-	-
4-Chloro-3-methylphenol	59-50-7	-	-	-	-	-	-	-	-	2.00E+01	mg/m³	5.83E+00	2.00E+01
Aromatic Nonhalogenated Hydrocarbons													
2-Nitrotoluene	88-72-2	-	-	-	-	-	-	-	-	-	-	-	-
4-Nitrobiphenyl	92-93-3	-	-	-	-	-	-	-	-	7.50E-01	mg/m³	8.14E+00	7.50E-01
Benzaldehyde	100-52-7	-	-	-	-	-	-	-	-	1.50E+01	mg/m³	4.34E+00	1.50E+01
Benzene	71-43-2	-	-	-	-	5.00E+01	ppm	1.30E+00	mg/m³	1.60E+02	mg/m³	3.19E+00	1.60E+02
Benzyl alcohol	100-51-6	-	-	-	-	-	-	-	-	5.53E+01	mg/m³	4.42E+00	5.53E+01
Ethyl benzene	100-41-4	-	-	-	-	-	-	-	-	5.42E+02	mg/m³	4.34E+00	5.42E+02
m-Xylene	108-38-3	-	-	-	-	-	-	2.20E+01	mg/m³	6.51E+02	mg/m³	4.34E+00	2.20E+01
o-Xylene	95-47-6	-	-	-	-	-	-	2.20E+01	mg/m³	6.51E+02	mg/m³	4.34E+00	2.20E+01
p-Xylene	106-42-3	-	-	-	-	-	-	2.20E+01	mg/m³	6.51E+02	mg/m³	4.34E+00	2.20E+01
Styrene	100-42-5	-	-	-	-	5.00E+01	ppm	2.10E+01	mg/m³	2.13E+02	mg/m³	4.26E+00	2.13E+02
Toluene	108-88-3	-	-	-	-	5.00E+01	ppm	3.70E+01	mg/m³	1.88E+02	mg/m³	3.77E+00	1.88E+02
Non-aromatic Nonhalogenated Hydrocarbons													
1,2-Epoxybutane	106-88-7	-	-	-	-	-	-	-	-	2.95E+02	mg/m³	2.95E+00	2.95E+02
1,3-Butadiene	106-99-0	-	-	-	-	1.00E+01	ppm			2.21E+01	mg/m³	2.21E+00	2.21E+01
1,4-Dioxane	123-91-1	-	-	-	-	-	-	3.00E+00	mg/m³	2.70E+02	mg/m³	3.60E+00	3.00E+00
1-Methylpropyl alcohol	78-92-2	-	-	-	-	-	-	-	-	4.55E+02	mg/m³	3.03E+00	4.55E+02
1-Nitropropane	108-03-2	-	-	-	-	-	-	-	-	2.73E+02	mg/m³	3.64E+00	2.73E+02
2,2,4-Trimethylpentane	540-84-1	-	-	-	-	-	-	-	-	3.50E+02	mg/m³	4.67E+00	3.50E+02
2-Butanone	78-93-3	-	-	-	-	-	-	1.30E+01	mg/m³	8.84E+02	mg/m³	2.95E+00	1.30E+01
2-Butenaldehyde (2-Butenal)	4170-30-3	-	-	1.90E-01	ppm	2.00E+00	ppm	-	-	5.73E+00	mg/m³	2.86E+00	5.44E-01
2-Ethoxyethanol	110-80-5	-	-	-	-	-	-	3.70E-01	mg/m³	5.53E+01	mg/m³	3.68E+00	3.70E-01
2-Heptanone	110-43-0	-	-	-	-	-	-	-	-	7.00E+02	mg/m³	4.67E+00	7.00E+02
2-Hexanone	591-78-6	-	-	-	-	-	-	-	-	4.09E+01	mg/m³	4.09E+00	4.09E+01
2-Methoxyethanol	109-86-4	-	-	-	-	-	-	9.30E-02	mg/m³	4.67E+01	mg/m³	3.11E+00	9.30E-02
2-Methyl-2-propanol	75-65-0	-	-	-	-	-	-	-	-	4.55E+02	mg/m³	3.03E+00	4.55E+02
2-Methyl-2-propenenitrile	126-98-7	-	-	-	-	-	-	-	-	2.74E+00	mg/m³	2.74E+00	2.74E+00
2-Methylaziridine	75-55-8	-	-	-	-	-	-	-	-	1.40E+01	mg/m³	2.33E+00	1.40E+01
2-Methylpropyl alcohol	78-83-1	-	-	-	-	-	-	-	-	4.55E+02	mg/m³	3.03E+00	4.55E+02
2-Pentanone	107-87-9	-	-	-	-	-	-	-	-	8.80E+02	mg/m³	3.52E+00	8.80E+02
2-Propanone (Acetone)	67-64-1	-	-	-	-	-	-	-	-	2.37E+03	mg/m³	2.37E+00	2.37E+03
2-Propene-1-ol	107-18-6	-	-	1.80E+00	ppm	-	-	-	-	9.50E+00	mg/m³	2.37E+00	4.27E+00
2-Propyl alcohol	67-63-0	-	-	-	-	-	-	3.20E+00	mg/m³	9.83E+02	mg/m³	2.46E+00	3.20E+00
3-Heptanone	106-35-4	-	-	-	-	-	-	-	-	-	-	-	-
3-Methyl-1-butanol	123-51-3	-	-	-	-	-	-	-	-	4.50E+02	mg/m³	3.60E+00	4.50E+02
3-Methyl-2-butanone	563-80-4	-	-	-	-	-	-	-	-	-	-	-	-

Table 4-4. Acute Inhalation Exposure Criteria for COPCs

COPC	CAS Number	NCEA Provis. Value[a]		AEGL-1[b]		ERPG-1[c]		AREL[d]		TEEL-1[e]		Conv. Factor[f]	AIEC[g] (mg/m³)
3-Pentanone	96-22-0	-	-	-	-	-	-	-	-	-	-	-	-
4-Heptanone	123-19-3	-	-	-	-	-	-	-	-	-	-	-	-
4-Methyl-2-pentanone	108-10-1	-	-	-	-	-	-	-	-	3.07E+02	mg/m³	4.09E+00	3.07E+02
4-Methyl-3-penten-2-one	141-79-7	-	-	-	-	-	-	-	-	1.00E+02	mg/m³	4.01E+00	1.00E+02
5-Methyl-2-hexanone	110-12-3	-	-	-	-	-	-	-	-	2.80E+03	mg/m³	4.67E+00	2.80E+03
Acetaldehyde	75-07-0	-	-	-	-	1.00E+01	ppm	-	-	1.80E+01	mg/m³	1.80E+00	1.80E+01
Acetamide	60-35-5	-	-	-	-	-	-	-	-	7.50E+01	mg/m³	2.41E+00	7.50E+01
Acetic acid	64-19-7	-	-	-	-	-	-	-	-	3.68E+01	mg/m³	2.45E+00	3.68E+01
Acetic acid ethyl ester	141-78-6	-	-	-	-	-	-	-	-	4.32E+03	mg/m³	3.60E+00	4.32E+03
Acetic acid n-butyl ester	123-86-4	-	-	-	-	-	-	-	-	-	-	-	-
Acetonitrile	75-05-8	-	-	-	-	-	-	-	-	1.01E+02	mg/m³	1.68E+00	1.01E+02
Acrolein	107-02-8	-	-	3.00E-02	ppm	1.00E-01	ppm	1.90E-04	mg/m³	2.29E-01	mg/m³	2.29E+00	6.87E-02
Acrylonitrile	107-13-1	-	-	-	-	1.00E+01	ppm	-	-	2.17E+01	mg/m³	2.17E+00	2.17E+01
Bis(isopropyl)ether	108-20-3	-	-	-	-	-	-	-	-	-	-	-	-
Butane	106-97-8	-	-	-	-	-	-	-	-	5.70E+03	mg/m³	2.38E+00	5.70E+03
Carbon disulfide	75-15-0	-	-	-	-	1.00E+00	ppm	6.20E+00	mg/m³	3.73E+01	mg/m³	3.11E+00	3.11E+00
Cyanogen	460-19-5	-	-	-	-	-	-	-	-	6.38E+01	mg/m³	2.13E+00	6.38E+01
Cyclohexane	110-82-7	-	-	-	-	-	-	-	-	3.10E+03	mg/m³	3.44E+00	3.10E+03
Cyclohexanone	108-94-1	-	-	-	-	-	-	-	-	3.01E+02	mg/m³	4.01E+00	3.01E+02
Cyclohexene	110-83-8	-	-	-	-	-	-	-	-	1.01E+03	mg/m³	3.36E+00	1.01E+03
Cyclopentane	287-92-3	-	-	-	-	-	-	-	-	-	-	-	-
Ethyl alcohol	64-17-5	-	-	-	-	-	-	-	-	5.65E+03	mg/m³	1.88E+00	5.65E+03
Ethyl ether	60-29-7	-	-	-	-	-	-	-	-	1.52E+03	mg/m³	3.03E+00	1.52E+03
Ethyl methacrylate	97-63-2	-	-	-	-	-	-	-	-	-	-	-	-
Formaldehyde	50-00-0	-	-	-	-	1.00E+00	ppm	9.40E-02	mg/m³	1.23E+00	mg/m³	1.23E+00	1.23E+00
Formamide	75-12-7	-	-	-	-	-	-	-	-	2.76E+01	mg/m³	1.84E+00	2.76E+01
Formic acid	64-18-6	-	-	-	-	-	-	-	-	1.88E+01	mg/m³	1.88E+00	1.88E+01
Formic acid, methyl ester	107-31-3	-	-	-	-	-	-	-	-	-	-	-	-
Glycidylaldehyde	765-34-4	-	-	-	-	-	-	-	-	2.21E-01	mg/m³	2.95E+00	2.21E-01
Methyl acetate	79-20-9	-	-	-	-	-	-	-	-	-	-	-	-
Methyl alcohol	67-56-1	-	-	-	-	2.00E+02	ppm	2.80E+01	mg/m³	2.62E+02	mg/m³	1.31E+00	2.62E+02
Methyl isocyanate	624-83-9	-	-	-	-	2.50E-02	ppm	-	-	5.83E-02	mg/m³	2.33E+00	5.83E-02
Methyl methacrylate	80-62-6	-	-	-	-	-	-	-	-	1.23E+03	mg/m³	4.09E+00	1.23E+03
Methyl tert-butyl ether	1634-04-4	-	-	-	-	-	-	-	-	4.32E+02	mg/m³	3.60E+00	4.32E+02
Methylacetylene	74-99-7	-	-	-	-	-	-	-	-	2.78E+03	mg/m³	1.64E+00	2.78E+03
Methylcyclohexane	108-87-2	-	-	-	-	-	-	-	-	4.82E+03	mg/m³	4.01E+00	4.82E+03
N,N-Dimethylacetamide	127-19-5	-	-	-	-	-	-	-	-	-	-	-	-
n-Butyl alcohol	71-36-3	-	-	-	-	-	-	-	-	1.52E+02	mg/m³	3.03E+00	1.52E+02
n-Heptane	142-82-5	-	-	-	-	-	-	-	-	1.80E+03	mg/m³	4.10E+00	1.80E+03
n-Hexane	110-54-3	-	-	-	-	-	-	-	-	5.28E+02	mg/m³	3.52E+00	5.28E+02
Nitromethane	75-52-5	-	-	-	-	-	-	-	-	1.50E+02	mg/m³	2.50E+00	1.50E+02

Table 4-4. Acute Inhalation Exposure Criteria for COPCs

COPC	CAS Number	NCEA Provis. Value[a]		AEGL-1[b]		ERPG-1[c]		AREL[d]		TEEL-1[e]		Conv. Factor[f]	AIEC[g] (mg/m³)
n-Nonane	111-84-2	-	-	-	-	-	-	-	-	1.05E+03	mg/m³	5.24E+00	1.05E+03
n-Octane	111-65-9	-	-	-	-	-	-	-	-	-	-	-	-
n-Pentane	109-66-0	-	-	-	-	-	-	-	-	1.80E+03	mg/m³	2.95E+00	1.80E+03
n-Propionaldehyde	123-38-6	-	-	-	-	-	-	-	-	7.50E+01	mg/m³	2.37E+00	7.50E+01
n-Propyl alcohol	71-23-8	-	-	-	-	-	-	-	-	6.14E+02	mg/m³	2.46E+00	6.14E+02
n-Valeraldehyde	110-62-3	-	-	-	-	-	-	-	-	-	-	-	-
Oxirane	75-21-8	-	-	-	-	-	-	-	-	1.35E+01	mg/m³	1.80E+00	1.35E+01
p-Cymene	99-87-6	-	-	-	-	-	-	-	-	-	-	-	-
Phosgene	75-44-5	-	-	-	-	-	-	4.00E-03	mg/m³	4.04E-01	mg/m³	4.04E+00	4.00E-03
Propargyl alcohol	107-19-7	-	-	-	-	-	-	-	-	6.87E+00	mg/m³	2.29E+00	6.87E+00
Propionic acid	79-09-4	-	-	-	-	-	-	-	-	3.03E+01	mg/m³	3.03E+00	3.03E+01
Propionitrile	107-12-0	-	-	-	-	-	-	-	-	3.38E+01	mg/m³	2.25E+00	3.38E+01
Propylene gylcol monomethyl ether	107-98-2	-	-	-	-	-	-	-	-	5.52E+02	mg/m³	3.68E+00	5.52E+02
p-tert-Butyltoluene	98-51-1	-	-	-	-	-	-	-	-	-	-	-	-
Triethylamine	121-44-8	-	-	-	-	-	-	2.80E+00	mg/m³	1.24E+01	mg/m³	4.14E+00	2.80E+00
Trimethylamine	75-50-3	-	-	-	-	1.00E-01	ppm	-	-	3.63E+01	mg/m³	2.42E+00	2.42E-01
Vinyl acetate	108-05-4	-	-	-	-	5.00E+00	ppm	-	-	1.76E+01	mg/m³	3.52E+00	1.76E+01
Non-aromatic Halogenated Hydrocarbons													
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	-	-	-	-	-	-	-	-	-	-	-	-
1,1,1,2-Tetrachloroethane	630-20-6	-	-	-	-	-	-	-	-	5.14E+01	mg/m³	6.86E+00	5.14E+01
1,1,1-Trichloroethane	71-55-6	-	-	-	-	3.50E+02	ppm	6.80E+01	mg/m³	1.91E+03	mg/m³	5.45E+00	1.91E+03
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	-	-	-	-	-	-	-	-	-	-	-	-
1,1,2,2-Tetrachloroethane	79-34-5	-	-	-	-	-	-	-	-	2.06E+01	mg/m³	6.86E+00	2.06E+01
1,1,2,2-Tetrachloroethene	127-18-4	-	-	-	-	1.00E+02	ppm	2.00E+01	mg/m³	6.78E+02	mg/m³	6.78E+00	6.78E+02
1,1,2-Trichloroethane	79-00-5	-	-	-	-	-	-	-	-	1.64E+02	mg/m³	5.45E+00	1.64E+02
1,1,2-Trichloroethylene	79-01-6	-	-	-	-	1.00E+02	ppm	-	-	5.37E+02	mg/m³	5.37E+00	5.37E+02
1,1-Dichloroethane	75-34-3	-	-	-	-	-	-	-	-	1.21E+03	mg/m³	4.04E+00	1.21E+03
1,1-Dichloroethene	75-35-4	-	-	-	-	-	-	-	-	7.92E+01	mg/m³	3.96E+00	7.92E+01
1,2,2-Trichloro-1,1,2-trifluoroethane	76-13-1	-	-	-	-	-	-	-	-	9.57E+03	mg/m³	7.66E+00	9.57E+03
1,2,3-Trichloropropane	96-18-4	-	-	-	-	-	-	-	-	1.81E+02	mg/m³	6.03E+00	1.81E+02
1,2-Dibromo-3-chloropropane	96-12-8	-	-	-	-	-	-	-	-	7.24E-01	mg/m³	9.66E+00	7.24E-01
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	-	-	-	-	-	-	-	-	2.10E+04	mg/m³	6.99E+00	2.10E+04
1,2-Dichloroethane	107-06-2	-	-	-	-	-	-	-	-	8.09E+00	mg/m³	4.04E+00	8.09E+00
1,2-Dichloroethylene	540-59-0	-	-	1.30E+01	ppm[h]	-	-	-	-	2.38E+03	mg/m³	3.96E+00	5.15E+01
1,2-Dichloropropane	78-87-5	-	-	-	-	-	-	-	-	5.08E+02	mg/m³	4.62E+00	5.08E+02
1,3-Dichloropropene	542-75-6	-	-	-	-	-	-	-	-	1.36E+01	mg/m³	4.54E+00	1.36E+01
1,4-Dichloro-2-butene	764-41-0	-	-	-	-	-	-	-	-	7.66E-02	mg/m³	5.11E+00	7.66E-02
1-Chloroethene	75-01-4	-	-	-	-	-	-	1.80E+02	mg/m³	1.28E+01	mg/m³	2.55E+00	1.80E+02
2,2-Dichloropropionic acid	75-99-0	-	-	-	-	-	-	-	-	-	-	-	-
2-Chloropropane	75-29-6	-	-	-	-	-	-	-	-	-	-	-	-
3-Chloropropene (Allyl chloride)	107-05-1	-	-	-	-	3.00E+00	ppm	-	-	9.38E+00	mg/m³	3.13E+00	9.38E+00

Table 4-4. Acute Inhalation Exposure Criteria for COPCs

COPC	CAS Number	NCEA Provis. Value[a]		AEGL-1[b]		ERPG-1[c]		AREL[d]		TEEL-1[e]		Conv. Factor[f]	AIEC[g] (mg/m³)
Bromochloromethane	74-97-5	-	-	-	-	-	-	-	-	1.06E+03	mg/m³	5.29E+00	1.06E+03
Bromodichloromethane	75-27-4	-	-	-	-	-	-	-	-	4.00E+00	mg/m³	6.70E+00	4.00E+00
Bromoethene	593-60-2	-	-	-	-	-	-	-	-	6.56E+01	mg/m³	4.37E+00	6.56E+01
Bromoform	75-25-2	-	-	-	-	-	-	-	-	1.55E+01	mg/m³	1.03E+01	1.55E+01
Bromomethane	74-83-9	-	-	-	-	-	-	3.90E+00	mg/m³	5.82E+01	mg/m³	3.88E+00	3.90E+00
Carbon tetrachloride	56-23-5	-	-	1.20E+01	ppm	2.00E+01	ppm	1.90E+00	mg/m³	1.26E+02	mg/m³	6.29E+00	7.54E+01
Chlorodibromomethane	124-48-1	-	-	-	-	-	-	-	-	6.00E+00	mg/m³	8.51E+00	6.00E+00
Chlorodifluoromethane	75-45-6	-	-	-	-	-	-	-	-	4.42E+03	mg/m³	3.53E+00	4.42E+03
Chloroethane	75-00-3	-	-	-	-	-	-	-	-	7.91E+03	mg/m³	2.64E+00	7.91E+03
Chloroform	67-66-3	-	-	-	-	-	-	1.50E-01	mg/m³	9.76E+00	mg/m³	4.88E+00	1.50E-01
Chloromethane	74-87-3	-	-	-	-	-	-	-	-	2.06E+02	mg/m³	2.06E+00	2.06E+02
Chloropentafluoroethane	76-15-3	-	-	-	-	-	-	-	-	-	-	-	-
cis-1,2-Dichloroethene	156-59-2	-	-	-	-	-	-	-	-	7.92E+02	mg/m³	3.96E+00	7.92E+02
cis-1,3-Dichloropropene	10061-01-5	-	-	-	-	-	-	-	-	1.13E+01	mg/m³	4.54E+00	1.13E+01
Cyanogen bromide	506-68-3	-	-	-	-	-	-	-	-	-	-	-	-
Cyanogen chloride	506-77-4	-	-	-	-	-	-	-	-	1.51E-01	mg/m³	2.51E+00	1.51E-01
Dichlorodifluoromethane	75-71-8	-	-	-	-	-	-	-	-	1.48E+04	mg/m³	4.94E+00	1.48E+04
Dichlorofluoromethane	75-43-4	-	-	-	-	-	-	-	-	1.26E+02	mg/m³	4.21E+00	1.26E+02
Dichloromethane	75-09-2					2.00E+02	ppm			6.94E+02	mg/m³	3.47E+00	6.94E+02
Difluorodibromomethane	75-61-6	-	-	-	-	-	-	-	-	-	-	-	-
Hexafluoroacetone	684-16-2	-	-	-	-	-	-	-	-	2.04E+00	mg/m3	6.79E+00	2.04E+00
Iodomethane	74-88-4	-	-	-	-	2.50E+01	ppm	-	-	1.45E+02	mg/m³	5.80E+00	1.45E+02
Methylene bromide	74-95-3	-	-	-	-	-	-	-	-	2.50E+02	mg/m³	7.11E+00	2.50E+02
Pentachloroethane	76-01-7	-	-	-	-	-	-	-	-	3.00E+01	mg/m³	8.27E+00	3.00E+01
trans-1,2-Dichloroethylene	156-60-5	-	-	-	-	-	-	-	-	4.95E+01	mg/m³	3.96E+00	4.95E+01
trans-1,3-Dichloropropene	10061-02-6	-	-	-	-	-	-	-	-	-	-	-	-
Trichloroacetic acid	76-03-9	-	-	-	-	-	-	-	-	6.68E+00	mg/m³	6.68E+00	6.68E+00
Trichlorofluoroethane	27154-33-2	-	-	-	-	-	-	-	-	-	-	-	-
Trichlorofluoromethane	75-69-4	-	-	-	-	-	-	-	-	2.81E+03	mg/m³	5.61E+00	2.81E+03
Trifluorobromomethane	75-63-8	-	-	-	-	-	-	-	-	1.83E+04	mg/m³	6.09E+00	1.83E+04
Dioxin and Furan Compounds													
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	35822-46-9	-	-	-	-	-	-	-	-	-	-	-	-
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	-	-	-	-	-	-	-	-	-	-	-	-
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	-	-	-	-	-	-	-	-	-	-	-	-
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	39227-28-6	-	-	-	-	-	-	-	-	-	-	-	-
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	-	-	-	-	-	-	-	-	7.50E-03	mg/m³	1.53E+01	7.50E-03
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	57653-85-7	-	-	-	-	-	-	-	-	1.50E-02	mg/m³	1.60E+01	1.50E-02
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	-	-	-	-	-	-	-	-	2.50E-03	mg/m³	1.55E+01	2.50E-03
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	19408-74-3	-	-	-	-	-	-	-	-	-	-	-	-
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	-	-	-	-	-	-	-	-	-	-	-	-
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	40321-76-4	-	-	-	-	-	-	-	-	2.50E-03	mg/m³	1.46E+01	2.50E-03

Table 4-4. Acute Inhalation Exposure Criteria for COPCs

COPC	CAS Number	NCEA Provis. Value[a]		AEGL-1[b]		ERPG-1[c]		AREL[d]		TEEL-1[e]		Conv. Factor[f]	AIEC[g] (mg/m³)
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	-	-	-	-	-	-	-	-	-	-	-	-
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	-	-	-	-	-	-	-	-	1.50E-03	mg/m³	1.53E+01	1.50E-03
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	-	-	-	-	-	-	-	-	7.50E-05	mg/m³	1.39E+01	7.50E-05
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1746-01-6	-	-	-	-	-	-	-	-	3.50E-03	mg/m³	1.32E+01	3.50E-03
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	-	-	-	-	-	-	-	-	2.00E-03	mg/m³	1.25E+01	2.00E-03
Dibenzofuran	132-64-9	-	-	-	-	-	-	-	-	-	-	-	-
Octachlorodibenzo(p)dioxin	3268-87-9	-	-	-	-	-	-	-	-	-	-	-	-
Octachlorodibenzofuran	39001-02-0	-	-	-	-	-	-	-	-	-	-	-	-
PCBs													
2,2',3,3',4,4',5- Heptachlorobiphenyl	35065-30-6	-	-	-	-	-	-	-	-	-	-	-	-
2,2',3,4,4',5,5'- Heptachlorobiphenyl	35065-29-3	-	-	-	-	-	-	-	-	-	-	-	-
2,3,3',4,4',5'- Hexachlorobiphenyl	69782-90-7	-	-	-	-	-	-	-	-	-	-	-	-
2,3,3',4,4',5- Hexachlorobiphenyl	38380-08-4	-	-	-	-	-	-	-	-	-	-	-	-
2,3,3',4,4',5,5'- Heptachlorobiphenyl	no cas #	-	-	-	-	-	-	-	-	-	-	-	-
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	-	-	-	-	-	-	-	-	-	-	-	-
2,3,4,4',5- Pentachlorobiphenyl	74472-37-0	-	-	-	-	-	-	-	-	-	-	-	-
2',3,4,4',5- Pentachlorobiphenyl	no cas #	-	-	-	-	-	-	-	-	-	-	-	-
2,3',4,4',5- Pentachlorobiphenyl	31508-00-6	-	-	-	-	-	-	-	-	-	-	-	-
2,3',4,4',5,5'- Hexachlorobiphenyl	no cas #	-	-	-	-	-	-	-	-	-	-	-	-
3,3',4,4',5- Pentachlorobiphenyl	no cas #	-	-	-	-	-	-	-	-	-	-	-	-
3,3',4,4',5,5'- Hexachlorobiphenyl	32774-16-6	-	-	-	-	-	-	-	-	-	-	-	-
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	-	-	-	-	-	-	-	-	-	-	-	-
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	-	-	-	-	-	-	-	-	-	-	-	-
Polychlorinated biphenyls (PCBs)	1336-36-3	4.00E-02	mg/m3	-	-	-	-	-	-	3.00E+00	mg/m³	-	4.00E-02
Phthalates													
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	-	-	-	-	-	-	-	-	1.00E+01	mg/m³	1.60E+01	1.00E+01
Butylbenzyl phthalate	85-68-7	-	-	-	-	-	-	-	-	5.00E+02	mg/m³	1.28E+01	5.00E+02
Dibutyl phthalate	84-74-2	-	-	-	-	-	-	-	-	1.50E+01	mg/m³	1.14E+01	1.50E+01
Diethyl phthalate	84-66-2	-	-	-	-	-	-	-	-	1.50E+01	mg/m³	9.08E+00	1.50E+01
Dimethylphthalate	131-11-3	-	-	-	-	-	-	-	-	1.50E+01	mg/m³	7.94E+00	1.50E+01
n-Dioctyl phthalate	117-84-0	-	-	-	-	-	-	-	-	1.50E+02	mg/m³	1.60E+01	1.50E+02
Light Polycyclic Aromatic Hydrocarbons													
2-Chloronapthalene	91-58-7	-	-	-	-	-	-	-	-	6.00E-01	mg/m³	6.65E+00	6.00E-01
2-Methyl naphthalene	91-57-6	-	-	-	-	-	-	-	-	2.00E+01	mg/m³	5.81E+00	2.00E+01
5-Nitroacenaphthene	602-87-9	-	-	-	-	-	-	-	-	-	-	-	-
Acenaphthene	83-32-9	-	-	-	-	-	-	-	-	1.25E+00	mg/m³	6.30E+00	1.25E+00
Acenaphthylene	208-96-8	-	-	-	-	-	-	-	-	2.00E-01	mg/m³	6.22E+00	2.00E-01
Anthracene	120-12-7	-	-	-	-	-	-	-	-	6.00E+00	mg/m³	7.28E+00	6.00E+00
Fluorene	86-73-7	-	-	-	-	-	-	-	-	7.50E+01	mg/m³	6.79E+00	7.50E+01
Indene	95-13-6	-	-	-	-	-	-	-	-	1.42E+02	mg/m³	4.75E+00	1.42E+02
Naphthalene	91-20-3	-	-	-	-	-	-	-	-	7.86E+01	mg/m³	5.24E+00	7.86E+01

Table 4-4. Acute Inhalation Exposure Criteria for COPCs

COPC	CAS Number	NCEA Provis. Value[a]		AEGL-1[b]		ERPG-1[c]		AREL[d]		TEEL-1[e]		Conv. Factor[f]	AIEC[g] (mg/m³)
Phenanthrene	85-01-8	-	-	-	-	-	-	-	-	2.00E+00	mg/m³	7.28E+00	2.00E+00
Pyrene	129-00-0	-	-	-	-	-	-	-	-	1.50E+01	mg/m³	8.27E+00	1.50E+01
Heavy Polycyclic Aromatic Hydrocarbons													
3-Methylcholanthrene	56-49-5	-	-	-	-	-	-	-	-	1.50E+00	mg/m³	1.10E+01	1.50E+00
5-Methylchrysene	3697-24-3	-	-	-	-	-	-	-	-				
Benzo(a)anthracene	56-55-3	-	-	-	-	-	-	-	-	6.00E-01	mg/m³	9.33E+00	6.00E-01
Benzo(a)pyrene	50-32-8	-	-	-	-	-	-	-	-	7.50E+00	mg/m³	1.03E+01	7.50E+00
Benzo(b)fluoranthene	205-99-2	-	-	-	-	-	-	-	-	-	-	-	-
Benzo(e)pyrene	192-97-2	-	-	-	-	-	-	-	-	-	-	-	-
Benzo(g,h,i)perylene	191-24-2	-	-	-	-	-	-	-	-	3.00E+01	mg/m³	1.13E+01	3.00E+01
Benzo(j)fluoranthene	205-82-3	-	-	-	-	-	-	-	-	-	-	-	-
Benzo(k)fluoranthene	207-08-9	-	-	-	-	-	-	-	-	-	-	-	-
Benzo[a,i]pyrene	191-30-0	-	-	-	-	-	-	-	-	-	-	-	-
Chrysene	218-01-9	-	-	-	-	-	-	-	-	2.00E-01	mg/m³	9.33E+00	2.00E-01
Dibenz(a,h)anthracene	53-70-3	-	-	-	-	-	-	-	-	3.00E+01	mg/m³	1.14E+01	3.00E+01
Dibenz[a,h]acridine	226-36-8	-	-	-	-	-	-	-	-	-	-	-	-
Dibenz[a,j]acridine	224-42-0	-	-	-	-	-	-	-	-	-	-	-	-
Dibenzo(a,e)fluoranthene	5385-75-1	-	-	-	-	-	-	-	-	-	-	-	-
Dibenzo(a,h)fluoranthene	no cas #	-	-	-	-	-	-	-	-	-	-	-	-
Dibenzo[a,e]pyrene	192-65-4	-	-	-	-	-	-	-	-	-	-	-	-
Dibenzo[a,h]pyrene	189-64-0	-	-	-	-	-	-	-	-	-	-	-	-
Dibenzo[a,i]pyrene	189-55-9	-	-	-	-	-	-	-	-	-	-	-	-
Fluoranthene	206-44-0	-	-	-	-	-	-	-	-	3.00E-02	mg/m³	8.27E+00	3.00E-02
Hexachloronaphthalene	1335-87-1	-	-	-	-	-	-	-	-	2.00E-01	mg/m³	1.37E+01	2.00E-01
Indeno(1,2,3-cd)pyrene	193-39-5	-	-	-	-	-	-	-	-	-	-	-	-
Octachloronaphthalene	2234-13-1	-	-	-	-	-	-	-	-	3.00E-01	mg/m³	1.65E+01	3.00E-01
Pentachloronaphthalene	1321-64-8	-	-	-	-	-	-	-	-	-	-	-	-
Tetrachloronaphthalene	1335-88-2	-	-	-	-	-	-	-	-	-	-	-	-
Trichloronaphthalene	1321-65-9	-	-	-	-	-	-	-	-	-	-	-	-
Light Substituted Benzene Compounds													
1,2,3-Trichlorobenzene	87-61-6	-	-	-	-	-	-	-	-	5.00E+01	mg/m³	7.42E+00	5.00E+01
1,2,4-Trichlorobenzene	120-82-1	-	-	-	-	-	-	-	-	3.71E+01	mg/m³	7.42E+00	3.71E+01
1,2,4-Trimethyl benzene	95-63-6	-	-	-	-	-	-	-	-	1.80E+02	mg/m³	4.91E+00	1.80E+02
1,2-Dichlorobenzene	95-50-1	-	-	-	-	-	-	-	-	3.00E+02	mg/m³	6.01E+00	3.00E+02
1,3,5-Trimethyl benzene	108-67-8	-	-	-	-	-	-	-	-	3.68E+02	mg/m³	4.91E+00	3.68E+02
1,3-Dichlorobenzene	541-73-1	-	-	-	-	-	-	-	-	3.60E+01	mg/m³	6.01E+00	3.60E+01
1,3-Dinitrobenzene	99-65-0	-	-	-	-	-	-	-	-	3.00E+00	mg/m³	6.87E+00	3.00E+00
1,4-Dichlorobenzene	106-46-7	-	-	-	-	-	-	-	-	6.61E+02	mg/m³	6.01E+00	6.61E+02
1,4-Dinitrobenzene	100-25-4	-	-	-	-	-	-	-	-	-	-	-	-
2,4,5-Trichlorophenol	95-95-4	-	-	-	-	-	-	-	-	3.00E+01	mg/m³	8.07E+00	3.00E+01
2,4,6-Trichlorophenol	88-06-2	-	-	-	-	-	-	-	-	3.00E+01	mg/m³	8.07E+00	3.00E+01

Table 4-4. Acute Inhalation Exposure Criteria for COPCs

COPC	CAS Number	NCEA Provis. Value[a]		AEGL-1[b]		ERPG-1[c]		AREL[d]		TEEL-1[e]		Conv. Factor[f]	AIEC[g] (mg/m^3)
2,4-Dichlorophenol	120-83-2	-	-	-	-	-	-	-	-	3.00E+01	mg/m^3	6.66E+00	3.00E+01
2,4-Dimethylphenol	105-67-9	-	-	-	-	-	-	-	-	-	-	-	-
2,4-Dinitrophenol	51-28-5	-	-	-	-	-	-	-	-	7.50E+00	mg/m^3	7.53E+00	7.50E+00
2,4-Dinitrotoluene	121-14-2	-	-	-	-	-	-	-	-	6.00E-01	mg/m^3	7.44E+00	6.00E-01
2,6-Dinitrotoluene	606-20-2	-	-	-	-	-	-	-	-	6.00E-01	mg/m^3	7.44E+00	6.00E-01
2-Chlorophenol	95-57-8	-	-	-	-	-	-	-	-	5.25E+00	mg/m^3	5.25E+00	5.25E+00
2-Chlorotoluene	95-49-8	-	-	-	-	-	-	-	-	3.88E+02	mg/m^3	5.17E+00	3.88E+02
2-Nitrophenol	88-75-5	-	-	-	-	-	-	-	-	-	-	-	-
4,6-Dinitro-o-cresol	534-52-1	-	-	-	-	-	-	-	-	5.00E-01	mg/m^3	8.10E+00	5.00E-01
4-Chlorotoluene	106-43-4	-	-	-	-	-	-	-	-	3.88E+02	mg/m^3	5.17E+00	3.88E+02
4-Nitrophenol	100-02-7	-	-	-	-	-	-	-	-	3.00E+01	mg/m^3	5.69E+00	3.00E+01
alpha-Methylstyrene	98-83-9	-	-	-	-	-	-	-	-	-	-	-	-
Aniline	62-53-3	-	-	8.00E+00[h]	ppm	-	-	-	-	2.28E+01	mg/m^3	3.81E+00	3.05E+01
Benzotrichloride	98-07-7	-	-	-	-	-	-	-	-	1.00E-01	mg/m^3	7.99E+00	1.00E-01
Benzyl chloride	100-44-7	-	-	-	-	1.00E+00	ppm	2.40E-01	mg/m^3	5.17E+00	mg/m^3	5.17E+00	5.17E+00
Bromobenzene	108-86-1	-	-	-	-	-	-	-	-	4.81E+01	mg/m^3	6.42E+00	4.81E+01
Chlorobenzene	108-90-7	-	-	-	-	-	-	-	-	1.38E+02	mg/m^3	4.60E+00	1.38E+02
Cumene	98-82-8	-	-	-	-	-	-	-	-	7.37E+02	mg/m^3	4.91E+00	7.37E+02
m-Cresol	108-39-4	-	-	-	-	-	-	-	-	-	-	-	-
n-Butyl benzene	104-51-8	-	-	-	-	-	-	-	-	1.10E+02	mg/m^3	5.49E+00	1.10E+02
Nitrobenzene	98-95-3	-	-	-	-	-	-	-	-	1.51E+01	mg/m^3	5.03E+00	1.51E+01
n-Propyl benzene	103-65-1	-	-	-	-	-	-	-	-	3.68E+02	mg/m^3	4.91E+00	3.68E+02
o-Cresol	95-48-7	-	-	-	-	-	-	-	-	-	-	-	-
o-Dinitrobenzene	528-29-0	-	-	-	-	-	-	-	-	-	-	-	-
o-Nitroaniline	88-74-4	-	-	-	-	-	-	-	-	-	-	-	-
o-Toluidine	95-53-4	-	-	-	-	-	-	-	-	2.63E+01	mg/m^3	4.38E+00	2.63E+01
p-Chloroaniline	106-47-8	-	-	-	-	-	-	-	-	3.00E+01	mg/m^3	5.21E+00	3.00E+01
p-Cresol	106-44-5	-	-	-	-	-	-	-	-	-	-	-	-
Phenol	108-95-2	-	-	-	-	1.00E+01	ppm	5.80E+00	mg/m^3	3.85E+01	mg/m^3	3.85E+00	3.85E+01
p-Nitrochlorobenzene	100-00-5	-	-	-	-	-	-	-	-	2.00E+00	mg/m^3	6.44E+00	2.00E+00
p-Toluidine	106-49-0	-	-	-	-	-	-	-	-	-	-	-	-
sec-Butyl benzene	135-98-8	-	-	-	-	-	-	-	-	2.74E+01	mg/m^3	5.49E+00	2.74E+01
tert-Butyl benzene	98-06-6	-	-	-	-	-	-	-	-	1.10E+02	mg/m^3	5.49E+00	1.10E+02
Toluene-2,6-diamine	823-40-5	-	-	-	-	-	-	-	-	-	-	-	-
Trimethyl benzene	25551-13-7	-	-	-	-	-	-	-	-	-	-	-	-
Other Light Semivolatile Compounds													
1,1`-Biphenyl	92-52-4	-	-	-	-	-	-	-	-	4.00E+00	mg/m^3	6.30E+00	4.00E+00
1,1-Dimethylhydrazine	57-14-7	-	-	-	-	-	-	-	-	1.84E+00	mg/m^3	2.46E+00	1.84E+00
1,2-Dimethylhydrazine	540-73-8	-	-	-	-	-	-	-	-	3.69E+00	mg/m^3	2.46E+00	3.69E+00
1,2-Diphenylhydrazine	122-66-7	-	-	-	-	-	-	-	-	3.00E+01	mg/m^3	7.53E+00	3.00E+01
1,3-Propane sultone	1120-71-4	-	-	-	-	-	-	-	-	2.50E+00	mg/m^3	4.99E+00	2.50E+00

Table 4-4. Acute Inhalation Exposure Criteria for COPCs

COPC	CAS Number	NCEA Provis. Value[a]		AEGL-1[b]		ERPG-1[c]		AREL[d]		TEEL-1[e]		Conv. Factor[f]	AIEC[g] (mg/m³)
2,4-Toluene diisocyante	584-84-9	-	-	2.00E-02	ppm	-	-	-	-	1.42E-01	mg/m³	7.12E+00	1.42E-01
2-Chloroacetophenone	532-27-4	-	-	-	-	-	-	-	-	-	-	-	-
2-Propenoic acid	79-10-7	-	-	-	-	2.00E+00	ppm	6.00E+00	mg/m³	5.89E+00	mg/m³	2.95E+00	5.89E+00
4,4-Methylenedianiline	101-77-9	-	-	-	-	-	-	-	-	8.10E-01	mg/m³	8.10E+00	8.10E-01
Acetophenone	98-86-2	-	-	-	-	-	-	-	-	3.00E+01	mg/m³	4.91E+00	3.00E+01
Benzoic acid	65-85-0	-	-	-	-	-	-	-	-	1.25E+01	mg/m³	4.99E+00	1.25E+01
Bis(2-chloroethoxy)methane	111-91-1	-	-	-	-	-	-	-	-	-	-	-	-
Bis(2-chloroethyl) ether	111-44-4	-	-	-	-	-	-	-	-	5.85E+01	mg/m³	5.85E+00	5.85E+01
Chlorocyclopentadiene	41851-50-7	-	-	-	-	-	-	-	-	-	-	-	-
Cyclohexanol	108-93-0	-	-	-	-	-	-	-	-	-	-	-	-
Dichloroisopropyl ether	108-60-1	-	-	-	-	-	-	-	-	6.99E+01	mg/m³	6.99E+00	6.99E+01
Dichloromethyl ether	542-88-1	-	-	-	-	-	-	-	-	1.41E-02	mg/m³	4.70E+00	1.41E-02
Dichloropentadiene	no cas #	-	-	-	-	-	-	-	-	-	-	-	-
Dimethyl sulfate	77-78-1	-	-	-	-	-	-	-	-	1.55E+00	mg/m³	5.16E+00	1.55E+00
Dimethylaniline	121-69-7	-	-	-	-	-	-	-	-	-	-	-	-
di-n-Propylnitrosamine	621-64-7	-	-	-	-	-	-	-	-	2.00E-01	mg/m³	5.32E+00	2.00E-01
Diphenyl ether	101-84-8	-	-	-	-	-	-	-	-	-	-	-	-
Epichlorohydrin	106-89-8	-	-	5.00E+00	ppm	2.00E+00	ppm	1.30E+00	mg/m³	7.56E+00	mg/m³	3.78E+00	1.89E+01
Ethyl Carbamate (urethane)	51-79-6	-	-	-	-	-	-	-	-	-	-	-	-
Ethyl methanesulfonate	62-50-0	-	-	-	-	-	-	-	-	-	-	-	-
Ethylene dibromide	106-93-4	-	-	-	-	-	-	-	-	1.54E+02	mg/m³	7.68E+00	1.54E+02
Ethylene glycol	107-21-1	-	-	-	-	-	-	-	-	5.07E+01	mg/m³	2.54E+00	5.07E+01
Ethylene glycol monobutyl ether	111-76-2	-	-	-	-	-	-	1.40E+01	mg/m³	3.62E+02	mg/m³	4.83E+00	1.40E+01
Ethylene glycol monoethyl ether acetate	111-15-9	-	-	-	-	-	-	1.40E-01	mg/m³	8.10E+01	mg/m³	5.40E+00	1.40E-01
Ethylene thiourea	96-45-7	-	-	-	-	-	-	-	-	1.00E+01	mg/m³	4.18E+00	1.00E+01
Furfural	98-01-1	-	-	-	-	-	-	-	-	7.85E+00	mg/m³	3.93E+00	7.85E+00
Maleic hydrazide	123-33-1	-	-	-	-	-	-	-	-	-	-	-	-
Malononitrile	109-77-3	-	-	-	-	-	-	-	-	-	-	-	-
Methyl styrene (mixed isomers)	25013-15-4	-	-	-	-	-	-	-	-	-	-	-	-
Methylhydrazine	60-34-4	-	-	-	-	-	-	-	-	3.77E-01	mg/m³	1.88E+00	3.77E-01
N,N-Diphenylamine	122-39-4	-	-	-	-	-	-	-	-	3.00E+01	mg/m³	6.92E+00	3.00E+01
Nitric acid, propyl ester	627-13-4	-	-	-	-	-	-	-	-	-	-	-	-
N-Nitrosodi-n-butylamine	924-16-3	-	-	-	-	-	-	-	-	-	-	-	-
N-Nitrosomorpholine	59-89-2	-	-	-	-	-	-	-	-	3.00E+01	mg/m³	4.75E+00	3.00E+01
N-Nitroso-N,N-dimethylamine	62-75-9	-	-	-	-	-	-	-	-	2.50E+00	mg/m³	3.03E+00	2.50E+00
o-Anisidine	90-04-0	-	-	-	-	-	-	-	-	1.50E+00	mg/m³	5.03E+00	1.50E+00
Oxalic acid	144-62-7	-	-	-	-	-	-	-	-	2.00E+00	mg/m³	3.68E+00	2.00E+00
Phthalic anhydride	85-44-9	-	-	-	-	-	-	-	-	1.80E+01	mg/m³	6.05E+00	1.80E+01
p-Phthalic acid	100-21-0	-	-	-	-	-	-	-	-	-	-	-	-
Pyridine	110-86-1	-	-	-	-	-	-	-	-	4.85E+01	mg/m³	3.23E+00	4.85E+01
Quinoline	91-22-5	-	-	-	-	-	-	-	-	3.17E+00	mg/m³	5.28E+00	3.17E+00

Table 4-4. Acute Inhalation Exposure Criteria for COPCs

COPC	CAS Number	NCEA Provis. Value[a]		AEGL-1[b]		ERPG-1[c]		AREL[d]		TEEL-1[e]		Conv. Factor[f]	AIEC[g] (mg/m³)
Quinone	106-51-4	-	-	-	-	-	-	-	-	1.33E+00	mg/m³	4.42E+00	1.33E+00
Safrole	94-59-7	-	-	-	-	-	-	-	-	-	-	-	-
Tetrahydrofuran	109-99-9	-	-	-	-	-	-	-	-	7.37E+02	mg/m³	2.95E+00	7.37E+02
Other Heavy Semivolatile Compounds													
1,2,4,5-Tetrachlorobenzene	95-94-3	-	-	-	-	-	-	-	-	3.00E+01	mg/m³	8.82E+00	3.00E+01
1,3,5-Trinitrobenzene	99-35-4	-	-	-	-	-	-	-	-	3.00E+01	mg/m³	8.71E+00	3.00E+01
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	-	-	-	-	-	-	-	-	-	-	-	-
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	-	-	-	-	-	-	-	-	-	-	-	-
2-sec-Butyl-4,6-dinitrophenol	88-85-7	-	-	-	-	-	-	-	-	4.50E+00	mg/m³	9.82E+00	4.50E+00
3,3-Dichlorobenzidine	91-94-1	-	-	-	-	-	-	-	-	6.21E+00	mg/m³	1.03E+01	6.21E+00
3,3'-Dimethoxybenzidine	119-90-4	-	-	-	-	-	-	-	-	5.00E+00	mg/m³	9.99E+00	5.00E+00
4-Bromophenylphenyl ether	101-55-3	-	-	-	-	-	-	-	-	-	-	-	-
Ammonium perfluorooctanoate	3825-26-1	-	-	-	-	-	-	-	-	-	-	-	-
Azobenzene	103-33-3	-	-	-	-	-	-	-	-	-	-	-	-
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	-	-	-	-	-	-	-	-	-	-	-	-
Captan	133-06-2	-	-	-	-	-	-	-	-	1.50E+01	mg/m³	1.23E+01	1.50E+01
Chlorobenzilate	510-15-6	-	-	-	-	-	-	-	-	2.50E-01	mg/m³	1.33E+01	2.50E-01
Dibutylphosphate	107-66-4	-	-	-	-	-	-	-	-	1.72E+01	mg/m³	8.59E+00	1.72E+01
Dimethyl aminoazobenzene	60-11-7	-	-	-	-	-	-	-	-	7.50E+01	mg/m³	9.21E+00	7.50E+01
Hexachlorobenzene	118-74-1	-	-	-	-	-	-	-	-	7.50E-02	mg/m³	1.16E+01	7.50E-02
Hexachlorobutadiene	87-68-3	-	-	-	-	3.00E+00	ppm	-	-	3.20E+01	mg/m³	1.07E+01	3.20E+01
Hexachlorocyclopentadiene	77-47-4	-	-	-	-	-	-	-	-	2.23E-01	mg/m³	1.11E+01	2.23E-01
Hexachloroethane	67-72-1	-	-	-	-	-	-	-	-	2.90E+01	mg/m³	9.68E+00	2.90E+01
Hexachlorophene	70-30-4	-	-	-	-	-	-	-	-	3.00E+01	mg/m³	1.66E+01	3.00E+01
Hexamethylene-1,5-diisocyanate	822-06-0	-	-	-	-	-	-	-	-	1.03E-01	mg/m³	6.88E+00	1.03E-01
Mirex	2385-85-5	-	-	-	-	-	-	-	-	7.50E-02	mg/m³	2.23E+01	7.50E-02
Nitrofen	1836-75-5	-	-	-	-	-	-	-	-	-	-	-	-
Pentachlorobenzene	608-93-5	-	-	-	-	-	-	-	-	3.00E+01	mg/m³	1.02E+01	3.00E+01
Pentachloronitrobenzene	82-68-8	-	-	-	-	-	-	-	-	1.50E+00	mg/m³	1.21E+01	1.50E+00
Pentachlorophenol	87-86-5	-	-	-	-	-	-	-	-	1.50E+00	mg/m³	1.09E+01	1.50E+00
Picric acid	88-89-1	-	-	-	-	-	-	-	-	3.00E-01	mg/m³	9.36E+00	3.00E-01
Pronamide	23950-58-5	-	-	-	-	-	-	-	-	-	-	-	-
Strychnine	57-24-9	-	-	-	-	-	-	-	-	3.00E-01	mg/m³	1.37E+01	3.00E-01
Terphenyls	26140-60-3	-	-	-	-	-	-	-	-	1.25E+00	mg/m³	9.41E+00	1.25E+00
Tributyl phosphate	126-73-8	-	-	-	-	-	-	-	-	6.53E+00	mg/m³	1.09E+01	6.53E+00
Trifluralin	1582-09-8	-	-	-	-	-	-	-	-	7.50E-02	mg/m³	1.37E+01	7.50E-02
Triphenylamine	603-34-9	-	-	-	-	-	-	-	-	-	-	-	-
Herbicides and Organochlorinated Pesticides													
2,4,5-T	93-76-5	-	-	-	-	-	-	-	-	3.00E+01	mg/m³	1.04E+01	3.00E+01
2,4-D and esters	94-75-7	-	-	-	-	-	-	-	-	3.00E+01	mg/m³	9.03E+00	3.00E+01
4,4-DDD	72-54-8	-	-	-	-	-	-	-	-	3.00E+01	mg/m³	1.31E+01	3.00E+01

Table 4-4. Acute Inhalation Exposure Criteria for COPCs

COPC	CAS Number	NCEA Provis. Value[a]		AEGL-1[b]		ERPG-1[c]		AREL[d]		TEEL-1[e]		Conv. Factor[f]	AIEC[g] (mg/m³)
4,4-DDE	72-55-9	-	-	-	-	-	-	-	-	3.00E+01	mg/m³	1.30E+01	3.00E+01
4,4-DDT	50-29-3	-	-	-	-	-	-	-	-	3.00E+00	mg/m³	1.45E+01	3.00E+00
Aldrin	309-00-2	-	-	-	-	-	-	-	-	7.50E-01	mg/m³	1.49E+01	7.50E-01
alpha-BHC	319-84-6	-	-	-	-	-	-	-	-	7.50E+00	mg/m³	1.19E+01	7.50E+00
beta-BHC	319-85-7	-	-	-	-	-	-	-	-	2.00E+00	mg/m³	1.19E+01	2.00E+00
Chlordane	57-74-9	-	-	-	-	-	-	-	-	1.50E+00	mg/m³	1.67E+01	1.50E+00
Delta-BHC	319-86-8	-	-	-	-	-	-	-	-	-	-	-	-
Dieldrin	60-57-1	-	-	-	-	-	-	-	-	7.50E-01	mg/m³	1.56E+01	7.50E-01
Endothall	145-73-3	-	-	-	-	-	-	-	-	-	-	-	-
Endrin	72-20-8	-	-	-	-	-	-	-	-	3.00E-01	mg/m³	1.56E+01	3.00E-01
gamma-BHC (Lindane)	58-89-9	-	-	-	-	-	-	-	-	1.50E+00	mg/m³	1.19E+01	1.50E+00
Heptachlor	76-44-8	-	-	-	-	-	-	-	-	1.50E-01	mg/m³	1.53E+01	1.50E-01
Isodrin	465-73-6	-	-	-	-	-	-	-	-	-	-	-	-
Methoxychlor	72-43-5	-	-	-	-	-	-	-	-	3.00E+01	mg/m³	1.41E+01	3.00E+01
Silvex (2,4,5-TP)	93-72-1	-	-	-	-	-	-	-	-	3.00E+01	mg/m³	1.10E+01	3.00E+01
Toxaphene	8001-35-2	-	-	-	-	-	-	-	-	1.00E+00	mg/m³	1.69E+01	1.00E+00
INORGANICS													
Metals													
Aluminum	7429-90-5	-	-	-	-	-	-	-	-	3.00E+01	mg/m³	1.10E+00	3.00E+01
Antimony	7440-36-0	-	-	-	-	-	-	-	-	1.50E+00	mg/m³	4.98E+00	1.50E+00
Arsenic	7440-38-2	-	-	-	-	-	-	1.90E-04	mg/m³	3.00E-02	mg/m³	3.06E+00	1.90E-04
Barium	7440-39-3	-	-	-	-	-	-	-	-	1.50E+00	mg/m³	5.61E+00	1.50E+00
Beryllium	7440-41-7	-	-	-	-	-	-	-	-	5.00E-03	mg/m³	3.68E-01	5.00E-03
Bismuth	7440-69-9	-	-	-	-	-	-	-	-	5.00E+00	mg/m³	8.54E+00	5.00E+00
Boron	7440-42-8	-	-	-	-	-	-	-	-	7.50E+00	mg/m³	4.42E-01	7.50E+00
Cadmium	7440-43-9	-	-	-	-	-	-	-	-	3.00E-02	mg/m³	4.59E+00	3.00E-02
Calcium	7440-70-2	-	-	-	-	-	-	-	-	3.00E+01	mg/m³	1.64E+00	3.00E+01
Chromium (and VI)	18540-29-9	-	-	-	-	-	-	-	-	1.50E+00	mg/m³	2.13E+00	1.50E+00
Cobalt	7440-48-4	-	-	-	-	-	-	-	-	6.00E-02	mg/m³	2.41E+00	6.00E-02
Copper	7440-50-8	-	-	-	-	-	-	1.00E-01	mg/m³	3.00E+00	mg/m³	2.60E+00	1.00E-01
Iron	7439-89-6	-	-	-	-	-	-	-	-	3.00E+01	mg/m³	2.28E+00	3.00E+01
Lead	7439-92-1	-	-	-	-	-	-	-	-	1.50E-01	mg/m³	8.47E+00	1.50E-01
Lithium	7439-93-2	-	-	-	-	-	-	-	-	3.00E+01	mg/m³	2.84E-01	3.00E+01
Magnesium	7439-95-4	-	-	-	-	-	-	-	-	3.00E+01	mg/m³	9.94E-01	3.00E+01
Manganese	7439-96-5	-	-	-	-	-	-	-	-	3.00E+00	mg/m³	2.25E+00	3.00E+00
Mercury	7439-97-6	-	-	-	-	-	-	1.80E-03	mg/m³	1.00E-01	mg/m³	8.20E+00	1.80E-03
Molybdenum	7439-98-7	-	-	-	-	-	-	-	-	1.50E+01	mg/m³	3.92E+00	1.50E+01
Nickel	7440-02-0	-	-	-	-	-	-	6.00E-03	mg/m³	3.00E+00	mg/m³	2.40E+00	6.00E-03
Potassium	7440-09-7	-	-	-	-	-	-	-	-	2.00E+00	mg/m³	1.60E+00	2.00E+00
Rhodium	7440-16-6	-	-	-	-	-	-	-	-	3.00E+00	mg/m³	4.21E+00	3.00E+00

Table 4-4. Acute Inhalation Exposure Criteria for COPCs

COPC	CAS Number	NCEA Provis. Value[a]		AEGL-1[b]		ERPG-1[c]		AREL[d]		TEEL-1[e]		Conv. Factor[f]	AIEC[g] (mg/m³)
Selenium	7782-49-2	-	-	-	-	-	-	-	-	6.00E-01	mg/m³	3.23E+00	6.00E-01
Silicon	7440-21-3	-	-	-	-	-	-	-	-	3.00E+01	mg/m³	1.15E+00	3.00E+01
Silver	7440-22-4	-	-	-	-	-	-	-	-	3.00E-01	mg/m³	4.41E+00	3.00E-01
Sodium	7440-23-5	-	-	-	-	-	-	-	-	1.50E+02	mg/m³	9.40E-01	1.50E+02
Strontium	7440-24-6	-	-	-	-	-	-	-	-	3.00E+01	mg/m³	3.58E+00	3.00E+01
Tantalum	7440-25-7	-	-	-	-	-	-	-	-	-	-	-	-
Thallium	7440-28-0	-	-	-	-	-	-	-	-	3.00E-01	mg/m³	8.35E+00	3.00E-01
Tin	7440-31-5	-	-	-	-	-	-	-	-	6.00E+00	mg/m³	4.85E+00	6.00E+00
Tungsten	7440-33-7	-	-	-	-	-	-	-	-	1.00E+01	mg/m³	7.51E+00	1.00E+01
Uranium	7440-61-1	-	-	-	-	-	-	-	-	6.00E-01	mg/m³	9.73E+00	6.00E-01
Vanadium	7440-62-2	-	-	-	-	-	-	-	-	7.50E-02	mg/m³	2.08E+00	7.50E-02
Yttrium	7440-65-5	-	-	-	-	-	-	-	-	3.00E+00	mg/m³	3.63E+00	3.00E+00
Zinc	7440-66-6	-	-	-	-	-	-	-	-	3.00E+01	mg/m³	2.67E+00	3.00E+01
Zirconium	7440-67-7	-	-	-	-	-	-	-	-	1.00E+01	mg/m³	3.73E+00	1.00E+01
Non-metals and Anions													
Ammonia/Ammonium	7664-41-7	-		2.50E+01	ppm	2.50E+01	ppm	3.20E+00	mg/m³	1.74E+01	mg/m³	6.96E-01	1.74E+01
Bromide	24959-67-9	-	-	-	-	-	-	-	-	-	-	-	-
Chloride	16887-00-6	-	-	-	-	-	-	-	-	-	-	-	-
Cyanide	57-12-5	-	-	-	-	-	-	-	-	5.00E+00	mg/m³	1.06E+00	5.00E+00
Fluoride	16984-48-8	-	-	-	-	-	-	-	-	7.50E+00	mg/m³	7.76E-01	7.50E+00
Hydroxide	14280-30-9	-	-	-	-	-	-	-	-	-	-	-	-
Iodine	7553-56-2	-	-	-	-	1.00E-01	ppm	-	-	1.04E+00	mg/m³	1.04E+01	1.04E+00
Nitrate	14797-55-8	-	-	-	-	-	-	-	-	3.00E+01	mg/m³	-	3.00E+01
Nitrite	14797-65-0	-	-	-	-	-	-	-	-	-	-	-	-
Phosphate	14265-44-2	-	-	-	-	-	-	-	-	-	-	-	-
Phosphorus	7723-14-0	-	-	-	-	-	-	-	-	3.00E-01	mg/m³	5.06E+00	3.00E-01
Sulfate	14808-79-8	-	-	-	-	-	-	1.20E-01	mg/m³	-	-	-	1.20E-01
Total Sulfur	63705-05-5	-	-	-	-	-	-	-	-	7.50E-01	mg/m³	1.31E+00	7.50E-01
Criteria Pollutants													
Carbon dioxide	124-38-9	-	-	-	-	-	-	-	-	5.40E+04	mg/m³	1.80E+00	5.40E+04
Nitrogen dioxide	10102-44-0	-	-	5.00E-01	ppm	-	-	4.70E-01	mg/m³	3.76E+00	mg/m³	1.88E+00	9.40E-01
Ozone	10028-15-6	-	-	-	-	-	-	1.80E-01	mg/m³	1.96E-01	mg/m³	1.96E+00	1.80E-01
Particulate matter	no cas #	-	-	-	-	-	-	-	-	-	-	-	-
Sulfur dioxide	7446-09-5	-	-	-	-	3.00E-01	ppm	6.60E-01	mg/m³	7.85E-01	mg/m³	2.62E+00	7.85E-01

[a] Provisional acute 1-hour inhalation value, obtained from National Center for Environmental Assessment (NCEA).
[b] Acute exposure guideline level (AEGL-1) values (1-hour averaging time), obtained from EPA Region X, unless otherwise noted.
[c] Emergency Response Planning Guidelines (ERPG-1) values, obtained from http://www.scapa.bnl.gov/scapawl.htm.
[d] Acute reference exposure level (AREL) values, obtained from California EPA (see http://oehha.ca.gov/scientific/relnums.htm).
[e] Temporary Emergency Response Limit (TEEL-1) values, obtained from http://tis.eh.doe.gov/web/chem_safety/teel.html.
[f] Conversion Factor, for converting from ppm to mg/m³, from http://tis.eh.doe.gov/web/chem_safety/teel.html.
[g] Acute Inhalation Exposure Criteria (AIEC) in mg/m³, used to quantify hazard quotients for short term inhalation exposures to COPCs.

Table 4-4. Acute Inhalation Exposure Criteria for COPCs

COPC	CAS Number	NCEA Provis. Value[a]		AEGL-1[b]		ERPG-1[c]		AREL[d]		TEEL-1[e]		Conv. Factor[f]	AIEC[g] (mg/m^3)

The following hierarchy was used in selecting the AIEC:

1. Values from NCEA (as provided by EPA Region X)
2. AEGL-1
3. ERPG-1
4. AREL
5. TEEL-1
6. SCAPA-approach.

[h] Acute exposure guideline level (AEGL-1) values (1-hour averaging time), obtained from EPA Federal Register, October 30, 1997 (Volume 62, Number 210).

5. Quantification of Emissions

Following the identification of constituents of potential concern (COPCs) (section 4), emission rates must be estimated for each chemical and radionuclide source. These emission rates will be used along with air dispersion modeling to estimate exposure point concentrations for both human and ecological receptor populations. Emission rates will be estimated for the preliminary risk assessment PRA (section 5.2) and final risk assessment FRA (section 5.3) as described below and in section 3. The primary difference between the PRA and the FRA is the estimation of emission rates. In the PRA, estimation of emission rates relies upon modeling and engineering designs, while the FRA utilizes environmental performance demonstration results to estimate emission rates. Both assessments rely on modeling to estimate exposure point concentrations resulting from these emissions.

5.1. Emission Sources

The SLRA will evaluate potential emissions from three operating conditions as specified by EPA 1998a guidance and defined below:

Stack Emissions – Chemicals and radionuclides released from the waste treatment plant stacks as a result of normal (also known as routine) operations.

Process Upset Emissions – Chemicals and radionuclides released from the waste treatment plant stacks as a result of nonroutine operations (such as a process malfunction). Upset emissions are assumed to be at a greater rate than normal stack emissions because the upset condition is assumed to result in decreased destruction or removal efficiency (DRE) or increased PIC formation. However, upset emissions are for a shorter duration. For the PRA, the conservative assumption that all upset conditions result in increased emission rates for short durations will be used, as described in section 5.2.2.

Fugitive Emissions – Fugitive emissions include chemicals and radionuclides released from the waste treatment plant from all non-stack sources (such as chemical storage areas). EPA guidance includes leaks from the combustion chamber due to "puffs" as a source of fugitive emissions. A "puff" results when a slug of high British Thermal Unit (BTU) waste is burned causing a rapid increase in the formation of combustion gases that exceeds the volume of the combustion chamber. Waste processed by the waste treatment plant will be of uniformly low BTU content.

Potential accidental releases (such as failure of tanks or transportation accidents) will not be included in the SLRA. Accident scenarios are addressed in the hazards analysis and other nuclear and process safety documents.

5.2. Emission Assumptions for the Preliminary Risk Assessment

The PRA will be conducted prior to construction of the waste treatment plant; therefore, no site-specific emission data will be available for the plant. A combination of data from other facilities, process and design information, model data, laboratory studies, and default assumptions will be used to estimate emission rates for COPCs and ROPCs for the PRA. These data, when available, will be published in the Work Plan. Otherwise, they will be published in the PRA report.

5.2.1. Stack Emissions

The following sections describe the waste feed, assumptions associated with the waste feed, and the approaches used for estimating stack emissions for different categories of COPCs and ROPCs. For the PRA different approaches will be used to estimate stack emissions for different categories of COPCs/ROPCs (i.e., COPCs present in waste feed) as described below. Actual emission rate calculations will be developed in the Air Quality Permit Application and especially Appendix B of the three parts of the Application and summarized in section 3 of this Work Plan. Estimated emission rates will be tabulated in Appendix E.

5.2.1.1. COPCs/ROPCs in Waste Feed

The RPP-WTP plant includes four treatment plants: pretreatment, LAW pretreatment, LAW vitrification, and HLW vitrification plants. Offgas from the treatment plants may be combined before exiting the plant stack. The following four waste types will be treated in the RPP-WTP:

- Envelope A. This LAW feed envelope will contain cesium and technetium at concentrations high enough to warrant their removal during pretreatment, to ensure the ILAW glass meets applicable requirements.

- Envelope B. This LAW feed envelope will contain higher concentrations of cesium than Envelope A. Both cesium and technetium will be removed to comply with the specifications for ILAW. This envelope will contain higher concentrations of chlorine, chromium, fluorine and phosphates, and sulfates, than those found in Envelope A, which may limit the rate of waste incorporation into the glass.

- Envelope C. This LAW feed envelope will contain organically complexed strontium and transuranic (TRU) compounds, that will require removal in a processing step unique to this waste envelope. As with Envelopes A and B, cesium and technetium will also require removal in the pretreatment process to ensure that the ILAW specifications can be met.

- Envelope D. HLW feed envelope will be in the form of a slurry containing 10 to 200 grams of solids per liter of slurry. The liquid fraction of the slurry will be comprised on envelope A, B, or C waste, and the solid fraction will be Envelope D waste. Envelope D waste will be delivered via pipeline into a receipt vessel or vessels located in the pretreatment plant.

Waste in Envelopes A, B, and C will be treated by the LAW treatment train. As part of this treatment, solids, cesium, strontium/TRU (Envelope C only), and technetium will be removed from the waste during pretreatment (that is, prior to entering the melter). The amount of waste from each envelope will depend on the sodium content of the waste. Waste Envelope D, which includes the solids, strontium, transuranics, cesium, and technetium removed from Envelopes A, B, and C, will be treated by the HLW treatment train.

Waste feed may vary over the operating lifetime of the waste treatment plant. Initial plans call for the plant to process Envelope B (LAW) waste through the pretreatment system only (that is, without vitrification) and Envelope D (HLW) through the entire process for the first two years. During the next nine years of operation both LAW (Envelopes A, B, and C) and HLW (Envelope D) will be processed through the waste treatment plant.

As a conservative approach, the following assumptions will be used in estimating waste feed rates for the PRA.

- The waste treatment plant will operate at full capacity starting at year 1 (that is, the first two years of reduced throughput will not be accounted for).

- The waste treatment plant will operate at full capacity for the entire 40-year design life of the facility. This assumed operating period is longer than the operating period covered by the current permit application.

- Waste feed volume for years 1 and 2 will be equal to those expected for each year 3 through 40. This is a conservative assumption because the waste feed for years 1 and 2 will actually be less than for each year 3 through 11 and there is no reason to believe that the waste feed volume for the following years 12 through 40 will be any greater.

- Acceptance of waste feed by the waste treatment plant will be based on the feed parameters for which the pretreatment and vitrification processes are designed. The constituent concentrations defined in the waste envelopes represent the maximum anticipated constituent concentrations in the waste feed.

- The concentration of each inorganic COPC and ROPC in the LAW waste used to estimate emission rates for the PRA will be the maximum envelope concentration limits or the maximum concentration reported in the most recent characterization data, whichever is larger. For organics, the COPC concentrations will represent the estimated total mass of total organic carbon (TOC) assumed to be present in the feed less the mass known to be accounted for by chelating agents and simple organic acids.

- The concentration of each COPC and ROPC in the HLW waste used to estimate emission rates for the PRA will be the maximum feed composition limit for this waste. The maximum feed composition limit is conservatively estimated based on:

  - Envelope D concentration limits
  - A maximum solids content of 200 g/L
  - A minimum sodium concentration of 3M
  - An assumption that all of the solids in the waste are present in their oxide form

Measured constituent concentrations reported in the Tank Waste Information Network System (TWINS) Tank Characterization Database (PNL 1998) do not exceed the envelope concentration limits for any constituents.

Emission rate estimates for organic and inorganic COPCs and ROPCs in the waste feed will be estimated as described below.

Organic Chemicals in Waste Feed

Organic chemicals are not the primary component of the waste material to be processed by the waste treatment plant; however, a variety of organic chemicals will be present in the waste material. Table 4-1 identifies the organic COPCs potentially present in the waste feed. The COPCs that are not products of incomplete combustion (PICs) (that is, that are potentially present in the waste feed) will be modeled through the process and treatment train. The COPCs will be used to represent the TOC mass in the waste

feed, less the mass known to be chelating agents or simple organic acids. The TOC mass in the waste feed will be estimated in the following manner.

To maintain a conservative bias in the risk assessment, it is assumed that the waste feed will contain 9% organic carbon. This is more than twice the organic concentration allowable under the BNFL Inc. contract. Samples were collected and analyzed for organic chemicals from two tanks in August 1999 (Klinger et al., 1999). One sample and a duplicate were analyzed from each tank. These two tanks are estimated, by BNFL, to have some of the highest concentrations of organic chemicals in Hanford tank waste. The results of this analysis showed that tank AN107 had higher concentration of organic chemicals. Historical analyses of Hanford tank wastes have revealed that 78% to 99% of TOC can be accounted for by chelating agents and simple organic acids (Campbell et al. 1995). Assuming the most conservative contribution of organic constituents are hazardous constituents, other than chelating agents (22%), the most conservative TOC in the waste feed (9%), and a conservative density of the waste, the estimated worst-case concentration of organic constituents in the waste feed is 29,700 mg/L. This value was compared to the total mass of speciated organic chemicals identified by the analysis of the two samples from Tank AN107 (1 sample and 1 duplicate) by summing the maximum detected concentration of each detected chemical and the maximum detection limit for each non-detected chemical. Based on these results, the total mass of identified organic chemicals is 23.9 mg/L.

The organic COPCs in the waste feed will be used to represent the total TOC mass in the waste feed by using a scaling factor (that is, all detected and non-detected concentrations of each organic constituent are multiplied by a scaling factor of 29,700/23.9).

Once COPC concentrations in the waste feed are established, emission rates will be estimated as follows:

$$E_i = [F_{LAW} \times C_{LAWi} \times (1\text{-}DRE_i)] + [F_{HLW} \times C_{HLWi} \times (1\text{-}DRE_i)]$$

where

E_i = emission rate for chemical i (g/s)
F = LAW or HLW waste feed rate (L/s)
C_i = chemical i concentration in LAW or HLW feed (g/L)
DRE_i = destruction and removal efficiency for chemical i for the total treatment process (%/100)

The DRE will be estimated based on (1) results from bench-scale experiments of the melter technology conducted by The Catholic University of America, Vitreous State Laboratory beginning in November 1998 (Matlock and Pegg 1999), (2) thermodynamic modeling of chemicals not included in the bench-scale experiments, and (3) process and design specifications for the pretreatment process and post-treatment APCS. The DRE value to be used in the PRA will be selected following completion of preliminary thermodynamic modeling and facility design.

This approach for determining DRE is proposed for the following reason: the miscellaneous treatment unit regulations allow flexibility in selecting performance standards by recognizing that the units are not intended to perform all functions specified in the regulations for which specific regulations are applicable (e.g., incinerators). The primary function of vitrification melters is to incorporate dangerous and radioactive constituents into a stable glass form to the extent technically and economically feasible. Asking these vitrification units to perform other functions, such as destruction of organics to an extremely high percentage DRE, places a burden on the units that may impact the ability of the units to perform their primary functions (that is, make acceptable glass). Therefore, it is proposed that the quantities of hazardous organic constituents in the waste feed be investigated (during technology development testing) and the applicable DRE be established that is adequately protective of human health and the environment as defined by the risk assessment process.

Radionuclides in Waste Feed

Table 4-3 lists the ROPCs known or suspected to be present in the waste feed materials. Emission rates for these radionuclides will be estimated based on an assumed maximum radionuclide feed rate and an estimated system removal efficiency (SRE) for the waste treatment plant process. Emission estimates for normal operations will be calculated as:

$$E_i = [F_{LAW} \times C_{LAWi} \times (1\text{-}SRE_i)] + [F_{HLW} \times C_{HLWi} \times (1\text{-}SRE_i)]$$

where

E_i = emission rate for radionuclide *i* (pCi/s)
F = LAW or HLW waste feed rate (L/s)
C_i = radionuclide *i* concentration in LAW or HLW feed (pCi/L)
SRE_i = system removal efficiency for radionuclide *i* (%/100)

The SRE will be estimated based on (1) results from bench-scale experiments of the melter technology conducted by The Catholic University of America, Vitreous State Laboratory in November 1998 (Matlock and Pegg 1999), (2) thermodynamic modeling because experimental SREs will be based on cold surrogates (that is, no radionuclides are used), and (3) process and design specifications for the pretreatment process and post-treatment APCS. This SRE applies across the entire process (pretreatment, vitrification, and offgas). The SRE value to be used in the PRA will be selected following completion of preliminary thermodynamic modeling and facility design.

Inorganics in Waste Feed

Metals and other inorganics will be introduced into the system both as components of the waste stream and as process inputs to the treatment system (such as glass-forming materials including Al_2O_3, Fe_2O_3, and so on). Table 4-2 lists the inorganic COPCs known or suspected to be present in the waste feed and treatment feed materials. Emission rates for these inorganics will be estimated based on an assumed maximum feed rate and estimated SRE for the waste treatment plant process. Emission estimates for normal operations will be calculated as:

$$E_i = [F_{LAW} \times C_{LAWi} \times (1\text{-}SRE_i)] + [F_{HLW} \times C_{HLWi} \times (1\text{-}SRE_i)]$$

where

E_i = emission rate for metal *i* (g/s)
F = LAW or HLW waste feed rate (L/s)
C_i = metal *i* concentration in LAW or HLW feed (g/L)
SRE_i = system removal efficiency for metal *i* (%/100).

Concentrations of the glass formers will be specified by the process design. The SRE will be estimated based on (1) results from bench-scale experiments of the melter technology conducted by The Catholic University of America, Vitreous State Laboratory beginning in November 1998 (Matlock and Pegg 1999), (2) thermodynamic modeling of metals not included in the bench-scale experiments, and (3) process and design specifications for the pretreatment process and post-treatment APCS. This SRE applies across the entire process (pretreatment, vitrification, and offgas). The SRE value to be used in the PRA will be selected following completion of preliminary thermodynamic modeling and facility design.

In the case of mercury, assumptions must be made regarding the chemical speciation of the emissions as well as the total emission rate. The following conservative EPA 1998a default assumptions regarding mercury speciation will be used for the PRA:

Chemical state	% of total mercury exiting the stack
Vapor phase	80
Elemental Mercury (Hg)	(20)
Divalent Hg^{+2}	(60)
Particulate phase	20
Elemental Hg	(0.2)
Divalent Hg^{+2}	(19.8)

It is assumed that elemental mercury would be the form that leaves the stacks. Once settled in moist areas, mercury could be methylated. The fraction of methylmercury in soil is assumed to be 2% and 15%, respectively, in wetland soil and sediment (EPA Region X, 1999). Methylmercury fractions based on data for mesic soils are expected to overestimate methylmercury fractions in Hanford's xeric soils. Under the most conducive conditions for methylation of mercury in the environment, methylmercury seldom is more than a few percent of the inorganic mercury. For example, two different and independent methods concluded that methylmercury represented less than 0.01% of the total mercury in floodplain soils at East Fork Poplar Creek (EFPC) near Oak Ridge, Tennessee. Therefore, a conservative estimate of methylmercury in Hanford soils for the PRA could be 0.1% of total mercury. Measurements taken from wetland soils at EFPC showed methylmercury did not exceed 2.3% of total mercury. Therefore, 15% of total mercury is a conservative estimate of the fraction of methylmercury in sediment and wetland soil at Hanford for the PRA. Methylmercury was less than 0.1% (filtered) and 0.03% (unfiltered) of the total mercury in surface water. Therefore, it is assumed for the PRA that the fraction of methylmercury in Hanford surface water is 1% of total dissolved mercury. If measurements of methylmercury can be found for Hanford media, then they may be used in lieu of the EPA guidance of 2% for soil, 15% for sediment and wetland soil, and 1% for surface water.

Metals may originate from the following sources in addition to the glass forming chemicals and tank waste to be processed:

- Chromium may be associated with refractory material lining the melter
- Copper, nickel, palladium, and zinc may be used as catalysts in the APCS

These metals will be addressed to the extent possible in a quantitative fashion in the PRA.

5.2.1.2. PCDDs/PCDFs

Combustion Emissions Technical Resource Document (CETRED) (EPA 1994b) provides a technical analysis of emissions of polychlorinated dibenzo-p-dioxin (PCDDs)/polychlorinated dibenzofurans (PCDFs) from four types of hazardous waste combustion facilities: cement kilns, lightweight aggregate kilns, incinerators, and industrial boilers. The results of this analysis indicate that the best controlled hazardous waste combustion facilities can generally achieve a PCDD/PCDF emission level of 0.17 ng TEQ/dry standard cubic meter (DSCM) at 7% O_2. The RPP-WTP differs from these combustion facilities in both technology and waste feed. The anticipated MACT standard is 0.2 ng TEQ/DSCM at 7% O_2.

The RPP-WTP will be required to meet the dioxin/furan MACT standard once promulgated. Therefore, the anticipated MACT standard of 0.2 ng TEQ/DSCM at 7% O_2 will be used as the assumed stack concentration for the PRA and converted to an emission rate based on the designed volumetric flow rate of the melter.

Under the current industry-standard dioxin analytical method (EPA method 1613), dioxins in water can be analyzed at a part-per-quadrillion level, whereas dioxins in solid media (such as waste mixtures, soil, fish tissue) can be analyzed at a part-per-trillion level. These analytical limits are higher than many of the human health risk preliminary remediation goals (PRGs) and equivalent to ecological toxicity reference values (TRVs). For example, using data based on the dioxin, 2,3,7,8 TCDD, the PRGs for agricultural land-use scenarios are one order of magnitude lower than the detection limit using EPA method 1613. Thus, under the best analytical conditions (few interferences) the analyses for dioxins in some matrices would not be below levels that might cause unacceptable risk. It should be noted that the PRGs in the example above are based on movement of the dioxin through the food chain into beef, milk, and vegetables. The PRGs based on more direct exposure pathways (such as dermal contact to or ingestion of dioxin-contaminated soils) are higher (that is, part-per-trillion) and are the same order of magnitude that can be analytically achieved. This situation is because the food chain models use bioaccumulation factors and tend to be conservative in the estimation of concentrations after transfers.

Alternatively, ecological TRVs are at or near the same order of magnitude as the analytical detection limits. The lowest TRVs (4E-6 microgram/L) are for freshwater environments, whereas the analytical detection limit in water is around 10E-6 microgram/L. Adherence to laboratory procedures that reduce interferences and do not increase the detection limits would result in dioxin analytical measurements that are comparable to units in the risk characterization for ecological receptors.

Toxics Best Available Control Technology (TBACT) standards (based on a top down assessment of control options as required for the Clean Air Act permitting) will be addressed in separate documents but not the RAWP.

5.2.1.3. Other Products of Incomplete Combustion

Many organic COPCs are not present in the RPP-WTP waste stream. However, they may be present in stack emissions as PICs. Based on emissions testing during the laboratory studies, it is assumed some PICs may be emitted from the waste treatment plant. Indicator PICs, such as chlorinated benzenes, et al., will be modeled using thermodynamic assumptions and models, and results from the laboratory studies. Maximum Achievable Control Technology (MACT) standards may also be used to develop PIC emissions rates. The EPA is committed to establishing technology-based regulations for hazardous air pollutants (HAPs) under the *Clean Air Act* (CAA Amendments of 1990). The EPA process for establishing MACT standards is based on measured emission levels achieved by the best performing facilities.

5.2.1.4. Unknown Emissions

Unknown emissions are the percentage of organic emissions that cannot be quantitatively analyzed during the environmental performance demonstration. Unknown emissions are nearly impossible to estimate for the PRA. However, use of the COPCs to represent the total organic fraction in the waste will address risk concerns associated with unknown emissions.

This methodology results in scaling up estimated waste feed concentrations of each organic COPC by a factor of 1242, which will result in a conservative estimate of emissions and risk. This will also be discussed in the uncertainty section.

5.2.2. Process Upset Conditions

Process upset conditions include periods of startup, shutdown, process malfunction (that is, the unit is operating outside the permitted operating conditions), or equipment failure. Process upset conditions are generally assumed to result in greater than normal stack emissions during the short period of the upset. However, the EPA 1998a has indicated that upsets are not generally expected to significantly increase stack emissions over the lifetime of a facility.

The potential for increased emissions during upset events will be addressed through the application of an upset factor. This upset factor is based on the amount of time the facility is expected to operate in an upset condition, and the estimated magnitude of stack emissions during upset relative to routine operating conditions. The preferred method for estimating this upset factor is through the use of data from existing facilities that have operating conditions similar to the proposed waste treatment plant. The frequency and duration of upset events may be estimated based on EPA 1998a:

- Data from emissions monitoring systems that measure operating parameters such as stack carbon monoxide or oxygen
- Data on combustion chamber, air pollution control system (APCS), or stack gas temperature
- Ratio of automatic waste feed cut-off frequency and duration to operating time
- Variations in the APCS operating conditions

The potential magnitude of emissions during upset events may be estimated based on stack test data collected during upset conditions.

EPA default upset factors represent worst-case conditions and will be used for the PRA unless sufficient process information is available to estimate conservative, site-specific upset factors. These default upset factors are based on the data described above from operating hazardous waste combustion facilities. These default upset factors are expected to over-predict upset emissions from the waste treatment plant for several reasons including:

- Carbon monoxide is frequently used as an indicator of upset conditions, and automatic waste feed cut-offs are often triggered by increased stack gas concentrations of carbon monoxide. However, routine operations such as adjusting waste feed or air intake rates will cause brief spikes in carbon monoxide concentration.
- Test data used for these defaults are based on hazardous waste combustion facilities designed for the destruction of liquid and/or solid organic waste. The technology and waste feed of the RPP-WTP melter are different and less subject to upset than these facilities.
- The waste treatment plant will include a standby offgas treatment system designed to treat emissions during an upset event.

EPA 1998a default upset factors are 2.8 for organic chemicals and 1.45 for metals calculated as shown below:

Organics. A default facility is assumed to operate under upset conditions 20% of the time and stack emissions are assumed to be 10 times greater than normal during this time.

$$\text{Upset Factor} = (0.80)\,(1) + (0.20)\,(10) = 2.8$$

Metals. A default facility is assumed to operate under upset conditions 5% of the time with stack emissions 10 times greater than normal during this time.

$$\text{Upset Factor} = (0.95)\,(1) + (0.05)\,(10) = 1.45$$

EPA has not determined a default upset factor for radionuclides. For the PRA, radionuclides are assumed to behave similarly to metals with an upset factor of 1.45. The same upset factors will be used for both the plausible and worst-case scenarios in the PRA and the FRA.

Site-specific upset factors may be derived based on the following:

- Data describing upset frequency, duration, and impact on emissions from similar melter facilities
- Process design data describing the potential for, and potential impact of, upset conditions at the waste treatment plant

The potential for increased emissions during upset conditions is addressed in section 3. The most likely condition that would result in a process upset in the LAW and HLW melters is a pressure surge in the melters resulting in a "bypass event". Potential emissions associated with bypass events will be evaluated separately from upsets that are due to deteriorated operations. The melter is designed to avoid bypass events through the use of an alternative route for emissions. This alternative route will prevent bypass events unless power is lost. In the event of a complete power failure, a bypass event could occur for a brief time (1 minute or less) until backup power is initiated. In the event of a power failure, the offgas would be released to the confinement cell and exhausted through the cell ventilation HEPA filter or returned to the melter and, then, exhausted through the vessel vent offgas system when power is restored.

5.2.3. Fugitive Emissions

Fugitive emissions include emissions of COPCs to air from any source except the stack. Because of the stringent controls designed to prevent radiation exposures and especially the negative pressures of the processing facilities, it is not anticipated that fugitive emissions will be emitted from either the pretreatment, LAW treatment, or HLW treatment processes. Insignificant amounts of fugitive emissions are anticipated from the materials storage areas that support processing plants as described below.

5.2.3.1. Waste Treatment Processing Facilities

The waste treatment plant ventilation system is being designed to confine all sources of radiation and radiological contamination and, thus, fugitive releases of COPCs to the environment are not anticipated. Confinement will be achieved through a combination of barriers or barrier systems, coupled with a cascade ventilation system that will be designed to conform to stringent nuclear facility ventilation standards.

The key to eliminating fugitive emissions is that the more contaminated areas are maintained at a greater negative pressure with respect to the less contaminated areas. Because of this pressure gradient, no fugitive emissions should be released from the processing plants (pretreatment, LAW pretreatment, LAW vitrification, and HLW vitrification). The air flow from this containment system is HEPA filtered and routed into the LAW and HLW offgas streams for treatment before emission to the environment.

5.2.3.2. Material Storage Systems

There are two storage facilities not directly associated with the main processing plant that have the potential to be minor sources of fugitive emissions: the glass former storage system and the nitric acid storage system.

The glass former storage bins will provide bulk storage for the alumina, boric acid, iron oxide, and similar materials that will be used as part of a recipe to support glass formation in the melters. These materials have no radiological contamination and do not contain COPCs, but very small levels of particulate emissions may occur from periodic refilling of the bins by outside vendors. The storage systems will have baghouses installed to minimize emissions during filling and normal operations.

The wet chemical storage system will be maintained to provide the nitric acid and other bulk chemicals used in operating the process plant. Emissions from this facility are not expected to be significant.

5.3. Measured Emission Rates for the Final Risk Assessment

The FRA will be conducted following construction of the waste treatment plant and completion of the environmental performance demonstration (see BNFL 1999c). Emission rates for the FRA will be estimated based on the environmental performance demonstration results.

The environmental performance demonstration program will be conducted in support of the *RPP-WTP Dangerous Waste Permit Application* (BNFL 1999a) using the "Universal Approach" to permitting described in EPA 1989a. With this approach, a single set of operating conditions is sought for treating a relatively broad range of waste. To establish the set of operating conditions, the following worst-case environmental performance demonstrations will be performed on the LAW vitrification and offgas treatment systems (LAW treatment system), and the HLW vitrification and offgas treatment systems (HLW treatment system):

- Low-temperature test designed to maximize the potential emissions of organic compounds
- High-temperature test designed to maximize the potential emissions of metals

Given that the waste treatment plant will not operate at the extreme limits of its operating envelope all the time, EPA 1998a provides for an additional test condition consisting of normal or average system operating parameters that may be specified in the testing program. If, as a result of the PRA, it appears that estimating potential risks based on emissions from worst-case DRE and/or SRE conditions are overly conservative, the normal operations test will be incorporated into the environmental performance demonstration. The normal operations test condition would be representative of long-term facility operations with equipment settings equivalent to those used during actual waste treatment. If the normal operations test condition is included in the environmental performance demonstration, these results will be compared to those derived under worst-case conditions and will be used for estimating emission rates in the FRA.

5.3.1. Stack Emissions

To minimize the dose to sampling and analytical personnel and to obtain useful detection limits in the environmental performance demonstration, a surrogate waste (performance demonstration feed) will be used to represent tank waste. The performance demonstration feed is a nonradioactive blend of chemicals (including radioactive surrogates), water, and glass-formers designed to mimic waste to be treated by the waste treatment plant spiked with specific principal organic dangerous constituents (PODCs), metals, and

chlorinated compounds. The performance demonstration feed will be formulated to represent worst-case blends of difficult-to-destroy PODCs, metals, and chlorinated compounds. One performance demonstration feed will be developed that will consist of the worst-case constituent concentration conditions for the LAW Envelopes A, B, and C for testing the LAW melter. A second performance demonstration feed will be developed that consists of the worst-case constituent concentration conditions for the HLW Envelope D for testing the HLW melter. The specific makeup of performance demonstration feed will be identified in the *Environmental Performance Test Plan* (BNFL 1999c) and will include surrogates for primary radionuclides.

Sufficient quantities of PODCs, metals, and chlorinated compounds will be part of the performance demonstration feed to measure the following:

- DRE for each PODC
- SRE for the selected metals and radionuclide surrogates
- Gaseous HCl and Cl_2 emissions
- Particulate matter emissions
- Emission rates for PCDDs/PCDFs
- Emission rates of other organic constituents

Chemicals to be spiked in the performance demonstration feed will be selected based on the compound's heat of combustion, thermal stability-based incinerability ranking, predicted behavior based on thermodynamic calculations, the results of small-scale testing and technology demonstrations, and Hanford waste analytical data as described in the *Environmental Performance Demonstration Plan*.

Different approaches will be used to estimate stack emissions for different categories of COPCs as described below.

5.3.1.1. COPCs Identified in Waste Feed

Chemicals and radionuclides present in the waste feed may be present in stack emissions and will be evaluated in the FRA as described below. Emission rates calculated from the measured removal efficiency across the waste treatment plant will be estimated based on the chemical feed rate and DRE/SRE as described previously for the PRA. Chemical feed rates will be estimated as described previously using any new information that may become available after completion of the PRA (such as new concentration data for tank waste, and waste feed information based on updated process designs). The DRE/SRE to be used for the FRA will be based on the performance demonstration results.

Spiked Chemicals (PODCs, metals, radionuclide surrogates). Stack emissions will be estimated from the DRE/SRE demonstrated for these chemicals. The DRE/SRE will be calculated as:

$$\text{DRE/SRE} = [(Q_{feed} - Q_{emission})/Q_{feed}] \times 100$$

where

Q_{feed} = feed rate of the chemical (g/s)

$Q_{emission}$ = emission rate of the chemical at the stack (g/s). The upper confidence limit on the average stack gas concentration will be used. If the chemical is detected in some but not all three of the runs, the detected concentration(s) will be used along with the detection limit for the chemical that was not detected.

The detection limit being used in a situation where chemicals are not detected is the method detection limit (MDL)-derived reported detection limit (RDL) for non-isotope dilution and is consistent with EPA 1998a. For isotope dilution methods where chemicals are not detected, the method or the method-defined estimated detection limit (EDL) is used and is consistent with EPA 1998a.

COPCs in Tank Waste and Included in Performance Demonstration Feed. If the chemical is detected in some but not all stack gas samples, the detected concentration(s) will be used along with the detection limit for the chemical that was not detected. Since the performance demonstration feed will contain worst-case concentrations of COPCs from all three LAW Envelopes, an adjustment may be made for the amount of each envelope to be included in the actual waste feed.

COPCs in Tank Waste but Not Included in Performance Demonstration Feed. Stack emissions will be estimated from the DRE/SRE demonstrated for PODCs or other COPCs. The DRE/SRE will be estimated as described above. The PODCs are intended to represent the most difficult to destroy chemicals. Therefore, the DRE/SRE estimated from other COPCs included in the performance demonstration feed may be used to estimate emissions of more easily destroyed COPCs.

5.3.1.2. PCDDs/PCDFs

Dioxin and furan congeners are not present in the waste treatment plant feed stream. However, they may be present in stack emissions as PICs. PCDDs/PCDFs identified in stack emissions will be evaluated as the upper confidence limit on the average from the three environmental performance demonstration runs. If a congener is detected in some but not all three of the runs, the detected concentration(s) will be used along with the detection limit for the non-detects. PCDDs/PCDFs not identified in stack emissions will be evaluated at their detection limit.

5.3.1.3. Other COPCs

Chemicals not present in waste may be present in stack emissions as PICs and will be evaluated in the FRA as described below.

COPCs Identified in Stack Emissions. For COPCs not present in the tank waste or performance demonstration feed but identified in stack emissions, the lesser of the 95th percentile or maximum stack gas concentration from the environmental performance demonstration will be used to develop the emission rate estimated used in the risk assessment (EPA 1998a). If the chemical is detected in some, but not all three of the runs, the detected concentration(s) will be used along with one-half the detection limit for the non-detect samples in these calculations.

COPCs That Are Potential PICs or Risk Drivers. COPCs that are not present in the tank waste or environmental performance demonstration feed and are not identified in stack emissions during the environmental performance demonstration, but which may be expected in the stack emissions as PICs (based on thermodynamic modeling) or are risk drivers will be evaluated at one-half their detection limit.

COPCs That Are Not Potential PICs or Risk Drivers. COPCs that are not present in the tank waste or environmental performance demonstration feed, are not identified in stack emissions during the environmental performance demonstration, are not expected in the stack emissions as PICs (based on thermodynamic modeling), and are not risk drivers will be assumed to be not present (that is, emission rate = 0).

5.3.1.4. Unidentified Organic Chemicals

The potential emission rate for unidentified organic chemicals will be based on environmental performance demonstration estimates of total organic emissions (TOE). The TOE results will be used to estimate a TOE factor as shown below:

$$F_{TOE} = (TOE_C/\Sigma_i C_i) \times (TOC_{waste}/TOC_{PTF})$$

where

F_{TOE} = total organic emission factor (unitless)
TOE_C = total organic emission concentration (mg/m^3)
C_i = stack concentration of organic $COPC_i$ (mg/m^3)
TOC_{waste} = total organic carbon concentration in waste feed (g/L)
TOC_{PTF} = total organic carbon concentration in performance demonstration feed (g/L).

The equation is from (EPA 1998a, p. 2-10) with a correction factor to account for the use of surrogate feed in the environmental performance demonstration. The TOE factor is corrected for the difference in organic carbon concentration between the actual waste feed and the performance demonstration feed used in the environmental performance demonstration. The TOE factor will be calculated from emission concentration data in the FRA because environmental performance demonstration data will be available at that time. TOC will be an analyte for the test burn. The $\Sigma_i C_i$ includes all compounds not detected at the same concentration assumed for emission rate estimates of these compounds as described above (that is, detection limit, 1/2 detection limit, or zero) and will include any TICs included with the COPCs.

5.3.2. Process Upset Conditions

The potential impact of upset conditions on emission rates will be addressed by using an upset factor as described in section 5.2.2. The upset factors used for the PRA will be reevaluated for the FRA based on any changes in the design of the waste treatment plant process made since the PRA was conducted or additional data than becomes available for similar facilities.

5.3.3. Fugitive Emissions

The potential sources and magnitude of fugitive emissions identified for the PRA (section 5.2.3) will be re-evaluated for the FRA based on the following:

- Changes in the design of the waste treatment plant process made since the PRA was conducted
- Observations during operation of the waste treatment plant during the operational testing period and environmental performance demonstration

Revised fugitive emissions factors will be determined based on the above evaluation.

5.4. Uncertainties in Estimating Emission Rates

The estimation of emissions from the waste treatment plant is an important initial step in the risk assessment. Uncertainties associated with this step will affect the results of the human health and ecological risk assessments.

5.4.1. Uncertainties in Estimating Emission Rates for the Preliminary Risk Assessment

The PRA will be conducted prior to the environmental performance demonstration and is based on the current facility design. This early PRA will result in a high level of uncertainty in the emission rate estimates. This uncertainty will be addressed two ways: first, conservative assumptions will be used in an attempt to ensure that the uncertainty will result in overestimating rather than underestimating emissions; and second, many of the assumptions used will be re-evaluated following construction of the waste treatment plant and environmental performance demonstration. The primary sources of uncertainty in estimating stack and fugitive emissions are described below.

5.4.1.1. Identification of COPCs/ROPCs

The COPC/ROPC list for the SLRA includes all chemicals known or suspected to be present in the tank waste as well as all chemicals known or suspected to be emitted from a wide variety of hazardous waste incinerators. Note that many of the COPCs/ROPCS included in this list will not be emitted from the waste treatment plant.

5.4.1.2. Estimated Concentration of COPCs and ROPCs in Waste Feed

Emission rates will be estimated from the concentration of COPCs and ROPCs in waste feed, waste feed rate, and the DRE/SRE achieved by the treatment process. Emission estimates will be based on high-end estimates of the waste feed concentrations.

5.4.1.3. Waste Feed Rate

The waste treatment plant is assumed to operate at its maximum design throughput capacity for its design life. It will take several years for the plant to reach its full capacity.

5.4.1.4. DRE/SRE

The DRE/SRE of the process will be estimated from the process design, vendor specifications for various process equipment, and laboratory-scale testing of the technology. The use of these data sources may result in an overestimation or underestimation of the actual DRE/SRE capability of the process.

5.4.1.5. Estimated Concentrations of PCDDs/PCDFs and Other PICs

The emission rate for PCDDs/PCDFs assumes that the waste treatment plant will meet the likely MACT standard for these compounds. This standard is based on data from hazardous waste incinerators burning large quantities of chlorinated organic chemicals. The waste treatment plant emissions are expected to be below this standard because the RPP-WTP is much different from an incinerator (that is, the organics in the feed are much lower) and there has been historical restriction on the amount of chlorides introduced to the tank system due to the carbon-steel construction of the tanks.

Emission rates of other PICs will be based on data from surrogate testing, process design, and thermodynamic modeling and mass balance. These data could result in an over or under estimation of PIC emissions.

5.4.1.6. Process Upset Conditions

Process upset conditions include periods of process malfunction. Process upset conditions are generally assumed to result in greater than normal stack emissions during the short period of the upset. However, EPA 1998a has indicated that upsets generally are not expected to significantly increase stack emissions over the lifetime of a facility.

The potential for increased emissions during upset events will be addressed by applying an upset factor. This upset factor is based on the amount of time the waste treatment plant is expected to operate in an upset condition and the estimated magnitude of stack emissions during the upset condition relative to routine operating conditions. The EPA default upset factors are designed to provide a conservative estimate of potential emissions at a variety of hazardous waste incinerators (EPA 1998a). The waste feed and technology used by the RPP-WTP differ from a hazardous waste incinerator, and these default values are expected to overestimate the potential contribution to emissions from process upsets. Conservative upset factors derived from the process design and similar facilities will be used if possible and appropriate.

5.4.1.7. Fugitive Emissions

The potential sources and magnitude of fugitive emissions are estimated based on the waste treatment plant design and conservative assumptions regarding the type and concentration of volatile chemicals in the tank waste. The plant is designed to minimize potential fugitive emissions with most potential emission sources directed toward the stack. The potential impact of fugitive emissions on the uncertainty of the assessment is low.

5.4.2. Uncertainties in Estimating Emission Rates for the Final Risk Assessment

The FRA will be conducted following construction and environmental performance demonstration of the waste treatment plant and will reduce the level of uncertainty in the emission rate estimates. The impacts of the FRA on the primary sources of uncertainty in estimating stack and fugitive emissions are described below.

5.4.2.1. Identification of COPCs

The COPC list will be re-evaluated based on the results of the PRA and the environmental performance demonstration.

5.4.2.2. Estimated Concentration of COPCs and ROPCs in Waste Feed

Emission rates will be estimated from the concentration of COPCs and ROPCs in waste feed, waste feed rate, and the DRE/SRE achieved by the treatment process. The emission estimates will be based on high-end estimates of the waste feed concentrations for the PRA and FRA.

5.4.2.3. Waste Feed Rate

The waste treatment plant is assumed to operate at its maximum design throughput capacity for its design life for both the PRA and FRA unless additional information becomes available to make a more realistic assumption.

5.4.2.4. DRE/SRE

The DRE/SRE of the process will be estimated from the results of the environmental performance demonstration. This estimate will reduce the uncertainty in the assessment. The environmental performance demonstration will utilize worst-case operating conditions to minimize the measured DRE/SRE and maintain the conservative bias of the assessment.

5.4.2.5. Estimated Concentrations of PCDDs/PCDFs and Other PICs

The emission rates of PCDDs/PCDFs and other PICs will be measured during the environmental performance demonstration. The performance demonstration feed will maximize the concentrations of hard-to-destroy and chlorinated organic chemicals to maximize the potential formation of PICs. This process will result in an overestimation of PIC emissions.

5.4.2.6. Unidentified Organics

The FRA will utilize measured concentrations of organic chemicals in stack emissions rather than waste feed concentrations, thus reducing the uncertainty in estimating unidentified organics.

5.4.2.7. Process Upset Conditions

The upset factors used for the PRA will be reevaluated for the FRA based on any changes in the design of the waste treatment plant process, thus reducing the uncertainty in this parameter.

5.4.2.8. Fugitive Emissions

Availability of the completed design and operating information will reduce the uncertainty in this parameter.

5.5. Summary for Quantification of Emissions

The SLRA uses estimated emission rates of the COPCs and ROPCs to calculate exposure concentrations in air and other media (section 6). Estimated emission rates will include potential emissions from the facility stack(s) during normal (routine) operations as well as emissions resulting from nonroutine (upset) events and the few fugitive emissions that may escape from parts of the waste treatment plant other than the stack(s).

The PRA will be conducted prior to construction of the waste treatment plant, therefore, COPC/ROPC emission rates will be estimated based on waste feed data, process and design information, emission data from other facilities, laboratory studies, and default assumptions.

The FRA will be conducted following completion of the environmental performance demonstration. Emission rates for the FRA will be estimated based on the environmental performance demonstration results. The primary difference between the PRA and FRA is the estimation of emission rates.

6. Environmental Modeling

Environmental modeling refers to four types of models that simulate the route of chemicals and radionuclides from the stack towards human and ecological receptors. Air dispersion modeling is defined first, followed by soil accumulation, biota uptake, and finally surface water and fish modeling.

6.1. Air Dispersion Modeling

Air dispersion modeling will be used to estimate the ambient air quality and deposition rates resulting from the emissions of chemicals and radionuclides during operations of the RPP-WTP. This section provides details of the approach that will be used in this task.

6.1.1. Model Selection

The Industrial Source Complex Short Term Model, Version 3 (ISCST3) (EPA 1995a and EPA 1995b) will be used to evaluate the air quality impacts in the vicinity of the RPP-WTP. This EPA-preferred model (*Guideline on Air Quality Models*, 40 *CFR* Part 51, Appendix W) is generally considered a conservative model for applications such as the SLRA. The model uses emissions source data and hourly meteorological data to estimate ambient air concentrations and deposition rates of gases and particles at locations (receptors) of interest in the vicinity of the facility (EPA 1995a, EPA 1995b, EPA 1998b). The most recent version of the ISCST3 (98226) will be used in this analysis and will be supplemented by EPA *Risk Assessment Guidelines* RAGs Part A and B models for radionuclides. This model has the capability of handling a large number of sources that could occur from a typical industrial source, including point sources (e.g., stacks) and area sources (e.g., fugitive emissions from an open area). However, it is expected that fugitive emissions will be so small that a qualitative evaluation of the near-field environment will suffice.

The ISCST3 model is capable of calculating ambient concentrations and deposition rates under variable terrain conditions. This capability is important in air modeling because the calculation of ambient air concentrations and deposition rates is greatly influenced by the height of the plume above local ground level. Terrain is defined as "complex" if the elevation of the surrounding terrain within the assessment area (up to 50 km for this risk assessment) is above the top of the stack, and "simple" if the terrain is at or below the stack height. EPA has developed specific guidance for use in complex terrain (to avoid the case where the source plume directly impacts elevated terrain), which was previously incorporated in the COMPLEX1 model. The ISCST3 model implements the algorithms from the COMPLEX1 air quality model when complex terrain occurs.

The ISCST3 model will be used to calculate ambient concentrations, and wet and dry deposition rates for COPCs and ROPCs at pre-determined exposure locations. The terrain elevation of each receptor will be included in the model input. Terrain elevations will be obtained from digitized maps of the Hanford Site for receptors located within the site or from U.S. Geological Survey digitized maps for receptors located outside of the site.

6.1.2. Model Parameters

This section discusses the modeling input parameters for the air dispersion and deposition modeling including: emissions data, meteorological data, exposure locations, calculations of deposition rates, and model variable settings.

6.1.2.1. Emissions Source Information

Identification of emission sources and quantification of emission rates for each specific COPC and ROPC was described in section 4.2, "Quantification of Emissions". Other data required for model execution, such as stack heights, stack diameters, stack gas flow velocities, and stack gas temperatures, will be provided in the PRA along with all model output data.

Unit Emission Rates. The ISCST3 model will be run with a unit (normalized) emission rate of 1.0 g/s, which is used as model input to preclude having to run the model with specific emissions for each COPC/ROPC. The relationship between the emission rate from a single source and the air parameter values (air concentrations and depositions) at a particular receptor is linear. Therefore, the normalized concentration and deposition outputs from ISCST3 can be adjusted to the COPC/ROPC-specific air concentrations and deposition rates by using COPC/ROPC-specific emission rates, as illustrated below:

$$COPC_{conc} = ISCST3_{conc} \times (COPC_{er}/Unit_{er})$$

where

$COPC_{conc}$ = COPC/ROPC-specific air concentration at a receptor ($\mu g/m^3$ for COPCs; pCi/m^3 for ROPCs)
$ISCST3_{conc}$ = modeled concentration based on the unit emission rate ($\mu g/m^3$ for COPCs; pCi/m^3 for ROPCs)
$COPC_{er}$ = COPC/ROPC-specific emission rate (g/sec)
$Unit_{er}$ = unit emission rate of 1 g/sec.

Both radiological and chemical decay will be accounted for by use of a decay term:

$$COPC_{conc(t)} = COPC_{conc(0)} \times e^{-\lambda t}$$

where:

COPCconc(t) = COPC/ROPC-specific air concentration at time t (μg/m3)
COPCconc(0) = initial COPC/ROPC-specific air concentration (μg/m3)
e = the base of the natural logarithm, approximately 2.71828
λ = radiological or chemical decay constant (day-1)
t = time elapsed in days since emission of the COPC/ROPC.

where

λ = 0.693/T1/2 and T1/2 is the half-life in days of chemical decay of COPCs (e.g., photooxidation) or radiological decay of ROPCs.

Analysis of Multiple Stacks. Based on the preliminary design of the RPP-WTP, there may be numerous separate stacks (LAW and HLW Vitrification System Stacks, Pretreatment Stack, and separate abated fugitive emissions stacks). If the stacks are located close together and have identical characteristics (stack heights, operating parameters, COPC/ROPC-specific emission rates), then a single model run may be used to represent the stacks. However, it is expected that each stack will need to be modeled separately using the ISCST3 model, because the COPC/ROPC-specific emission rates are likely to vary between stacks, and the stacks may have different elevations or emission characteristics. Thus, at a given receptor, one stack may be the main contributor of ambient concentrations and deposition rates of some COPCs/ROPCs, while another stack would be the primary contributor of other COPCs/ROPCs.

Assuming that the RPP-WTP has more than one stack, each source will be modeled separately using the ISCST3 model. The contribution from each source to the total COPC/ROPC-specific concentration and deposition rate at a receptor will be combined in a spreadsheet to determine the total COPC/ROPC-specific ambient concentration and deposition rate at each exposure location.

6.1.2.2. Meteorological Data

The ISCST3 model uses hourly meteorological data, along with the emissions data, as one of the two primary input data sets. Five years of comprehensive Hanford Site surface meteorological data collected by the Hanford Site Meteorological Station during the period 1992 to 1997 will be used as input into the model. The Hanford Meteorological Monitoring Network includes a station located at the 200-E area, within 1 mile of the location of the proposed RPP-WTP facility. The air-dispersion modeling uses wind speed, wind direction, and temperature data from that station. Other, less site-specific parameters used by the model, such as precipitation and solar radiation, are collected at the main Hanford Meteorological Station, located approximately 5 miles west-northwest of the RPP-WTP site. The terrain in the vicinity of the RPP-WPP site is relatively flat, with no major topographic features. The close proximity of the monitoring stations to the proposed facility and the lack of terrain obstacles ensure that the collected meteorological data are representative of the site.

The meteorological monitoring station at the 200-E area (near the site of the proposed BNFL facility) is part of the 30-station Hanford Meteorological Monitoring Network, operated by Pacific Northwest National Laboratory (PNNL) for the Department of Energy. The 200-E Station consists of a 10-meter tower, with wind speed and wind direction data collected at the standard 10-meter height and ambient temperature monitored at 2 meters. An on-site data logger collects the data for each parameter, computes 15-minute averages, and transmits these averages to the main Hanford Meteorological Station, where the data are permanently archived. Other data that are less site-specific, such as barometric pressure, precipitation, and solar radiation, are monitored at the main 410-foot Hanford Meteorological Station.

PNNL operates the stations on a daily basis and maintains a comprehensive quality assurance program to ensure the quality of the data collected in the Hanford Meteorological Monitoring Network. Because the risk assessment will be conducted on the concentrations and deposition rates calculated by using the meteorological data from the worst-case year, the ISCST3 model will be run independently with each year of meteorological data.

In addition to the surface meteorological data, upper air data collected during the same period are required to run the ISCST3 model. The upper air data, typically collected twice-daily at selected National Weather Surface (NWS) stations, are used to provide an estimate of the morning and afternoon mixing heights as an input to the model. Although mixing heights computed from -Hanford Site meteorological data would be desirable, the nearest valid upper air data are collected at Spokane, Washington, approximately 125 miles to the northeast of the Hanford Site. The Spokane upper air data should be representative of the Hanford Site, because Spokane is in the same regional climatic area as the Hanford Site. The upper air data are available from the National Climatic Data Center in Asheville, North Carolina, USA.

The screening-level risk assessment will use ISCST3 air quality model to determine the ambient concentrations and deposition rates of COPCs due to the emissions from the RPP-WTP. This model only allows the input of meteorological data from one station, which may result in some uncertainty in the results due to wind direction variations across a large area such as the Hanford Site. This limitation is one of the shortcomings of the ISCST3 model.

However, this is a screening level risk assessment, and EPA has approved the use of ISCST3 as the default model in similar applications. If the results of this risk assessment produce results that are near unacceptable risk levels, then it may be warranted to use a more comprehensive model such as CALPUFF, that allows the use of multiple meteorological data sets.

The *Hanford Site Climatological Data Summary 1997, With Historical Data* (PNNL-11794 UC-603) summarizes the precipitation data collected over the period of record for the Hanford meteorological monitoring stations. The document lists the average number of days per year with precipitation of a specified amount. These data are presented below.

- **Average annual number of days with 0.10 inch of precipitation or more: 23 days**
  [12 of these days (approximately 50 percent) occurred during the months of November through February, in which the precipitation is likely to be in the form of snow]

- **Average annual number of days with 0.25 inch of precipitation or more: 7 days**
  [4 of these days (approximately 50 percent) occurred during the months of November through February, in which the precipitation is likely to be in the form of snow]

- **Average annual number of days with 0.50 inch of precipitation or more: 1 day**

Thus, the extremely dry climate at the Hanford Site (average precipitation is just over 6 inches per year), means that there are very few days in which there is a potential for entrainment of soil due to rainwater splash. Investigation of this mechanism therefore does not appear to be appropriate for this screening level risk assessment.

The ISCST3 model requires the use of additional computer programs, referred to as "preprocessing" programs to format the meteorological data and other input variables into a file that can be used by ISCST3. The two preprocessing programs to be used for this modeling effort are described below.

The MPRM Preprocessing Program. A meteorological preprocessor program must be used to prepare the meteorological data for input into the ISCST3 program. For on-site data, the Meteorological Processor for Regulatory Models (MPRM) is recommended by EPA, and will be used in this risk assessment (EPA 1996a). This model performs the required meteorological formatting, and meshes the Hanford Site data with the NWS upper air data.

In addition to formatting the meteorological data, the MPRM model requires an additional series of input parameters, which are representative of the application site. These are site-specific parameters that can influence the calculated deposition rates, and include:

- Monin-Obukhov length (a measure of atmospheric stability)

- Surface roughness height (a measure of the height of obstacles to wind flow, used to determine how close a particle must be above the ground before it is subject to deposition)

- Noon-time albedo (the fraction of incoming solar radiation that is reflected from the ground at noon, which affects the hourly net heat balance)

- Bowen ratio (a measure of the amount of moisture at the surface, which affects the heat balance)

- Anthropogenic heat flux (surface heating caused by human activity)

- Fraction of net radiation absorbed at the ground (the final component of the radioactive heat balance)

EPA has provided guidance for estimating values for these parameters in the user guides for the MPRM meteorological preprocessing program and the ISCST3 air quality model (EPA 1996a and EPA 1995a and EPA 1995b). Estimated values are provided in these user guides for different land use categories or different vegetative types, depending on the parameter. Values also vary for some parameters by season. The most appropriate value for each parameter by season will be entered into the model input file.

The ISCST3 model requires that the meteorological data set contain no missing records. Any missing data will be substituted using EPA-recommended procedures (Atkinson and Lee 1992).

Building Profile Input Program (BPIP) Preprocessing Program. Building wake effects can have a significant impact on the dispersion of emissions near a stack. The turbulence induced by buildings produces a phenomenon, known as building downwash, in which a stack plume can be brought downward toward the ground much sooner than if the buildings were not there, resulting in localized areas of high emission concentrations.

Consequently, the ISCST3 model contains algorithms for evaluating the potential for downwash. Following EPA guidance (EPA 1998a), all nearby structures with heights of at least 40% of the height of the shortest stack to be modeled will be evaluated for building downwash. The BPIP calculates the maximum crosswind widths of the buildings, which are then input into the ISCST3 model to estimate building wake effects on air dispersion. The required building dimensions (length, height, and width) will be obtained from engineering drawings and used as input in the BPIP preprocessor. The BPIP output file is formatted for direct input into the ISCST3 model.

6.1.2.3. Receptor Data

The impact assessment area, for which COPC/ROPC-specific concentrations and deposition rates will be calculated, will extend out to 50 km in all directions from the RPP-WTP.

Receptor Grid. The ISCST3 model will be run using a Cartesian grid of exposure locations extending out to 50 km from the center of the RPP-WTP. The receptor grid points will be identified using universal transverse mercator coordinates. Receptor grid spacing will be established at 100-m intervals from the RPP-WTP up to a distance of 3 km, at 500-m intervals from 3 to 10 km, at 2-km intervals from 10 to 30 km, and at 4-km intervals from 30 to 50 km (Figure 6-1). As needed, additional modeling runs will be made with a finer grid resolution (50 m) to more precisely identify the locations of maximum impacts.

Discrete Receptors. Additional exposure locations will be evaluated at specific locations identified as potentially sensitive receptors in the human health and ecological risk assessments.

Receptor Elevations. The elevation for each exposure location will be used as input into the ISCST3 model. Elevations for all exposure locations within the Hanford Site will be obtained from digitized maps of the area. Elevations for exposure locations outside the site are available in digital form from the U.S. Geological Survey.

6.1.2.4. Calculation of Deposition Rates

The determination of deposition rates is an important input into the human health and ecological risk assessments being conducted for the RPP-WTP. The ISCST3 model will be used to calculate both wet and dry deposition rates, in addition to ambient concentrations, at each exposure location.

Dry deposition occurs in the absence of precipitation, while wet deposition is influenced by precipitation type and rate. The two types of deposition result from different physical processes and, therefore, must be considered separately.

Dry Deposition. The dry deposition flux, F_d, is calculated as follows:

$$F_d = \chi_d \times v_d$$

where

F_d = dry deposition flux ($\mu g/m^2$-sec)
χ_d = ambient concentration value calculated in ISCST3 ($\mu g/m^3$)
V_d = deposition velocity (m/sec).

The ISCST3 model calculates the deposition velocity from particle diameter, mass fraction, and particle density, which are the data input into the model for each particle size fraction. The calculation of deposition velocities within the ISCST3 model includes the effects of Brownian motion, inertial impaction, and gravitational settling. Particularly for the larger particles, the key parameter governing the rate of dry deposition is the terminal settling velocity. The terminal velocity, in turn, is affected primarily by the particle size and density: large particles have the highest terminal velocities (and therefore the highest deposition rates), and small particles have lower terminal velocities.

It is important to note that particles have a positive terminal settling velocity and, therefore, are subject to dry deposition. The revised (May 15, 1999) ISCST3 model issued by EPA in June 1999 now includes a gas dry deposition algorithm, and consequently the analysis of dry deposition will be extended to gasses through the application of the enhanced ISCST3 model. Solar radiation is a required parameter in the gas dry deposition algorithm and will be obtained from on-site data sets collected by the Hanford Meteorological Monitoring Network.

Wet Deposition. The wet deposition flux, F_w, is calculated by using a scavenging ratio to model the wet removal of particles and gases in the atmosphere. The flux of material to the surface through wet deposition is the product of a scavenging ratio times the ambient air concentration, integrated through the vertical:

$$F_w(x, y) = \int_0^\infty \Lambda \, \chi(x, y, z) \, dz$$

where

F_w = wet deposition flux ($\mu g/m^2$-s)
Λ = scavenging ratio (S^{-1})
χ = concentration ($\mu g/m^3$)

The scavenging ratio is calculated by the following equation:

$$\Lambda = \lambda \times R$$

where

Λ = scavenging ratio (S^{-1})
λ = scavenging coefficient (hr/s-mm)
R = precipitation rate (mm/hr)

The scavenging coefficient appears to depend on a complex combination of the characteristics of the COPC/ROPC (e.g., solubility and reactivity for gases, size distribution for particles) as well as the nature of the precipitation (e.g., liquid or frozen). However, the ISCST3 User's Guide (EPA 1995a) reports that the best fit to the data is a function of particle size. Particle size estimates for emissions through HEPA filters will be used. The two HEPA filters are designed to trap 99.97% of all particulates $\geq$ 0.3 microns (as well as many smaller particles). It will be assumed that particle size is 1 micron. The user guides also provide a figure with the scavenging coefficient, which will be used as one of the wet deposition data inputs into the ISCST3 model. Following EPA guidance, the scavenging coefficient for gases will be based on the scavenging coefficient for the smallest category of particle sizes.

Deposition Rate Calculations. COPC/ROPC emissions can occur in either the vapor or particle phase, and COPCs/ROPCs in both phases are subject to wet and dry deposition utilizing the revised (May 1999) EPA ISCST3 model. Due to the limitations of the ISCST3 model, estimates of deposition rates for all COPCs/ROPCs cannot be made in a single computer run. Three ISCST3 runs must be made: one for vapor phase COPCs/ROPCs, one for particle phase COPCs/ROPCs, and one for particle-bound COPCs/ROPCs. The appropriate phase for each COPC/ROPC will be designated prior to the disposition modeling task.

Particle size is a primary influence on the calculation of both dry and wet deposition of COPCs/ROPCs in the particle phase, as discussed above. Therefore, distribution of particle sizes in the stack emissions at the RPP-WTP is an important input parameter in the ISCST3 model for determining deposition rates. The primary source for obtaining the particle size distributions of emissions will be consultation with the RPP-WTP design engineers and reference to published specifications from control equipment manufacturers. In addition, a review of data from other hazardous waste combustion facilities processes which have similar control equipment may be used to help refine particle size distribution estimates. A final source of particle size distribution data that will be used if necessary, is EPA's generalized particle size distribution for hazardous waste combustion facilities (EPA 1998a), or such information that can come from the HEPA filter literature; e.g., particles < 0.3 μm. It is now planned that a single particle size of $\leq$ one micron will be assumed to be representative for all particles released from the stacks because of the use of HEPA filtration.

6.1.2.5. Model Variable Settings

Model settings used to run the ISCST3 model will follow recommendations in the ISCST3 User's Guide (EPA 1995b), as well as guidance from EPA Region X and the Washington State Departments of Ecology and Health. The standard default regulatory options, unless other options are agreed to, as specified in the ISCST3 User's Guide will be used in the model execution.

Land use information is important to air dispersion modeling. The ISCST3 model has the option to use "rural dispersion coefficients" or "urban dispersion coefficients," depending on the land use categories within 3 km of the RPP-WTP. The Auer method specified in the *Guideline on Air Quality Models* (40 *CFR* Part 51, Appendix W) is used to define local land use for purposes of specifying the appropriate dispersion model coefficients in ISCST3. The land use category for the RPP-WTP PRA will be "rural."

6.1.2.6. Resuspension of Particulate Matter

Wind erosion may cause the resuspension of particulate matter into the atmosphere, which could result in increased inhalation exposure at the site. The methodology used to address this potential pathway was first described in *Rapid Assessment of Exposure to Particulate Emissions from Surface Contamination*

Sites, EPA Office of Health and Environmental Assessment, by C. Cowherd, Jr., G. Muleski, P. Englehart, and D. Gillette (1985, EPA/600/8-85/002) and further discussed in *Soil Screening Guidance: Technical Background Document,* EPA Office of Solid Waste and Emergency Response (1996, EPA/540/R95/128).

The technical approach to address the resuspension of particulate matter requires three steps: (1) the estimation of emissions from the site, (2) the estimation of ambient concentrations of resuspended particulate matter, and (3) the estimation of exposure. The estimation of exposure is addressed elsewhere in this work plan; the following outlines the technical approach used to conduct the first two steps.

1 The Estimation of Emissions Resulting from the Resuspension of Particulate Matter

The approach suggested by Cowherd et. al. (1985) assumes that the soil surface at a site is characterized by a large number ("unlimited reservoir") of erodible particles. These surfaces tend to have a relatively low wind speed threshold for wind erosion. Field experiments of erodible soils have shown that the magnitude of particulate emissions was most sensitive to wind speed and surface soil characteristics. The best fit to the experimental data was a function of the cube of the mean annual wind speed, divided by the threshold friction velocity, which is the wind velocity necessary to initiate soil erosion. The threshold friction velocity depends upon characteristics of the local soil, such as the presence of a surface crust, soil moisture content, size distribution of the exposed material, and the presence of nonerodible elements (e.g., vegetation, rocks).

Particulate emissions per square meter due to resuspension are calculated by determining a representative mean wind speed and threshold friction velocity for that site, following guidance provided in the two references cited above. Total emissions due to resuspension at the site are determined by multiplying the emissions per unit area by the total exposed area.

2 The Estimation of Ambient Particulate Concentrations Resulting from the Resuspension of Particulate Matter.

The particulate emissions rate will be used as input into the ISCST3 model to determine the dispersion of resuspended particulate. The areas with emissions will be treated as area sources in the model.

6.1.3. Model Output

The modeled output from ISCST3 will be combined from each source (stack), if available, at each exposure location, so total emissions from the RPP-WTP will be presented at each exposure location. Tables of the results will be provided for use in the risk assessment, both in tabular and electronic form. In addition, plots will be used to graphically represent the concentrations and deposition rates of emissions from the RPP-WTP.

6.1.3.1. Chronic Output

Chronic output from the RPP-WTP, to be evaluated in the risk assessment, will be based on the annual average ambient air concentrations and deposition rates for each COPC/ROPC at each exposure location, as calculated by the ISCST3 model. Five complete sets of annual average concentrations and deposition rates will be calculated, one for each of the five years of meteorological data used as input into the ISCST3 model. The worst-case year, with the highest concentration and deposition, will be used in the risk assessment.

6.1.3.2. Acute Output

The acute output from the RPP-WTP, to be evaluated in the risk assessment, will be based on the highest 1-hour average air concentrations as required by EPA guidance (EPA 1998a) for each COPC/ROPC at each exposure location, as calculated by the ISCST3 model. The use of 1-hour average air concentrations is to support the analysis of worst-case acute effects in the risk assessment. An acute inhalation scenario is recommended in EPA guidance because it is possible for air concentrations of COPCs/ROPCs to significantly exceed the annual average concentration for a brief time and, thus, result in acute effects to receptor populations via inhalation. Because the acute effects are only due to direct inhalation, deposition rates are not important in determining the acute risk. Concentrations in soil and other media reflect long-term deposition of COPCs/ROPCs. The long-term cumulative concentration in these media will be greater than the concentration resulting from any single acute event. Therefore, the acute exposure scenario is only applicable to the inhalation pathway.

The highest 1-hour average concentration will be calculated for the worst-case year. Acute emissions estimates include process upset and fugitive emissions in addition to normal stack emissions as described in section 4.2.1. Acute emission modeling does not include accidental (i.e., catastrophic) releases. Accident scenarios are included in the risk management program under the Clean Air Act.

6.1.3.3. Exposure Point Concentrations

Exposure point concentrations used for estimating dose of COPCs and ROPCs depends on the location of the receptor. The location of the various receptor populations identified for the quantitative risk assessment will correspond to the receptor grid nodes defined during air dispersion modeling (section 6.1.2.3). In keeping with the protective approach used in this risk assessment, the location with the maximum concentration of COPCs/ROPCs will be used in estimating exposure point concentrations. Because the point of maximum concentration may be different for airborne COPCs/ROPCs and COPCs/ROPCs deposited via wet and dry deposition mechanisms, EPA 1998a recommends the following method for selecting the point of maximum concentration. Within each area to be evaluated, receptor grid nodes or exposure locations with the highest modeled unitized value will be identified for each of the following parameters:

- Vapor phase air concentration
- Vapor phase wet deposition rate
- Particle phase air concentration
- Particle phase wet deposition rate
- Particle phase dry deposition rate
- Particle-bound phase air concentration
- Particle-bound phase wet deposition rate
- Particle-bound phase dry deposition rate

The parameters are chosen because COPCs/ROPCs occur in three phases: vapor phase, particle phase, and particle-bound phase. Vapor phase COPCs/ROPCs are transported in the atmosphere as gases, particle phase COPCs/ROPCs are transported as particles, and particle-bound COPCs/ROPCs are transported primarily with the COPC/ROPC condensed (or adsorbed) on the surface of particles.

If all eight of these maxima occur at the same receptor grid node, the receptor will be evaluated at this location. If all eight of these parameters are not at the maxima at the same location, a receptor will be evaluated at the location where each maximum occurs. This could lead to the evaluation of each receptor at eight different locations. However, it is anticipated that all eight parameters will be maximized at three

grid locations or less because the location of highest air concentration and deposition generally occur at the same receptor grid. In summary, if all eight parameters are not at their maxima at the same receptor grid node or exposure location, multiple exposure locations will be evaluated. A single receptor will not be evaluated for concentration maxima occurring at different locations.

6.2. Soil Accumulation Modeling

COPC/ROPC concentrations in soil will be calculated by summing the vapor phase and particle phase of COPCs/ROPCs to the soil (Figure 6-2). Both wet and dry deposition of particles and deposition of vapors will be considered. The calculation of soil concentration also incorporates a term that accounts for loss of COPCs/ROPCs by several mechanisms, including leaching, erosion, runoff, degradation (biotic and abiotic), radiological decay, and volatilization. Loss of COPCs/ROPCs from the soil will be modeled using rates that depend on the physical and chemical characteristics of the soil, such as pH, structure, organic matter content, and moisture content. Equations presented in this section are based on those presented in EPA 1998a. Values for the equation variables are presented in Appendix B-1.

Because soil concentrations may require many years to reach steady state, the equations used to calculate the average soil concentration over the period of deposition are derived by integrating the instantaneous soil concentration equation over the period of deposition. The following equations are the basis for calculating the cumulative soil concentration for carcinogenic COPCs and ROPCs:

For exposure durations that are less than the operating life of the RPP-WTP ($T_x \leq tD$)

$$Cs = \frac{Ds}{ks \times (tD - T_1)} \times \left[\left(tD + \frac{\exp(-ks \times tD)}{ks}\right) - \left(T_1 + \frac{\exp(-ks \times T_1)}{ks}\right)\right]$$

For exposure durations longer than the operating life of the RPP-WTP ($T_1 < tD < T_2$)

$$Cs = \frac{\left(\frac{Ds \times tD - Cs_{tD}}{ks}\right) + \left(\frac{Cs_{tD}}{ks}\right) \times \left(1 - \exp[-ks \times (T_2 - tD)]\right)}{(T_2 - T_1)}$$

The following equation is the basis for calculating the cumulative soil concentration for noncarcinogenic COPCs/ROPCs:

$$Cs = \frac{Ds \times [1 - \exp(-ks \times tD)]}{ks}$$

where

- Cs = Average soil concentration over exposure duration (mass COPC/kg soil)
- Ds = Deposition term (mass COPC/kg soil/yr)
- T_1 = Time period at the beginning of combustion (yr)
- ks = COPC soil loss constant due to all processes (1/yr)
- tD = Time period over which deposition occurs or time period of combustion (yr)
- Cs_{tD} = Soil concentration at time tD (mass COPC/kg soil)
- T_2 = Length of exposure duration (yr).

Calculating the COPC/ROPC soil loss constant (ks) includes consideration of several removal processes that may not occur simultaneously. The constant (ks) will be determined by using the soil's physical, chemical, radiological, and biological characteristics to consider the loss resulting from the following:

- Leaching
- Runoff
- Erosion
- Biotic and abiotic degradation
- Radiological decay
- Volatilization.

The following equation will be used to calculate the soil loss constant ks:

$$ks = ksg + kse + ksr + ksl + kdecay + ksv$$

where

ks	=	COPC/ROPC soil loss constant due to all processes (1/yr)
ksg	=	COPC/ROPC soil loss constant due to biotic and abiotic degradation (1/yr)
kse	=	COPC/ROPC soil loss constant due to soil erosion (1/yr)
ksr	=	COPC/ROPC soil loss constant due to surface runoff (1/yr)
ksl	=	COPC/ROPT soil loss constant due to leaching (1/yr)
kdecay	=	ROPC radiological decay constant (1/yr)
ksv	=	COPC/ROPC soil loss constant due to volatilization (1/yr).

The soil loss constants listed above will be compiled from available sources for each COPC and ROPC. In some cases, the soil loss constant must be calculated based on additional parameters. The equations that describe these processes are presented in Appendix B-2.

The soil loss processes will be considered in the calculation of soil concentrations. A number of soil loss parameters are dependent on the available water, calculated as ($P+I-RO-E_v$), which is related to precipitation (P), irrigation (I), surface runoff (RO), and evapotranspiration (E_v) in the Hanford Site area. Climate in the region results in greater evapotranspiration than precipitation. Therefore, water added to the system to irrigate crops would not generally be available to generate surface runoff or leachate. Because little water is available for soil erosion, surface runoff, and leaching, these processes will have a negligible effect on the concentration of COPCs/ROPCs in soils. Exclusion of these processes is consistent with the conservative screening level approach being used.

The COPC/ROPC loss constant due to volatilization (ksv) will be assumed to be equal to zero as recommended by EPA 1998a. Any volatile COPCs sorbed to particulates would volatilize prior to deposition onto soil. Soil loss constant due to biotic and abiotic degradation (ksg) is presented for a number of COPCs in EPA 1998a and repeated in Appendix B. Where no value is presented in EPA 1998a, additional sources will be consulted prior to release of the final Work Plan. If no value can be obtained for this parameter, it will conservatively be assumed to be equal to zero. See Appendix B-1 for soil loss values for COPCs and ROPCs.

As described previously, soil erosion, surface runoff, leaching, volatilization, and chemical (biotic and abiotic) degradation are not expected to contribute significantly to chemical loss from soil at this site. The primary loss process at this site is radiological decay.

The deposition term (Ds) for the preceding equations will be calculated from the results of the dispersion modeling using the following equation:

$$Ds = \left[\frac{100 \cdot Q}{Z_s \cdot BD}\right] \times \left[F_v \times (315.36 \times Vdv \times Cyv + Dywv) + (Dydp + Dywp) \times (1 - F_v)\right]$$

where

Ds	=	Deposition term (mg COPC/kg soil/yr)
100	=	Units conversion factor (mg-m^2/kg-cm^2)
Q	=	COPC/ROPC emission rate (g/s)
Z_s	=	Soil mixing zone depth (cm)
BD	=	Soil bulk density (g soil/cm^3 soil),
F_v	=	Fraction of COPC air concentration in vapor phase (unitless)
315.36	=	Units conversion factor (m-g-s/cm-mg-yr)
Vdv	=	Dry deposition velocity (cm/s)
Cyv	=	Unitized yearly average air concentration from vapor phase (mg-s/g-m^3)
Dywv	=	Unitized yearly average wet deposition from vapor phase (s/m^2-yr)
Dydp	=	Unitized yearly average dry deposition from particle phase (s/m^2-yr)
Dywp	=	Unitized yearly average wet deposition from particle phase (s/m^2-yr).

In the equations presented above, a number of parameters require additional calculations, which are presented in Appendix B-2. The dry deposition velocity (Vdv) and COPC/ROPC emission rate (Q) will be provided by the air dispersion modeling. Site-specific factors such as soil mixing depth, soil bulk density, available water, and volumetric water content will have values as described below.

The soil mixing depth typically used in the calculations are:

- 1 cm for untilled soil
- 20 cm for tilled soil

These values are consistent with those presented in EPA 1998a. For this risk assessment, the soil mixing depth will conservatively be assumed to be 1 cm.

The soil dry bulk density (SD) used in the calculations will be the value of 1.5 g/cm^3 as recommended in EPA 1998a. Alternatively, if a Hanford Site-specific value can be determined using ASTM Method D2937 (ASTM 1994) and Method D2215 (ASTM 1998) by May 2000, the site-specific value will be used.

The values of the soil volumetric water content (θ_{sw}) will be 0.2 mL/cm^3 recommended by EPA 1998a. Alternatively, if a site-specific value can be determined by May 2000, the site-specific value will be used.

6.3. Terrestrial Plant and Animal Modeling

The modeled concentrations of contaminants in plants consumed by both humans and non-human receptors will be convergent. This is based upon previous stakeholder and tribal nations requests. This is not to say that receptors that ingest whole organisms will be exposed to the same contaminant concentration as receptors that eat only particular organs or tissues of the organism. Whole-body tissue concentrations will be calculated in the human health and ecological risk assessments using identical bioaccumulation factors, ingestion rates, and other parameters, and identical model equations. The

ecological risk assessment will use modeled whole-body concentrations to estimate doses to wildlife receptors.

6.3.1. Plants

The discussion in this section applies to plants for human consumption such as vegetables, as well as plants for animal consumption such as forage (including native plants consumed by browsing animals) and silage plants.

COPC and ROPC concentrations in plants will be estimated for aboveground produce as well as belowground produce. Aboveground produce will be exposed to particulate deposition and vapor phase contamination, as well as root uptake from soil and subsequent transfer to aboveground foliage (Figure 6-3). Aboveground plant parts are categorized as protected (i.e., the plant structure prevents accumulation of contaminants through the deposition and air-to-plant pathways) and unprotected. For example, corn kernels are protected by the husk. Protected plant parts will be limited to grain used as animal feed in this evaluation. All other plant parts for human and animal consumption will be considered unprotected; i.e., not physically shielded from deposition. Belowground produce will only be exposed to contaminants from root uptake.

Concentrations of COPCs/ROPCs in plants will be estimated using the equations presented below as recommended by EPA 1998a. Values for the chemical-specific parameters are presented in Appendix B-1.

6.3.1.1. Aboveground Plants/Direct Deposition

Concentrations of COPCs/ROPCs in aboveground produce due to direct deposition will be calculated using the following equation:

$$Pd = \frac{1{,}000 \times Q \times (1 - F_v) \times [Dydp + (Fw \times Dywp)] \times Rp \times [1 - \exp(-kp \times Tp)]}{Yp \times kp}$$

where

- Pd = Aboveground produce concentration due to wet and dry direct deposition (mass COPC/kg DW)
- 1,000 = Units conversion factor (mg/g)
- Q = COPC/ROPC emission rate (g/s)
- F_v = Fraction of COPC/ROPC in air concentration in vapor phase (unitless)
- Dydp = Unitized yearly average dry deposition from particle phase (s/m^2-yr)
- Fw = Fraction of COPC/ROPC wet deposition that adheres to plant surfaces (unitless)
- Dywp = Unitized yearly wet deposition from particle phase (s/m^2-yr)
- Rp = Interception fraction of the edible portion of plant (unitless)
- kp = Plant surface loss coefficient (1/yr)
- Tp = Length of plant exposure to deposition per harvest or growing season of the edible portion of the ith plant group (yr)
- Yp = Yield or standing crop biomass of the edible portion of the plant (kg DW/m^2).

Parameters representing the interception fraction (Rp), plant surface loss (kp), length of plant exposure (Tp), and crop yield (Yp) may require additional calculations. Values that will be used for each of these parameters are described below and are shown in Table 6-1.

Interception fraction (Rp) depends on the type and productivity of the plant being modeled. EPA 1998a reports interception fractions ranging from 0.053 for exposed fruits to 0.982 for exposed vegetables. The Rp is set to a value of 0.39 as recommended by EPA 1998a for fruits and vegetables consumed by human receptors. This value represents a weighted average Rp based on the relative ingestion rates of fruits and vegetables. Interception fractions of 0.499 for forage (including wild plants consumed by browsing animals) and 0.459 for silage were calculated from the correlation between Rp and productivity provided in EPA 1998a using the parameters below:

	Rp Interception Fraction	γ Constant	Yp Standing Biomass
Forage	0.499	2.88	0.24
Silage	0.459	0.769	0.8

These Rp values are used for estimating deposition onto forage and silage plants.

For chemical COPCs, EPA 1998a recommended value for the surface loss coefficient of 18 (1/yr) will be used in the calculations for chemicals. For radionuclides the surface loss coefficient will be calculated using the following equation:

$$kp = \left(\frac{\ln 2}{t_{1/2}}\right) \times CF$$

where

kp = Plant surface loss coefficient (1/yr)
$t_{1/2}$ = Half-life (days)
CF = Conversion factor of 365 (days/yr)

The value used for the length of edible plant exposure to deposition per harvest (Tp) will be 0.164 yr (60 days) as recommended by EPA 1998a. The EPA 1998a recommended value of 2.24-kg DW/m^2 for the standing crop biomass (Yp) will be used. Alternatively, if a Hanford site-specific length value for edible plant exposure to deposition can be determined by May 2000, the site-specific value will be used.

6.3.1.2. Aboveground Plants/Air-to-Plant Transfer

The concentration of COPCs/ROPCs in aboveground plants due to air-to-plant transfer will be calculated based on the following equation:

$$Pv = Q \times F_v \times \frac{Cyv \times Bv_{ag} \times VG_{ag}}{\rho_a}$$

where

Pv = Concentration of COPC/ROPC in the plant resulting from air-to-plant transfer (μg COPC/g DW)
Q = COPC/ROPC emission rate (g/s)
F_v = Fraction of COPC/ROPC air concentration in vapor phase (unitless)
Cyv = Unitized yearly average air concentration from vapor phase (μg-s/g-m^3)

Bv_{ag} = COPC/ROPC air-to-plant biotransfer factor [(mg COPC/g DW plant)/(mg COPC/g air)]
VG_{ag} = Empirical correction factor for aboveground produce (unitless)
ρ_a = Density of air (g/m^3).

6.3.1.3. Root Uptake

The concentration of COPCs/ROPCs in plants due to uptake from the soil will be calculated for aboveground and belowground plants. Concentrations in aboveground plants will be calculated using the following equation:

$$Pr = Cs \times Br$$

where

Pr = Concentration of COPC\ROPC in produce due to root uptake (mg/kg)
Cs = Average soil concentration over exposure duration (mg COPC/kg soil)
Br = Plant-soil bioconcentration factor for produce (unitless) from EPA 1998a and repeated in Appendix B

For concentrations in belowground plants the following equation will be used:

$$Pr = \frac{Cs \times RCF \times VG_{rootveg}}{Kd_s \times 1}$$

where

Pr = Concentration of COPC/ROPC in produce due to root uptake (mg/kg)
Cs = Average soil concentration over exposure duration (mg COPC/kg soil)
RCF = Root concentration to soil water concentration factor (unitless)
$VG_{rootveg}$ = Empirical correction factor for belowground produce (unitless)
Kd_s = Soil-water partition coefficient (L/kg)
1 = Conversion factor (kg/L)

6.3.2. Animal Tissue, Eggs, and Milk

The concentration of chemical and radiological COPCs/ROPCs in animal tissue (including beef, pork, poultry, and game animals), eggs, and milk products are estimated based on the amount of COPCs/ROPCs that each animal is assumed to consume through its diet from the following sources:

- Forage (primarily pasture grass and hay for domestic animals wild plants for game animals)
- Silage (forage that has been stored and fermented)
- Grain
- Incidental soil ingestion associated with grazing and browsing

The feed items are assumed to be contaminated through the following pathways:

- Direct deposition of particles – wet and dry deposition of particle phase COPCs/ROPCs onto forage, silage, and soil

- Vapor transfer – uptake of vapor phase COPCs/ROPCs in forage and silage

- Root uptake – root uptake of COPCs/ROPCs deposited in soil and subsequent transfer to aboveground plant portions of forage, silage, and grain

Figure 6-4, Figure 6-5 and Figure 6-6 present conceptual models of the sources and pathways that result in animal product contamination.

Concentrations of chemical and radiological COPCs/ROPCs in beef, pork, poultry, eggs, and milk will be estimated using the equations presented below as recommended by EPA 1998a. Values for the chemical-specific parameters are presented in Appendix B-1. Concentrations of COPCs/ROPCs in game animals will be estimated using the equations presented in section 8.2.2.1.

The concentration of COPCs in animal products, except wild game, will be estimated using the following equation:

$$A_j = \left(\sum\left(F_i \times Qp_i \times P_i\right) + Qs \times Cs \times Bs\right) \times Ba_j \times MF$$

where

A_j = Concentration of COPC/ROPC in animal product *j* (mass COPC/kg FW tissue)
F_i = Fraction of plant type *i* grown on contaminated soil and ingested by the animal (unitless)
Qp_i = Quantity of plant type *i* eaten by the animal per day (kg DW plant/day)
P_i = Concentration of COPC/ROPC in each plant type *i* eaten by the animal (mg/kg DW)
Qs = Quantity of soil eaten by the animal each day (kg/day)
Cs = Average soil concentration over exposure duration (mass COPC/kg soil)
Bs = Soil bioavailability factor (unitless)
Ba_j = COPC/ROPC biotransfer factor for animal product *j* (day/kg FW tissue)
MF = Metabolism factor (unitless).

The fraction of each plant type grown on contaminated soil and ingested by the animal (F_i) will be 1.0 (100%), as recommended by EPA 1998a. Values for P_i and *Cs* in the above equation will be calculated, using appropriate equations as previously discussed. Biotransfer factors (Ba) from EPA 1998a will be used. Values for Qp, Qs, Bs, and MF are shown in Table 6-2; these values are from EPA 1998a.

Animal products include tissue (beef, pork, poultry, game), dairy products, and eggs. The biotransfer factors used to estimate COPC/ROPC concentrations in meat (fat and muscle) generally do not include transfer to other organ tissues. Chemical-specific biotransfer factors are provided by EPA 1998a. Other parameters are summarized in Table 6-2.

The metabolism factor (MF) represents the estimated amount of COPC that remains in fat and muscle tissue. The MF only affects the concentration of bis(2-ethylhexyl)phthalate since evidence indicates this chemical is more readily metabolized and excreted by mammals than other COPCs. The MF does not apply to poultry or eggs.

6.4. Surface Water and Fish Bioaccumulation Modeling

Estimating the intake of COPCs/ROPCs due to bioaccumulation in fish from surface water contaminants will involve several steps (Figure 6-7). First, the concentration of COPCs/ROPCs in surface water is calculated based on the following:

- Direct deposition into surface water
- Runoff from impervious surfaces within the watershed
- Runoff from pervious surfaces within the watershed
- Soil erosion from the watershed
- Direct diffusion of vapor phase COPCs/ROPCs into the surface water
- Chemical, biological, or radiological transformation of compounds within the surface water body

COPCs/ROPCs in surface water and fish will be estimated using the equations presented below as recommended by EPA 1998a. Values for the chemical-specific parameters are presented in Appendix B-1. It should be noted that the Hanford Surface Environmental Surveillance Program collects and analyzes fish tissues in the Hanford Reach of the Columbia River. However, the SLRA will be conducted prior to release of emissions from the RPP-WTP; therefore, the data collected could not represent contamination contributed for the RPP-WTP and cannot be used to calibrate the fish model.

Finally, ROPCs will be evaluated using similar equations to those presented for COPCs.

The equation used to quantify the total COPC/ROPC load to the surface water body is as follows:

$$L_T = L_{dep} + L_{dif} + L_{ri} + L_{rp} + L_e + L_{transform}$$

where

L_T	=	Total COPC/ROPC load to the water body, including deposition, runoff, and erosion (g/yr)
L_{dep}	=	Total wet and dry particulate deposition and wet vapor deposition of COPC directly to water body (g/yr)
L_{dif}	=	Vapor phase COPC diffusion (dry deposition) load to water body (g/yr)
L_{ri}	=	Runoff load from impervious surfaces (g/yr)
L_{rp}	=	Runoff load from pervious surfaces (g/yr)
L_e	=	Load due to soil erosion (g/yr)
$L_{transform}$	=	Load due to biological, chemical, and/or radiological transformation within the water body (g/yr).

As discussed previously in section 6.2, evapotranspiration exceeds precipitation in the Hanford Site area, resulting in insufficient water available to cause significant erosion or runoff of COPCs/ROPCs. Therefore, the terms for runoff from pervious (L_{rp}), impervious surfaces (L_{ri}), and erosion (L_e) will not be quantified in the above equation. It should be noted that exclusion of these minor processes from the surface water calculations does not result in loss of contaminated material. This material is accounted for in the soil calculations where it is assumed no chemicals are lost due to erosion or runoff. Due to limited data and uncertainty associated with the chemical or biological internal transfer of compounds into daughter products, $L_{transform}$ will be assigned the default value of zero, as recommended by EPA 1998a. Therefore, the only terms quantified are L_{dep} and L_{dif}.

Total wet and dry particulate deposition and wet vapor deposition of COPCs/ROPCs directly to the waterbody (L_{dep}) will be calculated with the following equation:

$$L_{dep} = Q \times [F_v \times Dywwv + (1 - F_v) \times Dytwp] \times A_w$$

where

L_{dep} = Total (wet and dry) particle phase and wet vapor phase COPC/ROPC direct deposition load to water body (g/yr)
Q = COPC/ROPC emission rate (g/s)
F_v = Fraction of COPC/ROPC air concentration in vapor phase (unitless)
Dywwv = Unitized yearly (water body) average wet deposition from vapor phase (s/m^2-yr)
Dytwp = Unitized yearly (water body) average total (wet and dry) deposition from particle phase (s/m^2-yr)
A_w = Water body surface area (m^2).

Emission rates are modeled (see section 4.2); F_v is chemical-specific; A_w is site-specific; and Dywwv and Dytwp are modeled air parameters.

The vapor phase COPC/ROPC diffusion (dry deposition) load to the water body (L_{dif}) will be calculated with the following equation:

$$L_{dif} = \frac{K_v \times Q \times F_v \times Cywv \times A_w \times CF}{\frac{H}{R \times T_{wk}}}$$

where

L_{dif} = Vapor phase COPC/ROPC diffusion (dry deposition) load to water body (g/yr)
K_v = Overall COPC/ROPC transfer coefficient (m/yr)
Q = COPC/ROPC emission rate (g/s)
F_v = Fraction of COPC/ROPC air concentration in vapor phase (unitless)
Cywv = Unitized yearly (water body) average air concentration from vapor phase (μg-s/g-m^3)
A_w = Water body surface area (m^2)
CF = Conversion factor of 1×10^{-6} (g/μg)
H = Henry's Law constant (atm-m^3/mol)
R = Universal gas constant (atm-m^3/mol-K)
T_{wk} = Water body temperature (K).

Emission rates (Q) and Cywv are modeled values; A_w is site-specific; F_v, H, and R are chemical-specific values; and EPA 1998a recommends a value of 298°K for T_{wk}.

Total water body COPC/ROPC concentration (C_{wtot}) will be calculated with the following equation:

$$C_{wtot} = \frac{L_T}{Vf_x \times f_{wc} + k_{wt} \times A_W \times (d_{wc} + d_{bs})}$$

where

C_{wtot} = Total water body of COPC/ROPC concentration in g COPC/m^3 water body
L_T = Total COPC/ROPC load to the water body (g/yr)
Vf_x = Average volumetric flow rate through the water body (m^3/yr)
f_{wc} = Fraction of total water body COPC/ROPC concentration in the water column (unitless)
k_{wt} = Overall total water body COPC/ROPC dissipation rate constant (1/yr)
A_W = Water body surface area (m^2)

d_{wc} = Depth of water column (m)
d_{bs} = Depth of upper benthic sediment layer (m).

The total load to the water body (L_T) is calculated (see previous equation); and Vf_x, A_W, and d_{wc} are site-specific; d_{bs} will be 0.03, as recommended by EPA 1998a; equations for f_{wc} and k_{wt} appear in Appendix B-2.

Total COPC/ROPC concentration in the water column (C_{wctot}) will be calculated with the following equation:

$$C_{wctot} = f_{wc} \times C_{wtot} \times \frac{d_{wc} + d_{bs}}{d_{wc}}$$

where

C_{wctot} = Total COPC/ROPC concentration in water column (mg COPC/L water column)
f_{wc} = Fraction of total water body COPC/ROPC concentration, in the water column (unitless)
C_{wtot} = Total water body COPC/ROPC concentration including water column and bed sediment (mg COPC/L water body)
d_{wc} = Depth of water column (m)
d_{bs} = Depth of upper benthic sediment layer (m).

Equations for f_{wc} are found in Appendix B-2; C_{wtot} is calculated (see previous equation); d_{wc} is site-specific; and d_{bs} is 0.03 as recommended by EPA 1998a.

The dissolved phase COPC/ROPC water concentration (C_{dw}) will be calculated with the following equation:

$$C_{dw} = \frac{C_{wctot}}{1 + Kd_{sw} \times TSS \times CF}$$

where

C_{dw} = Dissolved phase water concentration (mg COPC/L water)
C_{wctot} = Total COPC/ROPC concentration in water column (mg COPC/Lwater column)
Kd_{sw} = Suspended sediment/surface water partition coefficient (L water/kg suspended sediment)
TSS = Total suspended sediment concentration (mg/L)
CF = Conversion factor of 1×10^{-6} (kg/mg).

C_{wctot} is calculated (see previous equation); Kd_{sw} is chemical-specific; and TSS is site-specific (see Appendix B-2 if measured data are not available).

The COPC/ROPC concentration sorbed to bed sediment (C_{sb}) will be calculated with the following equation:

$$C_{sb} = f_{bs} \times C_{wtot} \times \left(\frac{Kd_{bs}}{\theta_{bs} + Kd_{bs} \times C_{BS}} \right) \times \left(\frac{d_{wc} + d_{bs}}{d_{bs}} \right)$$

where

C_{sb} = COPC/ROPC concentration sorbed to bed sediment (mg COPC/kg sediment)
f_{bs} = Fraction of total water body COPC/ROPC concentration in benthic sediment (unitless)
C_{wtot} = Total water body COPC/ROPC concentration, including water column and bed sediment (mg COPC/L water body)
Kd_{bs} = Bed sediment/sediment pore water partition coefficient (L COPC/kg water body)
θ_{bs} = Bed sediment porosity (Lpore water/Lsediment)
C_{BS} = Bed sediment concentration (g/cm^3)
d_{wc} = Depth of water column (m)
d_{bs} = Depth of upper benthic layer.

The equation for f_{bs} and θ_{bs} are found in Appendix B-2; C_{wtot} is calculated (see previous equation); Kd_{bs} is chemical-specific; C_{BS} and d_{wc} are site-specific; and d_{bs} is 0.3, as recommended by EPA 1998a.

After calculating COPC/ROPC concentrations in the water column and benthic sediments, the concentration in fish tissue will be calculated. The fish concentration due to dissolved phase water concentrations will be calculated using bioconcentration factors (BCF) for COPCs with a log K_{ow} less than 4:

$$C_{fish} = C_{dw} \times BCF_{fish}$$

where

C_{fish} = Concentration of COPC in fish (mg COPC/kg FW tissue)
C_{dw} = Dissolved phase water concentration (mg COPC/L)
BCF_{fish} = Bioconcentration factor for COPC in fish (L/kg).

With the exception of extremely hydrophobic compounds such as dioxins, furans, and PCBs, fish tissue concentrations of COPCs with log K_{ow} greater than 4 will be calculated using bioaccumulation factors (BAF):

$$C_{fish} = C_{dw} \times BAF_{fish}$$

where

C_{fish} = Concentration of COPC in fish (mg COPC/kg FW tissue)
C_{dw} = Dissolved phase water concentration (mg COPC/L)
BAF_{fish} = Bioaccumulation factor for COPC in fish (L/kg FW tissue).

Extremely hydrophobic compounds such as dioxins, furans, and PCBs are expected to be sorbed to bed sediments but not suspended particulate and will bioaccumulate from the benthic sediment layer into fish tissue. The following equation will be used to estimate fish tissue concentrations (C_{fish}) for hydrophobic compounds:

$$C_{fish} = \frac{C_{sb} \times f_{lipid} \times BSAF}{OC_{sed}}$$

where

C_{fish} = Concentration of COPC in fish (mg COPC/kg FW tissue)
C_{sb} = Concentration of COPC sorbed to bed sediment (mg COPC/kg bed sediment)
f_{lipid} = Fish lipid content (unitless)
BSAF = Biota-to-sediment accumulation factors (kg sediment organic carbon/kg fish lipid) (unitless)
OC_{sed} = Fraction of organic carbon in bottom sediment (unitless)

Although there is no explicit guidance for ERAs or how to estimate fish-tissue concentrations from water concentrations (EPA 1992a, 1997c, 1998a; Wentzel et al. 1994), whole-body concentrations in fish also may be calculated using total water-column concentrations (C_{wctot}) and BCFs because fish are exposed to contaminants associated with suspended particulates, as well as dissolved contaminants. The use of whole water concentrations can be justified on procedural and technical grounds. Procedurally, screening-level risk assessments should be conservative. Technically, some BCFs are based on exposures to whole-water contaminant levels. Although dissolved contaminant is more available for uptake by fish than particle-bound contaminant, some uptake from ingestion of contaminated particulates is likely. Furthermore, some published BCFs are likely derived from field data. Field data for fish-tissue COPC concentrations include the transfer from ingested particles to fish. Therefore, calculating fish-tissue concentrations by multiplying the BCF by the whole water concentration gives a more accurate estimate in some cases, as well as a conservative estimate in all cases, as is appropriate to a screening-level risk assessment. Calculating fish tissue concentrations for fish consumed by ecological receptors is presented in section 8.2.3 and the BAFs and BCFs are discussed in section 8.2.4.

6.5. Uncertainties

Uncertainties will be presented in the risk assessment for each aspect of the environmental modeling. This includes air dispersion modeling, soil accumulation modeling, biota modeling, and surface water and fish bioaccumulation modeling. The uncertainty assessment will be presented in the form of both text and tables summarizing the primary contributors and potential magnitude of uncertainties.

A variety of conservative assumptions are used throughout the modeling process to compensate for uncertainties. One limitation of the air dispersion and deposition modeling will result in the overestimation of COPC/ROPC concentrations in all media. This is the double counting of COPCs/ROPCs in air and deposited on soil, plants, and surface water. Air dispersion and deposition components of the modeling are conducted separately. Therefore, when estimating ambient air concentrations for inhalation exposure, the model assumes no loss due to deposition. When estimating deposition, the deposition rate at each point on the receptor grid assumes no loss of COPCs/ROPCs in the air due to deposition at any other grid or receptor location.

6.5.1. Uncertainty in Air Dispersion Modeling

A number of sources of uncertainty exist in the air dispersion modeling, such as:

- input values, such as stack emission characteristics
- meteorological data

Some of these uncertainties are based on the limited data available, such as estimates of the emissions data. Other uncertainties become larger when the model is used at the limits of its normal application, for instance, in very complex terrain or at distances near 50 km.

Uncertainties in the application of the model are reduced when the model is used to calculate longer-term averages (annual averages), and when several years of meteorological data are available to use in the model.

6.5.2. Uncertainty in Soil Accumulation Modeling

Estimating soil concentrations incorporates numerous uncertainties, which are inherent in the assumptions that are the basis for the calculations. Examples of uncertainty in the parameters would be soil mixing depth, soil bulk density, and volumetric water content, which are assigned a single value, but may vary widely over a relatively small area. The concentration of COPCs/ROPCs in soil will be subject to loss due to biotic and abiotic degradation; however, transformation and subsequent increase of secondary COPCs are not considered in the assessment. Transformation of ROPCs and formation of daughter products is included in this assessment through the use of toxicity values that include daughter products.

6.5.3. Uncertainty in Terrestrial Plant and Animal Modeling

Calculation of COPC/ROPC concentration in biota incorporates the uncertainties inherent in calculation of soil concentrations because the soil is one of the sources of COPCs/ROPCs in biota. Uncertainties also arise from the assumption that the location of maximum soil concentration is the location of exposure to biota over a multiple-year period. Additionally, although COPCs/ROPCs are incorporated into plants and animal tissue, it is assumed that the COPC/ROPC concentration in soil does not decrease due to these processes. Assumptions of the animal feed ingestion rates introduces additional uncertainty because they are based on average rates.

Additional pathways such as fugitive dust emission or entrainment of soil in rainwater splash may contribute to COPC/ROPC concentrations in biota. However, no equations are available to quantify these pathways, and this contribution is considered negligible compared to direct deposition and uptake. In addition, the chemical transport through inedible portions of plants (e.g., corn husks) may contribute to COPC/ROPC concentrations in edible portions of plants (e.g., corn kernel). Transfer factors for this type of chemical transport are not available; however, if transfer does occur, it is likely to be negligible.

6.5.4. Uncertainty in Surface Water and Fish Modeling

Uncertainty in the estimation of surface water and fish concentrations of COPCs/ROPCs results from the assumptions used in the calculations. The equations assume the water body reaches a steady-state condition; however, for application to the Columbia River and any other surface water, this is extremely conservative. Additionally, many of the equations used to model the fate of COPCs/ROPCs deposited into the water body greatly simplify the mechanisms occurring within such a dynamic system and may overestimate or underestimate the concentration of COPCs/ROPCs in the surface water. It is also assumed that the maximum deposition of COPCs/ROPCs occurs over the entire depositional area of the water body which is a source of additional uncertainty and conservatism.

The calculation of fish tissue concentrations incorporates the uncertainty inherent in the calculation of surface water concentrations, as well as the following sources. It is assumed that steady-state bioconcentration factors (BCF) for transfer of COPCs/ROPCs from contaminated surface water to fish tissue apply to the Columbia River and any other surface water. Additionally, generic BCF values will be used in the calculations that do not account for variability among specific fish species associated with specific surface water systems. There is uncertainty in assigning COPCs exclusively to either water column or bed sediment for purposes of estimating fish-tissue concentrations as described in the EPA guidance for human health (EPA 1998a). The problem is that this approach to partitioning COPCs in the

aquatic environment may not reflect the multiple pathways by which different kinds of fish are potentially exposed to any given contaminant. The EPA approach estimates fish-tissue concentrations using BCFs and dissolved water concentrations for COPCs with low K_{ow}s; BAFs and whole-water concentrations for COPCs with moderate K_{ow}s; and BAFs and bed-sediment concentrations for COPCs with high K_{ow}s. The uncertainty lies in the source and meaning of the different biotransfer factors used for the different media.

If the EPA (1998a) biotransfer factors do not incorporate all the pathways to all fish from the single medium where each COPC is assumed to predominate, then the exposure will be underestimated. It is unlikely that the EPA transfer factors account for all pathways relevant to all fish. Fish take up contaminants into their tissue via the water in contact with their gills and via the ingestion of water, abiotic particulates, and biota. Some fish will be exposed by primarily one pathway, while others will be exposed over multiple pathways. Dissolved contaminants are primarily taken up across the gill membrane, thus, all fish living in the water column will be exposed to dissolved contaminants. Filter-feeding fish, which usually live in the water column, will also be exposed to contaminants bound to suspended particulates that they filter out of the water and ingest. Sediment-ingesting fish that live in the water column will also be exposed to sediment contaminants by direct ingestion. Predatory fish, which are also water-column dwellers, will also be exposed to dissolved, particulate-bound, and sediment contaminants by ingesting prey that were so exposed, as well as by direct uptake from the water column and ingestion of suspended particulates. In fact, there are probably few fish that are exposed to only dissolved contaminants; perhaps only those that live in the water column and selectively feed on planktonic animals to the exclusion of abiotic particulates. Therefore, assigning each contaminant to a particular class of media (dissolved water, whole water, and bed sediment), based on its tendency to adsorb to particles or organic carbon, potentially neglects pathways from other media. Further discussion of uncertainty related to these pathways is presented in the ecological section.

6.6. Summary for Environmental Modeling

Air dispersion modeling will be used to determine COPC/ROPC-specific concentrations and deposition rates resulting from emissions of the RPP-WTP. The assessment area will extend out to 50 km. These results will be used as input into the human health and ecological risk assessments.

The ISCST3 will be used for the air quality modeling task. The RPP-WTP emissions, as determined by the design engineers, and 5 years of Hanford Site meteorological data collected by the Hanford Site Meteorological Station will be used as input into the model. COPC/ROPC-specific concentrations and deposition rates will be calculated at a gridded network of receptors and at specific sensitive receptors identified by the risk assessment analysts.

Fate and transport modeling will be used to estimate COPC/ROPC concentrations in various exposure media (air, surface water, fish tissue, soil, animal tissue, eggs, milk, and plants). This modeling effort will utilize assumed emission rates with a combination of default and site-specific parameters to describe the movement of COPCs/ROPCs through the environment. This modeling is predictive and cannot be confirmed by sampling environmental media since the emissions source does not yet exist. The high level of uncertainty associated with this predictive modeling is addressed through the use of conservative assumptions whenever possible. Estimated media concentrations resulting from this modeling effort will be used in the exposure assessment for the Human Health (Chapter 7) and Ecological (Chapter 8) risk assessments. Environmental modeling will be the same for the PRA and FRA with the possible inclusion of additional site-specific modeling parameters in the FRA.

Figure 6-1. Exposure Grids in Each concentric Zone



Hanford ERA - 004a

Figure 6-2. Sources of COPC and ROPC Concentrations in Soil



(a) The only loss mechanism to be quantified for this conservative screening level analysis is radioactive decay.

(b) Production of daughter products will be evaluated through the use of toxicity values (+D) that include all potential daughter products at equalibrium with the parent radionuclide.

Hanford TWRS-P: 08

Figure 6-3. Sources of Constituents of Potential Concern Concentration in Plants



Loss mechanisms not included in the PRA

(a) The conservative screening level risk assessment does not account for loss between harvest and consumption

Hanford ERA - 005

Figure 6-4. Generic Sources of Constituents of Potential Concern Concentrations in Animal Tissue, Eggs, and Dairy Products



Figure 6-5. Sources of Constituents of Potential Concern Concentrations in Tissue, Eggs, and Dairy Products of Domestic Farm Animals



Figure 6-6. Sources of Constituents of Potential Concern Concentrations in Tissue of Game Animals



Figure 6-7. Sources of Constituents of Potential Concern Concentrations in Fish



(a) The only loss mechanism to be quantified for this conservative screening level analysis is radioactive decay.

Hanford ERA - 009

Table 6-1. Specific Variables for Plant Uptake Modeling[1]

Parameter	Plant Type	Value	Units
Interception Fraction (Rp)	Fruits	0.053	—
	Vegetables	0.932	—
	Fruits and Vegetables (weighted average)	0.39	—
	Forage	0.5	—
	Silage	0.46	—
Plant Surface Loss Coefficient (kp)	All (default)	18.0	1/yr
	All (calculated)	Applies to radionuclides	1/yr
Length of Plant Exposure (Tp)	Fruits and Vegetables	0.164	yr
	Forage	0.12	yr
	Silage	0.16	yr
Standing Crop Biomass (Yp)	Fruits	0.25	(kg DW/m^2)
	Vegetables	5.66	(kg DW/m^2)
	Fruits and Vegetables (weighted average)	2.24	(kg DW/m^2)
	Forage	0.24	(kg DW/m^2)
	Silage	0.8	(kg DW/m^2)
Empirical Correction Factor (VG_{ug})	Produce Log K_{ow} >4	0.01	—
	Produce Log K_{ow} <4	1.0	—
	Forage	1.0	—
	Silage	0.5	—

[1] All values from *Human Health Risk Assessment Protocol for Hazardous Waste Combustion Facilities* EPA 1998a. See document for justification of values. All values presented are default values; for radionuclides, plant surface loss (kp) includes radiological decay.

— = unitless

Table 6-2. Specific Variables of Uptake into Animal Products[a]

Parameter	Beef Cattle	Pork	Poultry	Dairy Cattle	Deer
Forage Ingestion Rate (kg DW/day) (Qp forage)	8.8	0.0	0.0	13.2	1.463[b]
Silage Ingestion Rate (kg DW/day) (Qp silage)	2.5	1.4	0.0	4.1	0.0
Grain Ingestion Rate (kg DW/day) (Qp grain)	0.47	3.3	0.2	3.0	0.0
Soil Ingestion Rate (kg/day) (Qs)	0.5	0.37	0.022	0.4	0.0
Soil Bioavailability (unitless) (Bs)	1.0	1.0	1.0	1.0	1.0
Metabolism Factor for Bis(2-ethylhexyl)phthalate (MF)	0.01	0.01	NA	0.01	0.01
Metabolism Factor all other COPCs (MF)	1.0	1.0	NA	1.0	1.0

[a] Parameter values are from *Human Health Risk Assessment Protocol for Hazardous Waste Combustion Facilities* (EPA 1998a), except where noted. See document for range and justification of values.

[b] Forage ingestion rate for deer is from Anderson, et al. 1974 and Allredge, et al. 1974, both presented in Higley and Kuperman 1996. Value presented has been calculated based on the body weight of 66.5 kg and an ingestion rate of 0.022 kg/kg BW/day.

NA = not applicable

7. Screening Human Health Risk Assessment

This section presents four fundamental components of the human health risk assessment (HHRA) process: (1) data evaluation (2) exposure assessment, (3) toxicity assessment, and (4) risk characterization. The data evaluation step includes the selection of COPCs/ROPCs (section 4) and the quantification of emissions (section 5). Each of the remaining three components is discussed below.

7.1. Exposure Assessment

Exposure assessment is the process of estimating the magnitude, frequency, duration, and type of potential exposures to COPCs/ROPCs. This section presents the exposure scenarios and approach for conducting the quantitative exposure assessment.

A human health conceptual exposure model has been developed that identifies potential receptors and potentially complete exposure pathways (i.e., exposure scenarios). The conceptual exposure model is shown as Figure 7-1. The end product of the conceptual model is identifying exposure scenarios that are defined by potentially exposed populations and exposure pathways. The conceptual model was developed from information obtained from and the *Screening Assessment and Requirements for a Comprehensive Assessment: Columbia River Comprehensive Impact Assessment* (DOE-RL 1998). The Native American scenario is taken from Harris and Harper 1997. The specific scenario contained in Harris and Harper 1997 represents a scenario developed for the Confederated Tribes of the Umatilla Indian Reservation. Exposure scenarios for other tribes may be different.

The conceptual exposure model focuses on identifying complete exposure pathways for potentially exposed populations. An exposure pathway is the means through which an individual may come in contact with a chemical in the environment. Exposure pathways are determined by environmental conditions (e.g., location of groundwater, vegetative cover, wind speed/direction), the potential for chemical migration among media (e.g., soil, surface water, or air), and by the lifestyles and work activities of potentially exposed populations. Although several potential pathways may exist, not all may be complete. For a pathway to be complete, the following four factors must exist:

- Source of COPC/ROPC release into the environment
- Release and transport mechanism (e.g., deposition to soil) that moves the COPC/ROPC from the source to other locations
- Point of contact with the affected medium
- Exposure route such as breathing vapors or ingesting affected media

These four factors were considered in the conceptual exposure model. The sources of COPC/ROPC release are the stack and any fugitive emissions from the RPP-WTP (see section 5). Transport processes, potential points of contact, and complete exposure pathways are identified to formulate exposure scenarios that will be the focus of the quantitative risk assessment. This process is summarized in the following section.

7.1.1. Characterization of Exposure Setting

Characterizing the exposure setting is the first step in identifying potentially exposed populations. This characterization includes identifying the location of human populations within the study area, and types of activities that can be expected under current and reasonable potential future land use.

The brief site characterization provided in this section is adequate to identify most possible receptors, especially the most significantly exposed receptors. A more detailed site characterization will be supplied in the PRA to describe all the populations evaluated.

The study area is defined as the area within a 50-kilometer radius of the RPP-WTP (Figure 7-2). This is considered to be the limit of the ISCST3 air-dispersion model. While it is possible for human populations to be exposed beyond this 50-km radius, the concentration of airborne and deposited emissions will be orders of magnitude less than those within the study area; essentially approaching zero. EPA 1998a reports that the most significant deposition of airborne emissions has been observed within a 3-km radius of a source (Figure 7-3). The Hanford Site boundary extends approximately 9 to 28 kilometers from the RPP-WTP. The Columbia River is located approximately 8 to more than 20 kilometers from the RPP-WTP. Therefore the potential for offsite impacts is likely to be minimal. Because of the importance of the Columbia River as a potable water and recreational resource, it will be included in the quantitative risk assessment despite its distance from the RPP-WTP. Currently, no residential receptors are present on the Hanford Site, nor are there likely to be any in the near future. Game animals that browse on Hanford Site property and plants that grow on Hanford Site property may be harvested by Native Americans living offsite.

Characterization of the exposure setting includes the following:

- Characterization of the physical setting – Including location of important physical features such as Gable Mountain, surface water bodies, and watersheds.

- Characterization of potentially exposed populations – Including identifying the location and activity patterns of current populations relative to the facility, determining plausible future land use, and identifying subpopulations of potential concern.

Characterization will concentrate on the areas potentially most impacted by emissions based on the results of the air-dispersion modeling.

<u>Current Land Use</u>

Current land use within the 50-km study area is characterized based on aerial photographs, zoning maps, land development plans, and information presented in:

- *Hanford Site Risk Assessment Methodology* (DOE-RL 1995)

- *Screening Assessment and Requirements for a Comprehensive Assessment: Columbia River Comprehensive Impact Assessment* (DOE-RL 1998)

- *The Future for Hanford: Uses and Cleanup* (DOE 1992)

- *Draft Hanford Remedial Action Environment Impact Study and Comprehensive Land Use Plan* (DOE 1996a).

Representatives of the Nez Perce Tribe, Yakama Nation, and Confederated Tribes of the Umatilla Indian Reservation are also being consulted in evaluating current land use in the study area.

Future Land Use

Plausible future land use is characterized based on land development plans, and information presented in:

- *Hanford Site Risk Assessment Methodology* (DOE-RL 1995)
- *Screening Assessment and Requirements for Comprehensive Assessment: Columbia River Comprehensive Impact Assessment* (DOE-RL 1998)
- *The Future for Hanford: Uses and Cleanup* (DOE 1992)
- *Draft Hanford Remedial Action Environmental Impact Study and Comprehensive Land Use Plan* (DOE 1996)

Representatives of the Nez Perce Tribe, Yakama Nation, and Confederated Tribes of the Umatilla Indian Reservation are also being consulted in evaluating plausible future land use in the study area. Worst-case future land use will be based on the assumption that the exposure locations could be used for any purpose regardless of development plans, or the actual potential for such use to occur.

7.1.2. Identification of Exposure Scenarios

EPA 1998a recommends the following exposure scenarios for evaluating potential risks from combustion facilities:

- Resident - Adult
- Resident - Child
- Subsistence Farmer - Adult
- Subsistence Farmer - Child
- Subsistence Fisher - Adult
- Subsistence Fisher - Child
- Nursing Infant
- Acute Risk

The subsistence farmer – adult and child, and subsistence fisher – adult and child, represent worst case scenarios.

The nursing infant scenario is recommended by EPA guidance (EPA 1998a) to address specific concerns regarding exposure to PCDDs and PCDFs. EPA guidance recommends inclusion of the nursing infant only for PCDDs/PCDFs; however, both coplanar and non-coplanar PCBs will also be evaluated for this pathway in the SLRA. Because radionuclides are the primary component of the waste to be treated at the facility, ROPCs will also be evaluated for this pathway. The ROPCs strontium (Sr-90), iodine (I-129), and cesium (Cs-134 and Cs-137) will be evaluated for the nursing infant scenario. These radionuclides are potentially present in the waste and are judged to have the highest potential for accumulation in milk and toxicity (EPA 1999d). No other COPCs/ROPCs will be evaluated for this scenario, which will be evaluated based on intakes for both the resident adult and the Native American resident adult.

In addition to the recommended exposure scenarios listed above, EPA recommends evaluation of special subpopulations by identifying their locations, and determining whether they are located in areas that are particularly conducive to impacts from RPP-WTP emissions. Special subpopulations are defined as human receptors or segments in the population that potentially may be at higher risk due to receptor sensitivity to COPCs. As noted by the EPA, the risk assessment approach used is designed to protect human health, including special subpopulations (e.g., through the use of RfDs developed to account for toxicity to sensitive receptors). However, to address actual characteristics of the exposure setting and potential community concerns, the following additional exposure scenarios will be evaluated: Hanford Site industrial worker, and Native American subsistence resident.

Workers employed at the RPP-WTP will not be included in the risk assessment because other regulations exist for occupational exposures within the RPP-WTP boundary (e.g., Occupational Safety and Health Administration or Department of Energy (DOE)-equivalent standard). However, because of its location within the Hanford Site, the closest and most likely receptors are other Hanford Site workers located outside the RPP-WTP boundary. Therefore, a Hanford Site industrial worker scenario will be included in the risk assessment.

Native American tribes (Yakama Nation, Confederated Tribes of Umatilla Reservation, Nez Perce Tribe) ceded the land currently occupied by the US Government's Hanford Site. However, these tribes claim retained rights to use resources on this land. A wide range of possible Native American activities related to traditional life styles exists. The Native American scenarios will address a variety of potential exposures associated with food gathering (including hunting and fishing), as well as cultural and social activities.

7.1.3. Description of Land Use Scenarios

Exposure scenarios to be addressed by the risk assessment are described in more detail below and summarized in Table 7-1. The scenarios to be carried through the quantitative risk assessment will be determined based on the results of the air-dispersion modeling. For example, if air-dispersion modeling indicates that significant emissions will not reach the Columbia River, the subsistence fisher scenario will not be evaluated in the quantitative risk assessment. Because the location of sacred sites is confidential within the tribes, representatives of the three tribes will be consulted following completion of the air-dispersion modeling to determine potential impacts to sacred sites.

Four types of exposure scenarios will be quantified: plausible current, plausible future, worst-case future, and acute exposure. By examining these four types of exposure scenarios, a range of possible exposures will be provided, giving the permit and risk managers additional information on which to base decisions. Using this approach, it will not be necessary to rely solely on worst-case assumptions.

The primary difference between plausible and worst-case scenarios is the assumed location of receptor populations. For the plausible scenarios, potential receptors are assumed to be located in areas where similar receptors currently exist or may reasonably be expected to exist some time in the future. For the worst-case scenarios, receptor populations are assumed to be located at the point of maximum concentration regardless of the likelihood of this occurring. For example, no residential development is expected within 10 kilometers (Figure 7-2 and Figure 7-3) of the RPP-WTP; however, if the point of maximum concentration falls within this area, a residential receptor will be evaluated at this location. The four types of exposure scenarios are defined below.

7.1.3.1. Plausible Current Land Use Scenarios

The area within a 10-km radius of the RPP-WTP is completely within the current boundaries of the Hanford Site. Current activities in this area are associated with waste management, environmental restoration, and industrial and commercial land use (Figure 7-4). The potentially exposed population in this area is limited to Hanford Site industrial workers. The 50-km study area extends beyond the Hanford Site boundary and includes areas that are, or can be used, for residential development. Although Native American residents do not live within the site boundaries, they may engage in food gathering (plant gathering, hunting, fishing) and cultural activities on Hanford Site land.

Based on this information, the plausible current exposure scenarios are as follows:

- Hanford Site industrial worker
- Residential adult and child at the Hanford Site boundary
- Native American resident adult and child hunter/gatherer living at the Hanford Site boundary and conducting some food gathering on the Hanford Site
- Native American resident adult and child fisher living at the Hanford Site boundary and fishing the Columbia River

7.1.3.2. Plausible Future Land Use Scenarios

Plausible future land use includes continued use of the area for industrial and commercial purposes and residential development beyond the Hanford Site boundary (Figure 7-5). Although Native American residents do not live within the site boundaries, they may engage in food gathering (i.e., plant gathering, hunting, fishing) and cultural activities on Hanford Site land. Based on this information the plausible future exposure scenarios are as follows:

- Hanford Site industrial worker
- Residential adult and child at the Hanford Site boundary
- Native American resident adult and child hunter/gatherer living at the Hanford Site boundary and conducting some food gathering on the Hanford Site
- Native American resident adult and child fisher living at the Hanford Site boundary and fishing the Columbia River

Plausible current and future land use scenarios are very similar and may be the same for some receptors.

7.1.3.3. Worst-Case Future Land Use Scenarios

EPA 1998a recommended exposure scenarios (resident adult and child, subsistence farmer adult and child, subsistence fisher adult and child, and nursing infant) will be evaluated at the point of maximum concentration identified by air-dispersion modeling to provide a worst-case estimate of exposure and risk.

In addition to these EPA-recommended scenarios, resident subsistence Native American scenarios (including a nursing infant) will be evaluated for unrestricted land-use. Based on this information, worst-case future exposure scenarios are as follows:

- Resident adult and child (on-site)
- Subsistence farmer adult and child (on-site)
- Subsistence fisher adult and child (off-site, at the Columbia River)
- Nursing infant (on-site)

- Native American subsistence resident adult and child hunter/gatherer (on-site)
- Native American subsistence resident adult and child fisher (offsite, at the Columbia River)
- Native American nursing infant (on-site)

7.1.3.4. Acute Exposure

EPA 1998a recommends evaluating potential acute exposures in addition to the chronic exposures evaluated by previously described exposure scenarios. The acute exposure scenario includes direct inhalation of airborne emissions at the estimated maximum 1-hour concentration. The receptor for the acute exposure scenario is located at the point of maximum 1-hour concentration and is independent of land- use. Because the acute exposure scenario is based on the maximum-modeled concentration and assumes that a receptor will be present at the location of that maximum during the hour in which it occurs, this is considered a worst-case scenario.

7.1.4. Exposure Pathways

Exposure pathways to be evaluated for each of these exposure scenarios are summarized in Table 7-1 and the conceptual exposure model (Figure 7-1). Both direct and indirect pathways will be evaluated. Direct exposure pathways to be included in the quantitative risk assessment are as follows:

- COPCs and ROPCs
  - direct inhalation of emissions

- ROPCs only
  - external exposure to radionuclides in air.

Indirect exposure pathways to be included in the quantitative risk assessment are as follows:

- COPCs and ROPCs
  - ingestion of soil
  - inhalation of resuspended soil
  - ingestion of homegrown or wild gathered produce
  - ingestion of homegrown beef, milk, chicken, eggs, and pork
  - ingestion of drinking water
  - ingestion of fish
  - ingestion of wild game (including fowl and eggs)

- COPCs only
  - dermal absorption in the sweat lodge

- ROPCs only
  - external exposure to radionuclides in soil

- PCDDs/PCDFs, PCBs, and selected ROPCs (Sr-90, I-129, Cs-137, and Cs-137) only
  - ingestion of breast milk

- VOCs, SVOCs, and volatile ROPCs only
  - inhalation of vapors in the sweat lodge

External radiation exposure will be quantitatively evaluated for radionuclides in air and soil. External radiation exposure is not expected to be significant for surface water because of the following:

- Distance to the Columbia River will result in extremely low concentrations of ROPCs via deposition
- ROPC concentrations in air near the facility and in soil following deposition and accumulation will far exceed surface water concentrations
- Exposure to air and soil are continuous while potential exposure to surface water is intermittent

Thus, external radiation exposure is not expected to be significant for surface water because the distance to the Columbia River will result in extremely low concentrations of ROPCs via deposition. Thus, ROPC concentrations in air near the facility, and in soil following deposition and accumulation, are expected to far exceed far away surface water concentrations.

EPA 1998a has identified three exposure pathways that are generally insignificant contributors to risk at thermal treatment facilities; they are as follows:

- Groundwater pathways
- Resuspended dust
- Dermal contact

Groundwater pathways are generally not significant contributors to risk from airborne emissions because exposure concentrations in groundwater following air dispersion, deposition, leaching, and groundwater dispersion are much less than concentrations in air, soil, and other media. Conditions at the Hanford Site (i.e., very low precipitation) will make the contribution to groundwater even less than at other sites. Therefore, exposure to groundwater will not be included in the quantitative risk assessment. However, surface water concentrations will be used to evaluate the ingestion of drinking water, as well as inhalation of VOCs/SVOCs and dermal absorption for the Native American sweat lodge scenario.

Inhalation of resuspended dust can be an important exposure pathway at contaminated sites where the contaminant source is at the surface or in the soil. This was explained in the air dispersion modeling section (6). At these sites, dust resuspension generally represents the only source of inhalation exposure (unless significant volatiles are present). At sites such as the RPP-WTP where the source of COPCs/ROPCs is airborne emissions, direct, continuous inhalation of these emissions is a much more important exposure pathway than periodic inhalation of fugitive dust. However, because of the dry, dusty conditions at the Hanford Site, inhalation of resuspended dust will be evaluated in the PRA.

Dermal exposure pathways (to soil, surface water, or air) will not be included in the PRA with the exception of the Native American sweat lodge scenario. This is a non-conservative assumption, because dermal contact will occur when direct contact with soil or surface water is a potentially complete pathway, along with incidental ingestion. However, dermal exposure pathways have been identified as insignificant contributors to risk in numerous risk assessments prepared and/or reviewed by EPA for airborne emissions from thermal treatment facilities. If initial PRA results indicate that the ingestion pathway results in risks that are borderline for any plausible receptor, then the dermal exposure pathway may be included in the PRA. A discussion of the potential impact associated with exclusion of this minor pathway from the quantitative risk assessment will be included in the uncertainty assessment.

7.1.5. Quantification of Exposure

The following section provides the calculation algorithms that will be used to quantify intake (or dose) for each COPC and ROPC. Estimates of the lifetime average daily doses (LADDs) and average daily doses (ADDs) will be quantified. The LADD defines a dose level that is distributed (averaged) over an entire lifetime, rather than a specific incremental exposure period. Unlike the LADD, the ADD is not averaged over an entire lifetime.

The equations that will be used to quantify each of the exposure pathways are given below and are based on those presented in Appendix C of EPA 1998a; these equations are subject to change as the guidance is modified.

7.1.5.1. Inhalation of Vapor Phase and Particulate Emissions

Inhalation of vapor phase and particulate emissions will be quantified with the following equations:

COPCs:
$$I_{avg} = \frac{C_a \times IR \times ET \times EF \times ED \times 0.001}{BW \times AT \times 365}$$

ROPCs:
$$I_{avg} = C_a \times IR \times ET \times EF \times ED$$

where

I_{avg} = average daily COPC/ROPC intake via inhalation (mg COPC/kg-day for chemicals; pCi ROPC for radionuclides)
C_a = total COPC concentration in air ($\mu g/m^3$ for chemicals; pCi/m^3 for radionuclides)
IR = inhalation rate (m^3/hour)
ET = exposure time (hour/day)
EF = exposure frequency (day/year)
ED = exposure duration (year)
0.001 = conversion factor (mg/µg)
BW = body weight (kg)
AT = averaging time (year)
365 = conversion factor (day/year)

7.1.5.2. Ingestion of Soil

Ingestion of COPCs/ROPCs in soil will be quantified with the following equations:

COPCs:
$$I_{soil} = \frac{Cs \times CR_{soil} \times F_{soil} \times EF \times ED}{BW \times AT \times 365}$$

ROPCs:
$$I_{soil} = Cs \times CR_{soil} \times F_{soil} \times EF \times ED \times 1000$$

where

I_{soil} = daily intake of COPC/ROPC due to soil ingestion (mg/kg-day for chemicals; pCi for radionuclides)
Cs = average soil concentration of COPC/ROPC over exposure duration (mg COPC/kg soil for chemicals; pCi ROPC/g soil for radionuclides)

CR_{soil} = consumption rate of soil (kg/day)
F_{soil} = fraction of ingested soil that is contaminated (unitless)
EF = exposure frequency (day/year)
ED = exposure duration (year)
BW = body weight of receptor (kg)
AT = averaging time (year)
365 = conversion factor (day/year)
1000 = conversion factor (g/kg)

7.1.5.3. Ingestion of Produce

Ingestion of COPCs/ROPCs in homegrown or wild gathered produce will be quantified with the following equations:

COPCs:

$$I_{ag} = \frac{\{[(Pd + Pv + Pr) \times CR_{ag}] + (Pr \times CR_{pp}) + (Pr_{bg} \times CR_{bg})\} \times F_{ag} \times EF \times ED}{AT \times 365}$$

ROPCs:

$$I_{ag} = \{[(Pd + Pv + Pr) \times CR_{ag}] + (Pr \times CR_{pp}) + (Pr \times CR_{bg})\} \times F_{ag} \times EF \times ED \times 1000 \times BW$$

where

I_{ag} = daily intake of COPC/ROPC from produce (mg/kg-day for chemicals; pCi for radionuclides)
Pd = COPC/ROPC concentration in aboveground produce due to direct deposition onto plant surfaces (mg/kg for chemicals; pCi/g for radionuclides)
Pv = COPC/ROPC concentration in aboveground produce due to air-to-plant transfer (mg/kg for chemicals; pCi/g for radionuclides)
Pr = COPC/ROPC concentration in aboveground produce due to root uptake (mg/kg for chemicals; pCi/g for radionuclides)
CR_{ag} = consumption rate of aboveground unprotected produce (kg/kg-day DW or dry weight)
CR_{pp} = consumption rate of aboveground protected produce (kg/kg-day DW or dry weight)
Pr_{bg} = COPC/ROPC concentration in belowground produce due to root uptake (mg/kg for chemicals; pCi/g for radionuclides)
CR_{bg} = consumption rate of belowground produce (kg/kg-day DW or dry weight)
F_{ag} = fraction of ingested produce that is contaminated (unitless)
EF = exposure frequency (day/year)
ED = exposure duration (year)
AT = averaging time (year)
365 = conversion factor (day/year)
1000 = conversion factor (g/kg)
BW = body weight (kg)

7.1.5.4. Ingestion of Beef, Milk, Pork, Poultry, Eggs, and Wild Game

Ingestion of COPCs/ROPCs in homegrown beef, milk, pork, poultry, eggs and wild game will be quantified with the following equations.

$$\text{COPCs:}\quad I_i = \frac{A_j \times CR_j \times F_j \times EF \times ED}{AT \times 365}$$

$$\text{ROPCs:}\quad I_i = A_j \times CR_j \times F_j \times EF \times ED \times 1000 \times BW$$

where

I_i = daily intake of COPC/ROPC *i* from animal *j* (mg/kg-day for chemicals; pCi for radionuclides)
A_j = concentration of COPC/ROPC i in animal *j* (mg/kg FW or fresh (wet) weight for chemicals; pC*i*/g for radionuclides)
CR_j = consumption rate of animal *j* (kg/kg-day FW or fresh (wet) weight)
F_j = fraction of ingested animal tissue *j* that is contaminated (unitless)
EF = exposure frequency (day/year)
ED = exposure duration (year)
AT = averaging time (year)
365 = conversion factor (day/year)
1000= conversion factor (g/kg)
BW = body weight (kg)

For the metals mercury, selenium, and cadmium, the concentrations in beef, milk, and pork as well as the consumption rate are in kg DW per day and must be adjusted using wet weight to dry weight conversion factors.

7.1.5.5. Ingestion of Fish

Ingestion of COPCs/ROPCs in fish (e.g., chinook salmon, bass) will be quantified using the following equations:

$$\text{COPCs:}\ I_{fish} = \frac{C_{fish} \text{ x } (CR_{fish} + CR_{organs}) \text{ x } F_{fish} \text{ x } EF \text{ x } ED}{AT \text{ x } 365}$$

$$\text{ROPCs:}\ I_{fish} = C_{fish} \text{ x } (CR_{fish} + CR_{organs}) \text{ x } F_{fish} \text{ x } EF \text{ x } ED \text{ x } 1000 \text{ x } BW$$

where

I_{fish} = daily intake of COPC/ROPC from fish (mg/kg-day for chemicals; pCi for radionuclides)
C_{fish} = concentration of COPC/ROPC in fish (mg/kg for chemicals; pCi/g for radionuclides)
CR_{fish} = consumption rate of fish (fillets) (kg/kg-day FW)
CR_{organs} = consumption rate of fish parts (e.g., head, fins, etc.) (kg/kg-day FW)
F_{fish} = fraction of ingested fish that are contaminated (unitless)
EF = exposure frequency (day/year)
ED = exposure duration (year)
AT = averaging time (year)
365 = conversion factor (day/year)
1000 = conversion factor (g/kg)
BW = body weight (kg)

7.1.5.6. Ingestion of Water

Ingestion of COPCs/ROPCs in drinking water will be quantified using the following equation:

$$\text{COPCs: } I_{dw} = \frac{C_{dw} \times CR_{dw} \times F_{dw} \times EF \times ED}{BW \times AT \times 365}$$

$$\text{ROPCs: } I_{dw} = C_{dw} \times CR_{dw} \times F_{dw} \times EF \times ED$$

where

I_{dw}	=	daily intake of COPC/ROPC in drinking water (mg/kg-day for chemicals, pCi for radionuclides)
C_{dw}	=	dissolved phase COPC/ROPC water concentration (mg/L for chemicals; pCi/L for radionuclides)
CR_{dw}	=	drinking water consumption rate (L/day)
F_{dw}	=	fraction of ingested drinking water that is contaminated (unitless)
EF	=	exposure frequency (day/year)
ED	=	exposure duration (year)
BW	=	body weight (kg)
AT	=	averaging time (year)
365	=	conversion factor (day/year)

7.1.5.7. Infant Ingestion of Breast Milk

The infants ingestion of COPCs/ROPCs in breast milk will be quantified for dioxins/furans, coplanar dioxin-like PCBs and selected radionuclides (Sr-90, I-129, Cs-134, and Cs-137) based on the following equations:

$$\text{COPCs: } I_{infant} = \frac{\frac{m \times CF \times h \times f_1}{0.693 \times f_2} \times f_3 \times f_4 \times IR_{milk} \times ED}{BW_{infant} \times AT}$$

$$\text{ROPCs: } I_{infant} = \frac{m \times h \times f_1}{0.693 \times f_2} \times f_3 \times f_4 \times IR_{milk} \times ED \times EFI$$

where

I_{infant}	=	average infant intake of COPC/ROPC (pg/kg-day for chemicals; pCi for radionuclides)
m	=	average maternal intake of COPC/ROPC across all adult ingestion scenarios (mg/kg BW-day for chemicals; pCi/kg BW-day for radionuclides)
CF	=	conversion factor of 1×10^9 (pg/mg)
h	=	half-life of COPC/ROPC in adults (days)
f_1	=	fraction of ingested COPC/ROPC that is stored in fat (unitless)
f_2	=	fraction of mother's weight that is fat (unitless)
f_3	=	fraction of mother's breast milk that is fat (unitless)
f_4	=	fraction of ingested COPC/ROPC that is absorbed (unitless)
IR_{milk}	=	ingestion rate of breast milk by infant (kg/day)

ED = exposure duration (year)
BW_{infant} = body weight of infant (kg)
AT = averaging time (year)
EF = exposure frequency (day/year)

This equation is expected to overestimate chemical concentrations in the lipid portion of breast milk because it does not consider the time required to reach steady state or contaminant losses from breast feeding as described below EPA (1998a):

- Chemical concentrations in milkfat may be overestimated by a factor of 10 or more if the half-life of the chemical is large compared to the exposure duration (i.e., near steady state conditions are not reached).

- Contaminant losses can be significant. EPA (1998a) reports that the highest concentrations of PCDDs/PCDFs are excreted in breast milk within the first few weeks after delivery. After 1 year of breast feeding, PCDD/PCDF concentrations are about 30-50% of initial levels.

For dixoins/furans, as recommended in EPA 1998a, I_{infant} will be compared to an average infant intake target level of 60 pg/kg-day of 2,3,7,8-TCDD TEQ. As recommended by EPA Region X, for coplanar (dioxin-like) PCBs, I_{infant} will be compared to an average infant intake target level of 13 pg/kg-day of 2,3,7,8-TCDD TEQ. Cancer risks will be quantified for ROPCs, based on I_{infant}.

7.1.5.8. External Exposure to Radionuclides in Soil

External exposure to ionizing radiation in soil from ROPCs will be quantified using the following equation:

$$\text{ROPCs: } I_{irs} = Cs \times (1\text{-}Se) \times Te \times EF \times ED \times (1/365)$$

where

I_{irs} = intake of ROPC due to external exposure to ionizing radiation in soil [(pCi-year)/g]
Cs = average soil concentration of ROPC (pCi/g)
Se = shielding factor (unitless)
Te = exposure time factor (unitless)
EF = exposure frequency (day/year)
ED = exposure duration (year)
1/365 = conversion factor (year/day)

7.1.5.9. External Exposure to Radionuclides in Air

External exposure to ionizing radiation in air from ROPCs will be quantified using the following equation:

$$\text{ROPCs: } I_{ira} = Ca \times EF \times ED \times 86400 \times AF \times 0.037$$

where

I_{ira} = intake of ROPC due to external exposure to ionizing radiation in air [(Bq-secs)/m^3]

Ca = concentration of ROPC in air (pCi/m^3)
EF = exposure frequency (day/year)
ED = exposure duration (year)
86400 = conversion factor (secs/day)
AF = age factor of 1 for adults and 1.3 for children (unitless) (Saito, et al. 1998)
0.037 = conversion factor (Bq/pCi)

7.1.5.10. Inhalation of Vapors in Sweat Lodge Scenario

The inhalation of volatile and semivolatile organic compounds for Native American adults inside the sweat lodge will be quantified with the following equation which assumes a hemisphere with a 2 m diameter:

VOCs and SVOCs that are COPCs: $$I_{inh} = \frac{Cw \times \left(\frac{VW}{\frac{2}{3}\pi r^3}\right) \times IR \times EF \times ED \times ET \times \frac{1}{24}}{BW \times AT \times 365}$$

Volatile ROPCs: $$I_{inh} = Cw \times \left(\frac{VW}{\frac{2}{3}\pi r^3}\right) \times IR \times EF \times ED \times ET \times \frac{1}{24}$$

where

I_{inh} = average daily intake of VOCs/SVOCs that are COPCs/ROPCs, from adult inhalation within the sweat lodge (mg COPC/kg-day for chemicals; pCi for radionuclides)
Cw = average surface water concentration (mg/L for chemicals; pCi/L for radionuclides)
VW = volume of water used in sweat lodge (L)
r = radius of sweat lodge (m)
IR = inhalation rate (m^3/day)
EF = exposure frequency (day/year)
ED = exposure duration (year)
ET = exposure time (hour/day)
$\frac{1}{24}$ = conversion factor (day/hour)
BW = body weight (kg)
AT = averaging time (year)
365 = conversion factor (day/year)

7.1.5.11. Dermal Absorption in Sweat Lodge Scenario

The dermal absorption of water for Native American adults inside the sweat lodge (a hemisphere with a 2 m diamter) will be quantified with the following equation:

COPCs: $$I_d = \frac{Cw \times SA \times Kp \times ET \times EF \times ED \times 10}{BW \times AT \times 365}$$

where

I_d	=	average daily intake of COPCs, from adult dermal absorption within the sweat lodge (mg COPC/kg-day)
Cw	=	average surface water concentration (mg/L)
SA	=	body surface area available for contact (m^2)
Kp	=	permeability constant (COPC-specific) (cm/hour)
ET	=	exposure time (hour/day)
EF	=	exposure frequency (day/year)
ED	=	exposure duration (year)
10	=	conversion factor $\left[\left(\frac{L}{m^3}\right)\times\left(\frac{m}{cm}\right), \text{or}\left(\frac{L}{m^2-cm}\right)\right]$
BW	=	body weight (kg)
AT	=	averaging time (year)
365	=	conversion factor (day/year)

7.1.5.12. Inhalation of Resuspended Soil

The inhalation of resuspended soil will be quantified with the equation shown below. See section 6.1.2.6 for information regarding the estimation of air concentrations of resuspended dust.

$$\text{COPCs: } I_{inhsoil} = \frac{Ca_{soil} \times IR \times ET \times EF \times ED \times 0.001}{BW \times AT \times 365}$$

$$\text{ROPCs: } I_{inhsoil} = Ca_{soil} \times IR \times ET \times EF \times ED$$

where

$I_{inhsoil}$	=	average daily COPC/ROPC intake via inhalation of resuspended soil (mg/kg-day for chemicals; pCi for radionuclides)
Ca_{soil}	=	average air concentration of COPC/ROPC from resuspended soil (µg COPC/m^3 for chemicals; pCi ROPC/m^3 for radionuclides)
IR	=	inhalation rate (m^3/hour)
ET	=	exposure time (hour/day)
EF	=	exposure frequency (day/year)
ED	=	exposure duration (year)
0.001	=	conversion factor (mg/µg)
BW	=	body weight (kg)
AT	=	averaging time (year)
365	=	conversion factor (day/year)
1000	=	conversion factor (g/kg)

7.1.6. Exposure Parameters

The equations presented above are the basis for quantifying the exposure to COPCs/ROPCs experienced by a potential receptor. The values that will be used for each parameter identified in the equations are provided in Tables 7-2, 7-3, and 7-4. For residential scenarios in Table 7-2, exposure values were predominantly taken from the *Human Health Risk Assessment Protocol for Hazardous Waste Combustion*

Facilities (EPA 1998a). For Native American subsistence scenarios in Table 7-3, exposure values were predominantly taken from Harris and Harper 1997. For the Hanford Site worker scenario in Table 7-4, exposure values were predominantly taken from the *Hanford Site Risk Assessment Methodology* (DOE-RL 1995).

7.1.7. Exposure Point Concentrations

The exposure point concentrations used for estimating dose of both COPCs and ROPCs are dependent on the location of the receptor. The location of the various receptor populations identified for the quantitative risk assessment will correspond to the receptor grid nodes defined during air-dispersion modeling (section 6). In keeping with the protective approach used in this risk assessment, the location with the maximum concentration of COPCs/ROPCs will be used in estimating exposure point concentrations. Because the point of maximum concentration may be different for airborne COPCs/ROPCs and COPCs/ROPCs deposited via wet and dry deposition mechanisms, EPA 1998a recommends the following method for selecting the point of maximum concentration. Within each area to be evaluated, receptor grid nodes or exposure locations with the highest modeled unitized value will be identified for each of the following parameters:

- Vapor phase air concentration
- Vapor phase wet deposition rate
- Particle phase air concentration
- Particle phase wet deposition rate
- Particle phase dry deposition rate
- Particle-bound phase air concentration
- Particle-bound phase wet deposition rate
- Particle-bound phase dry deposition rate

At each of the unique locations, air concentrations from vapor, particle phase, and particle bound fractions will be summed, and concentrations of deposited materials will be summed to determine the point of maximum exposure. Air concentrations and deposition rates will not be summed together.

7.1.7.1. Exposure Point Concentrations for Plausible Exposure Scenarios

For the plausible current and future exposure scenarios, exposure point concentrations will be estimated at actual or likely exposure locations using models presented in section 6. Locations to be used in estimating exposure point concentrations are described below and summarized in Table 7-1.

Residential Scenarios. Residential use of the Hanford Site is prohibited. Therefore, the nearest potential residential receptor would be at the Hanford Site boundary. Current and future residential adult and child scenarios will be located at the point(s) of maximum concentration (as defined above) at the property boundary.

Hanford Site Industrial Worker Scenario. The Hanford Site industrial worker will be located at the point(s) of maximum concentration as identified by the air-dispersion modeling unless access to this location is and will remain restricted.

Native American Scenarios. The plausible current and future location of activities will be determined with input from the three tribes after completion of the air-dispersion modeling. Locations selected will represent the maximally impacted location at the minimum distance from the RPP-WTP for each activity.

7.1.7.2. Exposure Point Concentrations for Worst-case Exposure Scenarios

For the worst-case future land use scenarios, exposure will be evaluated at the point(s) of maximum concentration regardless of actual land use patterns.

Residential Scenarios. Residential scenarios, including resident adult and child, subsistence farmer adult and child, nursing infant, and the residential portion of the subsistence fisher adult and child will be evaluated at the point(s) of maximum concentration as identified by the air-dispersion modeling unless physical factors would prevent a residence at this location. COPC/ROPC concentrations in drinking water will be estimated from the point(s) of maximum concentration at the Columbia River.

Subsistence Fisher Scenario. The adult and child subsistence fisher will obtain fish from the point(s) of maximum concentration at the Columbia River.

Native American Scenarios. The potential future location of activities will be determined with input from the three tribes after completing the air-dispersion modeling. The locations selected will represent the maximally impacted location at the minimum distance from the RPP-WTP for each activity.

7.2. Toxicity Assessment

This section of the SLRA will present the chemical specific dose-response information to be used in the risk assessment. The toxicity assessment will be performed in accordance with EPA 1998a guidance. Chronic toxicity data will be obtained according to the following hierarchy:

1. **Integrated Risk Information System (IRIS)** EPA 1998c or most current. IRIS is an on-line database that provides toxicity values for chronic oral and inhalation exposures. All data contained in IRIS are verified by an EPA work group. As such, IRIS serves as the primary source of toxicity values for the risk assessment.

2. **Health Effects Assessment Summary Tables (HEAST)** (EPA 1995d and EPA 1997a or most current). HEAST is an EPA document that supplements IRIS by providing nonverified toxicity values, as well as values for evaluating the potential for noncancer effects following subchronic exposures. Information in HEAST (EPA 1997a) is used as a secondary source for chemicals when information is not available from IRIS. HEAST (EPA 1995d) is the source for all carcinogenic toxicity values (slope factors) for radionuclides. Ingrowth of radioactive daughters will be accounted for by using the +D (plus daughters) slope factors. This is a conservative approach since use of the +D assumes secular equilibrium.

3. **Provisional Values.** In the absence of established values from IRIS or HEAST, provisional toxicity values are used and are available from several sources (i.e., NCEA's Superfund Technical Support Center, EPA Regional Toxicologists, ATSDR Toxicological Profiles).

4. **Surrogate Values.** When toxicity values for a chemical are not available from the sources listed above, the use of a surrogate value may be necessary. This process involves applying a toxicity value established for one chemical to another chemical for which no value has been established. The application of surrogate values is based on similarities in structure, mechanism of action, and toxicity. The surrogate values will be identified in the SLRA when their need is identified

The same approach will be used for the toxicity assessment in both the PRA and FRA. Any new toxicity values that become available prior to development of the FRA will be incorporated in the final assessment.

7.2.1. Noncarcinogenic Health Effects

Potential noncarcinogenic health effects associated with exposure to COPCs will be evaluated using acceptable daily intake levels (i.e., reference doses and concentrations) established by EPA 1997a and EPA 1998c or as described above.

It is widely accepted that most biological effects of chemicals occur only after a threshold dose is exceeded (Klaassen et al. 1996 and Paustenbach 1989). For purposes of establishing noncarcinogenic health criteria, this threshold dose is usually estimated from the no-observed adverse effect level (NOAEL) or lowest-observed adverse effect level (LOAEL) determined in animal or human studies. NOAEL is defined as the exposure level at which no statistically or biologically significant increases are present in the frequency or severity of adverse effects (EPA 1989b). The LOAEL is the lowest exposure level at which there are statistically or biologically significant increases in frequency or severity of adverse effects (EPA 1989b). The LOAEL or NOAEL from the most sensitive animal or human study is used by the EPA to establish long-term health criteria, termed reference doses (RfDs). A RfD is defined as an estimate (with uncertainty spanning perhaps an order of magnitude) of the dose of a chemical (expressed in mg/kg-day) that is likely to be without an appreciable risk of deleterious effects during a lifetime (EPA 1989b). Similarly, a reference concentration (RfC) represents the concentration of a chemical in environmental media (expressed in μg/L for water or mg/m^3 for air) that is likely to be without an appreciable risk of deleterious effects during a lifetime (EPA 1989b). When deriving a RfD or RfC, a NOAEL value is used preferentially over a LOAEL value if both are available from the key study. EPA derives RfDs and RfCs by applying uncertainty factors to the NOAEL or LOAEL value to provide a margin of safety. The equation for deriving a RfD or RfC is shown below:

$$\text{RfD or RfC} = (\text{NOAEL or LOAEL}) / (\text{UF} \times \text{MF})$$

where

RfD = reference dose (mg/kg/day)
RfC = reference concentration (mg/m^3)
NOAEL = no-observed-adverse-effect-level
LOAEL = lowest-observed-adverse-effect-level
UF = uncertainty factor
MF = modifying factor.

Uncertainty factors can range from 1 to 10,000 and may include a factor of up to 10 to account for each of the following.

- Variation in sensitivity within human populations
- Extrapolation of effects observed in animals to humans
- Extrapolation from less-than-lifetime exposures in the critical study to lifetime exposures
- Extrapolation from a LOAEL to a NOAEL, if necessary

In some cases, a modifying factor [usually ranging from 1 to 10 or <1 for most essential nutrients (EPA 1989b)] also is applied to the NOAEL/LOAEL. This value reflects a qualitative professional assessment of additional uncertainties in the critical study and in the entire database for the chemical not explicitly addressed by the above uncertainty factors (EPA 1989b). RfDs and RfCs can be interconverted using default exposure assumptions (i.e., 70-kg body weight, 2 L/day water intake, and 20 m^3/day breathing

rate). EPA establishes RfDs and RfCs for evaluating both subchronic (less than 7 years) and chronic (7 years or more) exposures.

EPA generally reports only RfC values in IRIS and HEAST because the EPA observes that it is technically more accurate to base toxicity values directly on measured air concentrations than to make an estimate of the administered dose. Inhalation RfDs will be calculated from the corresponding RfC values when necessary using the following equation:

$$\text{Inhalation RfD} = (\text{Inhalation RfC}) \times (\text{Breathing Rate}) / (\text{Body Weight})$$

where

Inhalation RfD = chemical-specific inhalation reference dose in mg/(kg-day)
Inhalation RfC = chemical-specific inhalation reference concentration in mg/m^3
Breathing Rate = 20 m^3/day, default value for an adult
Body Weight = 70 kg, default value for an adult.

Uncertainties associated with this type of conversion include those surrounding deposition and absorption of the chemical in the lung, both of which depend on physico-chemical properties of the chemical, the phase of the chemical in air (i.e., vapor, particle, particle-bound), and characteristics of the exposed species. EPA recognizes the need for expressing toxicity values in terms of a dose (mg/kg-day) for risk assessment purposes and acknowledges that in many cases the conversion of a RfC to a dose does not add significant uncertainty to the risk assessment process (EPA 1997a). In addition, the appropriateness of this conversion depends on the toxicological endpoint observed in the key study. For example, it may be inappropriate to estimate an internal dose for compounds that act at the point of contact (i.e., sensitizers and irritants of the upper respiratory tract). In these cases, the toxicological endpoint depends only on the concentration of the chemical in air and not on the chemical dose expressed on a per body weight basis. For example, a chemical irritant will irritate nasal passages and lungs at a given concentration regardless of whether the exposed individual weighs 15 kg or 70 kg. In addition, this conversion might inappropriately imply effects to other organ systems or effects from other exposure routes.

Oral RfDs are adjusted for the evaluation of the dermal exposure pathway (EPA 1989c, 1996c). Most RfDs are expressed as the amount of substance administered per time and body weight; however, dermal exposure to chemicals in soil and water is expressed as absorbed dose.

For the dermal assessments in this risk assessment, the oral RfD for each COPC is adjusted by the percent gastrointestinal absorption efficiency (%GI) for that chemical. The %GI values have been obtained from the Risk Assessment Information System (RAIS), which is maintained by the Oak Ridge National Laboratory (ORNL 2000). The protocol used to obtain %GI values in the RAIS is based on guidance from EPA Region IV (EPA 1995e). This protocol involves obtaining %GI values from the literature, where available, and assigning default values for specific classes of chemicals where no literature values can be found. Specifically, the RAIS notes that measured %GIs are available for only a limited number of chemicals; for those chemicals for which no %GI is available in the literature, a value of 80 percent is used for VOCs, 50 percent for SVOCs, and 20 percent for inorganics. Wide ranges of %GI values can be found for some chemicals and, in the absence of chemical-specific absorption data, the values are often estimated from data for related chemical structures; most organic compounds are readily absorbed (i.e., %GI=100).

Adjustments, which favor conservatism, are made to the oral RfDs used in the dermal assessments. The dermal RfDs are calculated as follows:

Dermal (absorbed dose) RfD = (Oral RfD) x (%GI/100).

Oral RfDs, inhalation RfCs and RfDs, and dermal RfDs are provided in Tables 4-1 and 4-2.

Many constituents have oral or inhalation toxicity values available but not both. For these chemicals route-to-route extrapolation will be used to calculate toxicity values per USEPA (1998a). This conversion assumes that the toxicity of a constituent is equivalent over all routes of exposure and, therefore, introduces uncertainty into the risk assessment.

7.2.2. Carcinogenic Health Effects

Health risks from exposures to carcinogens are defined in terms of probabilities. These probabilities identify the likelihood of a carcinogenic response in an individual that receives a given dose of a particular compound. The slope factor, expressed in units of $(mg/kg\text{-}day)^{-1}$ multiplied by the lifetime ADD of the chemical, provides an estimate of the theoretical excess cancer risk. Slope factors represent an upper-bound estimate of the probability of developing cancer per unit dose (expressed as risk per mg/kg-day) of a chemical over a lifetime (EPA 1989b). Similarly, unit risks (URs) represent an upper-bound estimate of the probability of developing cancer per unit concentration [expressed as risk per (μg/L) for water; risk per ($\mu g/m^3$) for air] of a chemical over a lifetime. Slope factor and UR values can be interconverted using default exposure assumptions (i.e., 70-kg body weight, 2 L/day water intake, 20 m^3/day breathing rate).

The cancer weight-of-evidence classification is a qualitative descriptor that characterizes the quality and quantity of the data concerning the potential carcinogenicity of the chemicals. As defined by EPA 1989b, six weight-of-evidence groups exist to which a chemical may be assigned:

- Group A Human Carcinogen (sufficient evidence of carcinogenicity in humans)
- Group B1 Probable Human Carcinogen (limited evidence of carcinogenicity in humans)
- Group B2 Probable Human Carcinogen (sufficient evidence of carcinogenicity in animals, with inadequate or lack of evidence in humans)
- Group C Possible Human Carcinogen (limited evidence of carcinogenicity in animals, or lack of human data)
- Group D Not Classifiable as to Human Carcinogenicity (inadequate or no evidence)
- Group E Evidence of Noncarcinogenicity for Humans (no evidence of carcinogenicity in adequate studies)

Consistent with EPA guidelines (EPA 1989b), chemicals assigned a weight-of-evidence classification of A, B1, or B2 will be quantitatively evaluated for carcinogenic dose-response. All Group C carcinogens will also be quantitatively evaluated for carcinogenic effects.

In some cases, an oral SF must be calculated from the corresponding UR using the following equation:

$$\text{Oral SF} = (\text{Oral UR}) \times (\text{Body Weight}) \times (\text{Conversion Factor}) / (\text{Water Intake})$$

where

Oral SF	=	oral slope factor in $(mg/kg\text{-}day)^{-1}$
Oral UR	=	chemical-specific oral unit risk in $(\mu g/L)^{-1}$
Body Weight	=	70 kg, default value for an adult
Conversion Factor	=	1,000 μg/mg
Water Intake	=	2 L/day, default intake for an adult.

Expression of the UR in terms of a dose is necessary to evaluate cancer risk associated with exposure media other than drinking water (i.e., soil, sediment). EPA recognizes the need for expressing toxicity values in terms of a dose (mg/kg-day) for risk assessment purposes, and acknowledges that in many cases this conversion does not add significant uncertainty to the risk assessment process (EPA 1997b). However, it will be carried through to the discussion of uncertainties in the SLRA.

In some cases, an inhalation SF must be calculated from the corresponding UR using the following equation:

$$\text{Inhalation SF} = \frac{(\text{Inhalation UR}) \times (\text{Body Weight}) \times (\text{Conversion Factor})}{(\text{Breathing Rate})}$$

where

Inhalation SF	=	inhalation slope factor in $(mg/kg\text{-}day)^{-1}$
Inhalation UR	=	chemical-specific inhalation unit risk in $(\mu g/m^3)^{-1}$
Body Weight	=	70 kg, default for an adult
Conversion Factor	=	1000 μg/mg
Breathing Rate	=	20 m^3/day, default value for an adult.

The same uncertainties that apply to the conversion of inhalation RfC to RfD for noncarcinogens (see section 7.2.1) also apply to the conversion of inhalation unit risk (UR) to inhalation slope factor (SF) for carcinogens.

Oral slope factors are adjusted for the evaluation of the dermal exposure pathway (EPA 1989c, 1996c). Most slope factors are expressed as the amount of substance administered per time and body weight; however, dermal exposure to chemicals in soil and water is expressed as absorbed dose.

For the dermal assessments in this risk assessment, the oral slope factor for each COPC is adjusted by the percent gastrointestinal absorption efficiency (%GI) for that chemical. The source of %GI values used in this risk assessment has been described in section 7.2.1.

Adjustments, which favor conservatism, are made to the oral slope factors used in the dermal assessments. The dermal slope factors are calculated as follows:

Dermal (absorbed dose) Slope Factor = (Oral Slope Factor) / (%GI/100).

Oral, inhalation, and dermal SFs are provided in Tables 4-1 and 4-2. Many constituents have oral or inhalation slope factors available but not both. For these chemicals route-to-route extrapolation will be used to calculate slope factors. This conversion assumes that the carcinogenicity of a constituent is equivalent over all routes of exposure and, therefore, introduces uncertainty into the risk assessment.

7.2.3. Polychlorinated Dibenzo-p-dioxins and Polychlorinated Dibenzofurans

Chlorinated dioxins, chlorinated dibenzofurans, and chlorinated biphenyls are thought to act through a common mechanism of toxicity by binding to a protein known as the arylhydrocarbon receptor (AR) (for review see ATSDR 1997 or WHO 1998). The AR-ligand complex is responsible for the activation of genes that have a deleterious effect when they are not under proper regulation by the receptor's hormones. Interaction of dioxins and similar compounds with AR therefore can cause immunological, neurological, endocrine, embryotoxic, and many other effects.

The similarity in action of these compounds is thought to result from their structural similarity. Dioxin is composed of two benzene rings joined by two carbon-oxygen-carbon bonds on two adjacent carbons of each benzene ring. Dibenzofurans have two benzene rings joined by a carbon-oxygen-carbon bond and a carbon-carbon bond on two adjacent carbons of each benzene ring. Biphenyls consist of two benzene rings joined by a single carbon-carbon bond. To form the polychlorinated derivatives, chloro groups are attached at various locations, as designated in the names of the compounds. Benzene rings are planar, i.e., flat, in conformation. Because two adjacent carbons on each benzene ring are joined in dioxins and dibenzofurans, both benzene rings are held in the same plane, and the chloro groups are also in that plane. Therefore, these molecules are said to be coplanar. The coplanar structure appears to be essential for interaction with AR. The benzene rings in biphenyl can rotate relative to each other, unless there are added groups that interfere with rotation (such as 2,2',6,6'-chloro groups, which occupy the carbons immediately on both sides of the carbon-carbon bond joining the rings). PCB congeners that are able to form a coplanar molecule (and are called coplanar PCBs) can interact with AR when they are in that configuration. Therefore, coplanar PCBs are included among the COPCs with similar action to dioxins and dibenzofurans.

EPA 1998a recommends evaluating all polychlorinated dibenzo-p-dioxins and polychlorinated dibenzofurans (PCDD/PCDF) with chlorine molecules substituted in the 2,3,7, and 8 positions as carcinogens.

Because these contaminants have a common mechanism of action, it is assumed that their toxicity to biota is additive (WHO 1998, EPA 1998a). That is, the toxicity of all dioxins, dibenzofurans, and PCBs should be added. Furthermore, their relative potency as chronic toxins is assumed to be related to the degree of affinity for AR, which can be measured much more conveniently than chronic toxic effects. Toxicity equivalence factors (TEFs) have been proposed for several chlorinated dioxins, chlorinated dibenzofurans, and chlorinated biphenyls (WHO 1998, EPA 1998a), always using the toxicity of 3,4,7,8-tetrachlorodioxin (TCDD), the most potent chlorinated dioxin, as a TEF of 1.0.

Potential cancer risks associated with PCDDs/PCDFs will be evaluated using the toxicity equivalency approach. This approach assigns a relative toxicity to each of the seventeen 2,3,7,8-substituted PCDDs/PCDFs as compared to 2,3,7,8-TCDD. EPA 1998a recommends evaluating all polychlorinated dibenzo-p-dioxins and polychlorinated dibenzofurans (PCDD/PCDF) with chlorine molecules substituted in the 2, 3, 7, and 8 positions as carcinogens. Potential cancer risks associated with PCDDs/PCDFs will be evaluated using the toxicity equivalency approach. This approach assigns a relative toxicity, reported as a Toxicity Equivalency Factor (TEF), to each of the seventeen 2, 3, 7, 8-substituted PCDDs/PCDFs as compared to 2, 3, 7, 8-TCDD. Exposure concentrations are converted to equivalent concentrations of 2, 3, 7, 8-TCDD by multiplying the concentration by the appropriate TEF. This results in Toxicity Equivalent (TEQ) concentrations of each congener. The slope factor for 2,3,7,8-TCDD is used to evaluate risk from total TEQ concentration. TEFs are available from EPA (1989b) and the World Health Organization (WHO, 1997). Values from both agencies are listed below:

Dioxin /Furan Congener	TEF (unitless) USEPA	WHO
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1.0	1.0
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	0.5	1.0
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	0.1	0.1
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	0.1	0.1
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	0.1	0.1
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	0.01	0.01
1,2,3,4,5,7,8,9-Octachlorodibenzo(p)dioxin	0.001	0.0001
2,3,7,8-Tetrachlorodibenzofuran	0.1	0.1
1,2,3,7,8-Pentachlorodibenzofuran	0.05	0.05
2,3,4,7,8-Pentachlorodibenzofuran	0.5	0.5
1,2,3,4,7,8-Hexachlorodibenzofuran	0.1	0.1
1,2,3,6,7,8-Hexachlorodibenzofuran	0.1	0.1
1,2,3,7,8,9-Hexchlorodibenzofuran	0.1	0.1
2,3,4,6,7,8-Hexachlorodibenzofuran	0.1	0.1
1,2,3,4,6,7,8-Heptachlorodibenzofuran	0.01	0.01
1,2,3,4,7,8,9-Heptachlorodibenzofuran	0.01	0.01
1,2,3,4,6,7,8,9-Octachlorodibenzofuran	0.001	0.0001

The TEFs published by these two agencies are very similar. The primary differences are:

- The WHO assigns 1,2,3,7,8-Pentachlorodibenzo(p)dioxin a toxicity equal to 2,3,7,8-TCDD, while the EPA assigns this congener a TEF of 0.5 or half the toxicity of 2,3,7,8-TCDD.

- The WHO assigns 1,2,3,4,5,7,8,9-octachlorodibenzo(p)dioxin and furan a TEF of 0.0001, while the EPA assigns these congeners a TEF of 0.001.

At the direction of EPA Region 10 (1999e) potential cancer risks associated with PCDD/PCDF emissions will be estimated using both WHO and EPA TEFs. This evaluation will be presented as follows:

- WHO TEFs will be used to estimate potential cancer risks in the SLRA.

- WHO TEFS will be used to develop risk-based emission limits for the facility.

- Potential cancer risks estimated using EPA TEFs will be presented as part of the uncertainty evaluation presented in the SLRA.

Exposure concentrations are converted to equivalent concentrations of 2,3,7,8-TCDD by multiplying the concentration by the appropriate TEF. This results in TEQ concentrations of each congener. The slope factor for 2,3,7,8-TCDD is used to evaluate risk from total TEQ concentration. The TEF approach will be used along with the SF for 2,3,7,8-TCDD to estimate potential cancer risks from all exposure pathways (including nursing infant) except acute inhalation.

Toxicity values are not available for evaluating noncancer effects of PCDDs/PCDFs. EPA 1998a recommends evaluating potential noncancer hazards using the national average background exposure levels. As recommended, national average background exposure levels of 1 to 3 pg/kg-day for adult and child receptors and 60 pg/kg-day for nursing infants will be used to estimate potential noncancer hazards for PCDDs/PCDFs.

7.2.4. Polycyclic Aromatic Hydrocarbons (PAHs)

The toxicity equivalency approach will also be used for evaluating potentially carcinogenic PAHs. Seven PAHs [benzo(*a*)pyrene, benzo(*a*)anthracene, benzo(*b*)fluoranthene, benzo(*k*)fluoranthene, chrysene, dibenz(*a,h*)anthracene, indeno(1,2,3-*cd*)pyrene] are considered to be potentially carcinogenic. Adequate toxicity data are available to determine a cancer slope factor only for benzo(*a*)pyrene. Relative potency factors (RPFs) developed by EPA 1993a to be used in estimating cancer risks for all seven potentially carcinogenic PAHs are shown below:

Compound	RPF
Benzo(*a*)pyrene	1.0
Benzo(*a*)anthracene	0.1
Benzo(*b*)fluoranthene	0.1
Benzo(*k*)fluoranthene	0.01
Chrysene	0.001
Dibenz(*a,h*)anthracene	1.0
Indeno(1,2,3-*cd*)pyrene	0.1

This method will be applied to both oral and inhalation exposure pathways. RPFs are available from CalEPA for additional potentially carcinogenic PAHs and may be considered if PAHs are predicted to be a significant emission from the facility.

Noncancer effects will also be evaluated for PAHs that have available toxicity values. If PAHs are predicted to be a significant emission from the facility, surrogate toxicity values may be considered.

7.2.5. Polychlorinated Biphenyls (PCBs)

In addition to the TEFs for dioxin/furan congeners, EPA 1996b is implementing the use of TEFs for coplanar, dioxin-like PCBs. The same method of converting emissions for individual congeners to a 2,3,7,8-TCDD equivalent will be used for dioxin-like PCBs.

TEFs are available for coplanar PCBs from EPA (1998a) and WHO (1997). Values from both agencies are listed below:

	TEF (unitless)	
PCB Congener	EPA	WHO
3,4,4',5-Tetrachlorobiphenyl	--	0.0001
3,3',4,4'-Tetrachlorobiphenyl	0.0005	0.0001
2,3,3',4,4'-Pentachlorobiphenyl	0.001	0.0001
2,3,4,4',5-Pentachlorobiphenyl	0.0005	0.0005
2,3',4,4',5-Pentachlorobiphenyl	0.0001	0.0001
2',3,4,4',5-Pentachlorobiphenyl	0.001	0.0001
3,3',4,4',5-Pentachlorobiphenyl	0.1	0.1
2,3,3',4,4',5-Hexachlorobiphenyl	0.0005	0.0005
2,3,3',4,4',5'-Hexachlorobiphenyl	0.0005	0.0005
2,3',4,4',5,5'-Hexachlorobiphenyl	0.00001	0.00001

3,3',4,4',5,5'-Hexachlorobiphenyl	0.01	0.01
2,2',3,3',4,4',5-Heptachlorobiphenyl	0.0001	W
2,2',3,4,4',5,5'-Heptachlorobiphenyl	0.00001	W
2,3,3',4,4',5,5'-Heptachlorobiphenyl	0.0001	0.0001

-- no value W value withdrawn

The two sets of TEFs differ for six of the 14 coplanar PCBs. At the direction of EPA Region 10 (1999d) potential cancer risks associated with coplanar PCB emissions will be estimated using both WHO and EPA TEFs. This evaluation will be presented as follows:

- WHO TEFs will be used to estimate potential cancer risks in the SLRA.
- WHO TEFS will be used to develop risk-based emission limits for the facility.
- Potential cancer risks estimated using EPA TEFs will be presented as part of the uncertainty evaluation presented in the SLRA.

The TEF approach will be used along with the SF for 2,3,7,8-TCDD to estimate potential cancer risks from coplanar PCBs for all exposure pathways (including nursing infant) except acute inhalation.

Noncancer effects of coplanar PCBs will be evaluated for nursing infants using average background exposure concentrations. A reported range for the average background concentration of coplanar PCBs in human milk is 10-20 ng/L (higher levels have been measured) with a recommended point value of 13 ng/L (EPA 1999d).

Other PCBs will be evaluated using the most recent slope factor for PCBs. EPA 1996b recommends different slope factors for different exposure routes and chlorine contents. For this assessment, the most conservative toxicity values (slope factors from the high-risk persistence tier) will be used.

Noncancer effects will also be evaluated for PCBs according to EPA guidelines. Typically, the RfD for Aroclor 1254 will be used when congeners are > 4 chlorines comprise > 0.5% of total PCBs. Otherwise, the RfD for Aroclor 1016 will be used.

7.2.6. Chromium

Hexavalent chromium (Cr^{+6}) is the most toxic valence state of chromium and has been shown to be a human carcinogen via inhalation. Trivalent chromium (Cr^{+3}) has not been shown to be carcinogenic in either humans or laboratory animals. However, the mechanism of Cr^{+6} carcinogenically in the lung is believed to be its reduction to Cr^{+3} and its generation of reactive intermediates (Goyer 1996 in Klaassen et al. 1996). While chromium emitted from the melter is not likely to be in the hexavalent form, the PRA will conservatively assume that 100% of the facility emissions are hexavalent chromium (EPA 1998a).

For the FRA, chromium may be assumed to exist in the trivalent form. If this assumption is made in the FRA, the rationale will be provided at that time. Performance test data or design information may also be used to provide more realistic estimates of Cr^{+6}/Cr^{+3} emissions from the RPP-WTP.

7.2.7. Lead

The EPA has not derived a RfD/RfC or SF for lead. The potential for adverse health effects associated with exposure to lead will be characterized through comparison with the following media-specific health-based levels as recommended by EPA Region 6 (EPA 1998a).

Exposure Medium	EPA Region 6 Health-Based Target Level
Soil	100 mg/kg
Air	0.2 $\mu g/m^3$

The target level for soil is based on acceptable lead concentrations in blood. The EPA recommends that lead exposures be limited so that 95% of the sensitive subpopulation (children) will have blood lead concentrations below 10 μg/dL. Using the EPA (1994c) Integrated Exposure Uptake Biokinetic (IEUBK) Model to estimate lead levels in blood of children exposed to lead through diet, drinking water, air, and soil using standard default values, the model predicts that at least 95% of children exposed to a soil lead concentration of 400 mg/kg will have blood lead concentrations below 10 μg/dL. EPA Region 6 has incorporated a margin of safety by allowing only 25% of this 400 mg/kg threshold level as their recommended target level of 100 mg/kg.

The recommended target level of 0.2 $\mu g/m^3$ for air is 25% of the National Ambient Air Quality Standard (NAAQS) quarterly average air concentration of 1.5 $\mu g/m^3$ translated to an annual value of 0.9 $\mu g/m^3$.

7.2.8. Nickel

EPA (1998a) recommends that nickel be evaluated as an inhalation carcinogen because some forms of nickel, including nickel carbonyl, nickel subsulfide, and nickel refinery dust, are considered to be carcinogens. This is contrary to EPA's previous analysis of the toxicity of nickel emissions from hazardous waste combustion units because it was previously assumed that nickel can only be emitted as nickel oxide, which by itself, is not considered to be a carcinogen. Nickel oxide is a major component of nickel refinery dust (other major components include nickel subsulfide and nickel sulfate) which is identified as a potential human inhalation carcinogen. The components responsible for the carcinogenicity of nickel refinery dust have not been conclusively established. Therefore, nickel emissions will be evaluated as a potential carcinogen via the inhalation pathway using the inhalation unit risk factor for nickel refinery dust.

For exposure pathways other than inhalation, nickel has not been shown to be carcinogenic and will be evaluated as a noncarcinogen using the oral RfD for nickel soluble salts.

7.2.9. Particulate Assessment

Toxicity values are not available to quantitatively evaluate potential adverse health effects associated with inhaling particulates. Therefore, modeled annual average concentration of respirable particulates less than 10 microns in diameter (PM_{10}) will be compared to the average annual NAAQS for PM_{10} of 50 $\mu g/m^3$ (40 CFR 50.4-50.12). The 24-hour average PM_{10} concentrations will be compared with the NAAQS 24-hour level of 150 $\mu g/m^3$.

7.2.10. Radionuclide Assessment

HEAST (EPA 1995d) and EPA Federal Guidance Report 13 (EPA 1999b) are the sources for all carcinogenic toxicity values (slope factors) for radionuclides. Ingestion, inhalation, and external exposure slope factors for radionuclides are provided in Table 4-3. Risks will be calculated using slope factors that include the potential toxicity of daughter products.

7.2.11. Nursing Infant Assessment

Potential infant exposure to PCDD/PCDFs, coplanar PCBs, and selected radionuclides (Sr^{90}, I^{129}, Cs^{134}, Cs^{137}) in human breast milk will be evaluated in the SLRA. The interpretation of infant exposures is limited by the lack of infant dose-response data (EPA 1998a). Potential infant exposures will be evaluated in the SLRA two ways; by comparison to background exposures, and by the calculation of a cancer risk value.

7.2.11.1 Comparison to Background

Because the possible impact of a brief early life exposure to these chemicals is unknown, the most appropriate method for evaluating the potential for adverse health effects to exposed infants is through a comparison to average background exposures.

Polychlorinated Dibenzo-p-dioxins and Polychlorinated Dibenzofurans

EPA (1998d) recommends that risk assessments include a comparison of exposures to PCDDs and PCDFs with a national average background exposure level for infants of 60 pg TEQ/kg/day. This value is a modeled ADD based on a measured average concentration of 16 ppt 2,3,7,8-TCDD TEQ in the lipid portion of breast milk. This comparison to background will utilize EPA TEF (rather than WHO TEFs) because historic data available for background concentrations and exposures are based on the EPA TEF values.

Polychlorinated Biphenyls (PCBs)

Possible effects of coplanar PCBs will be evaluated for nursing infants using average background exposure concentrations. A reported range for the average background concentration of coplanar PCBs in human milk is 10-20 ng/L (higher levels have been measured) with a recommended point value of 13 ng/L (EPA 1999d).

Radionuclides

Background concentrations of the four radionuclides of interest are not currently available. If background values become available they will be included in the SLRA. Background data identified by September 2000 will be included in the draft PRA. Background data identified up to four weeks prior to subsequent submissions (e.g., final PRA, FRA) will be included in these later documents.

7.2.11.2 Estimated Cancer Risk

Incremental Lifetime Cancer Risks are generally estimated from the total lifetime exposure or dose of a chemical (LADD) and the cancer slope factor. Cancer slope factors represent an upper-bound estimate of the probability of developing cancer per unit dose of a chemical. The possible impact of a brief early life exposure alone to a potential carcinogen is unknown. Per EPA guidance (EPA 1998a), an infant

cancer risk will be calculated for a 1 year exposure (1 year ADD) to PCDDs/PCDFs, coplanar PCBs and the four potentially bioaccumulative ROPCs.

Polychlorinated Dibenzo-p-dioxins and Polychlorinated Dibenzofurans

Infant cancer risks will be estimated using the same TEF/TEQ approach described previously. Because PCDDs/PCDFs act as a cancer promoter rather than a cancer initiator, exposures early in life may not be as important as those that occur later (i.e., following initiation).

Polychlorinated Biphenyls (PCBs)

Infant cancer risks will be estimated using the same TEF/TEQ approach described for coplanar PCBs previously.

Radionuclides

Risks will be estimated using slope factors that include the potential toxicity of daughter products as described for adult exposures.

7.2.12. Acute Effects Assessment

In addition to chronic effects, direct inhalation of airborne COPCs and ROPCs (vapor and particulate) will be evaluated for potential acute effects. Modeled short-term, one-hour maximum air concentrations will be compared to acute inhalation exposure criteria (AIEC) to evaluate the potential for COPCs/ROPCs emitted from the RPP-WTP to pose an acute health hazard to potentially exposed receptors. Acute toxicity data for COPCs/ROPCs will be obtained according to the following hierarchy:

1. Values from NCEA (as provided by EPA Region X)
2. Acute exposure guideline levels (AEGL-1)
3. Emergency response planning guidelines (ERPG-1)
4. Acute reference exposure levels (ARELs) from California EPA
5. Temporary emergency exposure limits (TEEL-1)
6. Subcommittee on consequence assessment and protective actions (SCAPA) toxicity-based approach (DOE 1997)

The most up-to-date references currently available for items 2-5 above are:

- AEGL-1: (EPA Region X and 10-30-99 *Federal Register)*
- ERPG-1: http://www.scapa.bnl.gov/scapawl.htm
- AREL: http://oehha.ca.gov/scientific/relnums.htm
- TEEL-1: http://tis.eh.doe.gov/web/chem_safety/teel.html

Updated references will be used for these acute toxicity data when the references become available.

For acute exposures to ROPCs, derived air concentrations based on the ALI will be used. These derived air concentrations for acute exposure comparison will be the concentration of the radionuclide in air that,

under continuous exposure for a one-hour period, would produce a total effective dose equivalent of 50 millirem (0.5 millisievert).

Federal guidance for occupational radiation protection specifies the following primary guides for limiting occupational exposure (USEPA, 1988):

Receptor	Primary Guide for Limiting Exposure
Adults	5000 mrem/yr
Children under 18	500 mrem/yr
Pregnant Women	50 mrem/gestation period

7.3. Risk Characterization

This section of the Work Plan presents how the potential noncarcinogenic and carcinogenic health risks for the exposure scenarios identified in the exposure assessment (see section 7.1) will be characterized. Potential health risks will be characterized separately for noncarcinogenic and carcinogenic endpoints, and chemical (nonradiological) and radiological cancer risks will be evaluated and presented separately. These endpoints will be characterized by comparing calculated dose levels to maximum "acceptable" doses. Potential noncarcinogenic health risks will be determined using the hazard quotient (HQ)/HI approach that defines the relative hazard based on the ratio of the estimated lifetime ADD (LADD) to the acceptable intake level (i.e., the RfD). Potential carcinogenic health risks will be determined based on the probability that an individual may contract cancer in a lifetime from the estimated LADD. COPCs having both carcinogenic and noncarcinogenic health effects will be evaluated for both endpoints. ROPCs will be evaluated for carcinogenic health effects.

The term risk means the probability of harm. More precisely, risk is the excess rate of cancer. By contrast, hazard refers to adverse noncarcinogenic health effects. In many cases, in this Work Plan, the use of risk generically means carcinogenic and/or noncarcinogenic consequences. Methodologies that will be used to characterize potential risks for the RPP-WTP are presented below.

7.3.1. Noncarcinogenic Effects

Noncarcinogenic health risks are typically characterized using an "HQ" and "HI" approach (EPA 1989b). The HQ is the ratio of the calculated ADD to the maximally allowable "safe" dose (i.e., EPA reference levels such as the RfD or similar value). The equation used to calculate the HQ for a chemical is presented below.

$$HQ = \frac{ADD}{RfD}$$

where

HQ = hazard quotient (unitless)
ADD = average daily dose (mg COPC/kg-day)
RfD = reference dose (mg COPC/kg-day)

An HQ of 0.25 or less indicates that the chemical-specific ADD for a particular pathway is below the level associated with an adverse health effect.

EPA 1998a recommends an HQ goal of 0.25 for evaluating emission rates and this risk goal will be applied here. Additive noncarcinogenic health effects can be evaluated when exposure to more than one chemical occurs by using the HI approach (see section 7.2.1).

Additivity of Noncarcinogenic Effects

Multiple chemical exposures can result in synergism, antagonism, and/or additivity of biological responses when the chemicals act on similar target organs or when they are metabolized by the same enzymatic pathways. It is appropriate in risk assessment to evaluate the possible health effects associated with multiple chemical exposures at a site. EPA risk assessment guidelines (EPA 1989b) state that additivity of noncarcinogenic health effects should only be considered if the chemicals have the same toxicological endpoint (e.g., organ or enzyme systems). Note that the same toxicological endpoint also implies the same mechanism of action. Additivity for all chemicals will initially be assumed to occur regardless of the toxicological endpoint. This approach is likely to overestimate the true human health risks associated with exposure to the COPCs/ROPCs since many chemicals may act on different target organs. If the target hazard index (HI) is exceeded, a segregation of the HI by toxicological endpoint will be considered. In addition to multiple chemicals, receptors may be exposed via more than one pathway. As the EPA (1989b) notes:

"There are two steps required to determine whether risks or hazard indices for two or more pathways should be combined for a single exposed individual or group of individuals. The first is to identify reasonable exposure pathway combinations. The second is to examine whether it is likely that the same individuals would consistently face the "reasonable maximum exposure" for more than one pathway."

The simplified equation for calculating a generic HI is presented below.

$$\text{Hazard Index} = \frac{ADD_1}{RfD_1} + \frac{ADD_2}{RfD_2} + \ldots + \frac{ADD_n}{RfD_n}$$

An HI of 0.25 or less indicates that levels of exposure are acceptable. HI will be calculated for all COPCs and all pathways for a given receptor.

7.3.2. Carcinogenic Risks

Carcinogenic health risks are defined in terms of a probability that an individual may develop cancer as a result of exposure to a given chemical or radionuclide at a given concentration (EPA 1989b). The incremental probability of developing cancer (i.e., the excess cancer risk) is the additional risk above and beyond the cancer risk an individual would face in the absence of exposures characterized in this assessment. The excess cancer risk will be determined for each potentially carcinogenic COPC or ROPC using the total LADD from all pathways and cancer slope factors as described below.

Excess Cancer Risk = LADD x SF for all pathways except external exposure to air

Excess Cancer Risk = LADD x RF for external exposure to air,

where

LADD = Lifetime average daily dose, with units of

- (mg/kg-day) for COPCs
- pCi for ingestion and inhalation of ROPCs

- (pCi-yr/g) for external exposure to ionizing radiation in soil (for ROPCs)
- (Bq-secs/m^3) for external exposure to ionizing radiation in air (for ROPCs)

SF = Cancer slope factor, with units of

- $(mg/kg\text{-}day)^{-1}$ for COPCs
- $(pCi)^{-1}$ for ingestion and inhalation of ROPCs
- $(pCi\text{-}yr/g)^{-1}$ for external exposure to ionizing radiation in soil (for ROPCs)

RF = Cancer incidence risk factor $(Bq\text{-}secs/m^3)^{-1}$ for external exposure to ionizing radiation in air (for ROPCs) (EPA 1999c).

Oral and inhalation slope factors are provided in Table 4-1 (organic COPCs), Table 4-2 (inorganic COPCs), and Table 4-3 (ROPCs). The slope factors provided for radionuclides are taken from EPA (1995d) and include toxicity due to daughter products. Use of these "+D" slope factors for radionuclides is conservative because it assumes the daughter species are in equilibrium with the parent. For many radionuclides this equilibrium may require thousands of years.

The LADD and slope factor are multiplied, yielding a dimensionless value that represents the probability of developing cancer within a lifetime (70 years) due to the chemical or radiological dose (LADD) received by a person. For example, a theoretical risk value of 0.0001 or 1×10^{-4} is referred to as a probability of 1 in 10,000 of developing cancer. As with the HIs, the theoretical excess cancer risk will be presented for each scenario on a chemical/radiological-specific, pathway-specific, and total risk basis. In addition, chemical and radiological risks will be presented separately. The theoretical excess cancer risk may be compared to both the *de minimus* lifetime cancer risk rate of 1×10^{-6} (i.e., negligible risk) and the acceptable regulatory risk range of 1×10^{-6} to 1×10^{-4} (OSWER Directive 9355.0-30 EPA 1991a). The target risk goal for emissions from the RPP-WTP is 1×10^{-5}.

7.4. Uncertainty Assessment

Numerous sources of uncertainty inherent in the risk assessment process exist. Some degree of uncertainty is introduced into the assessment each time an assumption is made. Many assumptions have valid and strong scientific bases, whereas others are estimates usually represented by a range of values.

The purpose of the uncertainty analysis is to identify and discuss areas of uncertainty associated with the quantitative estimates of risk for COPCs and ROPCs. This discussion serves to place the risk estimates in proper perspective by fully specifying the assumptions and qualitatively evaluating the uncertainties of the assessment.

A table providing a qualitative summary of all potential sources of uncertainty and the potential effect of that uncertainty on the risk estimates will be included in the PRA. The key variables and assumptions that contribute most to uncertainty will be identified and further addressed in the FRA. A qualitative discussion of the uncertainties associated with each component of the risk assessment will be provided including:

- **Site Characterization.** The degree of confidence in the current and future land use assumptions will be addressed.

- **Exposure Assessment.** Uncertainties surrounding fate and transport modeling, and the assignment of exposure parameter values will be addressed, including any excluded pathways such as dermal contact with soil and the handling of plants.

- **Toxicity Assessment**. Uncertainties surrounding the toxicity values (RfDs and SFs), weight-of-evidence classifications, toxicity value data gaps, conversions of toxicity values from concentration to dose (e.g., UR to SF, RfC to inhalation RfD), and route-to-route extrapolations will be addressed.

- **Risk Characterization**. Uncertainties surrounding the practice of summing HIs and risks across chemicals and pathways. The uncertainties associated with the evaluation of chemical and radionuclide emissions will also be addressed.

The four sources of uncertainty listed are unique to the human health risk assessment. Uncertainties associated with COPC/ROPC selection, emission rates, and environmental modeling are provided in previous sections.

To aid in the process of moving the dangerous waste permit process forward, special attention will be given to uncertainties associated with estimated risk that exceeds risk goals.

7.5. Summary for Screening Human Health Risk Assessment

Risks to human health from the potential emission of COPCs and ROPCs result from (1) exposure to the COPC/ROPC and (2) the toxicity of the COPC/ROPC. The screening HHRA utilizes the estimated emission rates (section 5) and results of fate and transport modeling (section 6) to calculate potential human exposure to COPCs and ROPCs. This exposure information is combined with toxicity data to estimate the potential for adverse effects to human populations in the vicinity of the RPP-WTP.

The PRA will use conservative exposure assumptions to compensate for the high level of uncertainty associated with conducting a risk assessment for a facility that is still in the design phase. The PRA will include a qualitative uncertainty analysis.

The FRA will focus on COPCs and ROPCs that exceed risk goals in the PRA and may utilize additional site-specific emission, fate and transport, and exposure data collected after completion of the PRA.

Figure 7-1. Human Health Conceptual Exposure Model



Hanford ERA - 014

Best Available Copy

Figure 7-2. Exposure Distances (Far-Field) from the RPP-WTP at the Hanford Site



Figure 7-3. Exposure Distances (Near-Field) from the RPP-WTP at the Hanford Site



Best Available Copy

Figure 7-4 Hanford Site Existing Land Use Map – 1996



WDFW	Washington Department of Fish and Wildlife
USFWS	U.S. Fish and Wildlife Service
LIGO	Laser Interferometer Gravitational Wave Observatory
HAMMER	Hazardous Materials Management and Emergency Response
WPPSS	Washington Public Power Supply System

Hanford ERA - 013

Best Available Copy

Figure 7-5 Hanford Site Projected Land Use Map – 2046



BHI:rpp 04/23/98 clup/prefalt.aml Database: 07-DEC-1998

Hanford ERA - 014

Best Available Copy

Table 7-1 Exposure Scenarios

Receptor	Receptor Location	Exposure Pathways	Reference Used to Define Exposures
Plausible Current Land Use Scenarios			
Hanford Site Industrial Worker[a]	Point of maximum exposure	Inhalation of emissions[b] Ingestion of soil Inhalation of resuspended soil Ingestion of drinking water[c] External exposure to radionuclides in air External exposure to radionuclides in soil	DOE-RL 1995
Resident Adult and Child	Point of maximum exposure at Hanford Site boundary	Inhalation of emissions[b] Ingestion of soil Inhalation of resuspended soil Ingestion of homegrown produce Ingestion of drinking water[c] External exposure to radionuclides in air External exposure to radionuclides in soil	EPA 1998a
Native American Subsistence Hunter/Gatherer Adult and Child	Point of maximum exposure at Hanford Site boundary	Inhalation of emissions[b] Ingestion of soil Inhalation of resuspended soil Ingestion of homegrown/wild produce Ingestion of wild game Ingestion of wildfowl and eggs[d] Ingestion of milk Ingestion of drinking water[c] External exposure to radionuclides in air External exposure to radionuclides in soil Inhalation of VOCs and SVOCs in sweat lodge[e] Dermal absorption in sweat lodge[e]	Harris and Harper 1997
Native American Subsistence Fisher Adult and Child	Point of maximum exposure at Hanford Site boundary	Inhalation of emissions[b] Ingestion of soil Inhalation of resuspended soil Ingestion of homegrown/wild produce Ingestion of fish (fillet and fish parts) Ingestion of drinking water[c] External exposure to radionuclides in air External exposure to radionuclides in soil Inhalation of VOCs and SVOCs in sweat lodge[e] Dermal absorption in sweat lodge[e]	Harris and Harper 1997
Plausible Future Land Use Scenarios			
Hanford Site Industrial Worker[a]	Point of maximum exposure	Inhalation of emissions[b] Ingestion of soil Inhalation of resuspended soil Ingestion of drinking water[c] External exposure to radionuclides in air External exposure to radionuclides in soil	DOE-RL 1995

Table 7-1 Exposure Scenarios

Receptor	Receptor Location	Exposure Pathways	Reference Used to Define Exposures
Resident Adult and Child	Point of maximum exposure at Hanford Site boundary	Inhalation of emissions[b] Ingestion of soil Inhalation of resuspended soil Ingestion of homegrown produce Ingestion of drinking water[c] External exposure to radionuclides in air External exposure to radionuclides in soil	EPA 1998a
Native American Subsistence Hunter/Gatherer Adult and Child	Point of maximum exposure	Inhalation of emissions[b] Ingestion of soil Inhalation of resuspended soil Ingestion of homegrown/wild produce Ingestion of wild game Ingestion of wildfowl and eggs[d] Ingestion of milk Ingestion of drinking water[c] External exposure to radionuclides in air External exposure to radionuclides in soil Inhalation of VOCs and SVOCs in sweat lodge[e] Dermal absorption in sweat lodge[e]	Harris and Harper 1997
Native American Subsistence Fisher Adult and Child	Point of maximum exposure	Inhalation of emissions[b] Ingestion of soil Inhalation of resuspended soil Ingestion of homegrown/wild produce Ingestion of fish (fillet and fish parts) Ingestion of drinking water[c] External exposure to radionuclides in air External exposure to radionuclides in soil Inhalation of VOCs and SVOCs in sweat lodge[e] Dermal absorption in sweat lodge[e]	Harris and Harper 1997
Worst-Case Future Land Use Scenarios			
Resident Adult and Child	Point of maximum exposure	Inhalation of emissions[b] Ingestion of soil Inhalation of resuspended soil Ingestion of homegrown produce Ingestion of drinking water[c] External exposure to radionuclides in air External exposure to radionuclides in soil	EPA 1998a
Subsistence Farmer Adult and Child	Point of maximum exposure	Inhalation of emissions[b] Ingestion of soil Inhalation of resuspended soil Ingestion of homegrown produce Ingestion of homegrown beef and milk Ingestion of homegrown chicken and eggs Ingestion of homegrown pork Ingestion of drinking water[c] External exposure to radionuclides in air External exposure to radionuclides in soil	EPA 1998a

Table 7-1 Exposure Scenarios

Receptor	Receptor Location	Exposure Pathways	Reference Used to Define Exposures
Subsistence Fisher Adult and Child	Point of maximum exposure at Columbia River	Inhalation of emissions[b] Ingestion of soil Inhalation of resuspended soil Ingestion of homegrown produce Ingestion of drinking water[c] Ingestion of fish External exposure to radionuclides in air External exposure to radionuclides in soil	EPA 1998a
Resident Nursing Infant[a]	Point of maximum exposure	Ingestion of breast milk	EPA 1998a
Native American Subsistence Hunter/Gatherer Adult and Child	Point of maximum exposure	Inhalation of emissions[b] Ingestion of soil Inhalation of resuspended soil Ingestion of homegrown/wild produce Ingestion of wild game Ingestion of wildfowl and eggs[d] Ingestion of milk Ingestion of drinking water[c] External exposure to radionuclides in air External exposure to radionuclides in soil Inhalation of VOCs and SVOCs in sweat lodge[e] Dermal absorption in sweat lodge[e]	Harris and Harper 1997
Native American Subsistence Fisher Adult and Child	Point of maximum exposure	Inhalation of emissions[b] Ingestion of soil Inhalation of resuspended soil Ingestion of homegrown/wild produce Ingestion of fish (fillet and fish parts) Ingestion of drinking water[c] External exposure to radionuclides in air External exposure to radionuclides in soil Inhalation of VOCs and SVOCs in sweat lodge[e] Dermal absorption in sweat lodge[e]	Harris and Harper 1997
Native American Resident Nursing Infant[a]	Point of maximum exposure	Ingestion of breast milk	Harris and Harper 1997 and EPA 1998a
Acute Exposure Scenario			
Acute Exposure	Point of maximum short-term exposure	Inhalation of emissions[b]	EPA 1998a

[a] Special subpopulation to be evaluated.

[b] Includes direct inhalation of vapor phase and particulate emissions.

[c] Surface water concentrations will be used to evaluate the ingestion of drinking water pathway.

[d] Consumption rates for the ingestion of poultry and eggs for the Native American will be based on a partitioning of the consumption rate of wild fowl (which includes bird and eggs), as presented in Harris and Harper 1997.

[e] Inhalation and dermal absorption in the sweat lodge is evaluated for adult Native Americans only.

Table 7-2 Exposure Parameters for Residential Scenarios

Exposure Pathway	Parameter	Description	Units	Value[a]	Comment
Inhalation of particulates and vapors	IR	Inhalation rate of air	m^3/hr	0.63 0.3	Value for adults Value for children
	ET	Exposure Time	hr/day	24	Assumption
	EF	Exposure Frequency	day/year	350	Assumed 2 weeks away from contaminated area per year
	ED	Exposure Duration	year	40 30 6	Value for adult farmer Value for adult fisher/resident Value for child in all scenarios
	BW	Body Weight	kg	70 15	Value for adults Value for children
	AT	Averaging Time	year	70 ED	Value for carcinogenic effects Equal to exposure duration for noncarcinogenic effects
Soil Ingestion	CR_{soil}	Consumption rate of soil	kg/day	0.0001[b] 0.0002[b]	Value for adults Value for children
	F_{soil}	Fraction of soil ingested that is contaminated	unitless	1.0	Assumption
	EF	Exposure Frequency	day/year	350	Assumed 2 weeks away from contaminated area per year
	ED	Exposure Duration	year	40 30 6	Value for adult farmer Value for adult fisher/resident Value for child in all scenarios
	BW	Body Weight	kg	70 15	Value for adults Value for children
	AT	Averaging Time	year	70 ED	Value for carcinogenic effects Equal to exposure duration for noncarcinogenic effects

Table 7-2 Exposure Parameters for Residential Scenarios

Exposure Pathway	Parameter	Description	Units	Value[a]	Comment
Inhalation of Resuspended Soil	IR	Inhalation rate of air	m^3/hr	0.63 0.3	Value for adults Value for children
	ET	Exposure Time	hr/day	24	Assumption
	EF	Exposure Frequency	day/year	350	Assumed 2 weeks away from contaminated area per year
	ED	Exposure Duration	year	40 30 6	Value for adult farmer Value for adult fisher/resident Value for child in all scenarios
	BW	Body Weight	kg	70 15	Value for adults Value for children
	AT	Averaging Time	year	70 ED	Value for carcinogenic effects Equal to exposure duration for noncarcinogenic effects
External Exposure to Ionizing Radiation in Soil	Se	Shielding Factor	unitless	0.2[c]	Assume 20% shielding
	Te	Exposure Time Factor	unitless	1.0[c]	Assumption (24 hours/24 hours)
	EF	Exposure Frequency	days/year	350	Assumed 2 weeks away from contaminated area per year
	ED	Exposure Duration	years	40 30 6	Value for adult farmer Value for adult fisher/resident Value for child in all scenarios
External Exposure to Ionizing Radiation in Air	EF	Exposure Frequency	day/year	350	Assumed 2 weeks away from contaminated area per year
	ED	Exposure Duration	year	40 30 6	Value for adult farmer Value for adult fisher/resident Value for child in all scenarios
	AF	Age Factor	unitless	1[d] 1.3[d]	No adjustment for adults Adjustment factor for child

Table 7-2 Exposure Parameters for Residential Scenarios

Exposure Pathway	Parameter	Description	Units	Value[a]	Comment
Produce Consumption	CR_{ug}	Consumption rate of unprotected aboveground produce	kg/kg-day DW	0.0003 0.00042	Value for adults Value for children
	CR_{pp}	Consumption rate of protected aboveground produce	kg/kg-day DW	0.00057 0.00077	Value for adults Value for children
	CR_{bg}	Consumption rate of below ground produce	kg/kg-day DW	0.00014 0.00022	Value for adults Value for children
	F_{ug}	Fraction of ingested produce that is contaminated	unitless	0.25 1.0 0.25	Value for resident Value for farmer Value for fisher
	EF	Exposure Frequency	day/year	350	Assumed 2 weeks away from contaminated area per year
	ED	Exposure Duration	year	40 30 6	Value for adult farmer Value for adult fisher/resident Value for child in all scenarios
	BW	Body Weight	kg	70 15	Value for adults Value for children
	AT	Averaging Time	year	70 ED	Value for carcinogenic effects Equal to exposure duration for noncarcinogenic effects
Ingestion of beef, milk, poultry, eggs, and pork	CR_{beef}	Consumption rate of beef	kg/kg-day FW	0.00114 0.00051	Value for adults Value for children
	CR_{milk}	Consumption rate of milk	kg/kg-day FW	0.00842 0.01857	Value for adults Value for children
	$CR_{poultry}$	Consumption rate of poultry	kg/kg-day FW	0.00061 0.000425	Value for adults Value for children
	CR_{eggs}	Consumption rate of eggs	kg/kg-day FW	0.00062 0.000438	Value for adults Value for children
	CR_{pork}	Consumption rate of pork	kg/kg-day FW	0.00053 0.000398	Value for adults Value for children
	F	Fraction of ingested food item that is contaminated	unitless	1.0	Assumption
	EF	Exposure Frequency	day/year	350	Assumed 2 weeks away from contaminated area per year

Table 7-2 Exposure Parameters for Residential Scenarios

Exposure Pathway	Parameter	Description	Units	Value[a]	Comment
Ingestion of beef, milk, poultry, eggs, and pork (cont'd)	ED	Exposure Duration	year	40 30 6	Value for adult farmer Value for adult fisher/resident Value for child in all scenarios
	BW	Body Weight	kg	70 15	Value for adults Value for children
	AT	Averaging Time	year	70 ED	Value for carcinogenic effects Equal to exposure duration for noncarcinogenic effects
Ingestion of fish	CR_{fish}	Consumption rate of fish	kg/kg-day FW	0.00117 0.000759	Value for adults Value for children
	CR_{organs}	Consumption rate of fish parts (e.g., head, fins, tails, etc.)	kg/kg-day FW	0.0	No consumption of fish parts
	F_{fish}	Fraction of ingested fish that are contaminated	unitless	1.0	Assumption
	EF	Exposure Frequency	day/year	350	Assumed 2 weeks away from contaminated area per year
	ED	Exposure Duration	year	30 6	Value for adult fisher Value for child fisher
	BW	Body Weight	kg	70 15	Value for adults Value for children
	AT	Averaging Time	year	70 ED	Value for carcinogenic effects Equal to exposure duration for noncarcinogenic effects

Table 7-2 Exposure Parameters for Residential Scenarios

Exposure Pathway	Parameter	Description	Units	Value[a]	Comment
Ingestion of drinking water[e]	CR_{dw}	Consumption rate of drinking water	L/day	1.4 0.67	Value for adults Value for children
	F_{dw}	Fraction of ingested drinking water that is contaminated	unitless	1.0	Assumption
	EF	Exposure Frequency	day/year	350	Assumed 2 weeks away from contaminated area per year
	ED	Exposure Duration	year	40 30 6	Value for adult farmer Value for adult fisher/resident Value for child in all scenarios
	BW	Body Weight	kg	70 15	Value for adults Value for children
	AT	Averaging Time	year	70 ED	Value for carcinogenic effects Equal to exposure duration for noncarcinogenic effects
Infant Ingesting Breast Milk[f]	h	Half-life of dioxin/furan/PCB or isotope in adult	day	2,555 isotope-specific	Upper bound of 5- to 7-year range is selected for dioxins Half-life of isotopes, in days
	f_1	Fraction of ingested COPC/ROPC that is stored in fat	unitless	0.9	EPA OSW recommendation
	f_2	Fraction of mother's weight that is fat	unitless	0.3	EPA OSW recommendation
	f_3	Fraction of mother's breast milk that is fat	unitless	0.04	EPA OSW recommendation
	f_4	Fraction of ingested COPC/ROPC that is absorbed	unitless	0.9	EPA OSW recommendation
	IR_{milk}	Ingestion rate of breast milk by infant	kg/day	0.8	Midpoint of range from 677 to 922 ml/day with assumed density >1 g/ml
	ED	Exposure duration	year	1.0	EPA OSW recommendation

Table 7-2 Exposure Parameters for Residential Scenarios

Exposure Pathway	Parameter	Description	Units	Value[a]	Comment
Infant Ingesting Breast Milk[f] (cont'd)	BW_{infant}	Body weight of infant	kg	10	Mean of 6 mo. to 1 year old infant's weight ranging from 9.1 to 11.3 kg
	AT	Averaging Time	year	1	EPA OSW recommendation acknowledging that this value may overestimate carcinogenic risks by a factor of 70
	EF	Exposure Frequency	day/year	365	Infant ingesting breast milk every day

[a] EPA default exposure parameters (EPA 1998a), unless otherwise noted.

[b] EPA Region X personal communication (EPA 1999e).

[c] RAGS, Part B (EPA 1991b).

[d] Saito, et al., 1998.

[e] Surface water concentrations will be used to evaluate the ingestion of drinking water pathway.

[f] Dioxins, furans, coplanar PCBs, and radionuclides are evaluated for the infant's ingestion of breast milk.

Table 7-3 Summary of Exposure Parameters for Native American Subsistence Scenarios[a]

Exposure Pathway	Parameter	Description	Units	Value[b]	Comment
Inhalation of particulates and vapors	IR	Inhalation rate of air	m^3/hr	0.833[c] 0.417[c]	Value for adults Value for children
	ET	Exposure Time	hr/day	24[c]	Assumption
	EF	Exposure Frequency	day/year	365	Exposure every day
	ED	Exposure Duration	year	70 6[d]	Value for adults Value for children
	BW	Body Weight	kg	70 15[d]	Value for adults Value for children
	AT	Averaging Time	year	70 ED	Value for carcinogenic effects Equal to exposure duration for noncarcinogenic effects
Soil Ingestion	CR_{soil}	Consumption rate of soil	kg/day	0.0002[e]	Value for adults and children
	F_{soil}	Fraction of soil ingested that is contaminated	unitless	1.0	Assumption
	EF	Exposure Frequency	day/year	365	Exposure every day
	ED	Exposure Duration	year	70 6[d]	Value for adults Value for children
	BW	Body Weight	kg	70 15[d]	Value for adults Value for children
	AT	Averaging Time	year	70 ED	Value for carcinogenic effects Equal to exposure duration for noncarcinogenic effects

Table 7-3 Summary of Exposure Parameters for Native American Subsistence Scenarios[a]

Exposure Pathway	Parameter	Description	Units	Value[b]	Comment
Inhalation of Resuspended Soil	IR	Inhalation rate of air	m^3/hr	0.833[c] 0.417[c]	Value for adults Value for children
	ET	Exposure Time	hr/day	24[c]	Assumption
	EF	Exposure Frequency	day/year	365	Exposure every day
	ED	Exposure Duration	year	70 6[d]	Values for adults Values for children
	BW	Body Weight	kg	70 15[d]	Values for adults Values for children
	AT	Averaging Time	year	70 ED	Value for carcinogenic effects Equal to exposure duration for noncarcinogenic effects
External Exposure to Ionizing Radiation in Soil	Se	Shielding Factor	unitless	0.2[f]	Assumed 20% shielding
	Te	Exposure Time Factor	unitless	1.0[f]	Exposure for 24 hr each day
	EF	Exposure Frequency	day/year	365	Exposure every day
	ED	Exposure Duration	year	70 6[d]	Value for adults Value for children
External Exposure to Ionizing Radiation in Air	EF	Exposure Frequency	day/year	365	Exposure every day
	ED	Exposure Duration	year	70 6[d]	Value for adults Value for children
	AF	Age Factor	unitless	1[g] 1.3[g]	No adjustment for adults Adjustment factor for children
Produce Consumption	CR_{ug}	Consumption rate of unprotected aboveground produce	kg/kg-day DW	0.00575[h] 0.02683[h]	Value for hunter/gatherer adults, based on 402.5 g/day[h] Value for hunter/gatherer children, based on 402.5 g/day[h]
	CR_{pp}	Consumption rate of protected aboveground produce	kg/kg-day DW	0.0[i]	Value for hunter/gatherer adults and children[i]

Table 7-3 Summary of Exposure Parameters for Native American Subsistence Scenarios[a]

Exposure Pathway	Parameter	Description	Units	Value[b]	Comment
Produce Consumption (cont'd)	CR_{bg}	Consumption rate of belowground produce	kg/kg-day DW	0.00245[j] 0.01143[j]	Value for hunter/gatherer adults, based on 171.5 g/day[j] Value for hunter/gatherer children, based on 171.5 g/day[j]
	F_{ag}	Fraction of ingested produce that is contaminated	unitless	1.0	Assumed
	EF	Exposure Frequency	day/year	365	Exposure every day
	ED	Exposure Duration	year	70 6[d]	Value for hunter/gatherer adults Value for hunter/gatherer children
	BW	Body Weight	kg	70 15[d]	Value for hunter/gatherer adults Value for hunter/gatherer children
	AT	Averaging Time	year	70 ED	Value for carcinogenic effects Equal to exposure duration for noncarcinogenic effects
Ingestion of game, milk, poultry, and eggs	CR_{game}	Consumption rate of beef, elk, and other game tissue	kg/kg-day FW	0.00357 0.01667	Value for hunter/gatherer adults based on 250 g/day of meat Value for hunter/gatherer children based on 250 g/day of meat
	CR_{milk}	Consumption rate of milk	kg/kg-day FW	0.00842[k] 0.01857[k]	Value for hunter/gatherer adults[k] Value for hunter/gatherer children[k]

Table 7-3 Summary of Exposure Parameters for Native American Subsistence Scenarios[a]

Exposure Pathway	Parameter	Description	Units	Value[b]	Comment
Ingestion of game, milk, poultry, and eggs (cont'd)	$CR_{poultry}$	Consumption rate of poultry	kg/kg-day FW	0.000329[l]	Value for hunter/gatherer adults, based on 23 g/day[l]
				0.001533[l]	Value for hunter/gatherer children, based on 23 g/day[l]
	CR_{eggs}	Consumption rate of eggs	kg/kg-day FW	0.0003[l]	Value for hunter/gatherer adults, based on 21 g/day[l]
				0.0014[l]	Value for hunter/gatherer children, based on 21 g/day[l]
	F	Fraction of ingested food item that is contaminated	unitless	1.0	Assumption
	EF	Exposure Frequency	day/year	365	Exposure every day
	ED	Exposure Duration	year	70	Value for hunter/gatherer adults
				6[d]	Value for hunter/gatherer children
	BW	Body Weight	kg	70	Value for hunter/gatherer adults
				15[d]	Value for hunter/gatherer children
	AT	Averaging Time	year	70	Value for carcinogenic effects
				ED	Equal to exposure duration for noncarcinogenic effects

Table 7-3 Summary of Exposure Parameters for Native American Subsistence Scenarios[a]

Exposure Pathway	Parameter	Description	Units	Value[b]	Comment
Ingestion of fish	CR_{fish}	Consumption rate of fish	kg/kg-day FW	0.00771	Value for fisher adults, based on 540 g/day. This same consumption rate is assumed for fisher children.[m]
	CR_{organs}	Consumption rate of fish parts (e.g., head, fins, tails, etc.)	kg/kg-day FW	0.000771	Value for fisher adults, based on 54 g/day. This same consumption rate is assumed for fisher children.[m]
	F_{fish}	Fraction of ingested fish that are contaminated	unitless	1.0	Assumption
	EF	Exposure Frequency	day/year	365	Exposure every day
	ED	Exposure Duration	year	70 6[d]	Value for fisher adults Value for fisher children
	BW	Body Weight	kg	70 15[d]	Value for fisher adults Value for fisher children
	AT	Averaging Time	year	70 ED	Value for carcinogenic effects Equal to exposure duration for noncarcinogenic effects
Ingestion of drinking water[n]	CR_{dw}	Consumption rate of drinking water	L/day	3.0 1.5[o]	Value for adults Value for children
	F_{dw}	Fraction of ingested drinking water that is contaminated	unitless	1.0	Assumption
	EF	Exposure Frequency	day/year	365	Exposure every day
	ED	Exposure Duration	year	70 6[d]	Value for adults Value for children
Ingestion of drinking water[n]	BW	Body Weight	kg	70 15[d]	Value for adults Value for children
	AT	Averaging Time	year	70 ED	Value for carcinogenic effects Equal to exposure duration for noncarcinogenic effects

Table 7-3 Summary of Exposure Parameters for Native American Subsistence Scenarios[a]

Exposure Pathway	Parameter	Description	Units	Value[b]	Comment
Inhalation of Volatile and Semi-Volatile Organic Compounds Inside the Sweat Lodge	VW	Volume of Water Used	L	4	Assumed value
	D	Diameter of Sweat Lodge	m	2	Hemisphere
	IR	Inhalation Rate	m^3/day	20	Assumed rate
	EF	Exposure Frequency	day/year	365	Exposure every day
	ED	Exposure Duration	year	70	Assumed lifetime
	ET	Exposure Time	hr/day	1	1 hour of exposure each day
	BW	Body Weight	kg	70	Value for adults
	AT	Averaging Time	year	70	Assumed
Dermal Absorption Inside the Sweat Lodge	SA	Surface Area	m^2	2	Whole body for an adult
	ET	Exposure Time	hr/day	1	1 hour of exposure each day
	EF	Exposure Frequency	day/year	365	Exposure every day
	ED	Exposure Duration	year	70	Assumed lifetime
	BW	Body Weight	kg	70	Value for adults
	AT	Averaging Time	year	70	Assumed
Infant Ingesting Breast Milk[p]	h	Half-life of dioxin (and related compounds) or isotope in adult	day	2555[q] isotope-specific	Upper bound of 5 - to - 7 year range for dioxins Half-life of isotope in days
	f_1	Fraction of ingested dioxin or isotope that is stored in fat	unitless	0.9[q]	EPA OSW recommendation
	f_2	Fraction of mother's weight that is fat	unitless	0.3[q]	EPA OSW recommendation
	f_3	Fraction of mother's breast milk that is fat	unitless	0.04[q]	EPA OSW recommendation

Table 7-3 Summary of Exposure Parameters for Native American Subsistence Scenarios[a]

Exposure Pathway	Parameter	Description	Units	Value[b]	Comment
Infant Ingesting Breast Milk[p] (cont'd)	f_4	Fraction of ingested dioxin compounds or isotopes that are absorbed	unitless	0.9[q]	EPA OSW recommendation
	IR_{milk}	Ingestion rate of breast milk by infant	kg/day	0.742	Harris and Harper 1997
	ED	Exposure Duration	year	1.0[q]	EPA OSW recommendation
	BW_{infant}	Body Weight of Infant	kg	10[q]	Mean of 6 months to 1 year old infant's weight, ranging from 9.1 to 11.3 kg
	AT[i]	Averaging Time	year	1[q]	EPA OSW recommendation, acknowledging that this value may overestimate carcinogenic risks by a factor of 70
	EF	Exposure Frequency	day/year	365[q]	Infant ingesting breast milk every day

[a] The hunter/gatherer (child and adult) is not exposed to the ingestion of fish, but is exposed to all other non-infant pathways. The fisher (child and adult) is not exposed to the ingestion of produce, meat, game, milk, poultry, and eggs, but is exposed to all other non-infant pathways. Children are not exposed to the two sweat lodge scenarios (inhalation of VOCs/SVOCs and dermal absorption). The infant's ingestion of breast milk is based on the mother's total daily intake of COPCs/ROPCs via all indirect exposure pathways (i.e., ingestion of soil, produce, meat, game, milk, poultry, eggs, fish, and drinking water).

[b] Parameter values are from Harris and Harper 1997, unless otherwise noted.

[c] When the inhalation rate in m^3/hr is combined with the exposure time of 24 hr/day, the resulting inhalation rates are 20 m^3/day for adults and 10 m^3/day for children (EPA 1989b).

[d] BW and ED values for children are consistent with those found in EPA 1998a.

[e] Adult soil ingestion rate is from Harris and Harper 1997; this same rate (200 mg/day) is commonly used as a child's soil ingestion rate (EPA 1989c). Therefore, this same rate is used for the Native American child soil ingestion rate.

[f] RAGS Part B (EPA 1991b).

[g] Age modifying factors are from Saito, et al. 1998.

[h] Consumption rates for unprotected aboveground produce are based on the Native American intake rates of 3.3 g/kg-day (231 g/day for a 70 kg adult) for fruits and 4.9 g/kg-day (343 g/day for a 70 kg adult) for vegetables (Harris and Harper 1997), along with the following assumptions: (1) fruits are unprotected aboveground produce; (2) vegetables are a combination of unprotected aboveground produce and belowground produce; and (3) 50% of the vegetables are assumed to come from unprotected aboveground produce and 50% from belowground produce. Thus, the unprotected aboveground produce intake for a 70-kg adult is 231 g/day of fruits and 171.5 g/day of vegetables, for a total unprotected aboveground produce intake rate of 402.5 g/day. Adult and child consumption rates are calculated from the 402.5 g/day intake.

[i] Consumption rate for protected aboveground produce is 0.0 kg/kg-day, based on the assumption that neither fruits nor vegetables are protected aboveground produce. Also, see footnote h above.

[j] Consumption rates for belowground produce are based on the Native American intake rate of 4.9 g/kg-day (343 g/day for a 70 kg adult) for vegetables (Harris and Harper 1997), along with the following assumptions: (1) vegetables are a combination of unprotected aboveground produce and belowground produce; and (2) 50% of the vegetables are assumed to come from unprotected aboveground produce and 50% from belowground produce. Thus, 171.5 g/day of vegetables is used as the basis for belowground produce consumption rates for adults and children.

[k] Milk consumption rates for non-Native Americans; values used for Native Americans are consistent with those values used for non-Native Americans.

[l] Consumption rates for poultry and eggs are based on the Native American intake rate of 44 g/day of fowl, which includes both bird and eggs (Harris and Harper 1997). Based on the ratio of 0.54:0.491 for Native American intake of poultry to eggs (see Tables 11-5 and 11-7 in EPA 1997b), the fowl intake of 44 g/day is subdivided as 23 g/day of poultry and 21 g/day of eggs. These same intake rates are used to calculate consumption rates for hunter/gatherer adults and children.

[m] Personal communication (EPA 1999f).

[n] Surface water concentrations will be used to evaluate the ingestion of drinking water pathway.

[o] Value obtained from personal communication with B. Harper (Harper 1999).

[p] Dioxins, furans, coplanar PCBs, and radionuclides are evaluated for the infant's ingestion of breast milk.

[q] Values from infant ingestion of breast milk scenario in EPA 1998a.

Table 7-4 Summary of Exposure Parameters for Hanford Site Worker Scenario

Exposure Pathway	Parameter	Description	Units	Value[a]	Comment
Inhalation of particulates and vapors	IR	Inhalation rate of air	m^3/hr	2.5[b]	Value for adult worker
	ET	Exposure Time	hr/day	8[b]	8-hour work day
	EF	Exposure Frequency	day/year	250	Assumed 2 weeks away from contaminated area per year
	ED	Exposure Duration	year	20	Value for adult worker
	BW	Body Weight	kg	70	Value for adult worker
	AT	Averaging Time	year	70 ED	Value for carcinogenic effects Equal to exposure duration for noncarcinogenic effects
Soil Ingestion	CR_{soil}	Consumption rate of soil	kg/day	0.00005	Value for adult worker
	F_{soil}	Fraction of soil ingested that is contaminated	unitless	1.0	Assumption
	EF	Exposure Frequency	day/year	146	Derived from frequency of exposure (365 day/yr x 0.4)
	ED	Exposure Duration	year	20	Value for adult worker
	BW	Body Weight	kg	70	Value for adult worker
	AT	Averaging Time	year	70 ED	Value for carcinogenic effects Equal to exposure duration for noncarcinogenic effects
Inhalation of Resuspended Soil	IR	Inhalation Rate of Air	m^3/hr	2.5[b]	Value for adult worker
	ET	Exposure Time	hr/day	8[b]	Assumption
	EF	Exposure Frequency	day/year	146	Derived from frequency of exposure (365 day/yr x 0.4)
	ED	Exposure Duration	year	20	Value for adult worker
	BW	Body Weight	kg	70	Value for adult worker
	AT	Averaging Time	year	70 ED	Value for carcinogenic effects Equal to exposure duration for noncarcinogenic effects
External Exposure to Ionizing Radiation in Soil	Se	Shielding Factor	unitless	0.2	Assumed 20% shielding
	Te	Exposure Time Factor	unitless	8/24	8-hour workday
	EF	Exposure Frequency	days/year	146	Derived from frequency of exposure (365 day/yr x 0.4)

Table 7-4 Summary of Exposure Parameters for Hanford Site Worker Scenario

Exposure Pathway	Parameter	Description	Units	Value[a]	Comment
	ED	Exposure Duration	years	20	Value for adult worker
External Exposure to Ionizing Radiation in Air	EF	Exposure Frequency	day/year	146	Derived from frequency of exposure (365 day/yr x 0.4)
	ED	Exposure Duration	year	20	Value for adult worker
	AF	Age Factor	unitless	1[c]	No adjustment for adults
Ingestion of drinking water[d]	CR_{dw}	Consumption rate of drinking water	L/day	1	Value for adult worker
	F_{dw}	Fraction of ingested drinking water that is contaminated	unitless	1.0	Assumption
	EF	Exposure Frequency	day/year	250	Assumed 2 weeks away from contaminated area per year
	ED	Exposure Duration	year	20	Value for adult worker
	BW	Body Weight	kg	70	Value for adult worker
	AT	Averaging Time	year	70 ED	Value for carcinogenic effects Equal to exposure duration for noncarcinogenic effects

[a] Values are from DOE-RL 1995, unless otherwise noted.

[b] The combination of the inhalation rate, in m^3/hr, and the exposure time, in hr/day, produce the standard industrial inhalation rate of 20 m^3/day.

[c] Age modifying factor is from Saito, et al. 1998.

[d] Surface water concentrations will be used to evaluate the ingestion of drinking water pathway.

8. Screening Ecological Risk Assessment

The screening ecological risk assessment incorporates four fundamental components of the ERA process: (1) problem formulation, (2) exposure assessment, (3) effects assessment, and (4) risk characterization. Selection of COPCs (section 4), quantification of emissions (section 5), and dispersion modeling (section 6) feed critical information to this process.

8.1. Problem Formulation

This section of the Work Plan focuses on the conceptual exposure model, ecological setting, ecological receptor identification, and assessment/measurement endpoints. Each is defined below.

8.1.1. Ecological Conceptual Exposure Model

A conceptual exposure model has been developed that identifies potential ecological receptors and potentially complete exposure pathways (that is, exposure scenarios). The conceptual model is shown as Figure 8-1. The end product of the conceptual model is the identification of exposure scenarios that are defined by potentially exposed populations and exposure pathways. The conceptual model was developed from information obtained from EPA (1998a) and *Screening Assessment and Requirements for a Comprehensive Assessment: Columbia River Comprehensive Impact Assessment* (DOE-RL 1998).

The conceptual model focuses on identifying complete exposure pathways for potentially exposed receptor populations. An exposure pathway is the means through which an organism comes in contact with a chemical in the environment. Exposure pathways are determined by environmental conditions (e.g., location of habitat and home ranges and wind speed/direction), the potential for chemical migration among media (e.g., air, soil, or surface water), and by the behavior and diet of potentially exposed plant and animal populations. Although several potential pathways may exist, as was explained in the human health risk assessment, not all pathways may be complete. For a pathway to be complete, the following four factors must exist:

1. A source of COPC/ROPC release into the Hanford Site environment,
2. A release and transport mechanism (e.g., deposition to soil) that moves the COPC/ROPC from the source, such as stack, to other locations in the environment,
3. A point of contact with a contaminated medium, and
4. An exposure route to the receptor such as ingesting or inhaling affected media

These four factors were considered in the conceptual model. The sources of COPC and ROPC release are the stack and fugitive emissions from the RPP-WTP. Transport processes, potential points of contact, and complete exposure pathways are identified to formulate exposure scenarios that will be the focus of the quantitative risk assessment. The technical context for this is as follows.

8.1.2. Ecological Characterization

The Hanford Site and adjacent region are a shrub-steppe vegetation zone with a shrub overstory and an understory of grasses. Ecological resources at the Hanford Site are extensive, diverse, and important as explained in Neitzel et al. 1998. Because the Hanford Site has not been farmed or grazed for over

50 years, it has become a refuge for a variety of plant and animal species (Gray and Rickard 1989) containing one of the largest remaining undisturbed shrub-steppe ecosystems in Washington (see Appendix C-1 for a listing of plants and animals observed on the site). About 665 km^2 (257 mi^2) of undeveloped lands located on-site (almost half of the total area of the Hanford Site) have been designated as ecological study areas or refuges (Figure 8-2).

8.1.2.1. Physiographic Setting

The Hanford Site lies within the Intermountain Semidesert Province (USFS 1994). This province includes the plains and plateaus of the Columbia-Snake River Plateaus and the Wyoming Basin. The climate is cool, the average temperature being about 50°F, and semi-arid, with the average annual precipitation ranging from 10 to 20 inches across the province from west to east. At the Hanford Site, the average annual precipitation totals about six inches. This precipitation is evenly distributed throughout the fall, winter, and spring months, with little precipitation during the summer months.

The Hanford Site lies within the semi-arid Pasco Basin of the Columbia Plateau in southeastern Washington and occupies an area of approximately 1,450 km^2 (560 mi^2) north of the confluence of the Yakima River with the Columbia River. The Pasco Basin lies within the southwest corner of the larger Columbia Basin. The Hanford Site occupies approximately one-third of the land area within the Pasco Basin. The Columbia River flows through the northern part of the Hanford Site and forms part of the Hanford Site's eastern boundary after turning south. The Yakima River runs near the southern boundary. Rattlesnake Mountain, Yakima Ridge, and Umtanum Ridge form the southwestern and western boundaries of the Hanford Site. The Saddle Mountains form the northern boundary. Adjoining lands to the west, north, and east are principally range and agricultural land. The Hanford Site exhibits low relief, ranging from 120 m (390 ft) above mean sea level (MSL) at the Columbia River to 230 m (750 ft) MSL in the vicinity of the RPP-WTP sites.

The 200 Areas/RPP-WTP sites are located on the Central Plateau. The Central Plateau is characterized by generally low-relief hills with deeply incised river drainages. Gable Butte and Gable Mountain (small east–west ridges) are prominent features of the Central Plateau.

8.1.2.2. Regional Ecology

The region has been characterized as shrub-steppe. Shrub-steppe vegetation is dominated by a shrub overstory with an understory of grasses (Daubenmire 1970). Non-industrialized lands on the Hanford Site have not been farmed or grazed by livestock for over 50 years, allowing it to serve as a refuge for a variety of plant and animal species (Gray and Rickard 1989). As stated earlier, approximately 665 km^2 (257 mi^2) of undeveloped lands within the Hanford Site have been designated as refuges or ecological study areas. Shrub-steppe is considered a priority habitat by the State of Washington because of its importance to wildlife species of concern (Neitzel et al. 1998). The National Biological Service has identified native shrub and grassland steppes in Washington and Oregon as endangered ecosystems (DOE 1996).

Biodiversity on the Hanford Site is enhanced by the large, relatively undisturbed tract of native shrub-steppe habitat and by the Hanford Reach which is a stretch of the Columbia River below the Priest Rapids Dam (DOE 1996a). Additional factors influencing biodiversity include topographic features such as Rattlesnake Mountain, Gable Butte, Gable Mountain, and the presence of a variety of soils ranging from sand to silty and sandy loam. Unique terrestrial habitats include basalt outcrops, scarps (cliffs), scree slopes, and sand dunes. Aquatic habitats are mostly associated with the Columbia River and include open water habitat, wetlands, and riparian areas (Figure 8-3).

Cold Creek and a tributary, Dry Creek, are ephemeral streams within the Yakima River drainage system that roughly parallel State Route 240 through the Hanford Site. Both streams drain areas to the west of Hanford Site. Surface flow, when it occurs, infiltrates and disappears into the surface sediments in the western portion of the Hanford Site. Rattlesnake Springs, located on the western portion of the Hanford Site, forms a small surface stream that flows for about 3 km (1.8 mi) before disappearing into the ground.

West Lake is located north of the 200 East Area and is recharged from groundwater (Neitzel et al. 1998). West Lake has not received direct effluent discharges from any Hanford Site facilities. This water body is created by an elevated water table within a low surface area south of Gable Mountain. This artificially elevated water table occurs under much of the Hanford Site, reflecting the augmented recharge from Hanford Site operations. Currently, West Lake has been reduced to a collection of small pools and mudflats (Neitzel et al. 1998).

Gable Mountain Pond (also to the north of 200 Area East but south of West Lake) and the B Pond System (immediately east of 200 Area East) received cooling water discharges from several facilities at the Hanford Site (Rogers and Rickard 1977). These artificial water bodies, formed by the waste water discharges from the operation of the separation facilities, were decommissioned and covered with soil.

Vegetation

The Hanford Site is a shrub-steppe ecosystem characterized by bunchgrasses and sagebrushes (Figure 8-3). This ecosystem is also referred to as high desert, northern desert shrub, or desert scrub (Franklin and Dyrness 1973). Prior to Western European settlement, the dominant plant in the area was big sagebrush (Artemisia tridentata) with an understory of perennial bunchgrasses, especially Sandberg's bluegrass (Poa sandbergii) and bluebunch wheatgrass (Agropyron spicatum). Following settlement in the early 1800s, grazing and agriculture disrupted the native vegetation and opened the way for invasive species such as Russian thistle (Salsola kali) and cheatgrass (Bromus tectorum). Cheatgrass is now dominant in fields that were cultivated prior to the establishment of the Hanford Site. Cheatgrass also is well established on rangelands at elevations less than 244 m (800 ft) (DOE and Ecology 1996). Establishment of the Hanford Site as a nuclear complex in 1943 resulted in the creation of a secured area of mostly undeveloped land with scattered, small industrial facilities. Consequently, the Hanford Site is one of a small number of remaining shrub-steppe tracts in Washington that is relatively undisturbed. Wildfire is a common occurrence and can significantly alter the shrub component of the vegetation. The most recent extensive fire on the Hanford Site was in 1984. Trees were planted and irrigated on most of the pre-1943 farms to provide windbreaks and shade. Some of these trees have persisted and serve as nesting platforms for several species of birds, (that is, hawks, owls, ravens, magpies, and great blue herons), and as night roosts for wintering bald eagles (DOE and Ecology 1996) (Figure 8-4).

The Central Plateau and surrounding areas have been identified as predominantly shrub-steppe (Neitzel et al. 1998 and Duranceau 1995). This designation includes communities dominated by big sagebrush and bitterbrush (*Purshia tridentata*) with an understory of cheatgrass or Sandberg's bluegrass. Over 100 plant species occur on the Central Plateau and vicinity. Common plant species include big sagebrush, rabbitbrush (*Chrysothamnus nauseous*), cheatgrass, and Sandberg's bluegrass.

The RPP-WTP sites in the 200 East Area and the immediately surrounding area are approximately 40% big sagebrush and rabbitbrush (Figure 8-3). Another 20% is dominated by Russian thistle, with the remainder being disturbed vegetation or bare gravel (PNL 1994). Other vegetation in the 200 Areas includes introduced perennial grasses planted to revegetate and stabilize disturbed areas such as waste burial grounds. Siberian wheatgrass (*Agropyron sibericum*) has been used extensively and has proven to be drought tolerant and better adapted to sandy soil than other species (Stegen 1993).

Wetland species such as cattail, reeds, and various trees, such as willow, cottonwood, and Russian olive, are established around some of the man-made ditches and ponds that are in the area. The decommissioning of some facilities have eliminated the supply of industrial water feeding some ponds. Without this water supply, the artificially supported wetland habitats have also been eliminated.

Almost 600 species of plants have been identified on the Hanford Site (Neitzel et al. 1998). The dominant plants are big sagebrush, rabbitbrush, cheatgrass, and Sandberg's bluegrass, with cheatgrass providing half of the total plant cover on much of the Hanford Site. Cheatgrass and Russian thistle that are annuals introduced to the United States from Eurasia in the late 1800's, invade disturbed areas. Big sagebrush and bitterbrush are widely spaced and usually provide less than 20% canopy cover. Bitterbrush provides important browse for the resident mule deer herd. The important understory plants are grasses, especially cheatgrass, Sandberg's bluegrass, Indian ricegrass, June grass, and needle-and-thread grass.

Central Plateau. More than 100 species of plants have been identified on the Central Plateau (Cushing 1992). Common species include sagebrush, rabbitbrush, cheatgrass, and Sandberg's bluegrass. Dominant vegetation type consists of big sagebrush with an understory of cheatgrass and Sandberg's bluegrass (Neitzel et al. 1998). Cheatgrass provides approximately 50% of the total plant cover. Cheatgrass also is common where native plant communities have been disturbed by wildfire or past construction activities. Three vegetation subtypes occurring in the vicinity of the 200 West Area of the Central Plateau are sagebrush and Sandberg's bluegrass, sagebrush and needle-and-thread grass, and spiny hopsage and Sandberg's bluegrass. Past wildfires in the Central Plateau have opened up some areas, creating a mosaic of shrub- and grass-dominated areas. Sandberg's bluegrass and cheatgrass are the most common grass species. Other vegetation includes riparian species associated with man-made ditches and ponds and introduced perennial grass planted to revegetate disturbed areas (DOE and Ecology 1996).

These riparian species include herbaceous species, such as cattail and reeds, and trees, such as willow, cottonwood, and Russian olive (Neitzel et al. 1998). Several of these ponds recently have been decommissioned, resulting in the elimination of wetland habitat as the supply of industrial waste water feeding the ponds was terminated. Introduced perennial grasses (that is, Siberian wheatgrass) have been used extensively in the Central Plateau to revegetate and stabilize waste burial grounds against wind and water erosion. Siberian wheatgrass has proven to be drought tolerant and better adapted to sandy soils than other cultivars used in Central Plateau revegetation efforts (WHC 1993).

Columbia River. The two major vegetation types occurring along the Hanford Reach of the Columbia River are riparian and upland (NPS 1994). Riparian habitats are found along the shoreline, slack water and slough areas, and on islands in the river. Riparian vegetation at these locations includes both woody and herbaceous species maintained by the high water table immediately adjacent to the river. Common plant species occurring in the riparian zone include black cottonwood, mulberry, willow, dogbane, and a variety of grasses and forbs (Neitzel et al. 1998). Sensitive habitats within the riparian zone include islands and cobbled shorelines occurring as a narrow band along the Hanford Reach. Plant species occurring in these areas include perennial summer-blooming forbs adapted to seasonal changes in water levels (NPS 1994). Upland habitats along the Hanford Reach are composed of shrub-steppe vegetation similar to that found on the Central Plateau (DOE and Ecology 1996).

In summary, special topographic features include Gable Butte and Gable Mountain north of the Central Plateau and an extensive series of active sand dunes in the southeast portion of the area. The dominant plant communities are cheatgrass, sagebrush-bitterbrush and Sandberg's bluegrass, sagebrush and cheatgrass, Sandberg's bluegrass, and riparian plant communities (Sackschewsky et al. 1992). Depending on the location, many of the terrestrial plants occurring in this area are the same as those found in the adjacent Columbia River and Central Plateau. Big sagebrush, bitterbrush, rabbitbrush, cheatgrass, and

Sandberg's bluegrass are common species in the 300 and 400 Areas (Neitzel et al. 1998). Common plants growing in riparian areas along the Columbia River include reed canarygrass, common witchgrass, large barnyard grass, summer-blooming forbs, sandbar willow, poplar, white mulberry, and Russian olive (NPS 1994). Vegetation occurring on scree slopes, outcrops, and scarps on Gable Butte and Gable Mountain is limited to scattered individuals or groups of plants. Plant species include squaw currant, bluebunch wheatgrass, rock buckwheat, and thyme buckwheat. Rigid sagebrush occurs, at the Hanford Site, only on Gable Mountain and Umtanum Ridge (Downs et al. 1993).

Wildlife

Almost 300 species of terrestrial vertebrates have been observed at the Hanford Site. This number includes 41 species of mammals, 246 species of birds, four species of amphibians, and nine species of reptiles (Neitzel et al. 1998).

Mammals. Large herbivorous mammal species that are found on the Hanford Site include mule deer, elk, and white-tailed deer. Mule deer, with an on-site herd of several hundred, occur just about everywhere on the Hanford Site but are most often found near the Columbia River. White-tailed deer and elk began to appear on the Hanford Site during the early 1970s. White-tailed deer tend to remain in the riparian habitats along the rivers, while elk generally are restricted to the Fitzner-Eberhardt Arid Lands Ecology Reserve (FEALE Reserve). Elk frequently move off the reserve to private lands to the north and west, particularly during late spring, summer, and early fall. This herd grew from an estimated eight animals in 1975 to almost 600 animals in 1997 (Neitzel et al. 1998).

Common mammalian predators are the coyote, bobcat, and badger. These carnivores feed primarily on the several species of small mammals found on the Hanford Site, including the Great Basin pocket mouse, Western harvest mouse, grasshopper mouse, deer mouse, house mouse, Townsend's ground squirrel, mountain and sagebrush voles, black-tailed jackrabbit, brushy-tailed woodrat, and Northern pocket gopher. Of these small mammals, the Great Basin pocket mouse is the most abundant. Coyotes have been a major predator of Canada goose nests on Columbia River islands, especially upstream from the abandoned Hanford townsite (DOE and Ecology 1996).

Black-tailed jackrabbits are common on the Hanford Site and are most often found in mature stands of sagebrush. Cottontail rabbits also are common but are more closely associated with the developed areas of the Hanford Site. Townsend's ground squirrels occur in colonies of various sizes scattered across the Hanford Site. The most abundant mammal inhabiting the site is the Great Basin pocket mouse. This mouse occurs all across the Columbia River plain and on the slopes of the surrounding ridges. Other small mammals include the deer mouse, harvest mouse, grasshopper mouse, mountain vole, vagrant shrew, and Merriam's shrew (DOE and Ecology 1996).

Up to 14 species of bats are known or have the potential to occur on the Hanford Site. They include the pallid bat, hoary bat, silver-haired bat, little brown bat, California brown bat, Yuma brown bat, and the Pacific western big-eared bat (Fitzner and Gray 1991). The pallid bat, which roosts in abandoned buildings, is considered to be most abundant. All of these bat species feed on insects.

Birds. Nearly 250 species of birds occur on or near the Hanford Site as year-round residents, seasonal residents, migrants, and accidentals (Neitzel et al. 1998).

Eleven raptors have been documented as nesting on the Hanford Site. These include the northern harrier (*Circus cyaneus*), red-tailed hawk (*Buteo jamaicensis*), Swainson's hawk (*Buteo swainsoni*), golden eagle (*Aquila chrysaetos*), prairie falcon (*Falco mexicanus*), American kestrel (*Falco sparverius*), barn owl (*Tyto alba*), great horned owl (*Bubo virginianus*), long-eared owl (*Asio otus*), short-eared owl (*Asio*

flammeus), and burrowing owl (*Athene cunicularia*) (Fitzner and Gray 1991, Rickard et al. 1988). Raptors use a variety of habitats for nesting and foraging on the Hanford Site. Nesting habitats include outcrops, cliffs, trees, marshes, fields, and utility towers. Depending on raptor species, prey may include small mammals, birds, reptiles such as snakes, and insects.

A variety of songbird species occur in the shrub-steppe vegetation. These include the Western meadowlark (*Sturnella neglecta*), grasshopper sparrow (*Ammodramus savannarum*), horned lark (*Eremophila alpestris*), and sage thrasher (*Oreoscoptes montanus*) (Downs et al. 1993). The Western meadowlark and horned lark are the most abundant breeding bird species within the shrub-steppe habitat (Rickard and Poole 1989). These two species nest on the ground in the open, while other species e.g., sage sparrow, sage thrasher, and loggerhead shrike) require sagebrush or bitterbrush as nesting structures.

Common upland game bird species include the chukar partridge (*Alectoris chukar*), California quail (*Callipepla californicus*), and Chinese ring-necked pheasant (*Phasianus colchicus*). Sage grouse (*Centrocercus urophasianus*) and gray partridge (*Perdix perdix*) are less common and are rarely seen. Although once more common, sage grouse are now essentially absent from the Hanford Site, displaced after a major wildfire in 1984 (Brandt 1995). A 1997 inventory conducted by The Nature Conservancy did not record any sage grouse in the sagebrush-steppe habitat of the FEALE Reserve (Neitzel et al. 1998). None of the upland birds are native to the area except the sage grouse.

Reptiles and Amphibians. Nine species of reptiles and four species of amphibians are found at the Hanford Site (Neitzel et al. 1998). The most abundant reptile is the side-blotched lizard (*Uta stansburiana*) (Neitzel et al. 1998). The short-horned lizard (*Phrynosoma douglassii*) and northern sagebrush lizard (*Sceloporous graciosus*) are also common in mature sagebrush habitats with sandy soil. Commonly encountered snakes include the gopher snake (*Pituophis melanoleucus*), yellow-bellied racer (*Coluber constrictor*), and Pacific rattlesnake (*Crotalus viridis*). Less common are striped whipsnakes (*Masticophis taeniatus*) and desert night snakes (*Hyspiglena torquata*). Amphibians on the Hanford Site are associated with riparian habitats located along permanent water bodies or the Columbia River (Fitzner and Gray 1991). Species include the Great Basin spadefoot toad (*Spea intermontana*), Woodhouses toad (*Bufo woodhouseii*), and the Pacific tree frog (*Hyla regilla*).

Insects. Grasshoppers and darkling beetles represent some of the more conspicuous insect groups. The populations of both of these species of insects are subject to seasonal changes and weather variations (Rogers and Rickard 1977). Fifty percent of the known insect species are of the order Coleoptera (beetles) (ERDA 1975). Many of the insect species are important in the food web of birds and mammals found on the Hanford Site. Species like the darkling beetle play an important role in the decomposition process by feeding on decaying plant material, animal feces, fungi, and live plant tissue (Weiss and Mitchell 1992).

The Nature Conservancy has identified nearly 1,500 species of insects on the Hanford Site (Hall 1998). The Nature Conservancy identified 41 new species of insects including six new species of bees, six new species of flies, five new species of leafhopper and planthopper insects, one new species of wasp, and one new species of beetle (Neitzel et al. 1998). The Nature Conservancy focused on the FEALE Reserve, the North Slope, and along the Columbia River. Consequently, none of these new species has been reported from the 200 Areas.

Central Plateau. A characterization study of small mammals performed south of the 200 East Area resulted in the following five species being trapped: the Great Basin pocket mouse, deer mouse, northern grasshopper mouse, sagebrush vole, and western harvest mouse (Rogers and Rickard 1977). The Great Basin pocket mouse represented more than 90% of the individuals caught. Medium- and large-size

mammals that may occur in the Central Plateau include rabbits, coyotes, badgers, and mule deer (Rogers and Rickard 1977). Some of these organisms will be receptors in the ecological risk assessment. Other mammals potentially using areas associated with ponds and ditches in the 200 Areas include muskrats, porcupines, and raccoons. Many common bird species, such as the western meadowlark and sage sparrow, are likely to occur on the Central Plateau where suitable habitats exist. Thirty-seven species of terrestrial birds were recorded during surveys conducted in the 200 Areas in 1986 (Schuler et al. 1993).

Unique habitats can be found on Gable Butte and Gable Mountain situated north of the Central Plateau. These unique habitats include basalt outcrops, scarps, and scree slopes. Birds likely to occur in these habitats are the prairie falcon, rock wren, poorwill, and chukar; small mammals include the yellow-bellied marmot and wood rat; reptiles include rattlesnakes, gopher snakes, and horned lizards (Downs et al. 1993).

Columbia River. Terrestrial wildlife species use both shoreline riparian and shrub-steppe habitats occurring along the Columbia River and on the islands. Wildlife reported to use the Hanford Reach include 184 species of birds, 36 species of mammals, 9 species of reptiles, and 4 species of amphibians (NPS 1994). The Canada goose uses islands along the Hanford Reach extensively for nesting. Monitoring of nesting geese that use the Hanford Site have been ongoing since 1950. These studies indicate that Canada geese nest more frequently on islands in the downstream reach because of heavy predation by coyotes further upstream (Neitzel et al. 1998). Mule deer use the islands and other riparian areas for fawning habitat. Wildlife occurring in shoreline habitat includes 46 species that use willow communities and 49 species that use grass areas (NPS 1994).

8.1.2.3. Aquatic Ecosystems

Washington State has classified the stretch of the Columbia River that includes the Hanford Reach as Class A, Excellent (Neitzel et al. 1998). Class A waters must be suitable for essentially all uses, including raw drinking water, recreation, and wildlife habitat. Water from the Columbia River is used for both irrigation and municipal water supplies. Federal and state drinking water quality standards apply to the Columbia River and are currently being met (Neitzel et al. 1998). Water samples from the Columbia River and three ponds on the Hanford Site are routinely collected and analyzed.

The Columbia River supports an ecosystem of plankton, benthic invertebrates, fish, and other communities. Algae are abundant in the river and provide food for herbivores, such as immature insects, which are then eaten by carnivorous species, such as bass.

Aquatic plants in the Hanford Reach include water milfoil, waterweed, pondweed, Columbia yellowcress, watercress, and duckweed. Water milfoil is an aggressive introduced aquatic plant and is becoming a nuisance in the river. Other aquatic species found in the Hanford Reach include a variety of microflora, zooplankton, and benthic invertebrates. Microflora include both sessile types (periphyton) and free-floating types (phytoplankton). Microflora species include diatoms, golden or yellow-brown algae, green algae, blue-green algae, red algae, and dinoflagellates. Dominant zooplankton taxa include *Bosmina*, *Diaptomus*, and *Cyclops*. Benthic invertebrate taxa occurring in the Hanford Reach include; insect larvae such as caddisflies, midge flies, black flies, also snails, freshwater sponges, limpets, and crayfish (Neitzel et al. 1998).

The Hanford Reach supports over 40 species of fish. The anadromous Chinook salmon, sockeye salmon, coho salmon, and steelhead trout use the river to migrate to and from upstream spawning areas. Chinook salmon and steelhead trout also spawn in the Hanford Reach in the fall (Figure 8-6). Shad may also spawn in this stretch of river. Mountain whitefish, white sturgeon, smallmouth bass, crappie, catfish,

walleye, and yellow perch are important gamefish to sport fisherman and Native Americans. A healthy rough fish population includes carp, redside shiner, suckers, and northern squawfish (Neitzel et al. 1998).

West Lake near the 200 Areas is created by a rise in the water table under the Central Plateau and is not fed by surface flow. This results in the pond being highly saline as well as alkaline and having a low species diversity (DOE and Ecology 1996). West Lake, located southwest of Gable Mountain, fluctuates in size with changes in the water table and is currently reduced to a collection of small pools and mudflats (Neitzel et al. 1998). Unlike other ponds on the Hanford Site, West Lake does not receive direct effluent discharges from Hanford Site facilities (PNL 1993). Wetland vegetation found at West Lake is limited to scattered patches of emergent macrophytes such as cattails and bulrushes.

Other wetland habitats found on the Hanford Site are associated with man-made ponds and ditches occurring on the Hanford Site, including the B Pond system located near the 200 East Area and a small cooling and wastewater pond in the 400 Area. The B Pond system was constructed in 1945 to receive cooling water from facilities in that area. Since that time, effluent flow to the B Pond has halted. One lobe of the pond received cooling water until recently; the rest of the B Pond system is slowly reverting to a shrub-steppe ecosystem. Gable Mountain Pond (also to the north of 200 East Area but south of West Lake) also received cooling water discharges from several facilities on the Hanford Site (PNNL 1997). These artificial water bodies, formed by the wastewater discharges from the operation of the separation facilities, were decommissioned and covered with soil.

8.1.2.4. Threatened and Endangered Species

Species of concern on the Hanford Site include federally-listed threatened and endangered species, state-listed threatened or endangered species, state candidate species, state plant species of concern, and species of ethnobiological concern to Native Americans.

No federally-listed threatened or endangered plant or mammal species is documented as occurring on the Hanford Site (Neitzel et al. 1998). Three birds on the federal list occur regularly or incidentally on the site; two federally-listed species of fish occur within the Hanford Reach. The threatened bald eagle (*Haliaeetus leucocephalus*) is found regularly along the Hanford Reach, while the threatened Aleutian Canada goose (*Branta canadensis leucopareia*) and the endangered peregrine falcon (*Falco peregrinus*) occur only as incidental visitors to the Hanford Site (Neitzel et al. 1998). The two fish species are the anadromous chinook salmon (*Oncorhynchus tshawytscha*) and steelhead (*Oncorhynchus mykiss*). These species are regulated as Evolutionarily Significant Units (ESU) by the National Marine Fisheries Service based on historical geographic spawning areas. One ESU of the chinook salmon, the Upper Columbia River spring-run ESU, is listed as endangered (64 FR 14308). Two ESUs of the steelhead have been listed: the Upper Columbia River steelhead is listed as endangered (Neitzel et al. 1998) and the steelhead Middle Columbia River ESU is listed as threatened (64 FR 14517).

Washington State lists the peregrine falcon, Aleutian Canada goose, American white pelican (*Pelecanus erythrorhynchos*), sandhill crane (*Grus canadensis*), and pygmy rabbit (*Brachylagus idahoensis*) as endangered and the ferruginous hawk (*Buteo regalis*) and the bald eagle as threatened. The peregrine falcon is a casual migrant to the Hanford Site. The bald eagle is a regular winter resident along the Columbia River (Neitzel et al. 1998).

Eight species of plants listed by Washington State as threatened or endangered are found on the Hanford Site. Columbia milk-vetch (*Astragalus columbianus*), loeflingia (*Loeflingia squarrosa*), Hoover's desert parsley (*Lomatium tuberosum*), and dwarf evening primrose (*Oenothera pygmaea*), are designated as threatened. Columbia yellowcress (*Rorippa columbiae*), Umtanum desert buckwheat (*Erigonium codium*), and White Bluffs bladderpod (*Lesquerella tuplashensis*) are listed as endangered. Columbia

milkvetch occurs on upland terraces along the Columbia River and on Yakima Ridge within the FEALE Reserve. Dwarf evening primrose has been found north of Gable Mountain and on disturbed areas near the Wye Barricade. Umtanum desert buckwheat and White Bluffs bladderpod are indigenous to the Hanford Site (Neitzel et al. 1998).

Piper's daisy (*Erigeron piperianus*), a state sensitive species, has been found at B Pond near the 200 East Area and at Pit 30. Crouching milkvetch, stalked-pod milkvetch, and scilla onion, all state plant species of concern, are also found in the 200 East Area. Wildlife state candidate species observed or considered likely to be found on or near the Central Plateau include the sage sparrow (*Amphispiza belli*) and the loggerhead shrike (*Lanius ludovicianus*). Both of these birds commonly nest in undisturbed shrub-steppe habitat. The sage sparrow is one of the most common nesting birds on the Hanford Site (Downs et al. 1993). Other bird species of concern that may be found include the burrowing owl, ferruginous hawk, golden eagle, sage thrasher, and merlin (Neitzel et al. 1998). Another wildlife species of concern inhabiting the Central Plateau and vicinity is the striped whipsnake (*Mastocophis taeniatus*), a state candidate species.

Central Plateau. No federal or state listed threatened or endangered plant or animal species occur in the Central Plateau (DOE and Ecology 1996). Several state-listed plant species are found on the Central Plateau. Piper's daisy has been found at B Pond near the 200 East Area, and may occur in sagebrush-steppe habitat elsewhere on the Hanford Site (WHC 1992). Dwarf evening primrose has been found on disturbed areas near the Wye Barricade (Neitzel et al. 1998) and might also be found on the Central Plateau.

Wildlife species of state concern occurring in the 200 Areas include the loggerhead shrike and sage sparrow. Both species nest in undisturbed sagebrush habitat in the Central Plateau (PNL 1993). Other bird species of concern that may occur in shrub-steppe habitat in the Central Plateau are the burrowing owl, golden eagle, long-billed curlew, and Swainson's hawk. Reptile species of concern using the Central Plateau include the striped whipsnake (Rogers and Rickard 1977, Neitzel et al. 1998).

Columbia River. No federally listed threatened or endangered plant species occur on the Hanford Reach (DOE and Ecology 1996). State endangered plant species occurring along the Hanford Reach include the Columbia yellowcress. Preferred habitat for Columbia yellowcress is shoreline areas with gently sloping, cobbly, or sandy substrate (PNL 1993b). State plant species of concern that have been found along the shoreline and on islands of the Hanford Reach between the Vernita Bridge and the 300 Area include the southern mudwort, dense sedge, and shining flatsedge (WHC 1992, Neitzel et al. 1998).

Federally listed threatened and endangered birds include the Aleutian Canada goose, peregrine falcon, and bald eagle. The Aleutian Canada goose and the peregrine falcon are incidental migrants on the Hanford Site (Neitzel et al. 1998). State-listed bird species that occur along the Hanford Reach that are considered relatively common include the American white pelican (endangered), bald eagle (threatened), and sandhill crane (endangered). The common loon (*Gavia immer*), a state candidate species, is also found within the Hanford Reach.

8.1.2.5. Sensitive Environments

Sensitive habitats on the Hanford Site include wetlands and riparian habitats. Wetlands include those transitional lands occurring between terrestrial and aquatic ecosystems where the water table is usually close to the surface or where shallow water covers the surface (Cowardin et al. 1979). The primary wetlands found onsite occur along the Hanford Reach of the Columbia River and include the riparian habitats located along the river shoreline. Other wetland habitats found on the Hanford Site are associated with man-made ponds and ditches. These include B Pond and its associated ditches located

near the 200 East Area. The B Pond Complex was constructed in 1945 to receive cooling water from facilities in that area. Wetland plants occurring along the shoreline of B Pond include herbaceous species such as showy milkweed (*Asclepias speciosa*), western goldenrod (*Solidago occidentalis*), three square bulrush (*Scirpus americanus*), horsetails (*Equisetum* spp.), rushes (*Juncus* spp.), and common cattail (*Typha latifolia*), and woody species such as mulberry (*Morus alba*), silver poplar (*Populus alba*), black cottonwood (*Populus trichocarpa*), and willow (*Salix* spp.) (Sackschewsky et al. 1992). Wildlife species observed at B Pond include a variety of mammals and waterfowl species (Meinhardt and Frostenson 1979).

8.1.3. Receptor Identification

Food webs represent the transfer of matter among the components of an ecosystem. This transfer occurs through the uptake and absorption of substances from abiotic media or consumption of animal and plant tissue. In a contaminated environment, the transfer of contaminants may also occur during uptake or consumption of contaminated media.

The food web representing the terrestrial organisms of the Hanford Site and their trophic relationships is. The food web representing the aquatic organisms of the Hanford Site and their trophic relationships is shown in Figure 8-8.

8.1.3.1. Terrestrial Receptors

A simplified food web for selected terrestrial receptors is presented in Figure 8-9. The receptors selected for use in the ecological risk assessment are shown in bold in figure:

- Plants: cheatgrass, rabbitbrush
- Terrestrial Invertebrates: earthworms, darkling beetles
- Herbivorous Mammals: mule deer
- Omnivorous Mammals: Great Basin pocket mouse
- Omnivorous Birds: Western meadowlark
- Carnivorous Mammals: coyote
- Carnivorous Birds: burrowing owl, red-tailed hawk.

Species profiles are not required for plants (cheatgrass and rabbitbrush) and terrestrial invertebrates (earthworms and darkling beetles). Single-page species profiles for mammals and birds containing a short narrative and a table of receptor parameters are provided.

8.1.3.2. Aquatic Receptors

A simplified food web of selected aquatic receptors is presented in Figure 8-10. The receptors selected for use in the ecological risk assessment are shown in bold on the figure and are listed below:

- Benthic Invertebrates (sediment-dwelling organisms): clams, insects
- Aquatic Organisms, Fish, and Other: bass, salmon, channel catfish; water fleas, other invertebrates
- Waterfowl and Shorebirds: Canada goose, spotted sandpiper
- Piscivorous Terrestrial Carnivores: great blue heron, bald eagle, mink

Species profiles are not required for the very numerous benthic invertebrates (clams, insects, snails, and worms), the planktivorous fish and small invertebrates (small carp, small Northern squaw fish, small suckers, water fleas, and other invertebrates), and fish (bass, salmon, and channel catfish). Single-page

species profiles of birds and mammals containing a short narrative and a table of receptor parameters are provided.

8.1.4. Assessment and Measurement Endpoints

Environmental statutes govern the maintenance of ecological resources, including:

1 preservation and conservation of threatened and endangered organisms
2 maintenance and protection of terrestrial organism populations and ecosystems
3 maintenance and protection of aquatic organism populations and ecosystems

An assessment endpoint is defined by EPA as "an explicit expression of the environmental value that is to be protected" (EPA 1992a). This applies to ecological receptors (population, community, or individual in the case of a threatened or endangered species). Measures of effect are defined as a change in an attribute of an assessment endpoint or its surrogate in response to a stressor to which it is exposed. Decision criteria prescribe how the endpoints are evaluated.

Policy goals, assessment endpoints, measures of effect and exposure, and decision rules used for the ecological risk assessment are presented in Table 8-1.

Mule Deer (*Odocoileus hemionus*)

Mule deer, with an on-site herd of several hundred, occur just about everywhere on the Hanford Site, but are most often found near the Columbia River. Mule deer use the islands and other riparian areas as fawning habitat. Bitterbrush provides important browse for the resident mule deer herd. Summer browse is chiefly herbaceous plants and the young shoots of woody plants, while winter browse includes twigs of woody plants and trees, including cedar, yew, aspen, willow, dogwood, juniper, and sage. Coyotes are a major predator, along with bobcats to a lesser extent. Mule deer are most active in the mornings and evenings.



Parameter	Definition	Value	Reference/Notes
BW	Body weight (kg)	66.5	Average of males and females, north central Colorado (Anderson et al. 1974 in Higley and Kuperman 1996)
HR	Home range (ha)	285	Utah, Arizona, Montana (Harestad and Bunnell 1979 in Higley and Kuperman 1996)
TUF	Temporal use factor	1	Will be 1 unless specific value exists.
IR_F	Food ingestion rate (g/g-d=kg/kgBW/day)[a]	0.022	Colorado (Alldredge et al. 1974 in Higley and Kuperman 1996)
PF	Plant fraction	1	Estimated
AF	Animal fraction	0	Estimated
SF	Soil fraction	0.02	(Arthur-Alldredge 1979 in DOE and Ecology 1996)
IR_W	Water ingestion rate (g/g-d=L/kgBW/day)	44	(Bissell et al. 1955 in Higley and Kuperman 1996)
IR_i	Inhalation rate (m^3/day)	15.7	Allometric, 0.5458 $BW^{0.80}$ (EPA 1993b, Eq. 3-20)

[a] Food ingestion rate (g/g – d) reexpressed as kg/kg BW/day is assumed not to include ingested soil; therefore, PF + AF = 1.0.

Great Basin Pocket Mouse (*Perognathus parvus*)

The pocket mouse eats mostly seeds, but also eats insects (Fitzner and Gray 1991). It is the principal prey of the burrowing, great horned, long-eared, and barn owls at the Hanford Site (Downs et al. 1993) and serves as a vector for contaminant movement through the food chain from plants to mammalian and avian carnivores. The pocket mouse is a nocturnal, burrowing mammal, with most burrows being between 35 and 193 cm (1.2 and 6.3 ft deep) (Gano and Rickard 1982). The mouse has no requirement for drinking water, obtaining all its water from its food. Its small home range could cause it to spend all of its time within a contaminated area and obtain all of its food there (DOE and Ecology 1996).



Parameter	Definition	Value	Reference/Notes
BW	Body weight (kg)	0.0235	Burt and Grossenheider 1976 in DOE and Ecology 1996
HR	Home range (ha)	0.209	O'Farrell et al. 1975 in DOE and Ecology 1996
TUF	Temporal use factor	1	DOE and Ecology 1996
IR_F	Food ingestion rate (g/g-d=kg/kgBW/day)[a]	0.285	Calder 1984 in DOE-RL 1995
PF	Plant fraction	0.62[b]	Annual average, Colorado, short-grass prairie (Flake 1973 in EPA 1993b)
AF	Animal fraction	0.38[b]	Annual average, Colorado, short-grass prairie (Flake 1973 in EPA 1993b)
SF	Soil fraction	0.262	Beyer et al. 1991 in DOE and Ecology 1996
IR_W	Water ingestion rate (g/g-d=L/kgBW/day)	0.00	DOE and Ecology 1996
IR_i	Inhalation rate (m^3/day)	0.027	Allometric, 0.5458 $BW^{0.80}$ (EPA 1993b, Eq. 3-20)

[a] Food ingestion rate (g/g – d) reexpressed as kg/kg BW/day is assumed not to include ingested soil; therefore, PF + AF = 1.0.
[b] Values used for the Great Basin pocket mouse taken from values established for the deer mouse (*Peromyscus maniculatus*). (Cronin and Bradley 1988 in EPA 1993b)

Western meadowlark (*Sturnella neglecta*)

The Western meadowlark is a ground-nesting bird that nests in cheatgrass and sagebrush-bunchgrass communities (Rickard et al. 1988, Schuler et al. 1988). This species exhibits resistance to plant community structure changes resulting from grazing or wildfires (Rickard et al. 1988). The Western meadowlark is a common, omnivorous bird of open habitats in southeastern Washington and is abundant in the shrub-steppe ecosystem (Schuler et al. 1988). It feeds on a variety of items, which includes both insects and plant material, mostly seeds. One study (Bent 1958 in Sample et al. 1997) reports that the Western meadowlark's diet consists of roughly 70% insects and 30% plant material. Studies conducted in southeastern Washington indicate that it is the main bird prey item in the diets of the red-tailed, ferruginous, and Swainson's hawks (Rickard et al. 1988). Adult female Western meadowlarks average 94.2 grams in weight and lay three to seven eggs in dome-shaped nests concealed in the grass or weeds and constructed of the same materials.

Parameter	Definition	Value	Reference/Notes
BW	Body weight (kg)	0.094	Adult female
HR	Home range (ha)	0.42	Pitts 1984 in EPA 1993b; Bull and Farrand 1994
TUF	Temporal use factor	1	Will be 1 unless specific value exists
IR_F	Food ingestion rate (g/g-d=kg/kgBW/day)[a]	0.028	ATG 1998
PF	Plant fraction	0.30	Bent 1958 in Sample et al. 1997
AF	Animal fraction	0.70	Bent 1958 in Sample et al. 1997
SF	Soil fraction	0	No data
IR_W	Water ingestion rate (g/g-d=L/kgBW/day)	0.186[b]	Estimated (EPA 1993b)
IR_i	Inhalation rate (m^3/day)	0.066	Allometric, 0.4089 $BW^{0.77}$ (EPA 1993b, Eq. 3-19)

[a] Food ingestion rate (g/g – d) reexpressed as kg/kg BW/day is assumed not to include ingested soil; therefore, PF + AF = 1.0.

Coyote (*Canis latrans*)

The coyote is the most common carnivore on the Hanford Site. They are nocturnal, but may be active at any time of day. Primarily carnivorous, coyotes feed mainly on birds and small mammals, but also feed on insects and fruits in season. The typical hunting range is 10 miles, but may extend to 100 miles, reflecting the coyote's variable home range. Being an upper-trophic-level receptor, the coyote could be particularly susceptible to chemicals that bioaccumulate. Coyotes living in the shrub-steppe feed on pocket mice, Northern pocket gopher, Nuttall's cottontail, and black-tailed jackrabbit. Favored den sites are riverbanks and the sides of canyons or gulches.



Parameter	Definition	Value	Reference/Notes
BW	Body weight (kg)	12.75	Average of adult male and female (Bekoff 1977 in Higley and Kuperman 1996)
HR	Home range (ha)	11,300	Residents, southeast Colorado (Gese et al. 1988 in Higley and Kuperman 1996)
TUF	Temporal use factor	1	Will be 1 unless specific value exists
IR_F	Food ingestion rate (g/g-d=kg/kgBW/day)[a]	0.047	Adults (Gier 1975 in Higley and Kuperman 1996)
PF	Plant fraction	0.1	(Bekoff 1977 in Higley and Kuperman 1996)
AF	Animal fraction	0.9	(Bekoff 1977 in Higley and Kuperman 1996)
SF	Soil fraction	0	No data
IR_W	Water ingestion rate (g/g-d=L/kgBW/day)	0.077	(Calder and Braun 1983 in Higley and Kuperman 1996)
IR_i	Inhalation rate (m^3/day)	4.2	Allometric, 0.5458 $BW^{0.80}$ (EPA 1993b, Eq. 3-20)

[a] Food ingestion rate (g/g – d) reexpressed as kg/kg BW/day is assumed not to include ingested soil; therefore, PF + AF = 1.0.

Burrowing Owl (*Athene cunicularia*)

The burrowing owl is the most abundant of the owls that nest on the Hanford Site. Burrowing owls nest in holes in the ground that are abandoned by burrowing mammals. Their diet consists of pocket mice, deer, pocket gophers, mountain voles, black-tailed jackrabbits, Nuttall's cottontail, rock doves, mallards, and American coots.

The burrowing owl is more diurnal than most owls. The female lays 5 to 7 eggs in a long, underground burrow lined with grasses, roots, and dung. The burrows are usually abandoned prairie dog or pocket gopher burrows, but burrowing owls are capable of digging their own.



Parameter	Definition	Value	Reference/Notes
BW	Body weight (kg)	0.15 [b]	Earhart and Johnson, 1970
HR	Home range (ha)	78.5 [b]	Estimated from Baumgartner 1939 in Higley and Kuperman 1996
TUF	Temporal use factor	1	Will be 1 unless specific value exists
IR_F	Food ingestion rate (g/g-d=kg/kgBW/day)[a]	0.092 [b]	Estimated from Craighead and Craighead 1956 in Higley and Kuperman 1996
PF	Plant fraction	0	Estimated
AF	Animal fraction	1	Estimated
SF	Soil fraction	0	No data
IR_W	Water ingestion rate (g/g-d=L/kgBW/day)	0	No data
IR_I	Inhalation rate (m^3/day)	0.08	Allometric, 0.4089 $BW^{0.77}$ (EPA 1993b, Eq. 3-19)

[a] Food ingestion rate (g/g – d) reexpressed as kg/kg BW/day is assumed not to include ingested soil; therefore, PF + AF = 1.0.
[b] Values used for the Burrowing Owl are estimates calculated from values established for the Great Horned Owl (*Bubo virginianus*).

Red-tailed Hawk (*Buteo jamaicensis*)

The red-tailed hawk may be found on the Hanford Site year-round (Fitzner and Gray 1991). Forty-one nesting pairs of hawks (red-tailed, Swainson's and ferruginous) were observed on-site during the 1994 breeding season (Neitzel et al. 1998). Nests were constructed in trees, cliffs, basalt outcrops, and high-voltage transmission line towers (Neitzel et al. 1998).

The red-tailed hawk is a diurnal predator of rodents and other small mammals, including mice, shrews, voles, rabbits, and squirrels. Generally opportunistic, the red-tailed hawk feeds on whatever is most abundant and readily available. Red-tailed hawks maintain a territory year-round (Brown and Amadon 1968).



Parameter	Definition	Value	Reference/Notes
BW	Body weight (kg)	1.055	Average of adult male and female, southwest Idaho (Steenhof 1983 in EPA 1993b)
HR	Home range (ha)	1,770	Adult, both , Colorado upland prairie (Andersen and Rongstad 1989 in EPA 1993b)
TUF	Temporal use factor	1	Will be 1 unless specific value exists
IR_F	Food ingestion rate (g/g-d=kg/kgBW/day)[a]	0.105	Average of adult male and female, Michigan, captive, outdoors (Craighead and Craighead 1956 in EPA 1993b)
PF	Plant fraction	0	Estimated
AF	Animal fraction	1	Estimated
SF	Soil fraction	0	No data
IR_W	Water ingestion rate (g/g-d=L/kgBW/day)	0	No data
IR_i	Inhalation rate (m^3/day)	0.43	Allometric, 0.4089 $BW^{0.77}$ (EPA 1993b, Eq. 3-19)

[a] Food ingestion rate (g/g – d) reexpressed as kg/kg BW/day is assumed not to include ingested soil; therefore, PF + AF = 1.0.

Canada Goose (*Branta canadensis*)

Canada geese forage primarily in open fields, feeding on grains, grass sprouts, and some aquatic vegetation. Breeding habitats includes tallgrass prairies and shortgrass prairies, marshes, ponds, and lakes. Most nesting sites are close to open water, often on islands (EPA 1993b). The Canada goose uses islands along the Hanford Reach extensively for nesting. Studies on the nesting habits of geese that use the Hanford Site have been ongoing since 1953. These studies indicate a general decline over the years in numbers of nests on islands in the Hanford Reach because of heavy predation by coyotes (Cushing et al. 1995).

Parameter	Definition	Value	Reference/Notes
BW	Body weight (kg)	2.620	Average of adult male and female, *B.c. parvipes* (Grieb 1970 in EPA 1993b)
HR	Home range (ha)	983+/- 822 SD	Adult female and brood, *B.c. mofitti*, river (Eberhardt et al. 1989a in EPA 1993b)
TUF	Temporal use factor	1	Will be 1 unless specific value exists
IR_F	Food ingestion rate (g/g-d=kg/kgBW/day)[a]	0.032	Average of adult male and female, winter and spring, *B.c. interior* (Joyner et al. 1984 in EPA 1993b)
PF	Plant fraction	1	Craven and Hunt 1984; Prevett et al. 1985 in EPA 1993b
AF	Animal fraction	0	< 1% invertebrates (Yelverton and Quay 1959; Prevett et al. 1985 in EPA 1993b)
SF	Soil fraction	0	No data
IR_W	Water ingestion rate (g/g-d=L/kgBW/day)	0.044	Average of adult male and female, *B.c. mofitti*, estimated (EPA 1993b)
IR_I	Inhalation rate (m^3/day)	0.86	Allometric, 0.4089 $BW^{0.77}$ (EPA 1993b, Eq. 3-19)

[a] Food ingestion rate (g/g – d) reexpressed as kg/kg BW/day is assumed not to include ingested soil; therefore, PF + AF = 1.0.

Spotted Sandpiper (*Actitus macularia*)

The spotted sandpiper requires open water for drinking, semi-open habitat for nesting, and dense vegetation for breeding (Bent 1929 and Oring et al. 1983). The nest is a grassy scrape near water or in brush with a determinate clutch size of four eggs. Several clutches may be laid during a given breeding season. The diet of the spotted sandpiper consists mostly of terrestrial and aquatic insects (Bent 1929), with adult flying insects making up the bulk of the diet (Oring et al. 1983).



Parameter	Definition	Value	Reference/Notes
BW	Body weight (kg)	0.0425	Maxson and Oring 1980 in EPA 1993b
HR	Home range (ha)	0.25	Territory, Maxson and Oring 1980 in EPA 1993b
TUF	Temporal use factor	0.4	Estimated, 5 months (May-September)
IR_F	Food ingestion rate (g/g-d=kg/kgBW/day)[a]	no data	Estimated metabolic rate of 448 kcal/kg-day (EPA 1993b)
PF	Plant fraction	0	Estimated
AF	Animal fraction	1	Estimated
SF	Soil fraction	0	No data yet; likely in excess of 0.2
IR_W	Water ingestion rate (g/g-d=L/kgBW/day)	0.044	Average of adult male and female, *B.c. mofitti*, estimated (EPA 1993b)
IR_I	Inhalation rate (m^3/day)	0.036	Allometric, 0.4089 $BW^{0.77}$ (EPA 1993b, Eq. 3-19)

[a] Food ingestion rate (g/g – d) reexpressed as kg/kg BW/day is assumed not to include ingested soil; therefore, PF + AF = 1.0.

Great Blue Heron (*Ardea herodius*)

Great blue herons are year-round residents of the Hanford Reach. This bird is relatively common along the Hanford Reach (Fitzner and Gray 1991). Some of the trees planted on pre-1943 farms have persisted and serve as nesting platforms for several species of birds, including the great blue herons (DOE-RL 1995). Its nest is a platform of sticks lined with finer material and is sometimes found on the ground or in a reedbed. Principal prey items of the great blue heron are fish and frogs, although it will also feed on small mammals, reptiles, and occasionally birds.



Parameter	Definition	Value	Reference/Notes
BW	Body weight (kg)	2.39	Arithmetic mean, adult, both sexes, location not stated (EPA 1993b)
HR	Home range (km)	3.1	Foraging distance, mean, adults, both sexes, South Dakota, stream (EPA 1993b)
TUF	Temporal use factor	1	Will be 1 unless a specific value exists
IR_F	Food ingestion rate (g/g-d=kg/kgBW/day)[a]	0.18	EPA 1993b
PF	Plant fraction	0	None listed as dietary intake in EPA 1993b
AF	Animal fraction	1	98% aquatic vertebrates, a river in lower Michigan (EPA 1993b)
SF	Soil fraction	0	Not reported in EPA 1993b; assumed to be negligible
IR_W	Water ingestion rate (g/g-d=L/kgBW/day)	0.045	Estimated (EPA 1993b)
IR_I	Inhalation rate (m^3/day)	0.080	Allometric, 0.4089 $BW^{0.77}$ (EPA 1993b, Eq. 3-19)

[a] Food ingestion rate (g/g – d) reexpressed as kg/kg BW/day is assumed not to include ingested soil; therefore, PF + AF = 1.0.

Bald Eagle (*Haliaetus leucocephalus*)

Bald eagles use trees during the day for perching and occasionally at night for communal roosts (DOE and Ecology 1996). The bald eagle is a common winter resident, usually arriving in October. These birds forage throughout the Hanford Reach. Wintering eagles tend to concentrate where food is abundant and with minimal human disturbance. The diet of bald eagles varies locally as well as seasonally. Food may vary from spawned salmon and waterfowl (often killed by other predators or disease) during the winter to fish, small mammals, carrion, and waterfowl during the breeding season (EPA 1993b). Although bald eagles exhibit nesting behavior at the Hanford Site, most leave before laying eggs (WHC 1994).

Parameter	Definition	Value	Reference/Notes
BW	Body weight (kg)	4.22	Adult average, male and female (Wiemeyer, pers. com., 1991 in EPA 1993b; Johnsgard 1990 in Higley and Kuperman 1996)
HR	Home range (ha)	1,880	Near Missouri Lake (Griffin and Baskett 1985 in Higley and Kuperman 1996)
TUF	Temporal use factor	1	Will be 1 unless specific value exists.
IR_F	Food ingestion rate (g/g-d=kg/kgBW/day)[a]	0.12	Adult, both sexes, Washington, free-flying (Stalmaster and Gessaman 1984 in EPA 1993b)
PF	Plant fraction	0	Estimated
AF	Animal fraction	1	Estimated
SF	Soil fraction	0	No data
IR_W	Water ingestion rate (g/g-d=L/kgBW/day)	0.036	(Calder and Braun 1983 in Higley and Kuperman 1996)
IR_i	Inhalation rate (m^3/day)	1.24	Allometric, 0.4089 $BW^{0.77}$ (EPA 1993b, Eq. 3-19)

[a] Food ingestion rate (g/g – d) reexpressed as kg/kg BW/day is assumed not to include ingested soil; therefore, PF + AF = 1.0.

Mink (*Mustela vision*)

The mink is the most abundant and widespread carnivorous mammal in North America. The home range of mink encompasses both their foraging areas around waterways and their dens along the Columbia River. The mink is found in aquatic habitats of all kinds, including waterways such as rivers, streams, lakes, and ditches, as well as swamps, marshes, and backwater areas (Linscombe et al, 1982 in EPA 1993b). Mink are particularly sensitive to certain chemicals. Minks are predominantly nocturnal hunters, although they are sometimes active during the day. They can often be found along the Columbia River. Mammals are the mink's most important prey year-round in many parts of their range (Eagle and Whitman, 1987), but mink also hunt aquatic prey (e.g., fish, amphibians, and crustaceans) and other terrestrial prey (e.g., bird, reptiles, and insects) depending on the season (Linscombe et al, 1982 in EPA 1993b).



Parameter	Definition	Value	Reference/Notes
BW	Body weight (kg)	0.85	Average of adult male and female (summer and fall) (Mitchell, 1961 in EPA, 1993b)
HR	Home range (ha)	2.24	Foraging distance, mean, adults, both sexes, Sweden/stream (Gerell, 1970 in EPA, 1993b)
TUF	Temporal use factor	1	Will be 1 unless specific value exists.
IR_F	Food ingestion rate (g/g-d=kg/kgBW/day)[a]	0.14	Michigan (farm raised); (Bleavins and Aulerich, 1981 in EPA, 1993b)
PF	Plant fraction	0.02	Michigan/stream, river (%wet wt; stomach contents) (Alexander, 1977 in EPA, 1993b)
AF	Animal fraction	0.98	Michigan/stream, river (%wet wt; stomach contents) Alexander, 1977 in EPA, 1993b)
SF	Soil fraction	0	To be determined
IR_W	Water ingestion rate (g/g-d=L/kgBW/day)	0.11	Estimated in EPA (1993b)
IR_i	Inhalation rate (m^3/day)	0.48	Allimetric, 0.5458 $BW^{0.80}$ (EPA 1993b, Eq. 3-20)

[a] Food ingestion rate (g/g – d) reexpressed as kg/kg BW/day is assumed not to include ingested soil; therefore, PF + AF = 1.0.

8.2. Exposure Assessment

Estimation of the risk to ecological receptors from COPCs and ROPCs in environmental media at an exposure location requires an estimate of exposure and a toxicity reference value (TRV), that is, an exposure associated with little or no adverse effect. Exposure locations at the Hanford Site are areas within the deposition grid at which ecological receptors come into contact with COPCs and ROPCs in media contaminated by stack emissions. Contamination at a given location is represented by modeled concentrations of COPCs and ROPCs in environmental media. The ecological hazard quotient (HQ) is a risk estimate for the receptor at the exposure location and is calculated as the ratio of the estimated exposure and the TRV for a specific COPC or ROPC. The hazard index (HI) is the sum of the HQs for all COPCs with similar modes of toxicity. HQs and HIs exceeding the EPA specified threshold value (0.25) indicate that the receptor has an unacceptable probability of experiencing an adverse effect from exposure to the COPC(s).

The equations used to estimate exposure for terrestrial and aquatic receptors near the RPP-WTP at the Hanford Site are described below (section 8.2.2). Two types of exposure estimates are required. One, the exposure estimate for a wildlife receptor that does not live in a medium containing COPCs but is exposed by ingestion, inhalation, and other routes is the estimated average daily dose (ADD). The exposure equations for wildlife are variations of wildlife exposure equations from EPA (1998a, p. 68, Text Box 4-10) and implied in other sources (EPA 1997b). The equations for ecological receptors are functionally equivalent to the equations in section 7.1 that are used to quantify exposure of humans by ingestion of contaminated food (EPA 1998a). All these equations calculate the amount of contaminant ingested or inhaled per unit biomass per unit time by multiplying the concentration of the contaminant in the ingested medium (abiotic medium or food item) by the receptor's ingestion or inhalation rate for that medium and, explicitly or implicitly, by the fraction of time the receptor is exposed (TUF), dividing where necessary by the body weight of the receptor. The equations for specific wildlife receptors in the RPP-WTP work plan are different from the human health exposure equations (EPA 1998a) because they include a term for the fraction of contaminated space used by the receptor (AUF). The wildlife equations also allow for the contaminant concentration in a food item to be calculated as the product of the contaminant concentration in an ingested abiotic medium and the bioaccumulation (transfer) factor for the medium and the food item being ingested. Two, for receptors living immersed in a medium containing COPC/ROPCs (e.g., vegetation and soil invertebrates living in soil, fish and other aquatic life living in surface water, and sediment-dwelling organisms living in sediment), the exposure estimate is the concentration of COPC/ROPC in the medium.

The modeled whole-body concentrations of contaminants in plants and fish consumed by both humans and non-human receptors will be convergent. This is not to say that receptors that ingest whole organisms will be exposed to the same contaminant concentration as receptors that eat only particular organs or tissues of the organism. For instance, the whole-body concentration in a fish eaten by the great blue heron may not equal the concentration in the fillets of that fish eaten by humans. Whole-body tissue concentrations will be calculated in the human health and ecological risk assessments using identical bioaccumulation factors, ingestion rates, and other parameters, and identical model equations. The ecological risk assessment will use these modeled whole-body concentrations to estimate doses to wildlife receptors.

The exposure assessment will evaluate an exclusive diet and an equal diet in the PRA and a realistic diet in the FRA. An "exclusive diet" is a diet consisting of a single type of prey or food. An "equal diet" is a diet where the fractions of all types of prey or food reportedly eaten by the receptor are set equal to the average fraction reported for that receptor or a similar species. A "realistic diet" is a diet where the fractions of different types of prey or food eaten are more or less the fractions reported to actually occur in one or more cases for the receptor or similar species. The exclusive-diet scenario will be evaluated

when it is a worst-case scenario (that is, it gives the most conservative risk estimate). The realistic-diet scenario will be evaluated because it is realistic. An equal-diet scenario will be evaluated in the PRA because it provides a "standard" diet that does not require quantitative information on the diet of receptors and, therefore, can be easily implemented.

For a given receptor, there is no one exclusive diet for all COPC/ROPC that is the worst-case diet. The worst-case diet for a given COPC/ROPC is an exclusive diet of the prey type that has the highest expected body burden of that COPC/ROPC among all the receptor's potential prey or food types. Which prey or food type will have the highest expected body burden varies by COPC/ROPC as a function of the diets of the prey types, and the magnitudes of the BAFs for the different prey or food types. For example, a COPC/ROPC that bioaccumulates more in plants than in worms may result in mule deer being the worst-case diet for a coyote, whereas a COPC/ROPC that bioaccumulates more in worms than plants may result in the pocket mouse being the worst-case diet for the coyote. In general, the fractions of prey or food types in a given animal's diet, the body burdens in each prey or food type, and the animal's BAF for the COPC/ROPC determines the animal's body burden and, thus, the exposure to its predator. Assuming an exclusive diet, the dietary fractions are either 1 or 0, so the worst-case diet for a given receptor is determined by the product of the uptake and bioaccumulation factors along each possible food chain from the source medium to the receptor. To identify the worst-case diet for each COPC/ROPC for all receptors, the product of the uptake and bioaccumulation factors can be calculated for all food chains of all lengths leading to all receptors.

The equal diet gives a less-accurate exposure estimate than the realistic diet. The equal-diet scenario overestimates the exposure from those dietary items that are actually less than the average dietary fraction and underestimates the exposure from those dietary items that are actually more than the average dietary fraction. Thus, while easier to model because fewer data are required, the equal- proportions diet is less accurate than a realistic diet. The equal-diet scenario is also not consistently less conservative than the realistic-diet scenario and, therefore, does not serve with the worst-case scenario to bracket the range of possible exposures. An equal-proportion diet is an appropriate, unbiased model when no dietary fraction information is available. The equal-proportion diet will be evaluated in the PRA. Dietary fractions are expected to be available for all receptors to be evaluated in the risk assessment. Therefore, the realistic-diet scenario will be evaluated in the FRA.

Concentrations used to estimate exposure for ecological receptors, exposure equations for terrestrial and aquatic receptors, and the variables and parameters used in these equations to estimate exposures for ecological receptors are provided in the following sections.

8.2.1. Exposure Point Concentration

Exposure of ecological receptors to COPCs and ROPCs in the vicinity of the RPP-WTP will be estimated from the distribution of concentrations predicted by the aerial dispersion and other fate and transport models (section 6). Dispersion model output concentrations will be used to calculate exposure concentrations for surface soil (mg/kg, pCi/g), gases and particulates in air (mg/m^3, pCi/m^3), sediment (mg/kg, pCi/g) and surface water (µg/L, pCi/L). For any specified exposure location on the dispersion grid, the modeled concentration will be used to estimate the exposure to ecological receptors. Use of area-weighted modeled concentrations represents a conservative estimate of potential exposure due to the RPP-WTP.

Exposure point concentrations used to estimate doses of both COPCs and ROPCs are dependent on the location of the receptor. The location of the various receptor populations identified for the quantitative, screening risk assessment will correspond to the receptor grid nodes defined during air dispersion modeling (section 6). It is expected that two important places are Gable Mountain and the Hanford Site

boundary. Other places are those of maximum contaminant concentration. In keeping with the protective approach used in this risk assessment, the location with the maximum concentration of COPCs and ROPCs will be used in estimating exposure point concentrations. Because the point of maximum concentration may be different for airborne contaminants deposited via wet and dry deposition mechanisms, EPA 1998a recommends the following method for selecting the point of maximum concentration. Within each area to be evaluated, receptor grid nodes or exposure locations with the highest modeled unitized value will be identified for each of the following parameters:

- Vapor phase air concentration
- Vapor phase wet deposition rate
- Particle phase air concentration
- Particle phase wet deposition rate
- Particle phase dry deposition rate
- Particle-bound phase air concentration
- Particle-bound phase wet deposition rate
- Particle-bound phase dry deposition rate

At each of the unique locations, air concentrations from vapor, particle phase, and particle bound fractions will be summed, and concentrations of deposited materials will be summed to determine the point or location of area-weighted maximum exposure. Air concentrations and deposition rates will not be summed together.

8.2.2. Terrestrial Receptors

For receptors living in soil (e.g., vegetation and soil invertebrates), the HQ is calculated as the ratio of the concentration of COPC in soil and the TRV for the receptor and the COPC. That is:

$$HQ_{ijk} = C_{jk}/TRV_{ij},$$

where

HQ_{ijk} = hazard quotient for receptor i at exposure location k for COPC j (unitless)
C_{jk} = concentration of COPC j in soil at exposure location k (mg/kg_{soil})
TRV_{ij} = toxicity reference value of receptor i for COPC j (mg/kg_{soil}).

The HQ for a wildlife receptor that does not live in the medium containing COPCs but is exposed by ingestion, inhalation, and other routes is calculated as the ratio of the ADD and the TRV. That is:

$$HQ_{ijk} = ADD_{ijk}/TRV_{ij}$$

where

ADD_{ijk} = average daily dose of COPC j to receptor i at exposure location k (mg/kg BW/day) calculated using concentration of COPC j at exposure location k (units vary)
TRV_{ij} = toxicity reference value of receptor i for COPC j (mg/kg BW/day).

The second equation is used to estimate risk for the wildlife receptors in the terrestrial food web: mule deer, Great Basin pocket mice, Western meadowlark, coyote, burrowing owl, and red-tailed hawk. The RME concentrations of COPCs/ROPCs in surface soil, plant and animal tissue, and air, ingestion and inhalation rates, body weights, and bioaccumulation factors (Appendix C-2) will be used to estimate the ADDs for receptors, as published in the receptor profiles and as described below.

8.2.2.1. Ingestion Exposure Calculations (Soil to Wildlife)

For wildlife receptors, such as mammals and birds, that are exposed to COPCs/ROPCs in soil by ingestion of soil, ingestion of terrestrial biota, or both, the ADD is calculated as follows:

Herbivores, Mid-level Predators, and Omnivores

$$ADD_{ijk} = AUF_{ik} \times TUF_{ik} \times [(Cr_{jk} \times IPr_i) + (Cv_{jk} \times IPv_i) + (Cd_{jk} \times IAd_i) + (C_{jk} \times IS_i \times AEs_j)]/BW_i$$
$$= AUF_{ik} \times TUF_{ik} \times C_{jk} \times [(IPr_i \times SPr_j) + (IPv_i \times SPv_j) + (IAd_i \times BAF\text{-}S_j) + (IS_i \times AEs_j)]/BW_i$$

where

AUF_{ik} = Area Use Factor for receptor i at exposure location k (unitless) =

$$\begin{cases} 1 & \text{if } A_k \geq HR_i \\ A_k / HR_i & \text{if } A_k < HR_i \end{cases}$$

where

A_k = Area of exposure location k (ha)
HR_i = Home range, foraging range, or territory size of receptor i (ha)
TUF_{ik} = Temporal Use Factor for receptor i at exposure location k (unitless) = 1 unless receptor-specific value is available
Cr_{jk} = Concentration of COPC j in plant reproductive tissue (fruit, nuts, flowers) at exposure location k (mg/kg-wet wt.) = RME tissue concentration, unless otherwise directed
IPr_i = Rate of ingestion of plant reproductive tissue by receptor i (kg/day)
Cv_{jk} = Concentration of COPC j in plant vegetative tissue (leaf, stem, root) at exposure location k (mg/kg-wet wt.) = RME tissue concentration, unless otherwise directed
IPv_i = Rate of ingestion of plant vegetative tissue of receptor i (kg/day)
Cd_{jk} = Concentration of COPC j in terrestrial invertebrate tissue at exposure location k (mg/kg-wet wt.) = RME tissue concentration, unless otherwise directed
IAd_i = Rate of ingestion of terrestrial invertebrate tissue by receptor i (kg/day)
C_{jk} = Concentration of COPC j in soil at exposure location k (mg/kg-dry wt.) = RME_{jk} soil concentration, unless otherwise directed
IS_i = Rate of soil ingestion (kg-dry wt./day) by receptor i = IF_i x SI_i, where IF_i is the total food ingestion rate (kg-dry wt./day) of receptor i and SI_i is the fraction of food ingested by receptor i that is soil, on a dry weight basis
AEs_j = Absorption efficiency factor for COPC j in ingested soil (unitless) = 1, unless chemical-specific value is available
BW_i = Body weight of receptor i (kg)
SPr_j = Soil-to-Plant transfer factor for reproductive tissue for COPC j (kg_{soil}-dry wt./kg_{tissue}-wet wt.)
SPv_j = Soil-to-Plant transfer factor for vegetative tissue for COPC j (kg_{soil}-dry wt./kg_{tissue}-wet wt.)
$BAF\text{-}S_j$ = Soil-to-soil invertebrate tissue transfer factor for COPC j (kg_{soil}-dry wt./kg_{tissue}-wet wt.)

For wildlife receptors that are exposed to COPCs/ROPCs in soil by ingestion of terrestrial biota, the ADD is calculated as follows:

Top Predators

$$ADD_{ijk} = AUF_{ik} \times TUF_{ik} \times [\Sigma_n(Cn_{jk} \times IAn_i) + (C_{jk} \times IS_i \times AEs_j)]/BW_i$$

where

$$Cn_{jk} = BAF\text{-}T_{nj} \; x \; C_{jk} \times [(SPr_j \times FPr_n) + (SPv_j \times FPv_n) + (BAF\text{-}S_j \times FA_n) + (AEs_j \times FS_n)]$$

and

Cn_{jk} = Concentration of COPC j in animal tissue of prey type n at exposure location k (mg/kg-wet wt.) = RME tissue concentration, unless otherwise directed

IAn_i = Rate of ingestion of tissue of animal prey type n by receptor i (kg/day)

$BAF\text{-}T_{nj}$ = Tissue-to-tissue transfer factor for prey type n for COPC j (kg_{tissue}-wet wt./kg)

FPr_n = Fraction of total daily ingestion by prey type n that is plant reproductive tissue (unitless) = IPr_n/IT_n

FPv_n = Fraction of total daily ingestion by prey type n that is plant vegetative tissue (unitless) = IPv_n/IT_n

FA_n = Fraction of total daily ingestion by prey type n that is terrestrial invertebrate tissue (unitless) = IAd_n/IT_n

FS_n = Fraction of total daily ingestion by prey type n that is soil (unitless) = IS_n/IT_n,

and

$$IT_n = IPr_n + IPv_n + IAd_n + IS_n$$

where

IT_n = the total daily ingestion by prey type n of receptor i (kg/day)

IPr_n = Rate of ingestion of plant reproductive tissue by prey type n of receptor i (kg/day)

IPv_n = Rate of ingestion of plant vegetative tissue by prey type n of receptor i (kg/day)

IAd_n = Rate of ingestion of terrestrial invertebrate tissue by prey type n of receptor i (kg/day)

IS_n = Rate of soil ingestion (kg-dry wt./day) by prey species n of receptor i = $IF_n \times SI_n$ where IF_n is the total food ingestion rate (kg-dry wt./day) of prey species n of receptor i and SI_n is the fraction of food ingested by prey species n that is soil, on a dry weight basis

Soil ingestion rates for Hanford receptors (Table 8-2) are required to calculate ADDs. Receptors are exposed to contaminants in soil by direct and indirect soil ingestion. Direct soil ingestion occurs during grooming, burrowing, and ingesting soil attached to the outside of the food. Indirect soil ingestion occurs during feeding on prey that contain soil in the gut. Soil ingestion fractions (direct and indirect soil ingestion) for two out of eleven Hanford receptors (Canada goose and mule deer) are documented in EPA (1993b). These values (Table 8-2) are derived from data in Beyer et al. (1994), EPA (1993), and Higley and Kuperman (1996). For three receptors (coyote, Great Basin pocket mouse, and spotted sandpiper), soil ingestion fractions for surrogate species documented in Beyer et al. (1994) are used to estimate direct and indirect soil ingestion. The indirect soil ingestion rates for the remaining six receptors without published values (bald eagle, red-tailed hawk, burrowing owl, western meadowlark, great blue heron, and mink) are estimated as described below. Direct soil ingestion is not estimated for these six receptors because direct soil ingestion is assumed to be a minor contribution compared to indirect soil ingestion.

Indirect soil ingestion rates for the six receptors without published values are calculated from the soil-to-body weight ratios (SBRs) of their prey and the proportions of different prey in their diets. The SBR of prey species with units kg_{soil}/kg is calculated as the product of the soil ingestion fraction (kg_{soil}/kg_{food}), the daily food ingestion rate per unit body weight ($kg_{food}/kg/d$) and the residence time (d) of soil in the prey gut (Table 8-3). To derive the indirect soil ingestion rate, a receptor's daily food ingestion rate per unit body weight is multiplied by the sum of the products of the dietary fraction and SBR of each prey type (Table 8-4). Indirect soil ingestion rates for the six Hanford receptors will also be calculated for the equal and exclusive diets evaluated in the PRA.

8.2.2.2. Inhalation Exposure Calculations (Air to Wildlife)

For wildlife receptors such as mammals and birds that are exposed to chemicals emitted from the stack(s) by inhalation of particulates and gases in air, the ADD is calculated as follows:

All Terrestrial Receptors

$$ADD_{ijk} = AUF_{ik} \times TUF_{ik} \times [(Dg_{jk} \times AEg_j) + (Dp_{jk} \times AEp_j)] \times IR_i / BW_i$$

where

AUF_{ik} = Area Use Factor for receptor i at exposure location k (unitless) =

$$\begin{cases} 1 & \text{if } A_k \geq HR_i \\ A_k / HR_i & \text{if } A_k < HR_i \end{cases}$$

where

A_k = Area of exposure location k (ha)
HR_i = Home range, foraging range, or territory size of receptor i (ha)
TUF_{ik} = Temporal Use Factor for receptor i at exposure location k (unitless) = 1, unless receptor-specific value is available
Dg_{jk} = Density of COPC j in air vapors or gases at exposure location k (mg/m^3)
Dp_{jk} = Density of COPC j in air particulates at exposure location k (mg/m^3)
AEg_j = Absorption efficiency factor for COPC j in inhaled gases or vapors (unitless) = 1, unless chemical-specific value is available
AEp_j = Absorption efficiency factor for COPC j in inhaled particulates (unitless) = 1, unless chemical-specific value is available
IR_i = Rate of inhalation of particulates and gases by receptor i (m^3/day).

EPA (1993) presents inhalation rates (m^3/d) for a few bird and mammal receptor species and regression equations to derive inhalation rates associated with standard metabolic rate from body weight. The inhalation rates used in the work plan are in the receptor parameter tables-profiles (Appendix Table C).

8.2.2.3. Radiological Exposure Equations

Ecological receptors are exposed internally and externally to ROPCs in terrestrial habitats within the deposition grid surrounding the RPP-WTP at the Hanford Site. The risk from exposure to ROPCs is calculated as the ratio of the estimated total-body dose rate (R_{ik}) and the TRV. That is:

$$HQ_{ik} = R_{ik}/TRV_i$$

where

HQ_{ik} = hazard quotient for receptor *i* at exposure location *k* for all ROPCs (unitless)
R_{ik} = total body dose rate to receptor *i* at exposure location *k* (rem/day)
TRV_i = toxicity reference value (dose rate) of receptor *i* for ROPCs (rem/day).

The total-body dose rate of terrestrial ecological receptors results from both internal and external exposures to all ROPCs, which are summed to estimate the combined exposure. That is:

$$R_{ik} = \Sigma_j\ [R_{ijk}(\text{internal}) + R_{ijk}(\text{external})].$$

where

R_{ijk} = total-body dose to receptor *i* from indicated exposure to ROPC *j* at location *k* (rem/day).

Internal and external dose rates for terrestrial receptors are calculated as described below.

8.2.2.4. Internal Exposure

The internal total-body dose rate to a terrestrial receptor from exposure to a single ROPC is

$$R_{ijk} = CFa \times [Co_{jk} \times (QF \times \Phi_\gamma E_\gamma n_\gamma + QF \times \Phi_\beta E_\beta n_\beta + QF \times \Phi_\alpha E_\alpha n_\alpha)]$$

where

R_{ijk} = total-body dose to receptor *i* from internal exposure to ROPC *j* at location *k* (rem/day)
CFa = unit conversion factor from pCi MeV/g disintegration to rad/day = 5.11×10^{-5} rad/day / (pCi MeV/g disintegration) [(3.70×10^{-2} dis/s /pCi) x (1.60×10^{-6} erg/MeV) x (1.00×10^{-7} Joule/erg) x (1.00×10^{2} rad x kg tissue/Joule) x (1.00×10^{3} g tissue / kg tissue) x (1 g soil/g tissue) x (8.64×10^{4} s/d)]
Co_{jk} = concentration of ROPC *j* in organism *i* at exposure location *k* (pCi/g wet weight)
QF = quality factor to account for relative effectiveness of radiation types: one rem/rad for β and γ and 20 rem/rad for α radiation
Φ_m = absorbed fraction of energy emitted by α, β, or γ radiation (unitless), where m = α, β, or γ
E_m = average effective energy emitted by α, β, or γ radiation (MeV per disintegration)
n_m = proportion of disintegrations producing α, β, or γ radiation (unitless).

It is assumed that n_m is contained along with E (as $E_\gamma n_\gamma$, $E_\beta n_\beta$, and $E_\alpha n_\alpha$) in the decay energy terms from Table A.1 of *External Exposure to Radionuclides in Air, Water, and Soil, Federal Guidance Report No. 12* (Eckerman and Ryman 1993). The values of Φ_β and Φ_γ come from *Methodology for Estimating Radiation Dose Rates to Freshwater Biota Exposed to Radionuclides in the Environment* (Blaylock et al. 1993) or *Methods and Tools for Estimation of the Exposure of Terrestrial Wildlife to Contaminants* (Sample et al. 1997). The value of Φ_α is 1.0 for all radionuclides (Blaylock et al. 1993).

The internal concentration in a terrestrial plant or terrestrial invertebrate from exposure to a single ROPC is calculated by using soil-to-biota transfer factors and the radionuclide activity in soil. For plants,

Co_{jk} = SPr_j x C_{jk}

and for terrestrial invertebrates,

Co_{jk} = $BAF\text{-}S_j$ x C_{jk}

where

SPr_j = Soil-to-plant transfer factor for reproductive tissue for ROPC *j* (kg_{soil}-dry wt./kg_{tissue}-wet wt.)

$BAF\text{-}S_j$ = Soil-to-tissue transfer factor for terrestrial invertebrates for ROPC *j* (kg_{soil}-dry wt./kg_{tissue}-wet wt.)

The concentration of a radionuclide in a mammal or bird receptor results from the inhalation or ingestion of material that contains radionuclides and the uptake and effective bioaccumulation in the receptor. For ingestion:

$$Co_{jk} = Cm_{ijk} \times BAF\text{-}T_{ij} \times SIR_i$$

where

Cm_{ijk} = the average concentration of ROPC *j* in the matter ingested by receptor *i* at location *k* (pCi/g)

$BAF\text{-}T_{ij}$ = the bioaccumulation factor representing the retention of ROPC *j* in tissue of receptor *i* resulting from the ingestion rate of ROPC *j* by receptor *i* [pCi retained / (pCi ingested / day)].

SIR_i = specific ingestion rate or receptor *i* (kg matter/kg body weight/day) = $IT_I + IN_i$

Values of $BAF\text{-}T_{ij}$ (Appendix Table C-2-2) were derived from ingestion-to-beef uptake factors presented by Baes et al. (1984) [in units of (pCi retained/kg tissue)/(pCi ingested/day)]. The Baes et al. (1984) values were multiplied by an average beef body weight at slaughter of 550 kg to calculate a whole-body retention rate in units of pCi retained/pCi ingested/day (see Section 8.2.4.3). The same values (Appendix Table C-2-2) were used for both ingestion exposure and inhalation exposure.

The average concentration of a radionuclide in the material ingested or inhaled by a receptor (C_{jk}) is calculated using the equation for the receptor's ADD for that substance in its prey and in soil or other abiotic media ingested or inhaled, that is:

$$Cm_{jk} = ADD_{ijk} / SIR_i$$

where

ADD_{ijk} = average daily dose of ROPC *j* to receptor *i* at location *k* (pCi/g/day) by inhalation or ingestion

SIR_i = specific ingestion rate or receptor *i* (kg matter/kg body weight/day) = $IT_I + IN_i$

For inhalation, the contribution to body burden is given by:

$$Co_{jk(inhaled)} = AUF_{ik} \times TUF_{ik} \times Dp_{jk} \times AEp_j \times BAF\text{-}T_j \times IR_i / (BW_i \times 1000)$$

where

AUF_{ik} = Area Use Factor for receptor i at exposure location k (unitless) =
1 if A_k HR_i
A_k/HR_i if $A_k < HR_i$

where

A_k = Area of exposure location k (ha)
HR_i = Home range, foraging range, or territory size of receptor i (ha)

TUF_{ik} = Temporal Use Factor for receptor i at exposure location k (unitless) = 1, unless receptor-specific value is available
Dp_{jk} = Density of ROPC j in air particulates at exposure location k (pCi/m^3)
AEp_j = Absorption efficiency factor for ROPC j in inhaled particles (unitless) = 1, unless radionuclide-specific value is available
$BAF\text{-}T_j$ = Bioaccumulation factor representing the retention of ROPC j in tissue of receptor i resulting from the inhalation rate of ROPC j [pCi retained / (pCi inhaled/day)]
IR_i = Rate of inhalation of particulates in air by receptor i (m^3/day)
BW_i = Body weight of receptor i (kg)
1000 = Conversion factor, g/kg

The total body burden of ROPC j is found by adding body burdens from inhaled and ingested radionuclides:

$$Co_{ijk(total)} = Co_{ijk(ingested)} + Co_{ijk(inhaled)}$$

8.2.2.5. External Exposure

The external total-body dose rate to a terrestrial receptor, including plants, from aboveground exposure to a single ROPC is:

$$R_{ijk} = AUF_{ik} \times TUF_{ik} \times F_{above\,i} \times F_{rufk} \times CFb \times C_{jk} \times DCFs_j \times h_i$$

where

R_{ijk} = total-body dose to soil surface receptor i from external exposure to ROPC j in soil at location k (rem/day)
AUF_{ik} = Area Use Factor for receptor i at exposure location k (unitless) =
1 if $A_k >= HR_i$
A_k/HR_i if $A_k < HR_i$

A_k = Area of exposure location k (ha)
HR_i = Home range, foraging range, or territory size of receptor i (ha)
TUF_{ik} = Temporal Use Factor for receptor i at exposure location k (unitless) = 1 unless receptor-specific value is available
$F_{above\,i}$ = fraction of time receptor i spends on the ground surface (unitless)
F_{rufk} = dose rate reduction factor accounting for roughness of the ground surface at location k = 0.7 (unitless)
CFb = unit conversion factor from Sv/s per Bq/m^3 to rem/d per pCi/g = 5.12×10^{11} rem/day per pCi/g) / (Sv/s per Bq/m^3)
C_{jk} = concentration of ROPC j in soil at location k (pCi/g)

$DCFs_{jk}$ = dose coefficient for ROPC *j* in soil contaminated to a depth of 15 cm at location *k* (Sv/s per Bq/m^3) (Eckerman and Ryman 1993, Table III.6)

h_i = factor to adjust for height of receptor *i* above the ground surface (Appendix Table C-2-4)

The external DCF values from Eckerman and Ryan (1993) are based on exposures at a height of 1 m (3.28 ft), but Hanford Site receptors are exposed at varying distances from the soil. The height adjustment factor (h) is assumed to be 1 for receptors exposed at approximately 1 m above the ground surface (such as mule deer); h is assumed to be 2 for receptors on the ground surface (such as western meadowlark); and h is assumed to be 2 plants growing close to the ground (that is, cheatgrass).

The external total-body dose rate to a terrestrial receptor from belowground exposure to a single ROPC is:

$$R_{ijk} = AUF_{ik} \times TUF_{ik} \times 1.05 \times F_{below} \times CFa \times C_{jk} \times QF \times (\Phi_\beta \times E_\beta n_\beta + \Phi_\gamma \times E_\gamma n_\gamma)$$

where

R_{ijk} = total-body dose to subsurface receptor *i* from external exposure to ROPC *j* in soil at location *k* (rem/day)

AUF_{ik} = Area Use Factor for receptor *i* at exposure location *k* (unitless) =

$$1 \text{ if } A_k >= HR_i$$
$$A_k/HR_i \text{ if } A_k < HR_i$$

A_k = Area of exposure location *k* (ha)

HR_i = Home range, foraging range, or territory size of receptor *i* (ha)

TUF_{ik} = Temporal Use Factor for receptor *i* at exposure location *k* (unitless) = 1 unless receptor-specific value is available

1.05 = conversion factor to account for immersion in soil rather than water

F_{below} = fraction of time receptor *i* spends below ground in tunnels or burrows (unitless)

CFa = unit conversion factor from pCi/g x MeV/disintegration to rad/day = 5.11×10^{-5} rad/day/[(pCi MeV/(g soil x disintegration)]

C_{jk} = concentration of ROPC *j* in soil at location *k* (pCi/g)

QF = quality factor to account for relative effectiveness of radiation types: 1 rem/rad for β and γ radiation

Φ_m = absorbed fraction of energy emitted by β or γ radiation (unitless), where m = β or γ

E_m = average effective energy emitted by β or γ radiation (MeV/disintegration)

n_m = proportion of disintegrations producing β or γ radiation (unitless)

The dose rate for external radiation from radionuclides suspended in air will be calculated as described by Eckerman and Ryman (1993). The dose rate equation is:

$$R_{ijk} = AUF_{ik} \times TUF_{ik} \times Dp_{jk} \times DCFa_j \times CFc$$

where

R_{ijk} = total-body dose to terrestrial receptor *i* from external exposure to ROPC *j* in air at location *k* (rem/day)

AUF_{ik} = Area Use Factor for receptor *i* at exposure location *k* (unitless) =

$$1 \text{ if } A_k >= HR_i$$
$$A_k/HR_i \text{ if } A_k < HR_i$$

A_k = Area of exposure location k (ha)
HR_i = Home range, foraging range, or territory size of receptor i (ha)
TUF_{ik} = Temporal Use Factor for receptor i at exposure location k (unitless) = 1 unless receptor-specific value is available
Dp_{jk} = density of $ROPC_j$ in suspended particles at location k (pCi/m^3)
$DCFa_j$ = dose conversion factor for exposure to ROPC j by immersion in air (Sv/sec per Bq/m^3) (Eckerman and Ryman 1993, Table III.1)
CFc = unit conversion factor from Sv/sec per Bq/m^3 to rem/day per pCi/m^3 = 3.20 x 10^5 (rem/day per pCi/m^3) / (Sv/s per Bq/m^3)

Concentrations of ROPCs in air (pCi/m^3) derived by fate and transport modeling will be used for the concentration term Dp_{jk}.

The total-body dose from external exposure is added to the total-body dose from internal exposure to calculate the combined total-body dose, R_{ik}.

8.2.3. Aquatic Receptors

For receptors living in surface water or sediment (e.g., aquatic life and salmon and other fish living in surface water, and sediment-dwelling organisms living in sediment), the HQ is calculated as the ratio of the measured concentration of COPC in the medium and the TRV. That is:

$$HQ_{ijk} = C_{jk}/TRV_{ij}$$

where

HQ_{ijk} = hazard quotient for receptor i at exposure location k for COPC j (unitless)
C_{jk} = concentration of COPC j in water or sediment at exposure location k (µg/L or mg/kg)
TRV_{ij} = toxicity reference value of receptor i for COPC j (µg/L or mg/kg).

The HQ for a wildlife receptor that does not live in the surface water or sediment containing the COPCs but is exposed from aquatic food webs by ingestion, inhalation, and other routes is calculated as the ratio of the estimated ADD (mg/kg BW/day) and the TRV (mg/kg BW/day). That is:

$$HQ_{ijk} = ADD_{ijk}/TRV_{ij}$$

The above equation is used to estimate risk for the wildlife receptors in the aquatic food web: Canada goose, spotted sandpiper, great blue heron, bald eagle and mink. The concentrations of COPCs in sediment or surface water, estimated COPC concentrations in plant and animal tissue, ingestion rates, and bio-accumulation factors (Appendix C-2), will be used to estimate the ADDs for these receptors, as described below.

8.2.3.1. Ingestion Exposure Calculation (Sediment to Wildlife)

For animal receptors such as invertebrates and birds that are exposed to chemicals in sediment by ingestion of sediment, ingestion of sediment-associated biota, or both, the exposure is calculated as follows:

$$ADD_{ijk} = AUF_{ik} \times TUF_{ik} \times [(Cp_{jk} \times IP_i) + \Sigma_m(C_{mjk} \times IAm_i) + (Cs_{jk} \times IS_i \times AEs_j)]/BW_i$$
$$= AUF_{ik} \times TUF_{ik} \times Cs_{jk} \times [(IP_i \times SP_j) + \Sigma_m(IAm_i \times BSF_{mj}) + (IS_i \times AEs_j)]/BW_i$$

where

Cp_{jk} = Concentration of COPC j in aquatic plant tissue at exposure location k (mg/kg-wet wt.) = RME tissue concentration, unless otherwise directed
IP_i = Rate of ingestion of aquatic plant tissue by receptor i (kg/day)
Cm_{jk} = Concentration of COPC j in tissue of prey type m at exposure location k (mg/kg-wet wt.) = RME tissue concentration unless otherwise directed
IAm_i = Rate of ingestion of prey type m by receptor i (kg/day)
Cs_{jk} = Concentration of COPC j in sediment at exposure location k (mg/kg-dry wt.) = RME_{jk} sediment concentration
IS_i = Rate of ingestion of sediment by receptor i (kg/day)
AEs_j = Absorption efficiency factor for COPC j in ingested sediment (unitless) = 1, unless chemical-specific value is available
SP_j = Sediment-to-Plant transfer factor for aquatic plant tissue for COPC j ($kg_{sediment}$-dry wt./kg_{tissue}-wet wt.)
BSF_{mj} = Sediment-to-tissue transfer factor for prey m for COPC j ($kg_{sediment}$-dry wt./kg_{tissue}-wet wt.).

Note that BSF in the literature (also referred to as BSAF) is often provided as a ratio of lipid-normalized tissue chemical residue to organic carbon-normalized sediment chemical concentration so that a conversion may be needed to determine BSF_{mj}.

8.2.3.2. Ingestion Exposure Calculation (Surface Water to Animals and Wildlife)

For animal receptors such as mammals and birds that are exposed to chemicals in surface water by ingestion of surface water, ingestion of aquatic biota, or both, the exposure is calculated as follows:

$$ADD_{ijk} = AUF_{ik} \times TUF_{ik} \times [(Cp_{jk} \times IP_i) + \Sigma_m(Cm_{jk} \times IAm_i) + (Cw_{jk} \times IW_i)]/BW_i$$
$$= AUF_{ik} \times TUF_{ik} \times Cw_{jk} \times [(IP_i \times WP_j) + \Sigma_m(IAm_i \times BCF_{mj}) + IW_i]/BW_i$$

where

Cw_{jk} = Concentration of COPC j in surface water at exposure location k (mg/L) = RME_{jk} water concentration, unless otherwise directed
WP_j = Water-to-plant transfer factor for aquatic plant tissue for COPC j (L/kg tissue wet wt., Table C-2-3)
IW_i = Rate of water ingestion by receptor i (L/day)
BCF_{mj} = Surface water-to-tissue transfer factor for prey type m for COPC j (L/kg_{tissue}-wet wt.).

8.2.3.3. Radiological Exposure Calculations

Ecological receptors are exposed internally and externally to ROPCs in aquatic habitats within the deposition grid surrounding the RPP-WTP at the Hanford Site. The risk from exposure to radionuclides is calculated as the ratio of the estimated total-body dose rate (R_{ik}) and the TRV. That is:

$$HQ_{ik} = R_{ik}/TRV_i$$

where

HQ_{ik} = hazard quotient for receptor i at exposure location k for all ROPCs (unitless)

R_{ik} = total body dose rate to receptor *i* at exposure location *k* (rem/day)
TRV_i = toxicity reference value (dose rate) of receptor *i* for ROPCs (rem/day).

The total-body dose rate of aquatic ecological receptors results from both internal and external exposure to all radionuclides, which are summed to estimate the combined exposure. That is:

$$R_{ik} = \Sigma_j \; [R_{ijk}(\text{internal}) + R_{ijk}(\text{external})]$$

Dose rates for aquatic receptors are calculated as in Blaylock et al. (1993).

8.2.3.4. Internal Exposure

The internal total-body dose rate to an aquatic receptor from exposure to a single ROPC is:

$$R_{ijk} = CFa \times [Co_{jk} \times (QF \times \Phi_\gamma E_\gamma n_\gamma + QF \times \Phi_\beta E_\beta n_\beta + QF \times \Phi_\alpha E_\alpha n_\alpha)]$$

where

R_{ijk} = internal total-body dose of ROPC *j* to receptor *i* at exposure location *k* (rem/day)
CFa = unit conversion factor from pCi/g x MeV/dis to rad/day, 5.11×10^{-5}
Co_{jk} = concentration of radionuclide ROPC *j* in organism *i* at exposure location *k* (pCi/g wet weight)
QF = quality factor to account for relative effectiveness of radiation types: 1 rem/rad for β and γ and 20 rem/rad for α radiation
Φ_m = absorbed fraction of each radiation-energy type *m* (dimensionless) where m = α, β , or γ
E_m = average effective energy emitted by α, β, or γ radiation (MeV per disintegration)
n_m = proportion of disintegrations producing α, β, or γ radiation.

According to Blaylock et al. (1993), internal beta radiation for large fish and internal alpha radiation for both small and large fish are completely absorbed (i.e., $\Phi = 1$).

8.2.3.5. External Exposure

The external total-body dose rate to a generic aquatic receptor from exposure to a single ROPC in aquatic habitat is calculated as in Blaylock et al. 1993, that is:

$$R_{ijk} = AUF_{ik} \times TUF_{ik} \times CFa \times [Cw_{jk} \times (1 - F_s - F_{in}) \times 1000 \text{ mL/L} / 1 \text{ g/mL} + Cs_{jk} \times (0.5F_s + F_{in})] \times QF \times [(1-\Phi_\gamma) \times E_\gamma n_\gamma + (1-\Phi_\beta) \times E_\beta n_\beta]$$

where

R_{ijk} = total-body dose to terrestrial receptor *i* from external exposure to ROPC *j* in surface water and sediment at location *k* (rem/day)
AUF_{ik} = Area Use Factor for receptor *i* at exposure location *k* (unitless) =
1 if $A_k >= HR_i$
A_k/HR_i if $A_k < HR_i$

A_k = Area of exposure location *k* (ha)
HR_i = Home range, foraging range, or territory size of receptor *i* (ha)
TUF_{ik} = Temporal Use Factor for receptor *i* at exposure location *k* (unitless) = 1 unless receptor-specific value is available

CFa = unit conversion factor from pCi/g x MeV/disintegration to rad/d = 5.11×10^{-5} rad/day/[pCi MeV/(g x disintegration)]
Cw_{jk} = concentration of ROPC *j* in water at location *k* (pCi/L)
F_s = fraction of time the receptor spends at the sediment-water interface (unitless) [exposure is assumed to be one-half as great at the interface as while immersed (Blaylock et al. 1993)]
F_{in} = fraction of time the receptor spends buried in sediment (unitless)
1000 mL/L = unit conversion factor for volume of water
1 g/mL = unit conversion factor for density of water
Cs_{jk} = concentration of ROPC *j* in sediment at location *k* (pCi/g)
QF = quality factor to account for relative effectiveness of radiation types: 1 rem/rad for β and γ radiation
Φ_m = absorbed fraction of energy emitted by β or γ radiation (unitless), where m = β or γ
E_m = average effective energy emitted by β or γ radiation (MeV/disintegration)
n_m = proportion of disintegrations producing β or γ radiation (unitless)

According to Blaylock et al. (1993), external beta radiation from water and sediment for large fish and external alpha radiation for both small and large fish from water and sediment are negligible.

Where ROPC concentrations are available for sediment but not surface water or surface water but not sediment, screening benchmarks presented by DOE (1998) will be used to screen exposures of fish and aquatic invertebrates.

Note that DOE (1998) has used the methods developed by, and TRUs published by, Blaylock et al (1993) to derive these sediment and surface water radiological benchmarks for aquatic biota.

8.2.4. Exposure Variables

The magnitude of exposure of ecological receptors to COPCs and ROPCs in environmental media depends on various parameters and variables in the above exposure equations. The exposure variables include factors correcting for the fraction of a receptor's total exposure originating at the exposure location, variables determining the rate of uptake into tissues and cells, and factors accounting for the accumulation in one medium of substances present in another medium. The exposure variables for ecological receptors are briefly discussed below.

8.2.4.1. Space and Time Factors for Exposure Calculations

For wildlife receptors that are exposed to COPCs and ROPCs by ingestion and inhalation, the calculation of exposure may require exposure factors that adjust for the fraction of a receptor's exposure obtained from the contaminated site. A receptor may obtain only a fraction of its exposure to a contaminant from the exposure location as a result of the receptor foraging over an area larger than the exposure location, spending only a fraction of its lifetime at the exposure location, or both. The rules for use and derivation of area-use and temporal-use factors follow.

Area-Use Factor

The area-use factor (AUF) estimates the fraction of a receptor's exposure that comes from the exposure location. The AUF is the smaller of 1 and the ratio of the area of the exposure location and the area in which a receptor lives or forages, whichever is more appropriate to the routes by which the receptor is exposed. The AUF is calculated as follows:

$$AUF_{ik} = 1, \text{ if } A_k \geq HR_i$$
$$= A_k/HR_i, \text{ if } A_k < HR_i$$

If the receptor has a home range large enough to comprise multiple exposure locations in the exposure grid, then the exposure location k can be an aggregate of smaller sites, in which case the AUF is calculated as above, using the area of the aggregate, which is the sum of the areas of the smaller units. Assuming that the receptor moves randomly over the exposure location, the area-weighted exposure concentrations of COPCs are used to calculate the average daily dose for that receptor that comes from the multiple nodes on the exposure grid.

Temporal-Use Factor

There are several approaches to dealing with the temporal aspect of exposure. The first approach is to assume conservatively, that receptors are exposed throughout their lifetime to COPCs and ROPCs present at the exposure location. The second approach is to estimate the temporal-use factor (TUF) as the fraction of time each year that a receptor is in the vicinity of the exposure location during which it forages or resides at the exposure location. The remaining time is assumed to be spent in either, one an area free of contamination, or two an area with 'background' levels of contaminants. The third approach is more complicated because it depends on receptor behavior and physiology, and the duration of the toxicity test on which the TRV is based. In general, the TUF is calculated as follows:

$$TUF_{ik} = (EF_{ik} \times ED_{ik})/AT_i$$

where

EF_{ik} = frequency of exposure of receptor i at exposure location k (days/year)
ED_{ik} = duration of exposure of receptor i at exposure location k (years)
AT_i = exposure averaging time for receptor i (days).

For the SLRA, the TUF will be assumed to be 1 for all receptors (unless explicitly stated otherwise). This is a reasonable assumption because the duration of toxicity tests is short compared to the duration of exposure of the receptor at the site.

8.2.4.2. Uptake Variables

The exposure equations for ecological receptors include parameters for the ingestion and inhalation of ecological receptors and the efficiency of absorption of COPCs and ROPCs from inhaled or ingested media.

Ingestion and Inhalation Rates

The magnitude of exposure of ecological receptors to COPCs and ROPCs in environmental media depends on the rate of intake of the contaminated media. For wildlife receptors exposed by ingestion and inhalation, receptor-specific ingestion and inhalation rates are required to estimate exposure. Published values for food ingestion (IP and IA), soil and sediment ingestion (IS), water ingestion (IW), and inhalation (IR) will be used to estimate exposure.

Absorption Efficiency

Substances ingested or inhaled by ecological receptors are absorbed and taken up into the receptors cells and organs to varying degrees. The efficiency of absorption depends on the relative affinity of the

substance for the environmental medium (soil, air, particulate, sediment, water, and tissue) and for the receptor's tissues. Published absorption values will be used for a COPC or ROPC when available. For both the PRA and the FRA, the absorption efficiency (AE) for inhaled and ingested media will be assumed to be the same or 100% of the actual absorption of the contaminant in the experiment or field observation used to derive the TRV. This is a conservative assumption for COPCs and ROPCs ingested as soil, sediment, or particulates in water, and inhaled as particulates.

8.2.4.3. Bioaccumulation Factors for Calculating Terrestrial Exposures

The calculation of exposure for ecological receptors may require one or more bioaccumulation and transfer factors. These factors are used to estimate the concentration in the tissue of an organism from the concentrations in the contaminated media to which it is exposed. Such factors are required to estimate exposure for wildlife receptors, such as mammals and birds, that are exposed to chemicals in soil by ingestion of plants, soil-dwelling invertebrates or other wildlife, when the concentration in the ingested organism is not measured directly. In each case, the numerator of the factor must have units corresponding to concentration in the medium taking up the substance (tissue) and the denominator the units of concentration in the 'source' medium (soil, tissue). The rules for use and derivation of bioaccumulation or transfer factors follow:

Soil-to-plant transfer factor

The soil-to-plant transfer factor (SP) is the ratio of the chemical concentration in plant tissue and in soil as follows: The SP is used to estimate the tissue concentration of plants exposed to chemicals in soil from the concentration of chemical in bulk soil. That is:

$$SP = [(mg/kg_{tissue}\text{-wet wt.})/(mg/kg_{soil}\text{-dry wt.})].$$

$$SPr_j = \text{Soil-to-Plant transfer factor for reproductive tissue for COPC } j = (kg_{soil}\text{-dry wt.}/kg_{tissue}\text{-wet wt.})$$

and

$$SPv_j = \text{Soil-to-Plant transfer factor for vegetative tissue for COPC } j = (kg_{soil}\text{-dry wt.}/kg_{tissue}\text{-wet wt.})$$

where reproductive tissues are flowers, seeds, and fruit, and vegetative tissues are leaf, stem, and root. Concentrations are estimated for plant tissues that are fed upon by wildlife receptors.

The first choice for terrestrial soil-to-plant (SP) transfer values will be EPA (1999) values. Per EPA (1999a), recommended SP values for inorganic elements are values published in Baes et al. (1984), and for those with no published field or laboratory data, the arithmetic average of the available SP values for the other inorganics will be used as the SP. EPA recommends using bioaccumulation equivalency factors (BEFs) to estimate the bioaccumulation of PCDD and PCDF congeners for which field or laboratory measurements are not available. The BEF is the predicted ratio of bioaccumulation of a PCDD or PCDF congener in soil to the bioaccumulation of TCDD (EPA 1999a). BEFs were used by EPA (1999a) to calculate the BAFs presented in Appendix C for PCDD and PCDF congeners. The EPA (1999a) recommended SP values for organic compounds with no field or laboratory data are estimated using the Travis and Arms (1988) regression on K_{ow}:

$$\log SP = 1.588 - (0.578 \log K_{ow})$$

For organic compounds, the same SP values are used for both SP_r and SP_v. Baes et al. (1984) and Travis and Arms (1988) values are reported in Appendix C-2, Table C-2-1. Where K_{ow} values are not available, SP_r and SP_v values for organic compounds will be values preferred and provided by the State of Washington. Where such values are not available, we will seek values from PNNL derived from Hanford site-specific data. Default SPs are 1 for COPCs/ROPCs without values from any of the above sources.

SP_r and SP_v values are also used for aboveground protected and unprotected plant parts for human health exposure (section 6.3.1).

Air-to-plant tissue transfer factor

The air-to-plant tissue transfer factor (APv) is the ratio of the chemical concentration in aboveground vegetative tissue of a plant and the concentration in air [(mg/kg)/(mg/L or m^3)]. The APv is used to estimate the tissue concentration of plants exposed to chemicals in air from the concentration of chemical in the air. That is,

$$AP\nu_j = \text{air-to-plant tissue transfer factor for COPC } j \text{ (L or m}^3\text{/kg)}$$

Tissue concentrations are estimated for plants that are fed upon by wildlife receptors, such as mule deer, because plant-tissue concentrations are needed to estimate the daily dose to receptors.

The first choice for air-to-plant tissue transfer factors (APv) will be EPA (1999a) values, where they are referred to as Bv. EPA (1999a) calculates the APv for organics (except PCDDs and PCDFs) using regressions on K_{ow} published by Bacci et al. (1990, 1992) and the calculated values are reduced by a factor of 100. For PCDDs and PCDFs, EPA (1999a) recommends the use of APv values from Lorber and Pinsky (1999). For metals except mercuric chloride, EPA (1999a) assumes that APv = 0. For mercuric chloride, EPA (1999a) recommended APv values are obtained from EPA (1997d).

Where K_{ow} values are not available, APv values for organic compounds will be values preferred and provided by the State of Washington. Where such values are not available, we will seek values from PNNL derived from site-specific field or laboratory data. Default APv values are 1 for organic COPCs without values from any of the above sources.

Soil-to-soil invertebrate tissue transfer factor

The soil-to-soil invertebrate tissue transfer factor (BAF-S) is the ratio of the chemical concentration in soil invertebrate tissue and that in soil [(mg/kg_{tissue}-wet wt.)/(mg/kg_{soil}-dry wt.)]. The BAF-S is used to estimate the tissue concentration of terrestrial invertebrates exposed to chemicals in soil by all exposure routes (ingestion, direct contact, and inhalation) from the concentration of a chemical in bulk soil. That is:

$$\text{BAF-S}_{mj} = \text{Soil-to-soil invertebrate tissue transfer factor for prey type } m \text{ for COPC } j \text{ (kg}_{soil}\text{-dry wt./kg}_{tissue}\text{-wet wt.)}$$

where prey type m is a soil-dwelling invertebrate, such as the earthworm, which is an important diet item of many mid-level predators, such as shrews and robins. Tissue concentrations are estimated for terrestrial invertebrates that are fed upon by wildlife receptors.

The first choice for terrestrial soil-to-earthworm bioaccumulation (BAF-S) values will be field or laboratory values for earthworms in EPA (1999a). Per EPA (1999a), recommended BAF-S values for inorganic elements (except lead) with no published field or laboratory data are arithmetic averages of the

BAF-S values available for other inorganics. For organic compounds with no field or laboratory data. EPA (1999a) uses values derived from K_{ow}s for daphnids, an aquatic macroinvertebrate, exposed to PAHs (Southworth et al 1978). For organic compounds with no field or laboratory data available, we will seek values from the State of Washington and Pacific Northwest National Laboratory (PNNL) derived from Hanford site-specific data. If values are not available from either of these sources, published values from published sources (e.g. HAZWRAP 1994) will be used. The Southworth et al (1978) regression for daphnids is not recommended for earthworms. Where no appropriate published surrogate data are available, the default BAF-S for organic compounds will be 10.

The earthworm BAF-S for lead (Pb) is calculated from the site-specific concentrations of calcium and lead (Corp and Morgan 1991), as follows:

$$\text{BAF-S}_{worm,Pb,k} = 0.2 \times C_{worm,Pb,k}/C_{soil,Pb,k}$$

where

$$\log C_{worm,Pb,k} = 1.16 + (0.916 \times \log C_{soilPb,k}) - (0.326 \times \log C_{soil,Ca,k})$$

and

$C_{worm,Pb,k}$ = the concentration of lead in worm tissue at exposure location k (mg/kg$_{tissue}$-dry wt.)
$C_{soil,Pb,k}$ = the concentration of lead in soil at exposure location k (mg/kg$_{soil}$-dry wt.)
$C_{soil,Ca,k}$ = the concentration of calcium in soil at exposure location k (mg/kg$_{soil}$-dry wt.)

The concentration of lead in the worm tissue (mg/kg$_{tissue}$-dry wt.) is converted to wet weight by multiplying by 0.2, on the assumption that worms are 80% water. No other substance is known to have an equation for predicting site-specific BAFs that depends on the concentration of a different substance.

BAF-S values are listed in Appendix C2. Note that the earthworm data (of which there are several values) serve as proxy for the darkling beetle and other desert terrestrial invertebrates (Figure 8-9) for which there are no known BAF-S values.

Tissue-to-tissue transfer factor

The tissue-to-tissue transfer factor (BAF-T) is the ratio of the chemical concentrations in predator tissue and that in the tissue of its prey [(mg/kg/(mg)/kg$_{tissue}$-wet wt.)]. The BAF-T is used to estimate the tissue concentration of animals exposed to chemicals by ingestion of prey from the concentration of chemical in the prey. That is:

$$\text{BAF-T}_{nj} = \text{Tissue-to-tissue transfer factor for prey type } n \text{ for COPC } j \text{ (kg}_{tissue}\text{-wet wt./kg)}$$

where, for example, prey type n is a mid-level predator of soil-dwelling invertebrates such as a Western meadowlark or Great Basin pocket mouse and also a prey of top predators, such as coyotes or owls and receptors such as deer and rabbits, which are considered wild game in the human health risk assessment. Tissue concentrations are estimated for animals that are fed upon by wildlife receptors.

For medium-to-tissue accumulation factors for mammals and birds, EPA (1999a) proposes the use of Baes et al. (1984) and Travis and Arms (1988) Ba values (d/kg) multiplied by the receptor's absolute ingestion rate for the medium (kg-medium/d). The EPA approach holds that the ratio of the receptor's tissue concentration (mg/kg body weight) to the receptor' contaminant intake rate (mg/d) is invariant. In other words, a deer (56.5 kg body weight, 1.74 kg/d ingestion rate) would have a seven-fold higher body

burden than a cottontail (1.2 kg body weight, 0.237 kg/d ingestion rate) eating the same vegetation because of the deer's seven-fold larger food ingestion rate.

It is proposed to calculate BAF-Ts from published biotransfer factors (Ba), as follows:

$$\text{BAF-T}_{nj} = \text{Ba}_A \times \text{BW}_A \times \text{SIR}_n$$

where

BAF-T_{nj} = animal-to-animal tissue transfer factor for prey species n and COPC j ($\text{kg}_{\text{tissue}}$-wet wt./kg)
Ba_A = biotransfer factor for animal A (d/kg)
BW_A = body weight of animal A (kg)
SIR_n = specific food ingestion rate of prey species n ($\text{kg}_{\text{tissue}}$-wet wt./kg/d)

This approach to estimating BAF-T assumes that the ratio of the amount of contaminant taken up per unit body weight and the amount of contaminant ingested per unit body weight per day (mg/kg / mg/kg/d) is constant among similar species, e.g., mammals.

The proposed approach can be derived from a simple model,

$$dC_{tissue}/dt = Z - \lambda C_{tissue},$$

where C_{tissue} (mg COPC/kg body weight) is the COPC-tissue concentration (amount of contaminant per unit tissue) at time t; Z is the amount of contaminant absorbed per unit tissue per unit time (mg COPC/ kgBW/day); and λ is the loss (depuration) rate constant (d^{-1}). A biouptake factor (BF) can be defined as the ratio of the equilibrium tissue concentration [$C_{tissue}(*)$] and the "input" of contaminant into the organism, that is,

$$BF = [C_{tissue}(*)]/Z.$$

BF has units mg COPC/kgBW per mg COPC/kgBW/d, or more simply days (d). BF is equal to the inverse of the loss rate,

$$BF = [C_{tissue}(*)]/Z = [Z/\lambda]\ /Z = 1/\lambda,$$

because $C_{tissue}(*)$ can be shown to be equal to Z/λ.

It is likely that the depuration rate constant varies in nature, but possibly not as much as food ingestion rates. Metabolic scaling theory suggests that rate constants or frequencies (time^{-1}) vary as the inverse of the 0.25 power of body weight while rates with units of mass or volume per unit time (e.g., kg/d) vary approximately as the 0.75 power of body weight (Schmidt-Nielsen 1984). If depuration rate constants are less variable than ingestion rates, then BF will be more invariant across species and individuals than EPA.

Given that the amount of contaminant absorbed per unit tissue per unit time (Z) can be estimated as

$$Z = C_{ingested} \times AE \times SIR,$$

then the relationship of BF to the BAF-T required in the RPP-WTP is as follows:

$$\text{BAF-T} = C_{tissue}(*)/C_{ingested}$$

$$= (Z/\lambda)/(C_{ingested})$$
$$= C_{ingested} \times AE \times SIR/(\lambda \times C_{ingested})$$
$$= (1/\lambda) \times AE \times SIR$$
$$= BF \times AE \times SIR,$$

where BAF has units (mgCOPC/kg-receptor tissue/mgCOPC/kg-medium ingested), AE is the absorption efficiency (mgCOPC-receptor tissue/mgCOPC-medium ingested), and the specific ingestion rate (SIR) has units kg-medium ingested/kg BW/d. Thus, BF isolates the storage component of the bioaccumulation process from the ingestion (SIR) and absorption (AE) components of the process.

The BAF will be relatively invariant with body weight if the BF, AE, and SIR vary as the 0.25, 0, and –0.25 powers of body weight, respectively. As stated above, the depuration constant likely varies as the inverse of the 0.25 power of body weight and BF is the inverse of the depuration constant, so it likely varies as the 0.25 power. To the extent that absorption is a tissue property, absorption efficiency should not vary with body weight (exponent = 0). If ingestion rate varies approximately as the 0.75 power of body weight (Schmidt-Nielsen 1984), then the body-weight specific ingestion rate will vary as approximately the –0.25 power of body weight ($W^{0.75}/W^{1} = W^{-0.25}$).

The proposed BAF values are calculated from Baes et al. (1984) and Travis and Arms (1988) Ba values because BF values are not available. Assuming that the ratio of the receptor-tissue concentration to specific ingestion rate of contaminant is invariant across species, that is,

$$BF_{cow} = BF_{receptor}$$

$$C_{cow}(*)/(C_{ingested} \times AE_{cow} \times SIR_{cow}) = C_{receptor}(*)/(C_{ingested} \times AE_{receptor} \times SIR_{receptor}),$$

the BAF for the receptor is calculated from the Ba value for the cow as described above.

Assuming that $AE_{receptor} = AE_{cow}$, it follows that,

$$BAF_{receptor} = C_{receptor}(*)/C_{ingested}$$
$$= [C_{receptor}(*)/(C_{ingested} \times AE_{receptor}/AE_{cow} \times SIR_{receptor})] \times SIR_{receptor}$$
$$= [C_{cow}(*)/(C_{ingested} \times SIR_{cow})] \times SIR_{receptor}$$
$$= [C_{cow}(*)/(C_{ingested} \times IR_{cow})] \times BW_{cow} \times SIR_{receptor}$$
$$= Ba_{cow} \times BW_{cow} \times SIR_{receptor},$$

because the biotransfer factor Ba (d/kg) is defined as follows (Baes et al. 1984):

$$Ba_{cow} = C_{cow}(*)/(C_{ingested} \times IR_{cow}),$$

where IR_{cow} is the ingestion rate (kg/d) of the cow.

When published field or laboratory values are not available, the first choices for Ba values are those recommended by EPA (1999a). For mammals, EPA (1999a) recommends using the following regression on K_{ow} (Travis and Arms 1988) for organic compounds (except dioxins/furans):

$$\log Ba = -7.6 + \log K_{ow}$$

For dioxins/furans, EPA (1999a) recommends for mammals the use of the Ba values presented in EPA (1995). For mammals and inorganic elements (except mercury), the EPA (1999a) recommended values are those reported in Baes et al. (1984). For mercury, EPA (1999a) recommends for mammals the use of the Ba value for total mercury reported for cows in EPA (1997d) converted to a wet weight basis and partitioned into methylmercury and mercuric chloride forms in the ratio of 0.13:0.87. For all mammal receptors, the Ba values are multiplied by the ratio of the receptor fat to the beef cattle fat.

For birds, EPA (1999a) recommends using for organic compounds (except dioxins/furans) the Travis and Arms (1988) Ba values for mammals adjusted for the lower fat content of birds. Bird Ba values are derived by multiplying the mammal value by the ratio of bird fat to mammal fat (0.15/0.19). For dioxins/furans, EPA (1999a) recommends for birds using the Ba values for chickens derived in Stephens et al. (1995). For birds and cadmium, selenium, and zinc, the EPA (1999a) recommended values are those reported in EPA (1992). For mercury, EPA (1999a) recommends for birds the use of the Ba value for total mercury for poultry reported in EPA (1997d) converted to a wet weight basis and partitioned into methylmercury and mercuric chloride forms in the ratio of 0.13:0.87.

The second choice for tissue-to-tissue bioaccumulation (BAF-T) values will be values preferred and provided by the State of Washington. Where such values are not available, we will seek values from PNNL derived from Hanford site-specific data. If values are not available from these sources, we will use values published in Baes et al. (1984) for elements and from published studies that have been compiled by HAZWRAP (1994). These values are found in Appendix C-2.

Default BAFs for mammals and birds are 1 for inorganic and 10 for organic COPCs.

Published BAF-Ts are listed in Appendix C2. Unless Hanford site-specific BAF-Ts become available, these and other BAF values will be used.

Abiotic medium-to-vertebrate tissue transfer factor

The abiotic medium-to-vertebrate animal tissue transfer factor (BAF-T) is the ratio of the chemical concentration in vertebrate animal tissue and the concentration in soil, drinking water or air (mg/kg/ mg/kg_{soil}-dry wt. or L or m^3). The BAF-T is used to estimate the tissue concentration of vertebrates exposed to chemicals by soil ingestion (direct or indirect), water ingestion, or inhalation from the concentration of chemical in the abiotic medium. That is,

$BAF\text{-}T_{nj}$ = abiotic medium-to-vertebrate animal tissue transfer factor for prey species n and COPC j (kg_{soil}-dry wt. or L or m^3/kg)

where, prey type n is a mid-level predator, such as the Western meadowlark or Great Basin pocket mouse, that feeds upon plants or soil-dwelling invertebrates. Tissue concentrations are estimated for mid-level predators that are fed upon by wildlife receptors, such as coyotes and burrowing owls, because prey-tissue concentrations are needed to estimate the daily dose to receptors.

The BAF-Ts for vertebrates ingesting soil, drinking water or inhaling air are the same as the BAF-Ts for tissue-to-tissue uptake. As with BAF-Ts for tissue-to-tissue uptake, BAF-Ts for uptake from abiotic media are calculated from published biotransfer factors (Ba), the beef cattle body weight (200 kg), and the prey's specific ingestion or inhalation rate for the abiotic medium. The Ba values used to calculate BAF-Ts for abiotic medium-to-tissue uptake are the same as the Ba values used to calculate BAF-Ts for tissue-to-tissue uptake. The default BAF-T values for abiotic media are the same as the default BAF-Ts for tissue.

Default bioaccumulation values

When a bioaccumulation factor for terrestrial receptors is not available, a default value will be used. Default values are conservative, order of magnitude estimates, based on an evaluation of available published values (HAZWRAP 1994). Default values are selected to be larger than most published values.

The default SP for terrestrial plants is 1 for all contaminants without published values. Default AP value for organic compounds is 1. Default BAF-S values are 1 for inorganic and 10 for organic contaminants without published values. Default BAF-Ts for terrestrial mammals and birds are 1 for inorganic and 10 for organic contaminants.

8.2.4.4. Bioaccumulation Factors for Calculating Aquatic Exposures

The calculation of exposure for ecological receptors may require one or more bioaccumulation or transfer factors that are used to estimate the concentration in the tissue of an organism from the concentrations in the contaminated media to which it is exposed (Figure 8-11). Such factors are required to estimate exposure for wildlife receptors, such as mammals and birds that are exposed to chemicals in sediment or surface water by ingestion of plants, sediment-dwelling invertebrates, or aquatic biota, when the concentration in the ingested organism is not measured directly. In each case, the numerator of the factor must have units corresponding to the units of the medium taking up the substance (tissue) and the denominator the units of the 'source' medium (sediment, water, tissue). The rules for use and derivation of these factors follow:

Water-to-plant transfer factor

The water-to-plant transfer factor (WP) is the ratio of the chemical concentration in aquatic plant tissue and that dissolved in water [(mg/kg$_{tissue}$-wet wt.)/(mg/L)]. The WP is used to estimate the tissue concentration of aquatic plants exposed to chemicals in sediment or surface water from the concentration of chemical dissolved in sediment pore water or surface water, respectively. That is:

WP_j = water-to-plant transfer factor for aquatic plant tissue for COPC j dissolved in water (L/kg$_{tissue}$-wet wt.).

Aquatic plants are assumed to be exposed only to the dissolved phase of contaminants in sediment pore water, or surface water, depending on the morphology of the particular type of aquatic plant (e.g., rooted, floating). Concentrations are estimated for aquatic plant tissues that are fed upon by terrestrial receptors (e.g., Canada goose).

If field-measured WPs at the Hanford Site are available, they are the first choice. Second choice are laboratory-measured WPs. The third choice is field or laboratory values in EPA (1999a). If no field or laboratory values are available, WPs will be calculated as the product of the SP and the Sediment Kd. We do not recommend that WP values for organic compounds be derived using the Southworth et al (1978) regression on K_{ow} for daphnids as described in EPA (1999a) because aquatic plants and aquatic microcrustacean invertebrate animals are expected to have different uptake and depuration. Default WPs are 1 for COPCs/ROPCs without more preferred values.

Sediment-to-plant transfer factor

The sediment-to-plant transfer factor (SP) is the ratio of the chemical concentration in aquatic plant tissue and that in sediment [(mg/kg$_{tissue}$-wet wt.)/(mg/kg$_{sediment}$-dry wt.)]. The SP is used to estimate the tissue concentration of aquatic plants exposed to chemicals in sediment. That is,

SP_j = Sediment-to-plant transfer factor for aquatic plant tissue for COPC *j* ($kg_{sediment}$-dry wt./kg_{tissue}-wet wt.)

Rooted aquatic plants are assumed to be exposed only to the dissolved phase of contaminants in sediment pore water. Therefore, if SP values are not available for a constituent, the SP can be estimated from the water-to-plant tissue transfer factor (WP) and the sediment Kd. That is,

$$SP_j = WP_j/Kd_j$$

where

WP_j = Water-to-plant transfer factor for aquatic plant tissue for COPC *j* (L/kg_{tissue}-wet wt.)

Kd_j = Soil-to-water partitioning coefficient for COPCj (L/$kg_{sediment}$-dry wt.)

This assumes equilibrium between sediment and sediment pore water.

The first choice for sediment-to-aquatic plant tissue (SP) values will be field or laboratory values preferred and provided by EPA (1999a). Where such values are not available, values will be obtained from the State of Washington or PNNL values derived from Hanford site-specific data. If values are not available from either of these sources, terrestrial soil-to-plant transfer values will be used for the vegetative parts of plants (SPv) published in Baes et al. (1984) for elements and values derived from regressions on K_{ow} (Travis and Arms 1998) for organics in accordance with EPA (1999a) (Appendix C.2). If SPv values are not available, calculate the SP as the ratio of the water-to-plant tissue value (WP) and soil-to-water partition coefficient (Kd), as described above. Default SP values are the same as the terrestrial SP_r, and SP_v values.

Surface water-to-aquatic invertebrate transfer factor

The surface water-to-aquatic invertebrate tissue transfer factor (BCF) is the ratio of the chemical concentrations in animal tissue and that in the surface water to which the animal is exposed [(mg/kg_{tissue}-wet wt.)/(mg/L)]. The BCF is used to estimate the tissue concentration of animals exposed to chemicals in surface water from the concentration of chemical in the surface water. The tissue concentration is estimated for animals that are fed upon by wildlife receptors. That is:

BCF_{invj} = Surface water-to-aquatic invertebrate tissue transfer factor for prey for COPC *j* (L/kg_{tissue}-wet wt.).

where *m* is an aquatic organism such as small fish and crustaceans, which are an important diet item of many terrestrial predators of aquatic biota, such as herons.

The first choice for BCFs for aquatic invertebrates are values preferred and provided by the State of Washington or EPA, including field or laboratory values reported in EPA (1999a). Where such values are not available, we will seek values from PNNL derived from site-specific field or laboratory data. If values are not available from either of these sources, we will use published values for surrogates derived from field or laboratory data and values for organics derived from regressions on K_{ow}.

If no more preferred values are available, BCFs for aquatic invertebrates are calculated per EPA (1999a) from the octanol water-partitioning coefficient (K_{ow}) using the regression equation of Southworth et al. (1978):

$$\log BCF_{inv,j} = 0.819 \times \log K_{ow,j} - 1.146$$

where

$BCF_{inv,j}$ = Surface water-to-tissue transfer factor for invertebrate aquatic biota for COPC j (L/kg_{tissue}-wet wt.).

$K_{ow,j}$ = Octanol-water partition coefficient of COPC j (Lyman et al.1982).

For BCFs derived from field data on tissue concentrations and total water column concentrations, modeled total COPC concentration in the water column (C_{wctot}) will be used to estimate aquatic invertebrate tissue concentrations. For BCFs derived from laboratory data or K_{ow} regressions, the modeled dissolved phase water concentration (C_{dw}) will be used to estimate the aquatic invertebrate tissue concentrations.

Published BCFs are listed in Appendix C2. The default BCFs are 500 for inorganics and 100,000 for VOCs, SVOCs, and pesticides/PCBs.

Water-to-fish tissue transfer factor

The water-to-fish tissue transfer factor (BCF) is the ratio of the chemical concentration in fish tissue and the concentration in water [(mg/kg_{tissue}-wet wt.)/(mg/L)]. The fish BCF is used to estimate the tissue concentration of fish from the concentration in the water to which the fish is exposed. That is,

$BCF_{fish,j}$ = Water-to-fish tissue transfer factor for COPC j (L/kg_{tissue}-wet wt.)

The fish tissue concentrations are estimated because fish are consumed by wildlife receptors, such as herons, bald eagles, and mink.

The first choice for BCFs for fish are values preferred and provided by the State of Washington or EPA, including field or laboratory values reported in EPA (1999a). Where such values are not available, we will seek values from PNNL derived from site-specific field or laboratory data. If values are not available from either of these sources, we will use published values for surrogates derived from field or laboratory data and values for organics derived from regressions on K_{ow}.

If preferred values are not available for organic compounds, BCFs for fish are calculated using the following regression on the K_{ow} (Bintein et al. 1993):

$$\log BCF_{fish} = 0.91 \times \log K_{ow} - 1.975 \times \log (6.8 \times 10^{-7} \times K_{ow} + 1.0) - 0.786$$

If preferred values are not available for inorganic COPCs and ROPCs, per EPA (1999a), the BCF will be estimated as the arithmetic average of available BCFs for other inorganics.

For BCFs derived from field data for fish-tissue concentrations, modeled total COPC concentrations in the water column (C_{wctot}) will be used to estimate fish-tissue concentrations. In this case, the BCF is equivalent to the BAF (EPA 1999a). For BCFs derived from laboratory data or K_{ow} regressions, the modeled dissolved phase water concentration (C_{dw}) will be multiplied by both BCFs and EPA (1995f) food-chain multipliers (FCMs) to estimate the fish-tissue concentrations (EPA 1999a). That is,

$$BAF = BCF_{fish} \times FCM,$$

where

BAF = Bioaccumulation factor (mg/kg_{tissue}-wet wt./ mg/L)
BCF_{fish} = Bioconcentration factor (mg/kg_{tissue}-wet wt./ mg/L)
FCM = trophic level food-chain multiplier (unitless)

EPA (1995) has calculated FCMs for log K_{ow} values ranging from 3.5 to 9.0.

Published BCFs for fish are listed in Appendix C-2, Table C-2-3. The default fish BCFs are 500 for inorganics and 100,000 for organic VOCs, SVOCs, and pesticides/PCBs.

Sediment-to-animal tissue transfer factor

The sediment-to-animal tissue transfer factor (BASF) is the ratio of the chemical concentrations in sediment-dwelling animal tissue and that in bulk sediment [(mg/kg_{tissue}-wet wt.)/(mg/$kg_{sediment}$-dry wt.)]. The BASF is used to estimate the tissue concentration of animals exposed to chemicals in sediment by all exposure routes (ingestion, direct contact) from the concentration of chemical in bulk sediment. The tissue concentration is estimated for animals that are fed upon by wildlife receptors. That is:

$BASF_{mj}$ = Sediment-to-animal tissue transfer factor for prey *m* for COPC *j* ($kg_{sediment}$-dry wt./kg_{tissue}-wet wt.)

where prey type *m* is typically a sediment-dwelling invertebrate, such as a burrowing amphipod or chironomid, which are important diet items of predators, such as the spotted sandpiper or certain fishes but may also be a sediment-ingesting fish such as carp and catfish.

BASFs are available in the literature for only a few chemicals. The first choice for BASFs are field or laboratory values preferred and provided by the EPA (1999a). However, we do not recommend that the BCF_{inv} derived using the Southworth et al (1978) regression for daphnids be used directly for the BASF as described in EPA (1999a). Where such values are not available, values from the State of Washington or PNNL will be obtained which are derived from Hanford site-specific data. If values are not available from either of these sources, published soil-to-soil invertebrate tissue transfer factors (BAF-S) will be used for inorganics and water-to-aquatic invertebrate tissue transfer factors (BCFs) will be derived for organics. For organics, the BASF will be estimated from the BCF as follows:

$$BASF_{mj} = BCF_{mj}/Kd_{jk}$$

where

BCF_{mj} = Surface water-to-aquatic invertebrate transfer factor for prey type *m* for COPC *j* (L/kg_{tissue}-wet wt.),

Kd_{jk} = Sediment-water partitioning coefficient for COPC *j* at exposure location *k* (L/$kg_{sediment}$-dry wt.),

Note that BASF in the literature (also referred to as BSAF) is often provided as a ratio of lipid-normalized tissue chemical residue to organic carbon-normalized sediment chemical concentration, so that a conversion may be needed to determine BASF.

For organic compounds, the Kd is a function of the K_{oc} and the organic matter content of the sediment, that is:

$$Kd_{jk} = K_{oc,j} \times FOC_k$$

where

$K_{oc,j}$ = the carbon-water partitioning coefficient for COPC j (L/kg$_{sediment}$-dry wt.)
FOC_k = mass fraction of organic carbon content of the sediment at exposure location k (unitless).

Because the K_{oc} is available for few compounds, the K_{ow}, the octanol-water partitioning coefficient, is used to estimate K_{oc} (EPA 1993c).

To estimate animal tissue concentrations, the BASFs derived from both field data and from aquatic invertebrate BCFs will be multiplied by the modeled bed-sediment COPC concentration (C_{bs}). Using bed-sediment concentrations with BASFs derived from BCFs is equivalent to using dissolved phase water concentrations and BCFs.

Published BASF values are listed in Appendix C2. The default BASF values are the same as the default BAF-S values (1 for inorganics, 10 for organics).

Default bioaccumulation values

When a transfer factor for aquatic receptors is not available because it was not measured in a field or laboratory investigation, it has no suitable surrogate value, or it could not be calculated, then a default value will be used. Default values are conservative, order of magnitude estimates, based on an evaluation of available published values (e.g., HAZWRAP 1994) and professional judgement. Default values are selected to be larger than most published values.

The default WP and SP values for aquatic plants is 1 and is the same as the default SPv for terrestrial plants. Default BCFs for aquatic animals are 500 for inorganics and 100,000 for VOCs, SVOCs, and pesticides/PCBs. The default BASF values are the same as the default BAF-S values (1 for inorganics, 10 for organics). Default BAF-Ts for mammals and birds ingesting aquatic biota are 1 for inorganic and 10 for organic COPCs.

8.3. Effects Assessment Calculations

TRVs are concentrations or doses of constituents that are associated with a specified level of adverse effect on ecological receptors. TRVs are used as the denominator in hazard quotients, as shown in the HQ equations (section 8.4).

8.3.1. Toxicity Reference Values for Terrestrial Receptors

TRVs for receptors dwelling in and thus, exposed by direct contact to COPCs/ROPCs in soil (plants, earthworms) are typically values from published sources, if field observations or site-specific toxicity tests of these media are not available.

8.3.1.1. Single Chemical TRVs

TRVs for plants and soil-dwelling invertebrates are derived values based on a review of published single-chemical laboratory studies (Efroymson et al. 1997a and 1997b).

For wildlife receptors, TRVs are required for both the ingestion and inhalation exposure pathways. Ingestion TRVs (Section 8.3.1.1) will be used to calculate HQs for the ingestion exposure pathway. Inhalation TRVs (Section 8.3.1.2) will be used to calculate HQs for the inhalation exposure pathway.

Inhalation and ingestion HQs for a given COPC can be added to calculate a HI for the combined pathways. This can be understood as adding the pathway-specific exposures, each weighted by its pharmacokinetic efficiency factor for the COPC, and comparing this combined "normalized" dose to the pathway-independent TRV (TRV'). For example, if the COPC is blood labile, there is a blood concentration associated with a particular level of adverse effect. This blood level corresponds to potentially different inhaled or ingested doses due to different pharmacokinetics of ingestion and inhalation. The factor that relates the dose to the corresponding blood concentration, with units kgBW/d/L, can be used to normalize the inhaled and ingested exposure doses to blood concentration so that they can be added together and compared on a common basis to the threshold blood concentration. That is,

$$\begin{aligned} HI &= ADD_{inh}/TRV_a + ADD_{ing}/TRV \\ &= ADD_{inh}/(TRV'/AE_{inh}) + ADD_{ing}/(TRV'/AE_{ing}) \\ &= (ADD_{inh} \times AE_{inh})/TRV' + (ADD_{ing} \times AE_{ing})/TRV' \\ &= (ADD_{inh} \times AE_{inh} + ADD_{ing} \times AE_{ing})/TRV' \end{aligned}$$

where

ADD_{inh} = the average daily dose by inhalation
ADD_{ing} = the average daily dose by ingestion
TRV_a = the inhalation TRV
TRV = the ingestion TRV
AE_{inh} = the inhalation absorption efficiency
AE_{ing} = the ingestion absorption efficiency

8.3.1.2. Ingestion TRVs

Ingestion TRVs for wildlife receptors are derived values based on published single-chemical laboratory toxicity tests (Sample et al. 1996). The outputs from these tests are subchronic or chronic NOAEL or LOAEL doses (mg/kg BW/day) for the test species. If the NOAEL or LOAEL is from a subchronic study, then the benchmark is adjusted downward by a factor of 10 to estimate the chronic benchmark. If the benchmark is a LOAEL for a mortality or reproduction endpoint, it is adjusted downward by a factor of 10 to estimate the NOAEL. To adjust a subchronic LOAEL for mortality or reproductive effect to a chronic NOAEL, divide by a factor of 100. As an alternative or addition to the above adjustments, an uncertainty factor can be applied to a NOAEL for a test species to adjust for extrapolating from a test species to a receptor species. The effect on HQs of adjusting benchmarks and alternative benchmarks will be discussed in the uncertainty section of the risk characterization.

If the desired TRV is that corresponding to the NOAEL, then the endpoint observed in the study should be mortality or reproduction. Non-lethal or non-reproductive NOAELs are conservative, that is, lower than necessary to protect the receptor, but will be used if a NOAEL for mortality or reproduction is not available. If the TRV is a LOAEL, then the endpoint observed in the study should be non-lethal or a non-reproductive effect. If the observed LOAEL endpoint is mortality or reproduction, then the non-conservative nature of the TRV should be considered in the risk characterization.

8.3.1.3. Inhalation TRVs

Appropriate inhalation toxicity data are those from chronic studies with ecologically significant endpoints. For example, mortality, impaired reproduction, organ dysfunction, and, in some cases, reduced growth are potentially ecologically significant endpoints, where as cancer is not. Portal effects on the respiratory tract (e.g., irritation) that do not lead to systemic effects (e.g., death) under chronic exposure are also not likely to be ecologically significant effects. The use of such data to derive TRVs is extremely conservative, which may be appropriate in certain situations, e.g., as a NOAEL in a screening ERA.

Chronic studies are usually more appropriate than acute data. In the absence of chronic data and assuming the endpoint is appropriate for the ERA, an acute benchmark is the conservative choice for the TRV, if it is less than the oral benchmark. In some cases, it may be appropriate to further reduce the acute benchmark by an acute-to-chronic uncertainty factor. No cases are known of any common contaminant causing ecologically significant effects from acute inhalation exposure at a dose lower than the dose causing significant effects from chronic oral exposure.

There are inhalation toxicity data from laboratory experiments reported in IRIS, RTECS and other databases. Air concentrations associated with varying levels of effect are reported for one or more laboratory animals, most if not all of which are mammals. These TRVs are reported as concentrations (mg/m^3) with a specified exposure duration and period. Assuming that the test duration and period is appropriately long for natural conditions, then test data can be used to derive inhalation TRVs for wildlife receptors of the same vertebrate class (mammal to mammal, bird to bird) given certain crucial assumptions. These assumptions are discussed below in Section 8.3.1.3.

When converting from published threshold-air concentrations to threshold doses, we assume that the values of the test species inhalation rate and body weight used to make the conversions are equal to those in the published study. A similar approach is used to interconvert threshold doses for ingestion and threshold dietary concentrations (Sample et al. 1996). See Appendix C2 for data.

8.3.1.4. Body-weight Scaling of TRVs

The starting point for deriving TRVs for receptors is thus the TRVs for one or more test species. TRVs for receptors should be derived from TRVs for test species from a similar taxonomic group. TRVs for birds should be derived from the TRV for a bird test species, and those for mammals from a mammal test species. The following equation is used to derive the TRV dose for wildlife receptors for COPCs in ingested matter (e.g., soil, sediment, water, biota) or inhaled air from published TRVs for test species:

$$TRV_{ij} = TRV_{tj} \times (BW_i/BW_t)^{0.25}$$

where

TRV_{ij} = Toxicity Reference Value of receptor i for COPC j (mg/kg BW/day)
TRV_{tj} = Toxicity Reference Value of test species t for COPC j (mg/kg BW/day)
BW_i = Body weight of receptor i (kg)
BW_t = Body weight of test species t (kg)

For ingestion TRVs, the body-weight ratio scaling exponent is 0.25 based on the weight of theoretical and empirical evidence (Travis et al. 1998, EPA 1992b) as presented in Sample et al (1996). Although there is some evidence that the body-weight ratio scaling exponent for acute toxicity in birds is 0 (Sample et al. 1996, Sample 1999), acute toxicity data (i.e., LC_{50}) may not be appropriate to predict body-weight scaling of chronic toxicity (Sample 1999). Acute toxicity likely operates by different mechanisms

than does chronic toxicity. Chronic toxicity depends on the balance of absorption and depuration, which probably scale to body weight. Therefore, an exponent of 0.25 will be used for both mammal and bird receptors for ingestion and inhalation TRVs.

Applying this body-weight scaling to inhalation TRVs for wildlife receptors involves two assumptions. First, it assumes that the ¾ power metabolic scaling law invoked by Sample et al. (1996) for toxicity of ingested contaminant applies equally well to inhaled contaminant. Second, it assumes that the absorption of contaminant per unit air inhaled is a constant fraction of body weight. This means that the amount of contaminant absorbed into the blood per unit body weight given a unit of inhaled contaminant is constant across animals of a given type.

The first assumption (inhalation toxicity scaling as ingestion toxicity) is reasonable because the Sample et al. (1996) approach is based on the observation that toxicity scales to the ¾ power of body weight (Chappell 1992, USEPA 1992, Travis and White 1988). Sample et al. (1996) attribute the scaling of toxicity to body weight to size-dependent variation in detoxification (depuration rates). The depuration rate for a blood-labile substance should be the same irrespective of the place of absorption, because depuration occurs primarily in the kidney, not at the site of absorption. Allometric scaling works best for substances processed by the kidney (Chappell 1992). Therefore, the scaling of toxicity with body weight should apply equally to ingested and inhaled substances.

The second assumption (constant specific lung-absorption efficiency) is reasonable for animals given the observed "constancy" with body weight of several related lung parameters: tidal volume, vital capacity, diffusion capacity, and especially, oxygen removal fraction (Schmidt-Nielsen 1989). The second assumption is analogous to the assumption of constant specific gut-absorption efficiencies (amount absorbed per unit ingested per unit body weight) that is required to extrapolate ingestion TRVs for test species to wildlife receptors (Sample et al. 1996), and which is also based on metabolic scaling arguments. In summary, we are assuming that inhalation toxicity scales to the ¾ power of body weight, for the same reason as does ingestion toxicity.

Finally, to apply the body-weight scaled inhalation thresholds derived from laboratory data for test species, one must assume that lab and the field conditions (e.g., temperature) are insignificantly different with respect to toxicity and metabolic function. This assumption is also made for toxicity via ingestion. Therefore, the application of the Sample et al. (1996) approach to deriving ingestion-based TRVs for wildlife receptors to the problem of deriving inhalation TRVs is defensible, and laboratory inhalation-toxicity data for test species can be used to derive inhalation TRVs for wildlife receptors.

If there is no published benchmark for a COPC, the lowest benchmark for a compound having like properties can be used as a surrogate. Also, when TEFs are available for a class of COPCs, such as dioxins and furans, then TEFs can be used to derive TRVs (see section 7.2.3 and following sections).

Published TRVs for chemical COPCs for plants, soil-dwelling invertebrates, and wildlife receptors (mammals and birds) are listed in Appendix C-3.

8.3.1.5. Toxicity Equivalence Factors for Dioxins, Dibenzofurans, and PCBs

Chlorinated dioxins, chlorinated dibenzofurans, and chlorinated biphenyls are thought to act through a common mechanism of toxicity by binding to a protein known as the arylhydrocarbon receptor (AR) (for review see ATSDR 1997 or WHO 1998). The AR-ligand complex is responsible for the activation of genes that have a deleterious effect when they are not under proper regulation by the receptor's hormones. Interaction of dioxins and similar compounds with AR therefore can cause immunological, neurological, endocrine, embryotoxic, and many other effects.

The similarity in action of these compounds is thought to result from their structural similarity. Dioxin is composed of two benzene rings joined by two carbon-oxygen-carbon bonds on two adjacent carbons of each benzene ring. Dibenzofurans have two benzene rings joined by a carbon-oxygen-carbon bond and a carbon-carbon bond on two adjacent carbons of each benzene ring. Biphenyls consist of two benzene rings joined by a single carbon-carbon bond. To form the polychlorinated derivatives, chloro groups are attached at various locations, as designated in the names of the compounds. Benzene rings are planar, i.e., flat, in conformation. Because two adjacent carbons on each benzene ring are joined in dioxins and dibenzofurans, both benzene rings are held in the same plane, and the chloro groups are also in that plane. Therefore, these molecules are said to be coplanar. The coplanar structure appears to be essential for interaction with AR. The benzene rings in biphenyl can rotate relative to each other, unless there are added groups that interfere with rotation (such as 2,2',6,6'-chloro groups, which occupy the carbons immediately on both sides of the carbon-carbon bond joining the rings). PCB congeners that are able to form a coplanar molecule (and are called coplanar PCBs) can interact with AR when they are in that configuration. Therefore, coplanar PCBs are included among the COPCs with similar action to dioxins and dibenzofurans.

EPA has recommended that toxicity equivalence factors (TEFs) be used to evaluate the cumulative toxicity of chlorinated dioxins, chlorinated dibenzofurans, and chlorinated biphenyls. Because these contaminants have a common mechanism of action, it is assumed that their toxicity to biota is additive (WHO 1998, EPA 1999b). That is, the toxicity of all dioxins, dibenzofurans, and PCBs should be added. Furthermore, their relative potency as chronic toxins is assumed to be related to the degree of affinity for AR, which can be measured much more conveniently than chronic toxic effects. TEFs have been proposed for several chlorinated dioxins, chlorinated dibenzofurans, and chlorinated biphenyls (WHO 1998, EPA 1999b), always using the toxicity of 3,4,7,8-tetrachlorodioxin (TCDD), the most potent chlorinated dioxin, as a TEF of 1.0. Separate lists were developed for mammals, birds, and fish, and these lists are presented in Table 8-5.

TEFs are reported in Table 8-5 for individual PCB congeners (e.g., 2,3,3',4,4',5-hexachlorobiphenyl), but analytical values for individual congeners in the exposure media are sometimes not available. It is also possible to calculate TEFs for Aroclors, which are mixtures of PCB congeners, using the typical composition of Aroclor mixtures (Hutzinger, Safe, and Zitko 1983).

Using TEFs, HQs can be calculated for chlorinated dioxins, chlorinated dibenzofurans, and PCBs for which TRVs are not available. The concentration of the COPC is multiplied by its TEF to calculate an equivalent concentration of TCDD. The TRV for TCDD is then used to calculate the HQ for the COPC. Because the mechanism of action of these compounds is thought to be the same, the TCDD-equivalent concentrations are added to determine the composite equivalent concentration of TCDD. This value is the same as the hazard index for the individual COPCs.

8.3.1.6. Radionuclide Benchmarks

The benchmark values for radiation given by IAEA (1992) are 1 mGy/d (0.1 rad/d) for terrestrial mammals and birds and 10 mGy/d (1 rad/d) for plants, invertebrates, and aquatic biota. These benchmarks are confirmed in *Effects of Ionizing Radiation on Terrestrial Plants and Animals: A Workshop Report* (Barnthouse 1995) and DOE Order 5400.5 (1990). Alpha radiation has a much higher effect on biological tissue than beta and gamma radiation because of the large mass of the alpha particle. When internal exposure is being evaluated, it is particularly important to consider the relative effectiveness of the radiation (P.M. Achey, U. of Fla, personal communication to C. Hadden, 9/20/99). *Rad* and *gray* are units of absorbed radiation (alpha, beta, and gamma) that do not adjust for the biological effectiveness of the radiation. *Rem* and *sievert* are units of absorbed radiation that include a quality factor

(QF rem/rad or sievert/gray) for the greater damage done by alpha particles than by beta and gamma radiation, and QF was included in the dose calculations used by IAEA (1992) to evaluate the benchmark. Inclusion of QF results in the calculation of effective dose in rem/day rather than absorbed dose in rad/day. Therefore, for this evaluation, radiological benchmarks for biota have been restated in units of rem/day rather than rad/day.

8.3.2. Toxicity Reference Values for Aquatic Receptors

TRVs for receptors dwelling in and thus exposed by direct contact to COPCs in sediment (sediment-dwelling biota) or surface water (fish, aquatic biota) are typically values from published sources if field observations or site-specific toxicity tests of these media are not available. The units of these values vary by source and medium, e.g., μg/L for surface water and mg/kg-dry wt. for sediment. TRVs for sediment-dwelling biota are derived values from published studies and reports, including field measured concentration and effect co-occurrence data and laboratory bioassays (Jones et al. 1997). TRVs for aquatic biota are values such as chronic National Ambient Water Quality Criteria (NAWQC) and Tier II values and lowest chronic values derived from published studies (Suter and Tsao 1996). TRVs for wildlife receptors exposed by ingesting biota exposed to surface water or sediments are derived as for terrestrial wildlife receptors (section 8.3.1).

Published TRVs for COPCs for sediment-dwelling invertebrates, aquatic biota, and wildlife receptors (mammals and birds) are listed in Appendix C-3.

For all sediment and aquatic biota, the TRV for total-body radiological dose from combined internal and external exposure for all ROPCs combined is 1.0 rem/day (IAEA 1992). However, the TRV for aquatic wildlife receptors (that is, birds and mammals) is 0.1 rem/day.

8.4. Risk Equations

Risk estimates for a receptor at an exposure location is calculated as the HQ, which is the ratio of the estimated exposure and the TRV. That is:

$$HQ = \text{Estimated Exposure/TRV}.$$

The HQ equation takes different forms depending on how the receptor is exposed, which also determines how the TRV is expressed (section 8.2.2). In the ecological risk assessment for the RPP-WTP, the exposure to ecological receptors is either a media concentration (C_{jk}), an average daily dose of a COPC (ADD_{ijk}), or a daily total-body radiological dose (R_{ik}).

The HQ is an index of the total risk to the receptor from exposure to the COPC if the COPC does not occur in the 'background' environment or if the home range of the receptor is smaller than the area of the exposure location, that is, if the AUF =1.

The HI is an index of the combined risk from exposure to multiple COPCs that have similar modes of action. A preliminary classification of inorganic COPCs groups arsenic, antimony, selenium, and vanadium as respiratory inhibitors; lead, manganese, and mercury as central nervous system inhibitors; and aluminum, chromium, and nickel as deoxyribonucleic acid (DNA) and protein reactors. Organic COPCs are typically grouped by chemical structure: VOCs, PAHs, organochloride pesticides, and PCBs. These chemical groupings are based on professional judgement and experience. For the PRA all HQs, regardless of mode of actions, will be grouped and summed because such summing represents the most conservative case. When the HI exceeds 0.25, additional HIs by mode of action will be developed. Thus,

the HI for a receptor at an exposure location is calculated from the HQs for the individual COPCs as follows:

$$HI_{ik} = \Sigma_j HQ_{ijk}$$

where

HI_{ik} = Hazard Index for receptor i at exposure location k (unitless)
HQ_{ijk} = Hazard Quotient for receptor i at exposure location k for COPC j (unitless).

Calculating HI assumes an additive effect on receptors from COPCs of all classes.

The HQ equation for receptors exposed to ROPCs is equivalent to an HI because the dose from all radionuclides is summed to estimate the total-body dose from internal and external exposures.

The threshold value for HQs and HIs for COPCs will be 0.25, unless a similar mode of action can be demonstrated and approved by Ecology. HIs for COPCs and ROPCs will not be added together.

8.5. Reporting of Major Ecological Risk Findings

Risk characterization will be reported in such a way as to capture all the various elements of the work. The following outline of headings is proposed:

- Current Risk at Exposure Location 1,2,3…n
  - Terrestrial Conditions: Central Plateau
    - Plausible Scenario
      - Chemicals
      - Radiologicals
    - Worst-case Scenario
      - Chemicals
      - Radiologicals
  - Aquatic Conditions: Columbia River
    - Plausible Scenario
      - Chemicals
      - Radiologicals
    - Worst-case Scenario
      - Chemicals
      - Radiologicals

- Future Risk at Exposure Locations 1,2,3…n
  - Terrestrial Conditions
    - Plausible Scenario
      - Chemicals
      - Radiologicals
    - Worst-case Scenario
      - Chemicals
      - Radiologicals
  - Aquatic Conditions
    - Plausible Scenario
      - Chemicals

Radiologicals
- Worst-case Scenario
  Chemicals
  Radiologicals

This will be done for the PRA as well as the FRA within the SLRA.

8.6. Uncertainties in Ecological Risk Assessment

Evaluation of uncertainties is part of the ERA process (EPA 1998e). Uncertainties in each of the four inter-related steps of the EPA approach to the screening the ERA will be discussed as follows:

- problem formulation
- exposure assessment
- effects assessment
- and risk characterization

Uncertainties about the data will be evaluated in the exposure assessment and the effects assessment steps.

8.6.1. Problem Formulation

Environmental concentrations of contaminants deposited on the soil and water at exposure locations will be based on many predictions. A degree of uncertainty exists about the predicted spatial distribution of contaminants. Exposure concentrations could be overestimated or underestimated, depending on how good the model is in predicting contaminant distribution.

Because the conservative exposure parameters (section 8.6.2) will be used to calculate HQs, the estimates of risk from ecological COPCs/ROPCs are conservative (that is, protective). Using conservative exposure concentrations decreases the likelihood of underestimating the risk posed by each ecological COPC/ROPC and increases the likelihood of overestimating the risk. Note that for wildlife receptors not living in soil, sediment, or surface water, HQ is a function of chemical dose (ADD) or radiological dose (R), which in turn, depends on a number of exposure factors (in addition to contaminant exposure concentration). Thus, several factors determine how conservative a HQ might be (in addition to contaminant exposure concentration).

The distribution and abundance of organisms comprising the ecological receptors at exposure locations have not been quantified by field studies. The lack of quantitative data introduces uncertainties concerning whether, and to what extent, the risk characterization based on the selected receptor species underestimates, or overestimates, the risk to organisms that are not used in the risk computations but are found at exposure locations. Additional reconnaissance on the Hanford Site at and near the exposure locations of maximum deposition will establish the nature and quality of habitat and confirm the presence of vegetation types and active, visible animal species. This is especially needed regarding bodies of water. Observations made during this reconnaissance will validate assumptions about the presence of unobserved organisms that are essential to normal ecosystem functioning, such as soil-dwelling worms and arthropods and herbivorous insects.

One (or more) unobserved species at exposure locations possibly is more sensitive than those ecological receptors for which toxicity data were available for use in setting TRVs. It does not necessarily follow that

these unevaluated species are at significantly greater risk of harmful ecological effects than that estimated in the SLRA. The reconnaissance will focus on places of predicted maximum ground concentrations.

8.6.2. Exposure Assessment

Movement of contaminants from the exposure locations through direct and indirect pathways to ecological receptors will not be measured for this SLRA. This introduces uncertainties about the actual modes and pathways of exposure and the actual exposure concentrations of these contaminants to the ecological receptors. Exposure concentrations can differ from the predicted environmental concentrations as a result of physical and chemical processes during transport from source to receptor and as a result of biomagnification through the food web. These processes will not be predicted quantitatively in this SLRA. Although bioaccumulation will be estimated for those receptors ingesting food for which toxicity thresholds were available, exposure of top predators possibly is underestimated because the biomagnification of certain contaminants in their prey was not considered.

The modes and pathways used to characterize the exposure of ecological receptors are the most important ones for the relatively large and active species in terrestrial habitats. Soil-dwelling terrestrial animals may be exposed to contaminants in soil by way of inhalation. However, it is expected that concentrations of VOCs will be very small and that gaseous concentrations in soil interstices, cavities, and burrows will not be available. Therefore, the exposure to burrowing organisms at the site from contaminated soil, and interstitial water in the soil, may be underestimated if gas concentrations are larger than soil concentrations. Overestimating exposure by using conservative exposure concentrations is thought to oppose the underestimation of exposure that results from neglecting certain exposure modes and pathways of lesser importance. Additional uncertainties are inherent in ingestion rates and dietary fractions of plants and animals. Likewise, uncertainties with dermal exposure will be considered. Exposure concentrations are likely overestimated because of conservative exposure factors. Sources of conservatism in the exposure factors include using published bioaccumulation factors, irrespective of species and environmental conditions.

8.6.3. Effects Assessment

Toxicity thresholds are based on concentrations reported to have no or little effect on the test organism or estimated conservatively from published toxicity data. TRVs for wildlife receptors exposed to soils derived from NOAELs or LOAELs reduced by safety factors of 10 for chronic LOAEFs and subchronic NOAELs or 100 for subchronic LOAELs (Sample et al. 1996). These thresholds would underestimate the risks only to organisms at the exposure locations that are considerably more sensitive than the receptor organisms for the specific toxicological endpoint. The thresholds are more likely to overestimate the risk to organisms that are equally or less sensitive than the receptor organisms. The possibility remains that some thresholds are set at levels at or above which some harm would occur to organisms at the exposure locations because receptors may be more sensitive to other toxicological endpoints.

There are limited data for developing inhalation TRVs and very few for developing dermal TRVs. Little is known about the actual absorption across the dermal layer of wildlife receptors. There is also uncertainty about the extrapolation of TRVs for ingestion to inhalation. Ingestion TRVs will not be extrapolated to dermal contact. The uncertainties associated with neglecting dermal contact and estimating inhalation toxicity will be discussed.

The calculated risks to the ecological receptors at the exposure locations are the risks of individual contaminants. The risks from exposure to multiple contaminants depend on contaminant interactions; effects could be greater or less than those from a single chemical. This screening ERA provides findings for ecological COPC-specific risk estimates and assumes additivity when calculating HIs. An evaluation of risk

from chemical mixtures cannot be conducted without additional data and evaluation of alternative models of contaminant interactions.

TRVs are not available for some COPCs/ROPCs. This lack of TRVs is especially true for organic COPCs Compounds. This situation likely results in underestimated risks.

Additional uncertainty exists as to the pertinence of individual organism toxicity for characterizing the risk to individuals, populations, and ecosystems. Populations possibly may compensate for the loss of large numbers of juveniles or adults with increased survival or birth rates, and habitats or ecosystems may possess functionally redundant species that are less sensitive to contaminants. Although the desert habitat at the exposure locations likely possesses some buffering mechanisms, a conservative approach is still justified to risk assessment based on organismal toxicity thresholds (i.e., NOAELs).

8.6.4. Risk Characterization

The uncertainties described above ultimately produce uncertainty in the quantification of current and future risks to plants and animals at the exposure locations. Three additional areas of uncertainty in the risk characterization exist: off-site risk, background risk, and cumulative risk. Each is briefly described below.

8.6.4.1. Risk Outside the Modeled Study Area

It is unlikely that the receptors outside the 50 km study area would have lower toxicity thresholds for contaminants than the thresholds used for receptors within the study area. In addition, there is little reason to expect that contaminants migrating outside the study area would be concentrated above predicted concentrations at the exposure locations unless a contaminant bioconcentrates in organisms that move extensively on and off the study area. In general, the risk to receptors outside the study area is likely to be overestimated rather than underestimated by the risk estimate for receptors within the 50 km radius of the site. (Spromberg et al. 1998).

8.6.4.2. Background Risk

Another source of uncertainty is ecological risk relative to background or predepositional conditions. Only incremental risk due to facility emissions will be evaluated because the concentration terms are modeled concentrations resulting from RPP-WTP operations. The HI< 0.25 threshold specified by EPA (1994a, 1994b, and 1994c), as cited in ATG (1998), are designed to "buffer" the potential for background contributions to risk (EPA 1998a). The amount of loading of various chemicals will be discussed. Background concentrations for the Hanford Site and the incremental loading rates will help to determine the portion of total risk coming just from stack emissions.

8.6.4.3. Cumulative Risk

The SLRA will estimate the risk to populations of ecological receptors from individual contaminants. In nature, receptors are exposed simultaneously to mixtures of chemicals. Generally, the methods used are sufficiently conservative, resulting in individual risks that are overestimated.

A second type of cumulative risk is possible when several living plants and animals are affected simultaneously. Harmful effects in ecosystems (including effects on individual organisms) may cascade throughout the system and have indirect effects on the ability of a population to persist in the area even though individual organisms are not sensitive to the given contaminants in isolation.

In addition, cumulative risk more broadly includes risk from multiple sources at the Hanford Site. Therefore, the ecological risk characterization for exposure locations may underestimate actual risks to plants and animals from cumulative risks.

8.6.5. Summary of Uncertainties

The most important uncertainties in the ecological portion of the SLRA for exposure locations are those surrounding the estimates of the contaminant concentrations to which ecological receptors are actually exposed (exposure point concentrations) and the concentrations that present an acceptable level of risk or harmful effects (toxicity thresholds or reference values). These uncertainties arise from multiple sources, especially from the lack of site-specific data on contaminant transport and transformation processes, organismal toxicity, animal behavior and diet, population dynamics, and the response of arid land plant and animal populations to stressors in the Hanford Site environments. Despite these uncertainties, the modeled exposure concentrations and published exposure and effects information will allow risks to be characterized for various exposure locations according to exposure/effects scenarios.

8.7. Summary for Screening Ecological Risk Assessment

Risks to ecological receptors from the potential emission of COPCs and ROPCs result from exposure to and ecological toxicity of the COPC/ROPC. The screening ecological risk assessment utilizes the estimated emission rates (section 5) and results of fate and transport modeling (section 6) to calculate potential ecological receptor exposure to COPCs and ROPCs. This exposure information is combined with toxicity data to estimate the potential for adverse effects to terrestrial and aquatic organisms and populations in the vicinity of the RPP-WTP.

The PRA will use conservative exposure assumptions to compensate for the high level of uncertainty associated with conducting a risk assessment for a facility that is still in the final design phases. The PRA will include a qualitative uncertainty analysis. Monte Carlo analysis or other quantitative methods may be used to identify and evaluate the primary sources of uncertainty in the FRA that may drive risk estimates. The exact procedures that may be used will be determined at a later time.

The FRA will focus on ecological COPCs that exceed risk goals in the PRA and may utilize additional site-specific emissions, fate and transport, and exposure data collected after the completion of the PRA.

Figure 8-1. Ecological Conceptual Exposure Model



Figure 8-2. Recreation and Wildlife Areas and the Hanford Reach



Hanford ERA - 016

Figure 8-3. Vegetation Types of the Hanford Site



BHI:rpp 04/22/96 clup/brmap d02.aml Database: 07-DEC-1998

Hanford ERA - 017

Best Available Copy

Figure 8-4. Selected Raptor Nesting and Perching Locations on the Hanford Site



Hanford ERA - 018

Figure 8-5. RPP-WTP Areas Vegetation Types (Simplified)



Hanford ERA - 019

Figure 8-6. Fall Chinook Salmon Spawning Areas Along the Columbia River



Figure 8-7. Trophic Levels and Receptor Species Considered in Development of the 200 Area and Vicinity Terrestrial Conceptual Exposure Model



Figure 8-8. Trophic Levels and Receptor Species at the Columbia River Considered in Development of the 200 Area and Vicinity Aquatic Conceptual Exposure Model



Figure 8-9. Trophic Levels and Selected Receptor Species (names in bold) from the 200 Area and Vicinity Terrestrial Conceptual Exposure Model



Figure 8-10. Trophic Levels and Selected Receptor Species (names in bold) at the Columbia River Considered in Development of the 200 Area and Vicinity Aquatic Conceptual Exposure Model



Figure 8-11. Relationship Between Sources and Biotransfer Factors for Calculating Terrestrial and Aquatic Exposures



1

Table 8-1. Policy Goals, Ecological Assessment Endpoints, Measures, and Decision Rules for 200 Areas and Surroundings

Policy Goals	Assessment Endpoint	Measures	Decision Rule
Policy Goal 1: The conservation of T&E species and their critical habitats.	**Assessment Endpoint 1:** Protection of individuals of state or federally designated threatened or endangered species. Endpoint Species: bald eagle and other species.	**Measure 1:** Modeled contaminant concentrations in prey (e.g., deer mouse, Western meadowlark, Great Basin pocket mouse, and house finch) based on modeled concentrations of vapors in air and particulates, depositions of contaminant particulates to soil, and measured concentrations of contaminants in abiotic media.	**Decision Rule for Assessment Endpoint 1:** If T&E species are not present, or RME concentrations in the media do not contribute to chronic NOAEL, possibly adjusted by a safety factor, exceedance (i.e., HQs <0.25), then it is indicated that the contaminant alone is unlikely to cause adverse ecological effects and, therefore, the T&E species should be preserved. If the HQ >0.25, lines of evidence will be evaluated to determine the potential for ecological risk and the need for any additional measurements or calculations.
Policy Goal 2: The protection of terrestrial populations and ecosystems.	**Assessment Endpoint 2:** Stable plant community for erosion control and energy production. Endpoint Species: cheatgrass, rabbitbrush.	**Measure 2:** Modeled concentrations of vapors in air and particulates and depositions of contaminant particulates to soil.	**Decision Rule for Assessment Endpoint 2:** If the HQ is <0.25, then it is indicated that the contaminant alone is unlikely to cause adverse ecological effects and, therefore, maintain the plant populations and communities. If the HQ >0.25, lines of evidence will be evaluated to determine the potential for ecological risk and the need for any additional measurements or calculations.
	Assessment Endpoint 3: Stable soil-dwelling invertebrate community for nutrient and energy processing. Endpoint Species: earthworms and darkling beetles.	**Measure 3:** Modeled concentrations of vapors in air and particulates and depositions of contaminant particulates to soil.	**Decision Rule for Assessment Endpoint 3:** If the HQ is <0.25, then it is indicated that the contaminant alone is unlikely to cause adverse ecological effects and, therefore, maintain the soil invertebrate community. If the HQ >0.25, lines of evidence will be evaluated to determine the potential for ecological risk and the need for any additional measurements or calculations.

Table 8-1. Policy Goals, Ecological Assessment Endpoints, Measures, and Decision Rules for 200 Areas and Surroundings

Policy Goals	Assessment Endpoint	Measures	Decision Rule
	Assessment Endpoint 4: Stable populations of herbivorous animals. Endpoint Species: mammals - mule deer.	**Measure 4:** Modeled contaminant concentrations in food chain, e.g., plants, based on modeled concentrations of vapors in air and particulates and depositions of contaminant particulates to soil.	**Decision Rule for Assessment Endpoint 4:** If the HQ is <0.25, then it is indicated that the contaminant alone is unlikely to cause adverse ecological effects and, therefore, populations of the herbivores, e.g., Great basin pocket mice, mule deer, and house finch, are maintained. If the HQ >0.25, lines of evidence will be evaluated to determine the potential for ecological risk and the need for any additional measurements or calculations.
	Assessment Endpoint 5: Stable populations of plant-eating and/or insectivorous animals (omnivores). Endpoint Species: bird-Western meadowlark.	**Measure 5:** Modeled contaminant concentrations in earthworms, plants, and other prey based on modeled concentrations of vapors in air and particulates and depositions of contaminant particulates to soil.	**Decision Rule for Assessment Endpoint 5:** If the HQ is <0.25, then it is indicated that the contaminant alone is unlikely to cause adverse ecological effects and, therefore, populations of omnivores, e.g., Western meadowlark, are maintained. If the HQ >0.25, lines of evidence will be evaluated to determine the potential for ecological risk and the need for any additional measurements or calculations.
	Assessment Endpoint 6: Stable populations of terrestrial predators. Endpoint Species: mammal-coyote; bird-burrowing owl and red-tailed hawk.	**Measure 6:** Modeled contaminant concentrations in prey (e.g., Western meadowlark and Great Basin pocket mice) based on modeled concentrations of vapors in air and particulates and depositions of contaminant particulates to soil.	**Decision Rule for Assessment Endpoint 6:** If the HQ is <0.25, then it is indicated that the contaminant alone is unlikely to cause adverse ecological effects and, therefore, populations of terrestrial predators are maintained. If the HQ >0.25, lines of evidence will be evaluated to determine the potential for ecological risk and the need for any additional measurements or calculations.

Table 8-1. Policy Goals, Ecological Assessment Endpoints, Measures, and Decision Rules for 200 Areas and Surroundings

Policy Goals	Assessment Endpoint	Measures	Decision Rule
Policy Goal 3: The protection of aquatic populations and ecosystems.	**Assessment Endpoint 7:** Stable populations of sediment-dwelling organisms. Endpoint Species: Clams, insects, snails, and worms.	**Measure 7:** Modeled sediment contaminant concentrations from dispersion and deposition.	**Decision Rule for Assessment Endpoint 7:** If the HQ is <0.25, then it is indicated that the contaminant alone is unlikely to cause adverse ecological effects and, therefore, populations of sediment-dwelling organisms are maintained. If the HQ >0.25, lines of evidence will be evaluated to determine the potential for ecological risk and the need for any additional measurements or calculations.
	Assessment Endpoint 8: Stable planktivorous fish and small invertebrate populations. Endpoint Species: water fleas and other invertebrates.	**Measure 8:** Modeled surface water contaminant concentrations.	**Decision Rule for Assessment Endpoint 8:** If the HQ is <0.25, then it is indicated that the contaminant alone is unlikely to cause adverse ecological effects and, therefore, populations of small invertebrates are maintained. If the HQ >0.25, lines of evidence will be evaluated to determine the potential for ecological risk and the need for any additional measurements or calculations.
	Assessment Endpoint 9: Stable waterfowl and shorebird populations. Endpoint Species: Canada goose, spotted sandpiper	**Measure 9:** Modeled contaminant concentrations in benthic invertebrates or aquatic plants based on modeled contaminant concentrations in surface water or sediments from dispersion and deposition.	**Decision Rule for Assessment Endpoint 9:** If the HQ is <0.25, then it is indicated that the contaminant alone is unlikely to cause adverse ecological effects and, therefore, populations of waterfowl and shorebirds are maintained. If the HQ >0.25, lines of evidence will be evaluated to determine the potential for ecological risk and the need for any additional measurements or calculations.

Table 8-1. Policy Goals, Ecological Assessment Endpoints, Measures, and Decision Rules for 200 Areas and Surroundings

Policy Goals	Assessment Endpoint	Measures	Decision Rule
	Assessment Endpoint 10: Stable populations of large carnivorous fish population for regulation. Endpoint Species: salmon, bass, channel catfish	**Measure 10:** Modeled surface water and sediment contaminant concentrations.	**Decision Rule for Assessment Endpoint 10:** If the HQ is <0.25, then it is indicated that the contaminant alone is unlikely to cause adverse ecological effects and, therefore, populations of large carnivorous fish are maintained. If the HQ >0.25, lines of evidence will be evaluated to determine the potential for ecological risk and the need for any additional measurements or calculations.
	Assessment Endpoint 11: Stable fish-eating terrestrial predator populations for population regulation. Endpoint Species: great blue heron, bald eagle, mink	**Measure 11:** Modeled contaminant concentrations in large carnivorous fish and planktivorous fish and small invertebrates based on modeled surface water and sediment concentrations.	**Decision Rule for Assessment Endpoint 11:** If the HQ is <0.25, then it is indicated that the contaminant alone is unlikely to cause adverse ecological effects and, therefore, populations of fish-eating terrestrial predators are maintained. If the HQ >0.25, lines of evidence will be evaluated to determine the potential for ecological risk and the need for any additional measurements or calculations.

RME = Reasonable maximum exposure
T&E = Threatened and endangered
NOAEL = No observed adverse effects level
HQ = Hazard (risk) quotient

1

Table 8-2. Soil Ingestion Rates For Hanford Receptors

Receptors	Soil Ingestion (kg/kgBW/d)	Source
Reported (Direct and Indirect)		
Canada goose	0.00254	Reported Soil fraction, 8.2% (EPA 1993, Table 4-4) *x* reported Food Ingestion Rate 0.031 (g/g-d) average adult both sexes (EPA 1993, Pg. 2-25)
Mule deer	0.00044	Reported Soil Fraction for Mule deer, 2.0% (Beyer et al.1994) *x* reported Food Ingestion Rate, 0.022 kg/kgBW/d (Alldredge et al. 1974 in Higley and Kuperman 1996)
Surrogate (Direct and Indirect)		
Great Basin pocket mouse	0.0057	Reported Soil Fraction for white-footed mouse, 2.0% (EPA 1993, Table 4-4) *x* reported Food Ingestion Rate for the Great Basin pocket mouse, 0.285 kg/kgbw/d (Calder 1984 in DOE-RL 1995)
Spotted sandpiper	0.056	Calculated from EPA (1993) using equations 3-39, 3-40, 4-11, and 4-12; Soil fraction for Stilt sandpiper 0.17 (Table 4-4), Diet of 100% Crustacea; Crustacea gross energy = 1.1 kcal/g wet wt. (Table 4-1); Waterfowl assimilation efficiency = 77% (Table 4-3); Body weight = 42.5g (average adult both sexes)
Calculated (Indirect only)		
Bald eagle	0.0000003	Calculated from soil-to-body weight ratio of prey
Red-tailed hawk	0.0004	Calculated from soil-to-body weight ratio of prey
Burrowing owl	0.0133	Calculated from soil-to-body weight ratio of prey
Western meadowlark	0.0082	Calculated from soil-to-body weight ratio of prey
Great blue heron	0.000005	Calculated from soil-to-body weight ratio of prey
Mink	0.000044	Calculated from soil-to-body weight ratio of prey
Coyote	0.00011	Calculated from soil-to-body weight ratio of prey

2

1

Table 8-3. Derivation of Soil-to-Body Weight Ratios for Prey of Hanford Receptors

Prey Species	SF	IR_f (kg/kgBW/d)	Rt (d)	SBR (kg/kgBW) =SFxIR_fxRt
Pronghorns[a]	0.057	0.015	1	0.0009
Meadow vole	0.024	0.81	1	0.0194
Black-tailed Jackrabbit[b]	0.063	0.047	1	0.003
Great Basin Pocket Mouse[c]	0.02	0.285	1	0.0057
Bullfrog	ND	0.0002	1	0[d]
Mallard	0.02	0.00025	1	0.000005

SF = Soil Ingestion Fraction (kg soil ingested/kg food ingested); all values for EPA (1993b) unless noted
IR_f =Food Ingestion Rate = ratio of amount of food ingested daily (kg/d) to body weight (kg/Bw)
Rt =Residence times (days); value assumed to be 1 day
SBR = Soil-to-body weight ratio
ND = No data

[a]Soil ingestion fraction and food ingestion rates from Arthur and Gates (1988)
[b]Food ingestion rate from Arthur and Gates (1988)
[c] Values for the white-footed mouse (Calder 1984 in EOE-RL 1995)
[d]Assumed to be zero

1

Table 8-4. Derivation of Indirect Soil Ingestion Rate for Hanford Receptors

Receptor	Receptor IR_f (kg/kgBW/d)	Prey														Receptor IS (kg/kgBW/d)
		Earthworm		Pronghorn[b]		Meadow vole		Black-tailed[c] jackrabbit		Great Basin[d] pocket mouse		Bullfrog[e]		Mallard[f]		
		DF	SBR[a]	DF	SBR	DF	SBR	DF	SBR	DF	SBR	DF	SBR	DF	SBR	$IS=IR_f x S_a(DF_a x SBR_a)$
Bald eagle[g]	0.12	0	0.53	0	0.0009	0	0.0194	0	0.003	0.0	0.0057	0	0	0.53	0.000005	0.0000003
Red-tailed hawk[h]	0.105	0	0.53	0	0.0009	0.024	0.0194	0.26	0.003	0.46	0.0057	0	0	0.26	0.000005	0.0004
Burrowing owl[i]	0.092	0.26	0.53	0	0.0009	0.31	0.0194	0	0.003	0.22	0.0057	0.21	0	0.00	0.000005	0.0133
Western meadowlark[j]	0.028	0.55	0.53	0	0.0009	0	0.0194	0	0.003	0	0.0057	0	0	0.00	0.000005	0.0082
Great blue heron	0.18	0	0.53	0	0.0009	0	0.0194	0	0.003	0.005	0.0057	0.04	0	0.005	0.000005	0.000005
Mink[k]	0.14	0	0.53	0	0.0009	0	0.0194	0	0.003	0.06	0.0057	0.05	0	0.06	0.000005	0.000044
Coyote[l]	0.047	0	0.53	0.52	0.0009	0	0.0194	0.21	0.003	0.21	0.0057	0	0	0.04	0.000005	0.00011

IR_f = Food Ingestion Rate; values taken from receptor parameter profiles
DF = Dietary fraction
SBR = Soil-to-Body weight ratio (see Table 8-3)
BW = body weight
IS = Indirect soil ingestion rate (kg/kgbw/d)

[a]Earthworm SBR values derived from soil concentrations and acid-insoluble residue data for earthworms (Stafford and McGrath 1986)
[b]Pronghorn includes deer and other ungulates
[c]Black-tailed jackrabbit include lagomorphs
[d]Great Basin pocket mouse includes mole, squirrel, porcupine, pocket gopher, deer mouse, and rat
[e]Bullfrog includes all herpetofauna; SBR assumed to be zero
[f]Mallard includes all birds
[g] Food ingestion rate and diets from EPA (1993b); values for the bald eagle (EPA 1993b); Washington, River; all birds assumed to be mallards
[h] Food ingestion rate and diets from EPA (1993b); summer, Alberta, Canada, farm and woodlands,
[i]Values for dietary fractions are values for the American Kestrel (EPA 1993b); California/hayfields, pasture plus; 0.925% added to each dietary fraction per prey represented
[j]Values for dietary fractions for the American Robin (EPA 1993b); Western United States % volume; stomach contents, average of all seasons; remainder dietary fraction 0.44 is fruit. We assume fruit SBR = 0
[k]Food ingestion rate and diets from EPA (1993b) pg 2-253; average all seasons; Michigan/stream, river year-round
[l]Food Ingestion Rate for Adults (Gier 1975 in Higley and Kuperman 1996), dietary fraction Table 1 in MacCracken (1984) Foods in the Black Hills, South Dakota (*J. Wildl. Manage.*48)
Herbivorous receptors such as mule deer and Canada goose are absent from table
Diets for bald eagle, great blue heron, and mink are exclusive of fish, therefore; we assume fish SBR = 0

Table 8-5 Toxicity Equivalence Factors for Chlorinated Dioxins, Chlorinated Dibenzofurans, and PCBs[a]

Congener	Receptor		
	Mammals	Birds	Fish
Dioxins[b]			
2,3,7,8-TCDD	1.0	1.0	1.0
1,2,3,7,8-PeCDD	1.0	1.0	1.0
1,2,3,4,7,8-HxCDD	0.1	0.05	0.5
1,2,3,6,7,8-HxCDD	0.1	0.01	0.01
1,2,3,7,8,9-HxCDD	0.1	0.1	0.01
1,2,3,4,6,7,8-HpCDD	0.01	<0.001	0.001
OCDD	0.0001	NA	NA
Dibenzofurans[c]			
2,3,7,8-TCDF	0.1	1.0	0.05
1,2,3,7,8-PeCDF	0.05	0.1	0.05
2,3,4,7,8-PeCDF	0.5	1.0	0.5
1,2,3,4,7,8-HxCDF	0.1	0.1	0.1
1,2,3,6,7,8-HxCDF	0.1	0.1	0.1
2,3,4,6,7,8-HxCDF	0.1	0.1	0.1
1,2,3,7,8,9-HxCDF	0.1	0.1	0.1
1,2,3,4,6,7,8-HpCDF	0.01	0.01	0.01
1,2,3,4,7,8,9-HpCDF	0.01	0.01	0.01
OCDD	0.0001	0.0001	0.0001
PCBs[d]			
3,4,4',5-TCB	0.0001	0.1	0.0005
3,3',4,4'-TCB	0.0001	0.05	0.0001
3,3',4,4',5-PeCB	0.1	0.1	0.005
3,3',4,4',5,5'-HxCB	0.01	0.001	0.00005
2,3,3',4,4'-PeCB	0.0001	0.0001	<0.000005
2,3,4,4',5-PeCB	0.0005	0.0001	<0.000005
2,3',4,4',5-PeCB	0.0001	0.00001	<0.000005
2',3,4,4',5-PeCB	0.0001	0.00001	<0.000005
2,3,3',4,4',5-HxCB	0.0005	0.0001	<0.000005
2,3',4,4',5,5'-HxCB	0.00001	0.00001	<0.000005
2,3,3',4,4',5,5'-HpCB	0.0001	0.00001	<0.000005
2,2',3,3',4,4',5'-HpCB	NA	NA	NA
2,2',3,4,4',5,5'-HpCB	NA	NA	NA

T = tetra, Pe = penta, Hx = hexa, Hp = hepta, O = octa, NA = not available
[a] Values from WHO 1997
[b] CDD = chlorinated dibenzodioxin
[c] CDF = chlorinated dibenzofuran
[d] CB = chlorinated biphenyl

9. Risk-Based Emissions Criteria

Risk-based emission criteria will be developed, as needed, for protection of human and ecological resources. This will be done in the PRA and updated in the FRA. Risk-based emission criteria will be developed for plausible scenarios.

9.1. Human Health Risk-Based Emission Levels

During the PRA, any COPCs, which do not meet the target risk goals, will be evaluated further to determine the driving factors behind the risk and the potential uncertainty associated with them. Risk-based emission limits will be calculated for any COPC which exceeds risk goals in the PRA and these compounds will be further evaluated in the FRA. For example, additional site-specific information will be available for the FRA. Risk goals that are exceeded will be addressed to the satisfaction of Ecology and EPA prior to approval of the FRA.

Risk-based emission levels will be calculated using the same exposure scenarios, pathways, toxicity values, and equations used to calculate plausible risk estimates. The risk equations will be used to back-calculate acceptable COPC concentrations in various media based on the following risk goals: for carcinogens a target incremental lifetime cancer risk level of 1×10^{-5} and for noncarcinogens with a HQ of 0.25. Air dispersion modeling results will then be used to convert these media concentrations to risk-based emission limits.

9.2. Ecological Receptor-based Emissions Levels

During the PRA, chemicals with ecological HQs greater than 0.25 for a given receptor will need development of a risk-based emission level. This emission level will be developed by setting the HQ to 0.25 and back-calculating through the same TRVs and exposure values in the food web, and the same dispersion modeling to the stack emission for plausible scenarios. The result will be risk-based emissions for protection of ecological resources.

10. Summary

A master summary will be provided in the SLRA that will capture the major points in each of the following areas:

- The SLRA approach, which includes identification of source COPCs
- The melter process, and quantification of chemical and radiological emissions
- Organization of environmental modeling to meet environmental requirements and to secure the dangerous waste permit from the State of Washington
- PRA and FRA human health risk assessment
- PRA and FRA ecological risk assessment
- Uncertainties
- Development of receptor-based emission levels

11. References

64 FR 14308. *Endangered and Threatened Species: Threatened Status for Three Chinook Salmon Evolutionarily Significant Units in Washington and Oregon, and Endangered Status of One Chinook Salmon ESU in Washington; Final Rule* and *Partial 6-Month Extension on Final Listing Determinations for Four Evolutionarily Significant Units of West Coast Chinook Salmon; Proposed Rule,* Volume 64, Number 56, pp 14308-14328, March 24, 1999. Federal Register.

64 FR 14517. *Endangered and Threatened Species: Threatened Status for Two ESUs of Chum Salmon in Washington and Oregon, for Two ESUs of Steelhead in Washington and Oregon, and for Ozette Lake Sockeye Salmon in Washington; Rules,* Volume 64, Number 57, pp 14507-14517, March 25, 1999. Federal Register.

ACGIH. 1998. *Guide to Occupational Exposure Levels,* Compiled by American Conference of Governmental Industrial Hygienists, Cincinnati, Ohio, USA.

Achey PM. 1999. Personal communication to Hadden C, 20 September 1999. University of Florida, USA.

AIHA. 1996. *Emergency Response Planning Guideline and Workplace Environmental Exposure Level Guides Handbook.* American Industrial Hygiene Association, Fairfax, Virginia, USA.

AIHA. 1998. *The AIHA Emergency Response Planning Guidelines and Workplace Environmental Exposure Level Guides Handbook.* American Industrial Hygiene Association Press, Fairfax, Virginia, USA.

Arthur WJ III, and Gates RJ. 1988. *Trace element intake via soil ingestion in pronghorns and black-tailed jackrabbits,* Journal of Range Management, volume 41, number 2, pp162-166.

ASTM. 1994. *D2937-94 Standard Test Method for Density of Soil in Place by the Drive-Cylinder Method,* 1994, American Society for Testing and Materials, West Conshohocken, Pennsylvania, USA.

ASTM. 1998. *D2216-98 Standard Test Method for Laboratory Determination of Water (Moisture) Content of Soil and Rock by Mass,* 1998. American Society for Testing and Materials, West Conshohocken, Pennsylvania, USA.

ATG. 1998. *Mixed Waste Facility RCRA/TSCA Permit Application.* Attachment 4 – Risk Assessment Work Plan. Rev. 4, 26 June 1998. Richland, Washington, USA.

Atkinson D, and Lee R. 1992. *Procedures for Substituting Values for Missing NWS Meteorological Data for Use in Regulatory Air Quality Models.* US Environmental Protection Agency, Office of Air Quality Planning and Standards, Research Triangle Park, North Carolina, USA.

.

Bacci E, Calamari D, Gaggi C, and Vighi M. 1990. *Bioconcentration of Organic Chemical Vapors in Plant Leaves: Experimental Measurements and Correlation,* Environmental Science and Technology. Volume 24, Number 6, pp 885-889.

Bacci E, Cerejeira C, Chemello G, Calamari D, and Vighi, M. 1992. *Chlorinated Dioxins: Volatilization from Soils and Bioconcentration in Plan Leaves,* Bullletin of Environmental Contamination and Toxicology, Volume 48, pp 401-408.

Baes CF, Sharp RD, Sjoreen AL and Shor RW. 1984. *A Review and Analysis of Parameters for Assessing Transport of Environmentally Released Radionuclides Through Agriculture*, ORNL-5786, September 1984. Martin Marietta Energy Systems, Oak Ridge, Tennessee, USA.

Baker DA, and Soldat JK. 1992. *Methods for Estimating Doses to Organisms from Radioactive Materials Released into the Aquatic Environment*, PNL-8150, June 1992. Pacific Northwest Laboratory, Richland, Washington, USA.

Barnthouse LW. 1995. *Effects of Ionizing Radiation on Terrestrial Plants and Animals: A Workshop Report*, ORNL/TM-13141, December 1995. Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA.

Bent AC. 1929. *Life Histories of North American Shore Birds*, Part 2. US National Museum, Bulletin 146, US Government Printing Office, Washington, D.C., USA.

Beyer WN, Connor EE, and Gerould S. 1994. *Estimates of Soil Ingestion by Wildlife*, Journal of Wildlife Management, volume 58, number 2 pp 375-382.

Beyer WN. 1990. *Evaluating Soil Contamination.* US Fish and Wildlife Service Biol. Report 90(2).

Bintein SJ, Devillers J, and Karcher W. 1993. *Nonlinear Dependence of Fish Bioconcentrations on n-Octanol/Water Partition Coefficients*, SAR and QSAR in Environmental Research, volume 1. pp 29-39.

Blaylock BG, Frank ML, and O'Neal BR. 1993. *Methodology for Estimating Radiation Dose Rates to Freshwater Biota Exposed to Radionuclides in the Environment*, ES/ER/TM-78. Martin Marietta Energy Systems, Oak Ridge, Tennessee, USA.

BNFL. 1999a. *TWRS-P Facility Dangerous Waste Permit Application*, BNFL-5193-RCRA-01, Draft Revision 0C, 7 May 1999. BNFL Inc., Richland, Washington, USA.

BNFL. 1999b. *Quality Assurance Project Plan for the Environmental Performance Test*, QP-W375-EN00001, Revision E, Draft, 13 December1999. BNFL Inc., Richland, Washington, USA.

BNFL. 1999c. *Environmental Performance Test Plan*, PL-W375-EN00003, Revision C, Draft, 7 May 1999. BNFL Inc., Richland, Washington, USA.

Boothe GF. 1996. *Predominant Radionuclides in Hanford Site Waste Tanks*, WHC-SD-WM-TI-731, Revision 0, January 1996. Westinghouse Hanford Company, Richland, Washington, USA.

Brandt C. 1995. Personal Communication with B. Cornaby, October 5, 1995, Biological Resources Specialist, Pacific Northwest National Laboratory, Richland, Washington, USA.

Brown L, and Amadon D. 1968. *Eagles, Hawks, and Falcons of the World*, Volume 1. McGraw-Hill Book Company, New York, New York, USA.

Bull J, and Farrand. 1994. *National Audubon Society Field Guide to North American Birds*. Alfred A. Knopf, New York, New York, USA.

Cal/EPA. 1995. Draft for Public Comment, *Technical Support Document for the Determination of Acute Toxicity Exposure Levels for Airborne Toxicants*. Office of Environmental Health Hazards Assessment, California Environmental Protection Agency.

Calder WA. 1984. *Size, Function, and Life History*. Harvard University Press, Cambridge, Massachusetts, USA.

Campbell JA and others. 1995. *Flammable Gas Safety Program. Organic Analysis and Analytical Methods Development: FY1995 Progress Report*, PNL-1077b, Pacific Northwest Laboratory, Richland, Washington, USA.

Corp N, and Morgan AJ. 1991. *Accumulation of Heavy Metals From Polluted Soils by the Earthworm, Lumbricus rubellus: Can laboratory exposure of "control" worms reduce biomonitoring problems?*, Environmental Pollution, volume 74, pp 39-52.

Cowardin LM, Carter V, Golet FC, and LaRoe ET. 1979. *Classification of Wetlands and Deepwater Habitats of the United States*, FWS/OBS-79/31, December 1979.

Cowherd C, Muleski, G., Engelhart P., and Gilete D. 1985. *Rapid Assessment of Exposure to Particulate Emissions from Surface Contamination.* Prepared for EPA Office of Health and Environmental Assessment.

Craven SR, and Hunt RA. *1984. Fall food habits of Canada geese in Wisconsin*, Journal of Wildlife Management, volume 48, pp 169-173. In: US Environmental Protection Agency 1993, *Wildlife Exposure Factors Handbook*, EPA/600/R-93/187. US Environmental Protection Agency, Office of Research and Development, Washington, D.C., USA.

Cushing CE. 1992. *Hanford Site National Environmental Policy Act (NEPA) Charact*erization, PNL-6415, Revision 5, December 1992. Pacific Northwest Laboratory, Richland, Washington, USA.

Cushing CE, Baker DA, Chamness MA, Crist ME, Fruland RM, Harvey DW, Hoitink DJ, Moore EB, Poston TM, Seely HE, and others. 1994. *Hanford Site National Environmental Policy Act (NEPA) Characterization*, PNL-6415, Revision 6, August 1994. Pacific Northwest Laboratory, Richland, Washington, USA.

Cushing CE, Baker DA, Chamness MA, Harvey DW, Hoitink DJ, Hostick DJ, Mazaika R, Moore EB, Poston TM, Seely HE, and others. 1995. *Hanford Site National Policy Act (NEPA) Characterization*, PNL-6415, Revision 7, September 1995. Pacific Northwest Laboratory, Richland, Washington, USA.

Daubenmire R. 1970. *Steppe Vegetation of Washington*, Technical Bulletin 62, Experimental Station. Washington State University, Pullman, Washington, USA.

Dempsey CR, and Oppelt ET. 1993. *Incineration of Hazardous Waste: A Critical Review Update*, Air & Waste, volume 43, pp. 25-73, January 1992. The Future for Hanford: Uses and Cleanup, US Department of Energy, Washington, D.C., USA.

DHHS. 1997. *Technical Support Document for ATSDR* Interim *Policy Guideline: Dioxin and Dioxin-like Compounds in Soil*. U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry, Atlanta, Georgia, USA.

DHHS. 1998. *Minimal Risk Levels for Hazardous Substances*, U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry, Atlanta, Georgia, USA

DOE. 1990. *Radiation Protection of the Public and the Environment*, DOE Order 5400.5. US Department of Energy, Washington, D.C., USA.

DOE. 1992. *The Future for Hanford: Uses and Cleanups*, 1992. US Department of Energy, Washington, D.C., USA.

DOE. 1996. *Draft Hanford Remedial Action Environmental Impact Statement and Comprehensive Land Use Plan*, Volume 1 of 4, DOE/EIS-0222D, August 1996. US Department of Energy, Washington, D.C., USA.

DOE. 1997. *Toxicity-Based Methodology, Text of Calc. Note, S-CLC-G-0074*, Facsimile Transmission from Craig D, DOE, to Monschein E, TetraTech EM Inc., 26 September 1997. US Department of Energy, Washington, D.C., USA.

DOE. 1998. *Radiological Benchmarks for Screening Contaminants of Potential Concern for Effects on Aquatic Biota at Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA*. BJC/OR-80. Bechtel Jacobs Company LLC, Oak Ridge, Tennessee, USA.

DOE and Ecology. 1996. *Final Environmental Impact Statement for the Tank Waste Remediation System, Hanford Site, Richland, Washington, USA*, DOE/EIS-0189, August 1996. US Department of Energy, Washington, D.C., USA, and Washington State Department of Ecology, Olympia, Washington, USA.

DOE-RL. 1991. *Double-Shell Tank System Dangerous Waste Permit Application*, DOE/RL-90-39, Revision 0, June 1991. US Department of Energy, Richland Operations Office, Richland, Washington, USA.

DOE-RL. 1995. *Hanford Site Risk Assessment Methodology*, DOE/RL-91-45, Revision 3, May 1995. US. Department of Energy, Richland Operations Office, Richland, Washington, USA.

DOE-RL. 1998. *Screening Assessment and Requirements for a Comprehensive Assessment: Columbia River Comprehensive Impact Assessment*, DOE/RL-96-16, Revision 1, March 1998. US Department of Energy, Richland Operations Office, Richland, Washington, USA.

Downs JL, Rickard WH, Brandt CA, Cadwell LL, Cushing CE, Geist DR, Mazaika RM, Neitzel DA, Rogers LE, Sackschewsky MR, and Nugent JJ. 1993. *Habitat Types on the Hanford Site: Wildlife and Plant Species of Concern*, PNL-8942, December 1993. Pacific Northwest Laboratory, Richland, Washington, USA.

Duranceau DA. 1995. *Site Evaluation Report for Candidate Basalt Quarry Sites*, BHI-00005, Revision 0, February 1995. Bechtel Hanford Inc., Richland, Washington, USA.

Eagle TC, and Whitman JS. 1987. *Wild Furbearer Management and Conservation,* Novak M, Baker JA, Obbarel ME, and others, eds., pp 615-624. University of Pittsburgh Press, Pittsburgh, Pennslyvania, USA.

Earhart and Johnson. 1970. *Size Dimorphism and Food Habits of North American Owls.* Condor, volume 72, pp 251-264.

Eckerman KF, and Ryman JC. 1993. *External Exposure to Radionuclides in Air, Water and Soil,* EPA 402-R-93-081, 1993, Federal Guidelines Report Number 12. US Environmental Protection Agency, Office of Radiation and Indoor Air, Washington, D.C., USA.

Efroymson RA, Will ME, Suter II GW, and Wooten AC. 1997a. *Toxicological Benchmarks for Screening Contaminants of Potential Concern for Effects on Terrestrial Plants: 1997 Revision*, ES/ER/TM-85/R3, Lockheed Martin Energy Systems, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA.

Eroymson RA, Will ME, and Suter II GW. 1997b. *Toxicological Benchmarks for Screening Contaminants of Potential Concern for Effects on Soil and Litter Invertebrates and Heterotrophic Process: 1997 Revision*, ES/ER/TM-126/R2. Lockheed Martin Energy Systems, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA.

EPA. 1986. *Test Methods For Evaluating Solid Waste, Physical and Chemical Methods*, SW-846, Third Edition (as amended by Updates I, II, IIA, IIB, and III), 1986. US Environmental Protection Agency, Washington, D.C., USA

EPA. 1988. *Federal Guidance Report No. 11: Limiting Values of Radionuclide Intake and Air Concentration and Dose Conversion Factors for Inhalation, Submersion, and Ingestion*, EPA/520/1-88/020. US Environmental Protection Agency, Washington D.C., USA.

EPA. 1989a. *Guidance on Setting Permit Conditions and Reporting Trial Burn Results*, EPA/625/6-89/019. US Environmental Protection Agency, Washington, D.C., USA.

EPA. 1989b *Risk Assessment Guidance for Superfund, Volume 1: Human Health Evaluation Manual (Part A),* EPA/540/1-89/002. US Environmental Protection Agency, Washington, D.C., USA.

EPA. 1989c. *Exposure Factors Handbook*, EPA/600/8-89/043. U.S. Environmental Protection Agency, Washington, D.C., USA.

EPA. 1991a. Memorandum from Don R. Clay, Assistant Administrator, *Role of the Baseline Risk Assessment in Superfund Remedy Selection Decisions*, OSWER Directive 9355.0-30, 22 April 1991. US Environmental Protection Agency, Washington, D.C., USA.

EPA. 1991b. *Risk Assessment Guidance for Superfund, Volume 1: Human Health Evaluation Manual (Part B, Development of Risk-Based Preliminary Remediation Goals).* US Environmental Protection Agency, Washington, D.C., USA.

EPA. 1992a. *Framework for Ecological Risk Assessment*, EPA/630/R-92-001. US Environmental Protection Agency, Washington, D.C., USA.

EPA. 1992b. *Draft Report: A Cross-species Scaling Factor for Carcinogen Risk Assessment Based on Equivalence of mg/kg 3/4/day, Notice,* volume 57, number 109, pp 24152-24173. *Federal Register.*

EPA. 1993a. *Provisional Guidance for Quantitative Risk Assessment of Polycyclic Aromatic Hydrocarbons*, EPA-600-R-93-089, July 1993. US Environmental Protection Agency, Washington D.C., USA.

EPA. 1993b. *Wildlife Exposure Factors Handbook*, EPA/600/R-93/187. US Environmental Protection Agency, Washington, D.C., USA.

EPA. 1993c. *Technical Basis for Establishing Sediment Quality Criteria for Non-Ionic Chemicals by Using Equilibrium Partitioning*, EPA-822-R-93-011. US Environmental Protection Agency, Washington, D.C., USA.

EPA. 1993d. *Review Draft Addendum to the Methodology for Assessing Health Risks Associated with Indirect Exposure to Combustor Emissions*, EPA-600-AP-93-003, November 10, 1993. US Environmental Protection Agency, Washington, D.C., USA.

EPA. 1994a. *Table 1 - Chemicals Recommended for Identification and Table 2 - Chemicals for Potential Identification, Draft Exposure Assessment Guidance for Resource Conservation and Recovery Act Hazardous Waste Combustion Facilities: Attachment*, April 15, 1994. US Environmental Protection Agency, Washington, D.C., USA.

EPA. 1994b. *Combustion Emissions Technical Resource Document* (CETRED), EPA-530-R-94-014. US Environmental Protection Agency, Washington, D.C., USA.

EPA. 1994c. *Guidance Manual for the Integrated Exposure Uptake Biokinetic Model for Lead in Children,* 9265, PB93-963510, Version 0.99D, February 1994, US Environmental Protection Agency, Washington, D.C., USA.

EPA. 1995a, *User's Guide for the Industrial Source Complex (ISC3) Dispersion Models*, Volume I - User Instructions, EPA-454/B-95-003a, US Environmental Protection Agency, Research Triangle Park, North Carolina, USA.

EPA. 1995b. *User's Guide for the Industrial Source Complex (ISC3) Dispersion Models*, Volume II - Description of Model Algorithms, EPA-454/B-95-003b. US Environmental Protection Agency, Research Triangle Park, North Carolina., USA.

EPA. 1995c. *Environmental Protection Agency Risk Characterization Policy,* March 21 memorandum from Carol Browner, Administrator to EPA Staff. US Environmental Protection Agency, Washington, D.C., USA.

EPA. 1995d. *Health Effects Assessment Summary Tables (HEAST) FY-1995 Annual,* EPA/540/R-95/036. US Environmental Protection Agency, Washington, D.C., USA.

EPA. 1995e. *Supplemental Guidance to RAGS: Region 4 Bulletins, Human Health Risk Assessment, Office of Health Assessment*, November 1995. US Environmental Protection Agency, Region 4, Atlanta, Georgia, USA.

EPA 1995f. *Final Water Quality Guidance for the Great Lakes System: Final Rule.* Federal Register 60(56)15366-15425. US Environmental Protection Agency, Washington, D.C., USA.

EPA. 1996a. *Meteorological Processor for Regulatory Models (MPRM) - User's Guide,* EPA-454/B-96-002. US Environmental Protection Agency, Research Triangle Park, North Carolina, USA.

EPA. 1996b. *PCBs: Cancer Dose-Response Assessment and Application to Environmental Mixtures,* EPA/600/P-96/001F. US Environmental Protection Agency, Washington, D.C., USA.

EPA. 1996c. *Soil Screening Guidance: Technical Background Document,* EPA/540/R95/128. US Environmental Protection Agency, Washington, D.C., USA.

EPA. 1997a. *Health Effects Assessment Summary Tables (HEAST) FY-1995 Annual,* EPA/540/R-95/036. US Environmental Protection Agency, Washington, D.C, USA.

EPA. 1997b. *Exposure Factors Handbook,* EPA/600/P-95/002Fb. US Environmental Protection Agency, Washington, D.C., USA.

EPA. 1997c. *Ecological Risk Assessment Guidance for Superfund: Process for Designing and Conducting Ecological Risk Assessments,* EPA/540/R/97/006, 9285.7-25, PB97-963211, Interim Final, June 1997. US Environmental Protection Agency, Washington D.C., USA.

EPA. 1997d. *Mercury Study Report to Congress, Volumes I through VIII,* EPA/452/R-97-001, December 1997. US Environmental Protection Agency, Washington, D.C., USA.

EPA. 1998a. *Human Health Risk Assessment Protocol for Hazardous Waste Combustion Facilities,* Peer Review Draft, EPA/530/D-98/001B. US Environmental Protection Agency, Washington, D.C., USA.

EPA. 1998b. *User's Guide for the Industrial Source Complex (ISC3) Dispersion Models,* Addendum for Volume I - User Instructions. US Environmental Protection Agency, Research Triangle Park, North Carolina, USA.

EPA. 1998c. *Integrated Risk Information System (IRIS) On-line Database of Toxicity Measures.* US Environmental Protection Agency, Cincinnati, Ohio, USA.

EPA. 1998d. *Guidelines for Ecological Risk Assessment,* EPA/630/R-95/002Fa. US Environmental Protection Agency, Washington, D.C., USA.

EPA. 1998e. *Methodology for Assessing Health Risks Associated with Multiple Pathways of Exposure to Combustor Emissions,* EPA 600/R-98/137, US Environmental Protection Agency, Washington, D.C., USA.

EPA. 1999a. *Screening Level Ecological Risk Assessment Protocol.* US Environmental Protection Agency, Region 6, USA.

EPA. 1999b. *Screening* Level *Ecological Risk Assessment Protocol for Hazardous Waste Combustion Facilities* (Peer Review Draft). EPA 530-D-99-001A. US Environmental Protection Agency, Washington, D.C., USA.

EPA. 1999c. *Cancer Risk Coefficients for Environmental Exposure to Radionuclides, Federal Guidance Report No. 13*, EPA 402-R-99-001, September 1999. US Environmental Protection Agency, Washington, D.C., USA.

EPA. 1999d. Personal communication with Marcia Bailey and Catherine Massimino of the US Environmental Protection Agency, October 18, 1999. US Environmental Protection Agency, Washington, D.C., USA.

EPA. 1999e. Personal communication with US Environmental Protection Agency, Region X, on September 16, 1999.

EPA 1999f. Personal communication with Marcia Bailey of the US Environmental Protection Agency, October 8 1999. US Environmental Protection Agency, Washington, D.C., USA.

ERDA. 1975. *Final Environmental Statement: Waste Management Operations, Hanford Reservation, Richland, Washington*, ERDA-1538, December 1975. Energy Research and Development Administration, Richland, Washington, USA.

Fitzner RE, and Gray RH. 1991. *The Status, Distribution, and Ecology of Wildlife on the US DOE Hanford Site: A Historical Overview of Research Activities*, Environmental Monitoring and Assessment, volume18 pp 173-202.

Franklin JF, and Dyrness CT. 1973. *Natural Vegetation of Oregon and Washington*, General Technical Report PNW-8. US Forest Service, Washington, D.C., USA.

Gano KA, and Rickard WH. 1982. *Small mammals of a bitterbrush-cheatgrass community*, Northwest Science, volume 56, pp 1-7.

Gish CD. 1970. *Organochlorine Insecticide Residues in Soils and Soil Invertebrates from Agricultural Lands*, Pesticides Monitor Journal, volume 3, number 71.

Gray RH, and Rickard WH. 1989. *The Protected Area of Hanford as a Refugium for Native Plants and Animals*, Environmental Conservation, volume 16, number 3, pp 251-260. The Foundation for Environmental Conservation, Switzerland.

Hall, John A. 1998. *Biodiversity Inventory and Analysis of the Hanford Site, 1997 Annual Report.* The Nature Conservancy of Washington, Seattle, Washington, USA.

Harper, BL. 1999. Personal communication with Barbara Harper, technical contact for Native-Americans. November 1999.

Harris SG, and Harper BL. 1997. *A Native American Exposure Scenario*, Risk Analysis, volume 17, number 6. Plenum Publishing Corporation, New York, New York, USA.

HAZWRAP. 1994. *Loring Air Force Base Ecological Risk Methodology,* August 1994. Hazardous Waste Remedial Actions Program, Lockheed Martin Energy Systems, Inc., Oak Ridge, Tennessee, USA.

Higley KA, and Kuperman R. 1996. *Radiological benchmarks for wildlife at Rocky Flats Environmental Technology Site*. Environmental Assessment Division, Argonne National Laboratory, Chicago, Illinois, USA.

Hutzinger O, Safe S, and Zitko V. 1983. *The Chemistry of PCB's.* Krieger Publishing Co., Malaber, Florida.

IAEA. 1992. *Effects of ionizing radiation on plants and animals at levels implied by current radiation protection standards*, Technical Report Series Number 332. International Atomic Energy Agency, Vienna, Austria.

Johnson. 1970. *Size dimorphism and food habits of North American owls*, Condor, volume 72, pp 251-265

Jones DS, Suter II GW, and Hull RN. 1997. *Toxicological Benchmarks for Screening Contaminants of Potential Concern for Effects on Sediment-Associated Biota: 1997 Revision*, ES/ER/TM-95/R4, November 1997. Lockheed Martin Energy Systems, Inc., Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA.

Klaassen CD, Amdur MO, and Doull J. 1996. *Casarett and Doull's Toxicology: The Basic Science of Poisons*, 5th Edition. MacMillan Publishing Co., Inc., New York, New York, USA.

Klinger GS and others. 1999. *Organic Analysis of AW-101 and AN-107 Tank Waste*, BNFL-RPT-001 Rev.0. BNFL Inc., Richland, Washington, USA.

Kupfer MJ, Boldt AL, Hodgson KM, Shelton LW, Simpson BC, Watrous RA, LeClair MD, Borsheim GL, Winward RT, Higley BA, and others. 1997. *Standard Inventories of Chemicals and Radionuclides in Hanford Site Tank Wastes*, HNF-SD-WM-TI-740, Revision 9, August 1997. Fluor Daniel Hanford, Inc., Richland, Washington, USA.

Lorber M and Pinsky P. 1999. *An Evaluation of Three Empirical Air-to-Leaf Models for Polychlorinated Dibenzo-p-Dioxins and Dibenzofurans*. National Center for Environmental Assessment (NCEA). US Environmental Protection Agency, Washington, D.C., USA.

Lyman WJ, Reehl WF, and Rosenblatt DH. 1982. *Handbook of Chemical Property Estimation Methods: Environmental Behavior of Organic Compounds*. McGraw-Hill Book Company, New York, New York, USA.

MacCracken JG. 1984. *Coyote foods in the Black Hills, South Dakota*. Journal of Wildlife Management, volume 4, number 4, pp1420-1423.

Matlock KS, and Pegg IL. 1999. *Determination of the Fate of Hazardous Organics During Vitrification of TWRS LAW and HLW Simulants*, VSL-99R3580-1, Revision 0, February 19, 1999. Vitreous State Laboratory, The Catholic University of America, Washington, D.C., USA.

Meinhardt CC, and Frostenson JC. 1979. *Current Status of the 200 Area Ponds*, RHO-CD-798, November 1979. US Department of Energy, Richland Hanford Operations, Richland, Washington, USA.

NAC. 1997. *National Advisory Committee for the Development of Acute Exposure Guideline Levels (AEGL) for Hazardous Substances 1997*, volume 62, number 210, pp 58839-58851, 30 October 1997. *Federal Register*.

Neitzel DA, Fosmire CJ, Fowler RA, Goodwin SM, Harvey DW, Hendrickson PL, Hoitink DJ, Poston TM, Rohay AC, Thorne PD, and Wright MK. 1998. *Hanford Site National Environmental Policy Act*

(NEPA) Characterization, PNL-6415, Revision 10, September 1998. Pacific Northwest National Laboratory, Richland, Washington, USA.

NPS 1994, *The Hanford Reach of the Columbia River Final River Conservation Study and Environmental Impact Statement*, June 1994. US Department of Interior, Pacific Northwest Regional Office, National Park Service, Seattle, Washington, USA.

NRC. 1992. *Residual Radioactive Contamination from Decommissioning, Technical Basis for Translating Contamination Levels to Annual Total Effective Dose Equivalent*, Final Report, NUREG/CR-5512, PNL-7794, Volume 1, 1992. US Nuclear Regulatory Commission, Washington, D.C., USA.

Opresko DM, Sample BE, and Suter II GW. 1994. *Toxicological Benchmarks for Wildlife: 1994 Revision*, ES/ER/TM-86/R1. Environmental Sciences Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA.

Oring LW, Lank DB, and Maxson SJ. 1983. *Population Studies of the Polyandrous Spotted Sandpiper*, Auk volume 100, pp 272-285.

ORNL 2000. Risk Assessment Information System, maintained on-line by the Oak Ridge National Laboratory at http://risk.lsd.ornl.gov/rap_hp.htm, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA.

Paustenbach DJ. 1989. *A Survey of Health Risk Assessment*, The Risk Assessment of Environmental and Human Health Hazards: A Textbook of Case Studies, Paustenbach DJ, ed. New York, New York, USA.

PNL. 1993. *Hanford Site Environmental Report for Calendar Year 1992*, PNL-8682, June 1993. Pacific Northwest Laboratory, Richland, Washington, USA.

PNL. 1994. *Biological Review of the Tank Waste Remediation System (TWRS) Sites*, Letter 94-WHC-142 from Brandt CA, Pacific Northwest Laboratory, to Granger JG. Westinghouse Hanford Corporation, Richland, Washington, USA.

PNNL. 1997a. *Hanford Site Environmental Report for Calendar Year 1996*, PNNL-11472, August 1997. Pacific Northwest National Laboratory, Richland, Washington, USA.

PNNL. 1997b. *Tank 241-Z-101 Criticality Assessment Resulting form Pump Jet Mixing Sludge Mixing Simulations*, PNNL-11486/UC-2030. Pacific Northwest National Laboratory, Richland, Washington, USA.

PNNL. 1998. *Regulatory Data Quality Objectives Supporting Tank Waste Remediation System Privatization Project*, PNNL-12040, Rev. 0. Pacific Northwest National Laboratory, Richland, Washington, USA.

Poston TM, Price KL, and Newcomer DR. 1991. *An Evaluation of the Chemical, Radiological, and Ecological Conditions of West Lake on the Hanford Site*, PNL-7662, March 1991. Pacific Northwest Laboratory, Richland, Washington, USA.

Rickard W, Rogers LE, Vaughn BE, and Liebetrau SF. 1988. *Shrub-Steppe Balance and Charge in a Semi-Arid Terrestrial Ecosystem*. Elsevier, Amsterdam, Holland.

Rickard WH, and Poole LD. 1989. *Terrestrial Wildlife of the Hanford Site: Past and Future*, Northwest Science, volume 63, number 4.

Rogers LE and Rickard WH. 1977. *Ecology of the 200 Area Plateau Waste Management Environs: A Status Report*, PNL-2253, October 1977. Pacific Northwest Laboratory, Richland, Washington, USA.

Sackschewsky MR, Landeen DS, Downs JL, Rickard WH, and Baird GI. 1992. *Vascular Plants of the Hanford Site*, WHC-EP-0554, Revision 1, July 1992. Westinghouse Hanford Company, Richland, Washington, USA.

Saito KN, Petoussi-Henss N, and Zanki M. 1998. *Calculation of the Effective Dose and Its Variation from Environmental Gamma Rays*, Health Physics, volume 74, number 6, pp 698-706.

Sample BE, Alpin MS, Efroymson RA, Suter II GW, and Welsh CJ. 1997. *Methods and tools for estimation of the exposure of terrestrial wildlife to contaminants*, ORNL/TM-13391, 1997. Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA.

Sample BE, Beauchamp JJ, Efroymson RA, Suter II GW, and Ashwood TL. 1998. *Development and Validation of Bioaccumulation Model's for Earthworms*. US Department of Energy Office of Environmental Management.

Sample BE, Opresko DM, and Suter II GW. 1996. *Toxicological Benchmarks for Wildlife: 1996 Revision*, ES/ER/TM-86/R3, 1996. Lockheed Martin Energy Systems, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA.

Sample, B. 1999. Personal communication to Thomas Burns (SAIC) at the 1999 Annual Meeting of the Society of Environmental and Toxicology and Chemistry, Charlotte, NC, USA.

SCAPA 1997b, *SCAPA ERPG Working List, Revised 2/26/97*. World wide web address http://www.sep.bnl.gov/es/scapa.htm. Accessed on October 1, 1997.

Schuler CA, Rickard WH, and Sargeant GA. 1988. *Bird Associations with Shrub-steppe Plant Communities at the Proposed Reference Repository Location in Southeastern Washington*, PNL-6493, March 1988. Pacific Northwest Laboratory, Richland, Washington, USA.

Schuler CA, Rickard WH, and Sargeant GA. 1993. *Conservation of Habitats for Shrub-Steppe Birds* Environmental Conservation, volume 20, number 1, pp 57-64.

Skarzinskas R. 1998. Notes from meeting of June 30th with Barbara Harper and Stuart Harris.

Southworth GR, Beauchamp JJ, and Schnieder PK. 1978a. *Bioaccumulation Potential of Polycyclic Aromatic Hydrocarbons in Daphnia pulex*, Water Res., volume 12, pp 973-977.

Southworth GR, Beauchamp JJ, and Schnieder PK. 1978b. *Bioaccumulation Potential and Acute Toxicity of Synthetic Fuels Effluents in Fresh Water Biota: azaarenes*, Environmental Science Technology, volume 12, pp1062-1066.

Spromberg JA, Bettina JM, and Landis WG. 1998. *Metapopulation Dynamic: Indirect Effects in Multidistinct Outcome in Ecological Risk Assessments*, Environ, Toxicol. Chem. 17:1640-1649.

Spromberg, JA, John, BM, and Landis, WG. 1998. *Metapopulation Dynamic: Indirect Effects in Multidistinct Outcome in Ecological Risk Assessments*, Environ, Toxicol, Chem. Volume 17, pp 1640-1649.

Stafford EA and McGrath SP. 1986. *The Use of Acid-insoluble Residue to Correct for the Presence of Soil-derived Metals in the Gut of Earthworms Used as Bio-Indicator Organisms*, Environ. Pollut. A. Ecol. and Biol, volume 42, pp 233-246.

Stegen JA. 1993. *Vegetation Communities Associated with the 100-Area and 200-Area Facilities on the Hanford Site*, WHC-SD-EN-TI-216, Revision 0, January 1994. Westinghouse Hanford Company, Richland, Washington, USA.

Stephens RD, Petreas M, and Hayward GH. 1995. *Biotransfer and Bioaccumulation of Dioxins and Furans from Soil: Chickens as a Model for Foraging Animals.* The Science of the Total Environment, volume 175, pp 253-273, 20 July 1995.

SCAPA. 1997a. *Emergency Response Planning Guideline (ErjPG) and TEELS for Chemical of Concern at the Savannah River Site: Rev. 12*, PEC-CAT-97-0044A, 9 July 1997. Subcommittee on Consequence Assessment and Protective Actions

SCAPA. 1997b. *SCAPA ERPG Working List, revised February 26, 1997.* World wide web address http://www.sep.bnl.gov/es/scapa.htm. Accessed on October 1, 1997. Subcommittee on Consequence Assessment and Protective Actions.

Suter II GW and Tsao CL. 1996. *Toxicological Benchmarks for Screening of Potential Contaminants of Concern for Effects on Aquatic Biota on Oak Ridge Reservation: 1996 Revision,* ES/ER/TM-96/R2. Lockheed Martin Energy Systems, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA.

Talmage SS, et al. (in press), "Nitroaromatic Munition Compounds: Environmental Effects and Screening Values", *Reviews of Environmental Contamination and Toxicology.*

Travis CC, and Arms AD. 1988. *Bioconcentration of Organics in Beef, Milk, and Vegetation.* Environmental Science Technology, volume 22, number 3, pp 271-274.

Travis CC, and White RK. 1988. *Interspecific scaling of toxicity data*, Risk Analysis, volume 8, number 1, pp 119-125.

USFS. 1994. *Ecological Subregions of the United States*, WO-WSA-5. U.S. Department of Agriculture, Forest Services, Washington, D.C, USA.

Weiss SG, and Mitchell RM. 1992. *A Synthesis of Ecological Data From the 100 Areas of the Hanford Site*, WHC-EP-0601, October 1992. Westinghouse Hanford Company, Richland, Washington, USA.

Wentsel RS, Checkai RT, LaPoint TW, Simini M, Ludwig D, and Brewer L. 1994. *Procedural Guidelines for Ecological Risk Assessments at US Army Sites*, volume 1, ERDEC-TR-221. Aberdeen Proving Ground, Maryland, USA.

Wentsel RS, LaPoint TW, Simini M, Checkai RT, Ludwig D, and Brewer L. 1996. *Tri-Service Procedural Guidelines for Ecological Risk Assessments.* Aberdeen Proving Ground, Maryland, USA.

WHC. 1992. *Biological Assessment for Rare and Endangered Plant Species Related to CERCLA Characterization Activities*, WHC-EP-0526, April 1992. Westinghouse Hanford Company, Richland, Washington, USA.

WHC. 1993. *100 Areas CERCLA Ecological Investigations*, WHC-EP-0620, September 1993. Westinghouse Hanford Company, Richland, Washington, USA.

WHC. 1994. *Threatened and Endangered Wildlife Species of the Hanford Site Related to CERCLA Characterization Activities*, WHC-EP-0513, June 1994. Westinghouse Hanford Company, Richland, Washington, USA.

WHO. 1997. *Meeting on the Derivation of Toxic Equivalency Factors (TEFs) for PCBs, PCDDs, PCDFs, and Other Dioxin-like Compounds for Human Health and Wildlife.* Institute for Environmental Medicine, Karolinska Institute. Stockholm, Sweden. June 15-18. Draft report, July 30. World Health Organization

WHO. 1998. *Assessment of the Health Risk of Dioxins: Re-*evaluation *of the Tolerable Daily Intake (TDI).* WHO Consultation, WHO European Centre for Environment and Health, International Programme on Chemical Safety, May 25-29. World Health Organization.

Wiemers KD, Lerchen ME, Miller M, and Meier K. 1998. *Regulatory Data Quality Objectives Supporting Tank Waste Remediation System Privatization Project*, PNNL-12040, Revision 0, December 1998. Pacific Northwest National Laboratory, Richland, Washington, USA.

Yelverton CS, and Quay TL. 1959. *Food habits of the Canada goose at Lake Mattamuskeet, North Carolina, USA*, North Carolina Wildlife Resources Commission.

Appendix A
Constituents of Potential Concern Lists

PDC
6-00

Table A-1. Organic Constituents of Potential Concern Retained by the Regulatory Data Quality Objective Process

CAS #	Constituent	CAS #	Constituent
50-32-8	Benzo(a)pyrene	106-93-4	Ethylene dibromide
53-70-3	Dibenz(a,h)anthracene	106-97-8	Butane
56-23-5	Carbon tetrachloride	106-99-0	1,3-Butadiene
57-14-7	1,1-Dimethylhydrazine	107-02-8	Acrolein
58-89-9	gamma-BHC (Lindane)	107-05-1	3-Chloropropene (Allyl chloride)
60-34-4	Methylhydrazine	107-06-2	1,2-Dichloroethane
60-57-1	Dieldrin	107-12-0	Propionitrile
62-75-9	N-Nitroso-N,N-dimethylamine	107-13-1	Acrylonitrile
64-17-5	Ethyl alcohol	107-87-9	2-Pentanone
64-18-6	Formic acid	108-10-1	4-Methyl-2-pentanone
64-19-7	Acetic acid	108-38-3	m-Xylene
67-56-1	Methyl alcohol	108-87-2	Methylcyclohexane
67-63-0	2-Propyl alcohol	108-88-3	Toluene
67-64-1	2-Propanone (Acetone)	108-90-7	Chlorobenzene
67-66-3	Chloroform	108-94-1	Cyclohexanone
71-23-8	n-Propyl alcohol	108-95-2	Phenol
71-36-3	n-Butyl alcohol	109-66-0	n-Pentane
71-43-2	Benzene	109-99-9	Tetrahydrofuran
71-55-6	1,1,1-Trichloroethane	110-12-3	5-Methyl-2-hexanone
72-20-8	Endrin	110-43-0	2-Heptanone
74-83-9	Bromomethane	110-54-3	n-Hexane
74-87-3	Chloromethane	110-82-7	Cyclohexane
75-00-3	Chloroethane	110-83-8	Cyclohexene
75-01-4	1-Chloroethene	110-86-1	Pyridine
75-05-8	Acetonitrile	111-65-9	n-Octane
75-09-2	Dichloromethane	111-84-2	n-Nonane
75-15-0	Carbon disulfide	118-74-1	Hexachlorobenzene
75-21-8	Oxirane	120-82-1	1,2,4-Trichlorobenzene
75-34-3	1,1-Dichloroethane	121-44-8	Triethylamine
75-35-4	1,1-Dichloroethene	122-39-4	N,N-Diphenylamine
75-43-4	Dichlorofluoromethane	123-19-3	4-Heptanone
75-45-6	Chlorodifluoromethane	123-38-6	n-Propionaldehyde
75-65-0	2-Methyl-2-propanol	123-86-4	Acetic acid n-butyl ester
75-69-4	Trichlorofluoromethane	123-91-1	1,4-Dioxane
75-71-8	Dichlorodifluoromethane	126-73-8	Tributyl phosphate
76-13-1	1,2,2-Trichloro-1,1,2-trifluoroethane	126-98-7	2-Methyl-2-propenenitrile
76-14-2	1,2-Dichloro-1,1,2,2-tetrafluoroethane	127-18-4	1,1,2,2-Tetrachloroethene
76-44-8	Heptachlor	128-37-0	2,6-Bis(tert-butyl)-4-methylphenol
78-87-5	1,2-Dichloropropane	141-78-6	Acetic acid ethyl ester
78-92-2	1-Methylpropyl alcohol	142-82-5	n-Heptane
78-93-3	2-Butanone	144-62-7	Oxalic acid
79-00-5	1,1,2-Trichloroethane	287-92-3	Cyclopentane
79-01-6	1,1,2-Trichloroethylene	309-00-2	Aldrin
79-10-7	2-Propenoic acid	319-84-6	alpha-BHC
79-34-5	1,1,2,2-Tetrachloroethane	319-85-7	beta-BHC
82-68-8	Pentachloronitrobenzene	465-73-6	Isodrin

Table A-1. Organic Constituents of Potential Concern Retained by the Regulatory Data Quality Objective Process

CAS #	Constituent	CAS #	Constituent
87-68-3	Hexachlorobutadiene	541-73-1	1,3-Dichlorobenzene
87-86-5	Pentachlorophenol	563-80-4	3-Methyl-2-butanone
88-85-7	2-sec-Butyl-4,6-dinitrophenol	591-78-6	2-Hexanone
88-89-1	Picric acid	624-83-9	Methyl isocyanate
92-52-4	1,1-Biphenyl	627-13-4	Nitric acid, propyl ester
95-47-6	o-Xylene	684-16-2	Hexafluoroacetone
95-50-1	1,2-Dichlorobenzene	1321-64-8	Pentachloronaphthalene
96-22-0	3-Pentanone	1335-87-1	Hexachloronaphthalene
98-86-2	Acetophenone	1335-88-2	Tetrachloronaphthalene
98-95-3	Nitrobenzene	1336-36-3	Polychlorinated biphenyls (PCBs)
100-00-5	p-Nitrochlorobenzene	2234-13-1	Octachloronaphthalene
100-25-4	1,4-Dinitrobenzene	3825-26-1	Ammonium perfluorooctanoate
100-41-4	Ethyl benzene	4170-30-3	2-Butenaldehyde (2-Butenal)
100-42-5	Styrene	8001-35-2	Toxaphene
106-35-4	3-Heptanone	10061-01-5	cis-1,3-Dichloropropene
106-42-3	p-Xylene	10061-02-6	trans-1,3-Dichloropropene
106-46-7	1,4-Dichlorobenzene		
PCBs			
31508-00-6	2,3',4,4',5- Pentachlorobiphenyl	69782-90-7	2,3,3',4,4',5'- Hexachlorobiphenyl
32598-13-3	3,3',4,4'-Tetrachlorobiphenyl	70362-50-4	3,4,4',5-Tetrachlorobiphenyl
32598-14-4	2,3,3',4,4'-Pentachlorobiphenyl	74472-37-0	2,3,4,4',5- Pentachlorobiphenyl
32774-16-6	3,3',4,4',5,5'- Hexachlorobiphenyl	no cas #	2,3,3',4,4',5,5'- Heptachlorobiphenyl
35065-29-3	2,2',3,4,4',5,5'- Heptachlorobiphenyl	no cas #	2',3,4,4',5- Pentachlorobiphenyl
35065-30-6	2,2',3,3',4,4',5- Heptachlorobiphenyl	no cas #	2,3',4,4',5,5'- Hexachlorobiphenyl
38380-08-4	2,3,3',4,4',5- Hexachlorobiphenyl	no cas #	3,3',4,4',5- Pentachlorobiphenyl

Table A-2. Inorganic Constituents of Potential Concern Retained by the Regulatory Data Quality Objective Process

CAS #	Metals	Constituent	CAS #	Ions	Constituent
7440-22-4	Ag	Silver	24959-67-9	Br	Bromide
7429-90-5	Al	Aluminum	16887-00-6	Cl	Chloride
7440-38-2	As	Arsenic	57-12-5	CN	Cyanide
7440-42-8	B	Boron	16984-48-8	F	Fluoride
7440-39-3	Ba	Barium	7553-56-2	I	Iodine
7440-41-7	Be	Beryllium	7664-41-7	NH4/NH3	Ammonia/Ammonium
7440-69-9	Bi	Bismuth	14797-65-0	NO2	Nitrite
7440-70-2	Ca	Calcium	14797-55-8	NO3	Nitrate
7440-43-9	Cd	Cadmium	14280-30-9	OH	Hydroxide
7440-48-4	Co	Cobalt	14265-44-2	PO4	Phosphate
18540-29-9	Cr	Chromium (and VI)	14808-79-8	SO4	Sulfate
7440-50-8	Cu	Copper	7440-36-0	Sb	Antimony
7439-89-6	Fe	Iron	7782-49-2	Se	Selenium
7439-97-6	Hg	Mercury	7440-21-3	Si	Silicon
7440-09-7	K	Potassium	7440-24-6	Sr	Strontium
7439-93-2	Li	Lithium	7440-31-5	Sn	Tin
7439-95-4	Mg	Magnesium	7440-25-7	Ta	Tantalum
7439-96-5	Mn	Manganese	7440-28-0	Tl	Thallium
7439-98-7	Mo	Molybdenum	7440-61-1	U	Uranium
7440-23-5	Na	Sodium	7440-62-2	V	Vanadium
7440-02-0	Ni	Nickel	7440-33-7	W	Tungsten
7723-14-0	P	Phosphorus	7440-65-5	Y	Yttrium
7439-92-1	Pb	Lead	7440-66-6	Zn	Zinc
7440-16-6	Rh	Rhodium	7440-67-7	Zr	Zirconium
63705-05-5	S	Total Sulfur			

Table A-3. Constituents of Potential Concern from the Environmental Protection Agency's Recommended Products of Incomplete Combustion List

CAS #	Constituent	CAS #	Constituent	CAS #	Constituent
			Organics		
50-00-0	Formaldehyde	99-35-4	1,3,5-Trinitrobenzene	218-01-9	Chrysene
51-28-5	2,4-Dinitrophenol	99-65-0	1,3-Dinitrobenzene	460-19-5	Cyanogen
51-79-6	Ethyl Carbamate (Urethane)	100-02-7	4-Nitrophenol	506-68-3	Cyanogen bromide
56-55-3	Benzo(a)anthracene	100-44-7	Benzyl chloride	506-77-4	Cyanogen chloride
57-24-9	Strychnine	100-52-7	Benzaldehyde	510-15-6	Chlorobenzilate
57-74-9	Chlordane	101-77-9	4,4-Methylenedianiline	528-29-0	o-Dinitrobenzene
60-11-7	Dimethyl aminoazobenzene	103-33-3	Azobenzene	532-27-4	2-Chloroacetophenone
62-50-0	Ethyl methanesulfonate	105-67-9	2,4-Dimethylphenol	534-52-1	4,6-Dinitro-o-cresol
62-53-3	Aniline	106-44-5	p-Cresol	540-73-8	1,2-Dimethylhydrazine
65-85-0	Benzoic acid	106-47-8	p-Chloroaniline	540-84-1	2,2,4-Trimethylpentane
67-72-1	Hexachloroethane	106-49-0	p-Toluidine	542-75-6	1,3-Dichloropropene
70-30-4	Hexachlorophene	106-51-4	Quinone	542-88-1	Dichloromethyl ether
72-43-5	Methoxychlor	106-89-8	Epichlorohydrin	584-84-9	2,4-Toluene diisocyanate
72-55-9	DDE	107-19-7	Propargyl alcohol	593-60-2	Bromoethene
74-95-3	Methylene bromide	107-21-1	Ethylene glycol	606-20-2	2,6-Dinitrotoluene
74-97-5	Bromochloromethane	107-98-2	Propylene gylcol monomethyl ether	608-93-5	Pentachlorobenzene
75-07-0	Acetaldehyde	108-05-4	Vinyl acetate	621-64-7	di-n-Propylnitrosamine
75-25-2	Bromoform	108-39-4	m-Cresol	630-20-6	1,1,1,2-Tetrachloroethane
75-27-4	Bromodichloromethane	108-60-1	Dichloroisopropyl ether	764-41-0	1,4-Dichloro-2-butene
75-29-6	2-Chloropropane	108-67-8	1,3,5-Trimethyl benzene	765-34-4	Glycidylaldehyde
75-44-5	Phosgene	109-77-3	Malononitrile	822-06-0	Hexamethylene-1,5-diisocyanate
76-01-7	Pentachloroethane	109-86-4	2-Methoxyethanol	823-40-5	Toluene-2,6-diamine
77-47-4	Hexachlorocyclopentadiene	110-80-5	2-Ethoxyethanol	924-16-3	N-Nitrosodi-n-butylamine
77-78-1	Dimethyl sulfate	111-15-9	Ethylene glycol monoethyl ether acetate	1120-71-4	1,3-Propane sultone
84-66-2	Diethyl phthalate	111-44-4	Bis(2-chloroethyl) ether	1634-04-4	Methyl tert-butyl ether
84-74-2	Dibutyl phthalate	111-76-2	Ethylene glycol monobutyl ether	1746-01-6	2,3,7,8-Tetrachlorodibenzo(p)dioxin
85-44-9	Phthalic anhydride	111-91-1	Bis(2-chloroethoxy)methane	3268-87-9	Octachlorodibenzo(p)dioxin
85-68-7	Butylbenzyl phthalate	117-81-7	Bis(2-ethylhexyl)phthalate (DEHP)	5385-75-1	Dibenzo(a,e)fluoranthene

Table A-3. Constituents of Potential Concern from the Environmental Protection Agency's Recommended Products of Incomplete Combustion List

CAS #	Constituent	CAS #	Constituent	CAS #	Constituent
88-06-2	2,4,6-Trichlorophenol	117-84-0	n-Dioctyl phthalate	19408-74-3	1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin
88-74-4	o-Nitroaniline	119-90-4	3,3'-Dimethoxybenzidine	23950-58-5	Pronamide
88-75-5	2-Nitrophenol	120-12-7	Anthracene	25013-15-4	Methyl styrene (mixed isomers)
90-04-0	o-Anisidine	120-83-2	2,4-Dichlorophenol	35822-46-9	1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin
91-20-3	Naphthalene	121-14-2	2,4-Dinitrotoluene	39001-02-0	Octachlorodibenzofuran
91-22-5	Quinoline	122-66-7	1,2-Diphenylhydrazine	39227-28-6	1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin
91-58-7	2-Chloronapthalene	123-33-1	Maleic hydrazide	40321-76-4	1,2,3,7,8-Pentachlorodibenzo(p)dioxin
91-94-1	3,3-Dichlorobenzidine	124-48-1	Chlorodibromomethane	41851-50-7	Chlorocyclopentadiene
94-59-7	Safrole	129-00-0	Pyrene	51207-31-9	2,3,7,8-Tetrachlorodibenzofuran
94-75-7	2,4-D and esters	131-11-3	Dimethylphthalate	55673-89-7	1,2,3,4,7,8,9-Heptachlorodibenzofuran
95-48-7	o-Cresol	131-89-5	2-Cyclohexyl-4,6-dinitrophenol	57117-31-4	2,3,4,7,8-Pentachlorodibenzofuran
95-53-4	o-Toluidine	132-64-9	Dibenzofuran	57117-41-6	1,2,3,7,8-Pentachlorodibenzofuran
95-57-8	2-Chlorophenol	133-06-2	Captan	57117-44-9	1,2,3,6,7,8-Hexachlorodibenzofuran
95-94-3	1,2,4,5-Tetrachlorobenzene	145-73-3	Endothall	57653-85-7	1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin
95-95-4	2,4,5-Trichlorophenol	156-60-5	trans-1,2-Dichloroethylene	60851-34-5	2,3,4,6,7,8-Hexachlorodibenzofuran
96-12-8	1,2-Dibromo-3-chloropropane	191-24-2	Benzo(g,h,i)perylene	67562-39-4	1,2,3,4,6,7,8-Heptachlorodibenzofuran
96-18-4	1,2,3-Trichloropropane	192-97-2	Benzo(e)pyrene	70648-26-9	1,2,3,4,7,8-Hexachlorodibenzofuran
96-45-7	Ethylene thiourea	193-39-5	Indeno(1,2,3-cd)pyrene	72918-21-9	1,2,3,7,8,9-Hexachlorodibenzofuran
97-63-2	Ethyl methacrylate	205-82-3	Benzo(j)fluoranthene	no CAS #	Dichloropentadiene
98-01-1	Furfural	205-99-2	Benzo(b)fluoranthene	no CAS#	Dibenzo(a,h)fluoranthene
98-07-7	Benzotrichloride	206-44-0	Fluoranthene		
98-82-8	Cumene	207-08-9	Benzo(k)fluoranthene		

Table A-4. Constituents of Potential Concern Identified During Bench-Scale Testing

CAS #	Constituent
74-88-4	Iodomethane
80-62-6	Methyl methacrylate
85-01-8	Phenanthrene
87-61-6	1,2,3-Trichlorobenzene
91-57-6	2-Methyl naphthalene
95-49-8	2-Chlorotoluene
95-63-6	1,2,4-Trimethyl benzene
98-06-6	tert-Butyl benzene
99-87-6	p-Cymene
100-51-6	Benzyl alcohol
103-65-1	n-Propyl benzene
104-51-8	n-Butyl benzene
106-43-4	4-Chlorotoluene
108-86-1	Bromobenzene
135-98-8	sec-Butyl benzene
156-59-2	cis-1,2-Dichloroethene

Table A-5. Organic Constituents of Potential Concern

Cas #	Constituent	Cas #	Constituent
Detected chemicals previously eliminated due to low toxicity		**Non-detected chemicals previously eliminated due to low toxicity (continued)**	
59-50-7	4-Chloro-3-methylphenol	75-99-0	2,2-Dichloropropionic acid
59-89-2	N-Nitrosomorpholine	76-03-9	Trichloroacetic acid
60-35-5	Acetamide	76-11-9	1,1,1,2-Tetrachloro-2,2-difluoroethane
74-99-7	Methylacetylene	76-12-0	1,1,2,2-Tetrachloro-1,2-difluoroethane
75-12-7	Formamide	76-15-3	Chloropentafluoroethane
75-50-3	Trimethylamine	88-72-2	2-Nitrotoluene
75-52-5	Nitromethane	92-93-3	4-Nitrobiphenyl
75-55-8	2-Methylaziridine	93-72-1	Silvex (2,4,5-TP)
78-83-1	2-Methylpropyl alcohol	93-76-5	2,4,5-T
79-09-4	Propionic acid	95-13-6	Indene
79-20-9	Methyl acetate	98-51-1	p-tert-Butyltoluene
83-32-9	Acenaphthene	100-21-0	p-Phthalic acid
86-73-7	Fluorene	101-55-3	4-Bromophenyl phenyl ether
96-69-5	Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	107-66-4	Dibutylphosphate
98-83-9	alpha-Methylstyrene	121-69-7	Dimethylaniline
101-84-8	Diphenyl ether	189-55-9	Dibenzo[a,i]pyrene
106-88-7	1,2-Epoxybutane	189-64-0	Dibenzo[a,h]pyrene
107-18-6	2-Propene-1-ol	191-30-0	Benzo[a,i]pyrene
107-31-3	Formic acid, methyl ester	192-65-4	Dibenzo[a,e]pyrene
108-03-2	1-Nitropropane	208-96-8	Acenaphthylene
108-20-3	Bis(isopropyl)ether	224-42-0	Dibenz[a,j]acridine
108-93-0	Cyclohexanol	226-36-8	Dibenz[a,h]acridine
110-62-3	n-Valeraldehyde	319-86-8	Delta-BHC
123-51-3	3-Methyl-1-butanol	540-59-0	1,2-Dichloroethylene
127-19-5	N,N-Dimethylacetamide	602-87-9	5-Nitroacenaphthene
141-79-7	4-Methyl-3-penten-2-one	603-34-9	Triphenylamine
Non-detected chemicals previously eliminated due to low toxicity		1321-65-9	Trichloronaphthalene
50-29-3	4,4-DDT	1582-09-8	Trifluralin
56-49-5	3-Methylcholanthrene	1836-75-5	Nitrofen
58-90-2	2,3,4,6-Tetrachlorophenol	3697-24-3	5-Methylchrysene
60-29-7	Ethyl ether	2385-85-5	Mirex
72-54-8	4,4-DDD	25551-13-7	Trimethyl benzene
75-61-6	Difluorodibromomethane	26140-60-3	Terphenyls
75-63-8	Trifluorobromomethane	27154-33-2	Trichlorofluoroethane

Table A-6. Radionuclide Constituents of Potential Concern

CAS #	Symbol	Constituent	CAS #	Symbol	Constituent
14952-40-0	^{227}Am	Actinium-227	14119-33-6	^{240}Pu	Plutonium-240
14596-10-2	^{241}Am	Americium-241	14119-32-5	^{241}Pu	Plutonium-241
14993-75-0	^{243}Am	Americium-243	13982-10-0	^{242}Pu	Plutonium-242
14234-35-6	^{125}Sb	Antimony-125	14331-85-2	^{231}Pa	Protactinium-231
13981-97-0	^{137m}Ba	Barium-137	13982-63-3	^{226}Ra	Radium-226
no CAS #	^{113m}Cd	Cadmium-113	15262-20-1	^{228}Ra	Radium-228
14762-75-5	^{14}C	Carbon-14	13967-48-1	^{106}Ru	Ruthenium-106
13967-70-9	^{134}Cs	Cesium-134	15715-94-3	^{151}Sm	Samarium-151
10045-97-3	^{137}Cs	Cesium-137	no CAS #	^{79}Se	Selenium-79
10198-40-0	^{60}Co	Cobalt-60	10098-97-2	^{90}Sr	Strontium-90
15510-73-3	^{242}Cm	Curium-242	14133-76-7	^{99}Tc	Technetium-99
15757-87-6	^{243}Cm	Curium-243	15594-54-4	^{229}Th	Thorium-229
13981-15-2	^{244}Cm	Curium-244	7440-29-1	^{232}Th	Thorium-232
14683-23-9	^{152}Eu	Europium-152	15832-50-5	^{126}Sn	Tin-126
15585-10-1	^{154}Eu	Europium-154	10028-17-8	^{3}H	Tritium
14391-16-3	^{155}Eu	Europium-155	14158-29-3	^{232}U	Uranium-232
15046-84-1	^{129}I	Iodine-129	13968-55-3	^{233}U	Uranium-233
13994-20-2	^{237}Np	Neptunium-237	13966-29-5	^{234}U	Uranium-234
14336-70-0	^{59}Ni	Nickel-59	15117-96-1	^{235}U	Uranium-235
13981-37-8	^{63}Ni	Nickel-63	13982-70-2	^{236}U	Uranium-236
7440-03-1	^{93m}Nb	Niobium-93	7440-61-1	^{238}U	Uranium-238
13981-16-3	^{238}Pu	Plutonium-238	10098-91-6	^{90}Y	Yttrium-90
15117-48-3	^{239}Pu	Plutonium-239	15751-77-6	^{93}Zr	Zirconium-93

Appendix B
Chemical-Specific Parameters and Supporting Equations for Human Health Risk Assessment

Appendix B1

Constituents of Potential Concern-Specific Parameters

Appendix B1
Constituents of Potential Concern-Specific Parameters

Database

A database will be developed for chemical-specific parameters that are used in the process of modeling concentrations for the screening level risk assessment. Values will be presented for the constituents of potential concern (COPCs), as identified in section 4 of this Work Plan, for the following parameters:

- MW: Molecular weight of COPC (g/mole)
- H: Henry's Law Constant (atm-m^3/mol)
- D_a: Diffusivity of COPC in air (cm^2/s)
- D_w: Diffusivity of COPC in water (cm^2/s)
- K_{ow}: Octanol/water partitioning coefficient (unitless)
- K_{oc}: Soil organic carbon-water partitioning coefficient (mL water/g soil)
- Kd_s: Soil-water partitioning coefficient (mL water/g soil)
- Kd_{sw}: Suspended sediment/surface water partitioning coefficient (L water/kg suspended sediment)
- Kd_{bs}: Bed sediment/sediment pore water partitioning coefficient (L COPC/kg water body)
- ksg: COPC soil loss due to biotic and abiotic degradation (year^{-1})
- $t_{1/2}$: Half time of COPC in soil (days)
- F_v: Fraction of COPC air concetration in vapor phase (unitless)
- RCF: Root concentration factor (ug COPC/g DW plant)/(ug COPC/ml soil water) - (unitless)
- $Br_{rootveg}$: Plant-soil bioconcentration factor for belowground produce (ug COPC/g DW plant)/(ugCOPC/ml soil) - (unitless)
- Br_{forage}: Plant-soil bioconcentration factor in forage and silage (ug COPC/g DW plant)/(ug COPC/ml soil) - (unitless)
- Br_{grain}: Plant-soil bioconcentration factor in grain (ug COPC/g DW plant)/(ug COPC/ml soil) - (unitless)
- Bv_{ag}: COPC air-to-plant biotransfer factor for aboveground produce ((ug COPC/g DW plant)/(ug COPC/ml air) - (unitless)

- Bv_{forage}: COPC air-to-plant biotransfer factor for forage and silage ((ug COPC/g DW plant)/(ug COPC/ml air) - (unitless)
- Ba_{milk}: Biotransfer factor for milk (mg COPC/kg FW tissue)/(mg COPC/day) OR (day/kg FW tissue)
- Ba_{beef}: Biotransfer factor for beef (mg COPC/kg FW tissue)/(mg COPC/day) OR (day/kg FW tissue)
- Ba_{pork}: Biotransfer factor for pork (mg COPC/kg FW tissue)/(mg COPC/day) OR (day/kg FW tissue)
- Ba_{egg}: Biotransfer factor for eggs (mg COPC/kg FW tissue)/(mg COPC/day) OR (day/kg FW tissue)
- $Ba_{chicken}$: Biotransfer factor for chicken (mg COPC/kg FW tissue)/(mg COPC/day) OR (day/kg FW tissue)
- BCF_{fish}: Bioconcentration factor in fish (L/kg FW) OR (unitless)
- BAF_{fish}: Bioaccumulati0n factor in fish (mg COPC/kg FW tissue)/(mg COPC/L total water column)
- $BSAF_{fish}$: Biota-sediment accumulation factor in fish (mg COPC/kg lipid tissue)/(mg COPC/kg sediment) - (unitless)

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Mol. Wt.		Henry's Constant			Diffusivity		
Constituent of Potential Concern	CAS Registry Number	MW (g/mol)	pH	H (Published Value)	H (atm*m^3/mol)	Source	D_a (cm^2/s)	Source	D_w (cm^2/s)
Aromatic Halogenated Hydrocarbons									
2,3,4,6-Tetrachlorophenol	58-90-2	231.9	NA	3.55E-01	3.51E-06	Properties a	2.17E-02	RAIS b	7.10E-06
4-Chloro-3-methylphenol	59-50-7	142.58	NA	2.53E-01	2.50E-06	Properties a	6.96E-02	EPA6 calc c	8.06E-06
Aromatic Nonhalogenated Hydrocarbons									
2-Nitrotoluene	88-72-2	137.13	NA	5.81E+00	5.75E-05	Properties a	4.75E-02	RAIS b	8.67E-06
4-Nitrobiphenyl	92-93-3	199.21	NA	NA	NA	NA	5.57E-02	EPA6 calc c	6.45E-06
Benzaldehyde	100-52-7	106.12	NA	4.18E-05	4.18E-05	EPA6 e	7.07E-02	EPA6 e	9.48E-06
Benzene	71-43-2	78.11	NA	5.49E-03	5.49E-03	EPA6 e	1.17E-01	EPA6 e	1.02E-05
Benzyl alcohol	100-51-6	108.13	NA	3.78E-07	3.78E-07	EPA6 e	6.89E-02	EPA6 e	9.38E-06
Ethyl Benzene	100-41-4	106.16	NA	7.73E-03	7.73E-03	EPA6 e	7.65E-02	EPA6 e	8.49E-06
m-Xylene	108-38-3	106.16	NA	9.26E-06	9.26E-06	EPA6 e	7.69E-02	EPA6 e	8.49E-06
o-Xylene	95-47-6	106.16	NA	6.73E-06	6.73E-06	EPA6 e	7.69E-02	EPA6 e	8.44E-06
p-Xylene	106-42-3	106.16	NA	6.89E-06	6.89E-06	EPA6 e	7.61E+02	EPA6 e	8.50E-06
Styrene	100-42-5	104.14	NA	3.33E-03	3.33E-03	EPA6 e	7.73E-02	EPA6 e	8.77E-06
Toluene	108-88-3	92.13	NA	6.13E-03	6.13E-03	EPA6 e	9.72E-02	EPA6 e	9.23E-06
Non-aromatic Nonhalogenated Hydrocarbons									
1,2-Epoxybutane	106-88-7	72.12	NA	1.89E-02	4.61E-04	RAIS b	1.34E-01	RAIS b	1.03E-05
1,3-Butadiene	106-99-0	54.091	NA	7.46E+00	7.39E-02	Henry's f	1.33E-01	EPA6 calc c	1.54E-05
1,4-Dioxane	123-91-1	88.1	NA	4.89E-06	4.89E-06	EPA6 e	9.20E-02	EPA6 e	1.05E-05
1-Methylpropyl alcohol	78-92-2	74.122	NA	1.03E+00	1.02E-05	Properties a	1.08E-01	EPA6 calc c	1.25E-05
1-Nitropropane	108-03-2	89.09	NA	NA	NA	NA	9.53E-02	EPA6 calc c	1.10E-05
2,2,4-Trimethlylpentane	540-84-1	114.23	NA	3.05E+05	3.02E+00	Properties a	8.07E-02	EPA6 calc c	9.34E-06
2-Butanone	78-93-3	72.1	NA	3.61E-05	3.61E-05	EPA6 e	1.35E-01	EPA6 e	1.03E-05
2-Butenaldehyde (2-Butenal)	4170-30-3	70.091	NA	NA	NA	NA	1.12E-01	EPA6 calc c	1.29E-05
2-Ethoxyethanol	110-80-5	90.12	NA	5.13E-02	5.13E-02	Organics g	9.32E-02	RAIS b	9.76E-06
2-Heptanone	110-43-0	114.19	NA	1.46E+01	1.45E-04	Properties a	8.07E-02	EPA6 calc c	9.35E-06
2-Hexanone	591-78-6	100.16	NA	9.70E+00	9.60E-05	Properties a	8.81E-02	EPA6 calc c	1.02E-05
2-Methoxyethanol	109-86-4	76	NA	3.38E-05	8.24E-07	RAIS b	1.29E-01	RAIS b	9.57E-06
2-Methyl-2-propanol	75-65-0	74.122	NA	1.21E+00	1.20E-05	Properties a	1.08E-01	EPA6 calc c	1.25E-05

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Mol. Wt.		Henry's Constant			Diffusivity		
Constituent of Potential Concern	CAS Registry Number	MW (g/mol)	pH	H (Published Value)	H (atm*m^3/mol)	Source	D_a (cm^2/s)	Source	D_w (cm^2/s)
2-Methyl-2-propenenitrile	126-98-7	67.09	NA	2.39E-04	2.39E-04	EPA6 e	1.15E-01	EPA6 e	1.33E-05
2-Methylaziridine	75-55-8	57.1	NA	NA	NA	NA	1.28E-01	EPA6 calc c	1.48E-05
2-Methylpropyl alcohol	78-83-1	74.12	NA	1.21E+00	1.20E-05	Properties a	8.60E-02	RAIS b	9.30E-06
2-Pentanone	107-87-9	86.133	NA	6.52E+00	6.46E-05	Properties a	9.74E-02	EPA6 calc c	1.13E-05
2-Propanone (Acetone)	67-64-1	58.08	NA	2.88E-05	2.88E-05	EPA6 e	1.87E-01	EPA6 e	1.15E-05
2-Propene-1-ol	107-18-6	58.08	NA	5.06E-01	5.01E-06	Properties a	1.14E-01	RAIS b	1.14E-05
2-Propyl alcohol	67-63-0	60.096	NA	8.20E-01	8.12E-06	Properties a	1.24E-01	EPA6 calc c	1.43E-05
3-Heptanone	106-35-4	114.19	NA	NA	NA	NA	8.07E-02	EPA6 calc c	9.35E-06
3-Methyl-1-butanol	123-51-3	88.15	NA	NA	NA	NA	9.59E-02	EPA6 calc c	1.11E-05
3-Methyl-2-butanone	563-80-4	86.133	NA	NA	NA	NA	9.74E-02	EPA6 calc c	1.13E-05
3-Pentanone	96-22-0	86.133	NA	8.13E+00	8.05E-05	Properties a	9.74E-02	EPA6 calc c	1.13E-05
4-Heptanone	123-19-3	114.19	NA	NA	NA	NA	8.07E-02	EPA6 calc c	9.35E-06
4-Methyl-2-pentanone	108-10-1	100.16	NA	1.25E-04	1.25E-04	EPA6 e	8.59E-02	EPA6 e	8.36E-06
4-Methyl-3-penten-2-one	141-79-7	98.14	NA	NA	NA	NA	8.93E-02	EPA6 calc c	1.03E-05
5-Methyl-2-hexanone	110-12-3	114.19	NA	NA	NA	NA	8.07E-02	EPA6 calc c	9.35E-06
Acetaldehyde	75-07-0	44.05	NA	7.89E-05	7.89E-05	Chemfate i	2.72E-01	EPA6 e	1.33E-05
Acetamide	60-35-5	59.07	NA	3.53E-04	3.50E-09	Properties a	1.25E-01	EPA6 calc c	1.45E-05
Acetic acid	64-19-7	60.052	NA	3.05E-02	3.02E-07	Properties a	1.24E-01	EPA6 calc c	1.43E-05
Acetic acid ethyl ester	141-78-6	88.1	NA	1.36E+01	1.35E-04	Properties a	7.32E-02	RAIS b	9.66E-06
Acetic acid n-butyl ester	123-86-4	116.16	NA	3.34E+01	3.31E-04	Properties a	7.98E-02	EPA6 calc c	9.24E-06
Acetonitrile	75-05-8	41.05	NA	3.79E+01	3.79E+01	EPA6 e	3.14E-01	EPA6 e	1.40E-05
Acrolein	107-02-8	56.06	NA	9.34E-05	9.34E-05	EPA6 e	1.92E-01	EPA6 e	1.22E-05
Acrylonitrile	107-13-1	53.06	NA	9.90E-05	9.90E-05	EPA6 e	2.11E-01	EPA6 e	1.23E-05
Bis(isopropyl)ether	108-20-3	102.18	NA	NA	NA	NA	8.69E-02	EPA6 calc c	1.01E-05
Butane	106-97-8	58.123	NA	9.59E+01	9.50E-01	Henry's f	1.27E-01	EPA6 calc c	1.47E-05
Carbon disulfide	75-15-0	76.14	NA	1.27E-02	1.27E-02	EPA6 e	1.04E-01	EPA6 e	1.29E-05
Cyanogen	460-19-5	52.04	NA	2.03E-01	4.96E-03	RAIS b	2.04E-01	RAIS b	1.37E-05
Cyclohexane	110-82-7	84.161	NA	1.97E+04	1.95E-01	Properties a	9.89E-02	EPA6 calc c	1.15E-05
Cyclohexanone	108-94-1	98.2	NA	1.22E+00	1.20E-05	Properties a	7.84E-02	RAIS b	8.62E-06

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Mol. Wt.		Henry's Constant			Diffusivity		
Constituent of Potential Concern	CAS Registry Number	MW (g/mol)	pH	H (Published Value)	H (atm*m^3/mol)	Source	D_a (cm^2/s)	Source	D_w (cm^2/s)
Cyclohexene	110-83-8	82.145	NA	4.02E+03	3.98E-02	Properties a	1.01E-01	EPA6 calc c	1.16E-05
Cyclopentane	287-92-3	70.134	NA	1.88E+04	1.86E-01	Properties a	1.12E-01	EPA6 calc c	1.29E-05
Ethyl alcohol	64-17-5	46.069	NA	6.29E-06	6.29E-06	Organics g	1.48E-01	EPA6 calc c	1.71E-05
Ethyl ether	60-29-7	74.12	NA	1.30E+02	1.29E-03	Properties a	7.40E-02	RAIS b	9.30E-06
Ethyl methacrylate	97-63-2	114.14	NA	1.38E-04	1.38E-04	EPA6 e	8.07E-02	EPA6 e	9.35E-06
Formaldehyde	50-00-0	30.03	NA	2.78E-04	2.78E-04	EPA6 e	5.00E-01	EPA6 e	1.74E-05
Formamide	75-12-7	45.04	NA	NA	NA	NA	1.50E-01	EPA6 calc c	1.74E-05
Formic acid	64-18-6	46.03	NA	2.49E-06	2.49E-06	EPA6 e	2.22E-01	EPA6 e	1.71E-05
Formic acid, methyl ester	107-31-3	60.05	NA	NA	NA	NA	1.24E-01	EPA6 calc c	1.43E-05
Glycidylaldehyde	765-34-4	72.07	NA	2.10E-05	5.11E-07	RAIS b	1.10E-01	EPA6 calc c	1.27E-05
Methyl acetate	79-20-9	74.08	NA	8.80E+00	8.71E-05	Properties a	1.04E-01	RAIS b	1.00E-05
Methyl alcohol	67-56-1	32.04	NA	1.44E-04	1.44E-04	EPA6 e	4.58E-01	EPA6 e	1.64E-05
Methyl isocyanate	624-83-9	57.052	NA	NA	NA	NA	1.28E-01	EPA6 calc c	1.48E-05
Methyl methacrylate	80-62-6	100.13	NA	3.21E+01	3.18E-04	Properties a	7.70E-02	RAIS b	8.60E-06
Methyl tert-butyl ether	1634-04-4	88.15	NA	5.95E+01	5.89E-04	Properties a	1.02E-01	RAIS b	1.05E-05
Methylacetylene	74-99-7	40.07	NA	NA	NA	NA	1.62E-01	EPA6 calc c	1.88E-05
Methylcyclohexane	108-87-2	98.188	NA	4.41E+04	4.36E-01	Properties a	8.93E-02	EPA6 calc c	1.03E-05
N,N-Dimethylacetamide	127-19-5	87.12	NA	NA	NA	NA	9.67E-02	EPA6 calc c	1.12E-05
n-Butyl alcohol	71-36-3	74.1	NA	8.60E-01	8.51E-06	Properties a	8.00E-02	RAIS b	9.30E-06
n-Heptane	142-82-5	100.2	NA	2.30E+05	2.28E+00	Properties a	8.81E-02	EPA6 calc c	1.02E-05
n-Hexane	110-54-3	86.177	NA	1.84E+05	1.82E+00	Properties a	9.74E-02	EPA6 calc c	1.13E-05
Nitromethane	75-52-5	61.04	NA	NA	NA	NA	1.23E-01	EPA6 calc c	1.42E-05
n-Nonane	111-84-2	128.26	NA	5.00E+05	4.95E+00	Properties a	7.47E-02	EPA6 calc c	8.65E-06
n-Octane	111-65-9	114.23	NA	3.27E+05	3.24E+00	Properties a	8.07E-02	EPA6 calc c	9.34E-06
n-Pentane	109-66-0	72.15	NA	1.27E+05	1.26E+00	Properties a	1.10E-01	EPA6 calc c	1.27E-05
n-Propionaldehyde	123-38-6	58.08	NA	7.49E+00	7.41E-05	Properties a	1.27E-01	EPA6 calc c	1.47E-05
n-Propyl alcohol	71-23-8	60.096	NA	NA	NA	NA	1.24E-01	EPA6 calc c	1.43E-05
n-Valeraldehyde	110-62-3	86.13	NA	NA	NA	NA	9.74E-02	EPA6 calc c	1.13E-05
Oxirane	75-21-8	44.05	NA	1.67E-04	1.67E-04	EPA6 e	2.71E-01	EPA6 e	1.44E-05

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Mol. Wt.		Henry's Constant			Diffusivity		
Constituent of Potential Concern	CAS Registry Number	MW (g/mol)	pH	H (Published Value)	H (atm*m³/mol)	Source	D_a (cm²/s)	Source	D_w (cm²/s)
p-Cymene	99-87-6	134.22	NA	8.00E-01	7.92E-03	Henry's f	7.25E-02	EPA6 calc c	8.39E-06
Phosgene	75-44-5	98.916	NA	NA	NA	NA	8.88E-02	EPA6 calc c	1.03E-05
Propargyl alcohol	107-19-7	56.06	NA	3.51E-04	8.56E-06	RAIS b	1.81E-01	RAIS b	1.33E-05
Propionic acid	79-09-4	74.08	NA	4.50E-02	4.46E-07	Properties a	1.08E-01	EPA6 calc c	1.25E-05
Propionitrile	107-12-0	55.079	NA	3.75E+00	3.71E-05	Properties a	1.31E-01	EPA6 calc c	1.52E-05
Propylene glycol monomethyl ether	107-98-2	90	NA	4.26E-05	1.04E-06	RAIS b	7.17E-02	RAIS b	9.73E-06
p-tert-Butyltoluene	98-51-1	148.25	NA	NA	NA	NA	6.78E-02	EPA6 calc c	7.85E-06
Triethylamine	121-44-8	101.19	NA	NA	NA	NA	8.81E-02	RAIS b	7.88E-06
Trimethylamine	75-50-3	59.11	NA	1.05E+01	1.04E-04	Properties a	1.25E-01	EPA6 calc c	1.45E-05
Vinyl acetate	108-05-4	86.09	NA	5.50E-04	5.50E-04	EPA6 e	9.94E-02	EPA6 e	1.00E-05
Non-aromatic Halogenated Hydrocarbons									
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	203.83	NA	NA	NA	NA	5.49E-02	EPA6 calc c	6.35E-06
1,1,1,2-Tetrachloroethane	630-20-6	167.85	NA	2.44E-03	2.44E-03	EPA6 e	3.15E-02	EPA6 e	9.30E-06
1,1,1-Trichloroethane	71-55-6	133.42	NA	1.86E-02	1.86E-02	EPA6 e	4.66E-02	EPA6 e	9.56E-06
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	203.83	NA	9.87E+03	9.77E-02	Properties a	5.49E-02	EPA6 calc c	6.35E-06
1,1,2,2-Tetrachloroethane	79-34-5	167.86	NA	3.72E-04	3.72E-04	EPA6 e	3.16E-02	EPA6 e	9.26E-06
1,1,2,2-Tetrachloroethene	127-18-4	165.85	NA	1.73E-02	1.73E-02	EPA6 e	7.20E-02	EPA6 e	8.20E-06
1,1,2-Trichloro-1,2,2-trifluoroethane	76-13-1	187.38	NA	5.33E+04	5.28E-01	Properties a	2.88E-02	RAIS b	8.07E-06
1,1,2-Trichloroethane	79-00-5	133.42	NA	1.00E-03	1.00E-03	EPA6 e	4.51E-02	EPA6 e	1.00E-05
1,1,2-Trichloroethylene	79-01-6	131.4	NA	1.06E-02	1.06E-02	EPA6 e	4.65E-02	EPA6 e	9.94E-06
1,1-Dichloroethane	75-34-3	98.97	NA	5.75E-03	5.75E-03	EPA6 e	7.42E-02	EPA6 e	1.05E-05
1,1-Dichloroethene	75-35-4	96.95	NA	2.55E-02	2.55E-02	EPA6 e	7.53E-02	EPA6 e	1.09E-05
1,2,3-Trichloropropane	96-18-4	147.43	NA	3.80E-04	3.80E-04	EPA6 e	3.99E-02	EPA6 e	9.24E-06
1,2-dibromo-3-chloropropane	96-12-8	236.36	NA	1.97E-04	1.97E-04	EPA6 e	1.79E-02	EPA6 e	8.79E-06
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	170.92	NA	NA	NA	NA	6.17E-02	EPA6 calc c	7.14E-06
1,2-Dichloroethane	107-06-2	98.96	NA	1.27E-03	1.27E-03	EPA6 e	7.19E-02	EPA6 e	1.10E-05
1,2-Dichloroethylene	540-59-0	96.94	NA	7.61E-01	7.53E-03	Henry's f	9.00E-02	EPA6 calc c	1.04E-05
1,2-Dichloropropane	78-87-5	112.99	NA	2.81E-03	2.81E-03	EPA6 e	6.21E-02	EPA6 e	9.71E-06
1,3-Dichloropropene	542-75-6	110.98	NA	2.94E-03	2.94E-03	EPA6 e	6.26E-02	EPA6 e	1.00E-05

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Mol. Wt.		Henry's Constant			Diffusivity		
Constituent of Potential Concern	CAS Registry Number	MW (g/mol)	pH	H (Published Value)	H (atm*m^3/mol)	Source	D_a (cm^2/s)	Source	D_w (cm^2/s)
1,4-Dichloro-2-butene	764-41-0	125	NA	1.06E-02	2.59E-04	RAIS b	7.60E-02	EPA6 calc c	8.80E-06
1-Chloroethene	75-01-4	62.5	NA	3.15E-01	3.15E-01	EPA6 e	1.58E-01	EPA6 e	1.19E-05
2,2-Dichloropropionic acid	75-99-0	142.97	NA	6.43E-08	6.43E-08	Chemfate i	6.95E-02	EPA6 calc c	8.05E-06
2-Chloropropane	75-29-6	78.541	NA	1.65E+03	1.64E-02	Properties a	1.04E-01	EPA6 calc c	1.20E-05
3-Chloropropene (allyl chloride)	107-05-1	76.526	NA	1.10E+00	1.09E-02	Henry's f	1.05E-01	EPA6 calc c	1.22E-05
Bromochloromethane	74-97-5	129.39	NA	1.06E+03	2.59E+01	RAIS b	4.74E-02	RAIS b	1.00E-05
Bromodichloromethane	75-27-4	163.83	NA	3.17E-03	3.17E-03	EPA6 e	2.98E-02	EPA6 e	1.06E-05
Bromoethene	593-60-2	106.95	NA	2.76E-01	6.74E-03	RAIS b	8.43E-02	EPA6 calc c	9.76E-06
Bromoform	75-25-2	252.77	NA	6.16E-04	6.16E-04	EPA6 e	1.41E-02	EPA6 e	1.03E-05
Bromomethane	74-83-9	94.95	NA	1.41E-02	1.41E-02	EPA6 e	7.28E-02	EPA6 e	1.21E-05
Carbon tetrachloride	56-23-5	153.84	NA	2.87E-02	2.87E-02	EPA6 e	3.56E-02	EPA6 e	9.77E-06
Chlordibromomethane	124-48-1	208.3	NA	1.21E-03	1.21E-03	EPA6 e	1.96E-02	EPA6 e	1.05E-05
Chlorodifluoromethane	75-45-6	86.47	NA	1.68E-01	1.68E-01	EPA6 e	9.72E-02	EPA6 e	1.13E-05
Chloroethane	75-00-3	64.52	NA	1.80E+00	1.80E+00	EPA6 e	1.27E-01	EPA6 e	1.53E-06
Chloroform	67-66-3	119.39	NA	4.03E-03	4.03E-03	EPA6 e	5.17E-02	EPA6 e	1.09E-05
Chloromethane	74-87-3	50.49	NA	4.52E-02	4.52E-02	EPA6 e	2.13E-01	EPA6 e	1.39E-05
Chloropentafluoroethane	76-15-3	154.47	NA	NA	NA	NA	6.60E-02	EPA6 calc c	7.64E-06
cis-1,2-Dichloroethene	156-59-2	96.94	NA	4.51E-03	4.51E-03	EPA6 e	7.36E-02	EPA6 e	1.13E-05
cis-1,3-Dichloropropene	10061-01-5	110.97	NA	NA	NA	NA	8.23E-02	EPA6 calc c	9.53E-06
Cyanogen bromide	506-68-3	105.92	NA	NA	NA	NA	8.49E-02	EPA6 calc c	9.83E-06
Cyanogen chloride	506-77-4	61.471	NA	NA	NA	NA	1.22E-01	EPA6 calc c	1.41E-05
Dichlorodifluoromethane	75-71-8	120.92	NA	2.58E+00	2.58E+00	EPA6 e	7.77E-02	EPA6 e	9.00E-06
Dichlorofluoromethane	75-43-4	102.92	NA	NA	NA	NA	8.65E-02	EPA6 calc c	1.00E-05
Dichloromethane	75-09-2	84.94	NA	2.38E-03	2.38E-03	EPA6 e	8.69E-02	EPA6 e	1.25E-05
Difluorodibromomethane	75-61-6	209.82	NA	NA	NA	NA	5.38E-02	EPA6 calc c	6.23E-06
Hexafluoroacetone	684-16-2	166.02	NA	NA	NA	NA	6.29E-02	EPA6 calc c	7.28E-06
Iodomethane	74-88-4	141.939	NA	5.55E+02	5.49E-03	Properties a	6.98E-02	EPA6 calc c	8.09E-06
Methylene bromide	74-95-3	173.86	NA	2.64E-02	2.64E-02	EPA6 e	6.10E-02	EPA6 e	7.06E-06
Pentachloroethane	76-01-7	202.29	NA	2.48E+02	2.45E-03	Properties a	5.51E-02	EPA6 calc c	6.38E-06

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Mol. Wt.		Henry's Constant			Diffusivity		
Constituent of Potential Concern	CAS Registry Number	MW (g/mol)	pH	H (Published Value)	H (atm*m^3/mol)	Source	D_a (cm^2/s)	Source	D_w (cm^2/s)
trans-1,2-Dichloroethene	156-60-5	96.95	NA	7.44E-03	7.44E-03	EPA6 e	8.16E-02	EPA6 e	9.75E-06
trans-1,3-Dichloropropene	10061-02-6	110.97	NA	NA	NA	NA	8.23E-02	EPA6 calc c	9.53E-06
Trichloroacetic acid	76-03-9	163.39	NA	1.80E-07	1.80E-07	EPA6 calc c	6.36E-02	EPA6 calc c	7.36E-06
Trichlorofluoroethane	27154-33-2	151.4	NA	NA	NA	NA	6.69E-02	EPA6 calc c	7.74E-06
Trichlorofluoromethane	75-69-4	137.38	NA	1.37E-01	1.37E-01	EPA6 e	4.27E-02	EPA6 e	1.00E-05
Trifluorobromomethane	75-63-8	148.91	NA	NA	NA	NA	6.76E-02	EPA6 calc c	7.83E-06
Dioxin and Furan Compounds									
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	35822-46-9	425.31	NA	7.50E-06	7.50E-06	EPA6 e	1.11E-02	EPA6 e	3.89E-06
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	409.31	NA	5.30E-05	5.30E-05	EPA6 e	1.55E-02	EPA6 e	3.99E-06
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	409.31	NA	5.30E-05	5.30E-05	EPA6 e	1.55E-02	EPA6 e	3.99E-06
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	39227-28-6	390.87	NA	1.20E-05	1.20E-05	EPA6 e	1.15E-02	EPA6 e	4.12E-06
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	374.87	NA	1.40E-05	1.40E-05	EPA6 e	1.62E-02	EPA6 e	4.23E-06
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	57653-85-7	390.87	NA	1.20E-05	1.20E-05	EPA6 e	1.15E-02	EPA6 e	4.12E-06
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	374.87	NA	6.10E-06	6.10E-06	EPA6 e	1.62E-02	EPA6 e	4.23E-06
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	19408-74-3	390.87	NA	1.20E-05	1.20E-05	EPA6 e	1.15E-02	EPA6 e	4.12E-06
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	374.87	NA	1.00E-05	1.00E-05	EPA6 e	1.62E-02	EPA6 e	4.23E-06
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	40321-76-4	356.42	NA	2.60E-06	2.60E-06	EPA6 e	1.21E-02	EPA6 e	4.38E-06
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	340.42	NA	6.20E-06	6.20E-06	EPA6 e	1.70E-02	EPA6 e	4.51E-06
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	374.87	NA	1.00E-05	1.00E-05	EPA6 e	1.62E-02	EPA6 e	4.23E-06
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	340.42	NA	6.20E-06	6.20E-06	EPA6 e	1.70E-02	EPA6 e	4.51E-06
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1746-01-6	321.98	NA	1.60E-05	1.60E-05	EPA6 e	1.27E-02	EPA6 e	6.81E-06
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	305.98	NA	8.60E-06	8.60E-06	EPA6 e	1.79E-02	EPA6 e	4.85E-06
Dibenzofuran	132-64-9	168.19	NA	1.30E-05	1.30E-05	EPA9 h	6.24E-02	EPA6 calc c	7.22E-06
Octachlorodibenzo(p)dioxin	3268-87-9	459.75	NA	7.00E-09	7.00E-09	EPA6 e	1.06E-02	EPA6 e	3.69E-07
Octachlorodibenzofuran	39001-02-0	444.76	NA	1.90E-06	1.90E-06	EPA6 e	1.48E-02	EPA6 e	3.78E-06
PCBs									
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	395.33	NA	1.93E+01	1.91E-04	Properties a	3.53E-02	EPA6 calc c	4.08E-06
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	395.33	NA	3.04E+01	3.01E-04	Properties a	3.53E-02	EPA6 calc c	4.08E-06
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	360.88	NA	NA	NA	NA	3.75E-02	EPA6 calc c	4.34E-06

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Mol. Wt.		Henry's Constant			Diffusivity		
Constituent of Potential Concern	CAS Registry Number	MW (g/mol)	pH	H (Published Value)	H (atm*m^3/mol)	Source	D_a (cm^2/s)	Source	D_w (cm^2/s)
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	395.33	NA	NA	NA	NA	3.53E-02	EPA6 calc c	4.08E-06
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	360.88	NA	NA	NA	NA	3.75E-02	EPA6 calc c	4.34E-06
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	326.44	NA	NA	NA	NA	4.01E-02	EPA6 calc c	4.64E-06
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	360.88	NA	NA	NA	NA	3.75E-02	EPA6 calc c	4.34E-06
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	326.44	NA	NA	NA	NA	4.01E-02	EPA6 calc c	4.64E-06
2',3,4,4',5-Pentachlorobiphenyl	no cas #	326.44	NA	NA	NA	NA	4.01E-02	EPA6 calc c	4.64E-06
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	326.44	NA	NA	NA	NA	4.01E-02	EPA6 calc c	4.64E-06
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	360.88	NA	5.98E+00	5.92E-05	Properties a	3.75E-02	EPA6 calc c	4.34E-06
3,3',4,4',5-Pentachlorobiphenyl	no cas #	326.44	NA	NA	NA	NA	4.01E-02	EPA6 calc c	4.64E-06
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	291.99	NA	NA	NA	NA	4.32E-02	EPA6 calc c	5.00E-06
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	291.99	NA	NA	NA		4.32E-02	EPA6 calc c	5.00E-06
Polychlorinated biphenyls (PCBs)**	1336-36-3	NA	NA	3.54E-02	8.64E-04	RAIS b	1.75E-02	RAIS b	8.00E-06
Phthalates									
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	390.54	NA	1.10E-08	1.10E-08	EPA6 e	1.32E-02	EPA6 e	4.22E-06
Butylbenzyl phthalate	85-68-7	312.39	NA	1.91E-06	1.91E-06	EPA6 e	1.65E-02	EPA6 e	5.17E-06
Dibutyl phthalate	84-74-2	278.34	NA	1.43E-06	1.43E-06	EPA6 e	4.38E-02	EPA6 e	7.86E-06
Diethyl phthalate	84-66-2	222.24	NA	5.48E-07	5.48E-07	EPA6 e	2.56E-02	EPA6 e	6.35E-06
Dimethyl phthalate	131-11-3	194.19	NA	1.01E-07	1.01E-07	EPA6 e	2.96E-02	EPA6 e	7.13E-06
n-Dioctyl phthalate	117-84-0	390.56	NA	7.68E-07	7.68E-07	EPA6 e	1.32E-02	EPA6 e	4.20E-06
Light Polycyclic Aromatic Hydrocarbons									
2-Chloronaphthalene	91-58-7	162.61	NA	1.43E-04	1.43E-04	EPA6 e	3.64E-02	EPA6 e	8.24E-06
2-Methyl naphthalene	91-57-6	142.2	NA	4.05E+01	4.01E-04	Properties a	6.97E-02	EPA6 calc c	8.08E-06
5-Nitroacenaphthene	602-87-9	199.21	NA	NA	NA	NA	5.57E-02	EPA6 calc c	6.45E-06
Acenaphthene	83-32-9	154.21	NA	2.00E-04	2.00E-04	EPA6 e	4.21E-02	EPA6 e	7.19E-06
Acenaphthylene	208-96-8	152.21	NA	1.19E+01	1.18E-04	Properties a	4.39E-02	RAIS b	7.53E-06
Anthracene	120-12-7	178.22	NA	1.11E-04	1.11E-04	EPA6 e	3.24E-02	EPA6 e	7.74E-06
Fluorene	86-73-7	166.22	NA	8.50E+00	8.42E-05	Properties a	3.63E-02	RAIS b	7.88E-06
Indene	95-13-6	116.16	NA	NA	NA	NA	7.98E-02	EPA6 calc c	9.24E-06
Napthalene	91-20-3	128.16	NA	4.82E-04	4.82E-04	EPA6 e	5.26E-02	EPA6 e	8.92E-06

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Mol. Wt.		Henry's Constant			Diffusivity		
Constituent of Potential Concern	CAS Registry Number	MW (g/mol)	pH	H (Published Value)	H (atm*m^3/mol)	Source	D_a (cm^2/s)	Source	D_w (cm^2/s)
Phenanthrene	85-01-8	178.22	NA	1.88E-01	1.88E-01	EPA6 e	3.33E-02	EPA6 e	7.47E-06
Pyrene	129-00-0	202.24	NA	1.14E-08	1.14E-08	EPA6 e	2.72E-02	EPA6 e	7.14E-06
Heavy Polycyclic Aromatic Hydrocarbons									
3-Methylcholanthrene	56-49-5	268.36	NA	1.24E-07	1.24E-07	EPA6 calc c	4.57E-02	EPA6 calc c	5.29E-06
5-Methylchrysene	3697-24-3	242.32	NA	NA	NA	NA	4.89E-02	EPA6 calc c	5.66E-06
Benzo(a)anthracene	56-55-3	228.28	NA	3.62E-06	3.62E-06	EPA6 e	2.47E-02	EPA6 e	6.21E-06
Benzo(a)pyrene	50-32-8	252.3	NA	8.36E-07	8.36E-07	EPA6 e	2.18E-02	EPA6 e	5.85E-06
Benzo(b)fluoranthene	205-99-2	252.32	NA	6.18E-06	6.18E-06	EPA6 e	2.28E-02	EPA6 e	5.49E-06
Benzo(e)pyrene	192-97-2	252.31	NA	4.62E-07	4.62E-07	EPA6 calc c	4.76E-02	EPA6 calc c	5.51E-06
Benzo(g,h,i)perylene	191-24-2	276.34	NA	7.09E-01	7.02E-06	Properties a	4.48E-02	EPA6 calc c	5.19E-06
Benzo(j)fluoranthene	205-82-3	252.31	NA	NA	NA	NA	4.76E-02	EPA6 calc c	5.51E-06
Benzo(k)fluoranthene	207-08-9	252.32	NA	4.15E-07	4.15E-07	EPA6 e	2.28E-02	EPA6 e	5.49E-06
Benzo[a,i]pyrene	191-30-0	302.37	NA	NA	NA	NA	4.22E-02	EPA6 calc c	4.88E-06
Chrysene	218-01-9	228.28	NA	1.21E-06	1.21E-06	EPA6 e	2.48E-02	EPA6 e	6.21E-06
Dibenz(a,h)anthracene	53-70-3	278.33	NA	1.12E-08	1.12E-08	EPA6 e	1.80E-02	EPA6 e	6.01E-06
Dibenz[a,h]acridine	226-36-8	279.34	NA	NA	NA	NA	4.45E-02	EPA6 calc c	5.15E-06
Dibenz[a,j]acridine	224-42-0	279.34	NA	NA	NA	NA	4.45E-02	EPA6 calc c	5.15E-06
Dibenzo(a,e)fluoranthene	5385-75-1	302.37	NA	NA	NA	NA	4.22E-02	EPA6 calc c	4.88E-06
Dibenzo(a,h)fluoranthene	no cas #	302.37	NA	NA	NA	NA	4.22E-02	EPA6 calc c	4.88E-06
Dibenzo[a,e]pyrene	192-65-4	302.37	NA	NA	NA	NA	4.22E-02	EPA6 calc c	4.88E-06
Dibenzo[a,h]pyrene	189-64-0	302.37	NA	NA	NA	NA	4.22E-02	EPA6 calc c	4.88E-06
Dibenzo[a,i]pyrene	189-55-9	302.37	NA	NA	NA	NA	4.22E-02	EPA6 calc c	4.88E-06
Fluoranthene	206-44-0	202.26	NA	9.33E-06	9.33E-06	EPA6 e	2.75E-02	EPA6 e	7.18E-06
Hexachloronaphthalene	1335-87-1	334.84	NA	NA	NA	NA	3.94E-02	EPA6 calc c	4.56E-06
Indeno(1,2,3-cd)pyrene	193-39-5	276.34	NA	4.86E-09	4.86E-09	EPA6 e	1.90E-02	EPA6 e	5.66E-06
Octachloronaphthalene	2234-13-1	403.73	NA	NA	NA	NA	3.48E-02	EPA6 calc c	4.03E-06
Pentachloronaphthalene	1321-64-8	300.4	NA	NA	NA	NA	4.24E-02	EPA6 calc c	4.90E-06
Tetrachloronaphthalene	1335-88-2	265.95	NA	NA	NA	NA	4.59E-02	EPA6 calc c	5.32E-06
Trichloronaphthalene	1321-65-9	231.51	NA	NA	NA	NA	5.04E-02	EPA6 calc c	5.84E-06

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Mol. Wt.		Henry's Constant			Diffusivity		
Constituent of Potential Concern	CAS Registry Number	MW (g/mol)	pH	H (Published Value)	H (atm*m^3/mol)	Source	D_a (cm^2/s)	Source	D_w (cm^2/s)
Light Substituted Benzene Compounds									
1,2,3-Trichlorobenzene	87-61-6	181.46	NA	2.84E-03	2.84E-03	EPA6 e	3.02E-02	EPA6 e	8.15E-06
1,2,4-Trichlorobenzene	120-82-1	181.46	NA	2.61E-03	2.61E-03	EPA6 e	3.00E-02	EPA6 e	8.23E-06
1,2,4-Trimethlyl benzene	25551-13-7	120.19	NA	NA	NA	NA	7.80E-02	EPA6 calc c	9.03E-06
1,2,4-Trimethyl benzene	95-63-6	120.19	NA	5.81E+02	5.75E-03	Properties a	7.80E-02	EPA6 calc c	9.03E-06
1,2-Dichlorobenzene	95-50-1	147.01	NA	2.11E-03	2.11E-03	EPA6 e	4.11E-02	EPA6 e	8.93E-06
1,3,5-Trimethyl benzene	108-67-8	120.19	NA	7.81E-03	7.81E-03	EPA6 e	6.48E-02	EPA6 e	7.86E-06
1,3-Dichlorobenzene	541-73-1	147.01	NA	1.11E+02	1.11E+02	EPA6 e	4.14E-02	EPA6 e	8.85E-06
1,3-Dinitrobenzene	99-65-0	168.11	NA	1.25E-07	1.25E-07	EPA6 e	3.18E-02	EPA6 e	9.15E-06
1,4-Dichlorobenzene	106-46-7	147.01	NA	2.80E-03	2.80E-03	EPA6 e	4.14E-02	EPA6 e	8.85E-06
1,4-Dinitrobenzene	100-25-4	168.11	NA	NA	NA	NA	6.24E-02	EPA6 calc c	7.22E-06
2,4,5-Trichorophenol	95-95-4	197.46	7	5.64E-06	5.64E-06	EPA6 e	2.91E-02	EPA6 e	7.03E-06
2,4,6-Trichlorophenol	88-06-2	197.46	7	4.06E-06	4.06E-06	EPA6 e	2.62E-02	EPA6 e	8.08E-06
2,4-Dichlorophenol	120-83-2	163.01	7	2.38E-07	2.38E-07	EPA6 e	2.69E-02	EPA6 e	7.79E-06
2,4-Dimethylphenol	105-67-9	122.17	7	4.27E-09	4.27E-09	EPA6 e	5.84E-02	EPA6 e	8.69E-06
2,4-Dinitrophenol	51-28-5	184.11	7	4.82E-09	4.82E-09	EPA6 e	2.73E-02	EPA6 e	9.06E-06
2,4-Dinitrotoluene	121-14-2	182.14	NA	1.46E-07	1.46E-07	EPA6 e	3.09E-02	EPA6 e	7.86E-06
2,6-Dinitrotoluene	606-20-2	182.15	NA	1.30E-07	1.30E-07	EPA6 e	3.11E-02	EPA6 e	7.76E-06
2-Chlorotoluene	95-49-8	126.59	NA	6.41E-02	6.35E-04	Henry's f	7.54E-02	EPA6 calc c	8.73E-06
2-Chorophenol	95-57-8	128.56	NA	1.66E-05	1.66E-05	EPA6 e	5.01E-02	EPA6 e	9.46E-06
2-Nitrophenol	88-75-5	139.11	NA	1.46E-05	1.46E-05	EPA6 e	4.44E-02	EPA6 e	9.19E-06
4,6-Dinitro-o-cresol	534-52-1	198.13	NA	5.76E-05	1.40E-06	RAIS b	2.93E-02	RAIS b	6.91E-06
4-Chlorotoluene	106-43-4	126.59	NA	NA	NA	NA	7.54E-02	EPA6 calc c	8.73E-06
4-Nitrophenol	100-02-7	139.11	NA	7.32E-09	7.32E-09	EPA6 e	4.30E-02	EPA6 e	9.61E-06
alpha-Methylstyrene	98-83-9	118.18	NA	2.42E-01	5.91E-03	RAIS b	2.64E-01	RAIS b	1.14E-05
Aniline	62-53-3	93.12	NA	2.28E-06	2.28E-06	EPA6 e	8.56E-01	EPA6 e	1.01E-05
Benzotrichloride	98-07-7	195.47	NA	4.02E-02	9.81E-04	RAIS b	2.75E-02	RAIS b	7.77E-06
Benzyl Chloride	100-44-7	126.58	NA	4.13E-04	4.13E-04	EPA6 e	5.43E-02	EPA6 e	8.80E-06
Bromobenzene	108-86-1	157.01	NA	2.11E+02	2.09E-03	Properties a	6.53E-02	EPA6 calc c	7.56E-06

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Mol. Wt.		Henry's Constant			Diffusivity		
Constituent of Potential Concern	CAS Registry Number	MW (g/mol)	pH	H (Published Value)	H (atm*m^3/mol)	Source	D_a (cm^2/s)	Source	D_w (cm^2/s)
Chlorobenzene	108-90-7	112.56	NA	4.38E-03	4.38E-03	EPA6 e	6.35E-02	EPA6 e	9.49E-06
Cumene	98-82-8	120.19	NA	1.29E-02	1.29E-02	EPA6 e	6.50E-02	EPA6 e	7.83E-06
m-Cresol	108-39-4	108.13	NA	8.93E-07	8.93E-07	EPA6 e	6.93E-02	EPA6 e	9.30E-06
n-Butyl benzene	104-51-8	134.22	NA	1.30E+03	1.29E-02	Properties a	7.25E-02	EPA6 calc c	8.39E-06
Nitrobenzene	98-95-3	123.11	NA	2.06E-05	2.06E-05	EPA6 e	5.43E-02	EPA6 e	9.43E-06
n-Propyl benzene	103-65-1	120.19	NA	1.11E+03	1.10E-02	Properties a	7.80E-02	EPA6 calc c	9.03E-06
o-Cresol	95-48-7	108.13	NA	1.62E-06	1.62E-06	EPA6 e	6.88E-02	EPA6 e	9.41E-06
o-Dinitrobenzene	528-29-0	168.11	NA	NA	NA	NA	6.24E-02	EPA6 calc c	7.22E-06
o-Nitroaniline	88-74-4	138.12	NA	1.17E-06	1.17E-06	EPA6 e	4.29E-02	EPA6 e	9.81E-06
o-Toluidine	95-53-4	107.15	NA	2.43E-06	2.43E-06	EPA6 e	7.14E-02	EPA6 e	9.12E-06
p-Chloroaniline	106-47-8	127.57	NA	1.17E-06	1.17E-06	EPA6 e	4.80E-02	EPA6 e	1.02E-05
p-Cresol	106-44-5	108.13	NA	7.99E-07	7.99E-07	EPA6 e	6.93E-02	EPA6 e	9.30E-06
Phenol	108-95-2	94.11	NA**	5.95E-07	5.95E-07	EPA6 e	8.27E-02	EPA6 e	1.03E-05
p-Nitrochlorobenzene	100-00-5	157.56	7	NA	NA	NA	6.51E-02	EPA6 calc c	7.54E-06
p-Toluidine	106-49-0	107.17	NA	6.56E-01	6.50E-06	Properties a	6.98E-02	RAIS b	9.43E-06
sec-Butyl benzene	135-98-8	134.22	NA	1.40E+03	1.39E-02	Properties a	7.25E-02	EPA6 calc c	8.39E-06
tert-Butyl benzene	98-06-6	134.22	NA	1.20E+03	1.19E-02	Properties a	7.25E-02	EPA6 calc c	8.39E-06
Toluene-2,6-diamine	823-40-5	122.17	NA	5.08E-10	1.24E-11	RAIS b	5.59E-02	RAIS b	9.23E-06
Other Light Semivolatile Compounds									
1,1 Dimethyl hydrazine	57-14-7	60.099	NA	6.72E-05	1.64E-06	RAIS b	1.06E-01	RAIS b	1.09E-05
1,1-Biphenyl	92-52-4	154.21	NA	2.80E+01	2.77E-04	Properties a	6.61E-02	EPA6 calc c	7.65E-06
1,2 Diphenylhydrazine	122-66-7	184.24	NA	1.28E-07	1.28E-07	EPA6 e	2.95E-02	EPA6 e	7.24E-06
1,2-Dimethylhydrazine	540-73-8	60.099	NA	NA	NA	NA	1.24E-01	EPA6 calc c	1.43E-05
1,3-Propane sultone	1120-71-4	122.1	NA	NA	NA	NA	7.72E-02	EPA6 calc c	8.94E-06
2,4-Toluene diisocyanate	584-84-9	174.16	NA	NA	NA	NA	6.09E-02	EPA6 calc c	7.05E-06
2-Chloroacetophenone	532-27-4	154.6	NA	3.57E-05	8.71E-07	RAIS b	3.83E-02	RAIS b	8.71E-06
2-Propenoic acid	79-10-7	72.06	NA	4.20E-02	4.16E-07	Properties a	9.80E-02	RAIS b	1.06E-05
4,4-Methylenedianiline	101-77-9	198.27	NA	1.15E-06	2.81E-08	RAIS b	3.56E-02	RAIS b	5.39E-06
Acetophenone	98-86-2	120.5	NA	1.03E-05	1.03E-05	EPA6 e	6.00E-02	EPA6 e	8.73E-06

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Mol. Wt.		Henry's Constant			Diffusivity		
Constituent of Potential Concern	CAS Registry Number	MW (g/mol)	pH	H (Published Value)	H (atm*m^3/mol)	Source	D_a (cm^2/s)	Source	D_w (cm^2/s)
Benzoic acid	65-85-0	122.13	7	6.31E-05	1.54E-06	RAIS b	5.36E-02	RAIS b	7.97E-06
bis(2-Chloroethoxy)methane	111-91-1	173.04	NA	NA	NA	NA	NA	NA	NA
bis(2-Chloroethyl) ether	111-44-4	143.02	NA	2.13E-05	2.13E-05	EPA6 e	4.40E-02	EPA6 e	8.70E-06
Chlorocyclopentadiene	41851-50-7	100.55	NA	9.18E-01	2.24E-02	RAIS b	8.79E-02	EPA6 calc c	1.02E-05
Cyclohexanol	108-93-0	100.16	NA	5.81E-01	5.75E-06	Properties a	8.81E-02	EPA6 calc c	1.02E-05
Dichloroisopropyl ether	108-60-1	171.07	NA	1.14E+01	1.13E-04	Properties a	3.50E-02	RAIS b	7.36E-06
Dichloromethyl ether	542-88-1	114.96	NA	3.70E-03	9.03E-05	RAIS b	5.73E-02	EPA6 calc c	9.38E-06
Dichloropentadiene	no cas #	137.01	NA	NA	NA	NA	7.15E-02	EPA6 calc c	8.28E-06
Dimethyl sulfate	77-78-1	126.14	NA	1.65E-04	4.01E-06	RAIS b	5.14E-02	RAIS b	9.57E-06
Dimethylaniline	121-69-7	121.18	NA	5.68E-04	1.39E-05	RAIS b	1.53E-01	RAIS b	1.41E-05
di-n-propylnitrosamine	621-64-7	130.19	NA	5.43E-08	5.43E-08	EPA6 e	5.67E-02	EPA6 e	7.75E-06
Diphenyl ether	101-84-8	170.21	NA	2.51E+01	2.49E-04	Properties a	6.19E-02	EPA6 calc c	7.16E-06
Epichlorohydrin	106-89-8	92.53	NA	3.08E-05	3.08E-05	EPA6 e	8.13E-02	EPA6 e	1.10E-05
Ethyl carbomate (urethane)	51-79-6	89.094	NA	NA	NA	NA	9.52E-02	EPA6 calc c	1.10E-05
Ethyl methane sulfanate	62-50-0	124.15	NA	8.87E-08	8.87E-08	EPA6 e	7.63E-02	EPA6 e	8.84E-06
Ethylene dibromide	106-93-4	187.88	NA	4.47E-04	4.47E-04	EPA6 e	2.17E-02	EPA6 e	1.19E-05
Ethylene glycol	107-21-1	62.08	NA	7.38E-08	1.80E-09	RAIS b	1.08E-01	RAIS b	1.22E-05
Ethylene glycol monobutyl ether	111-76-2	118	NA	2.15E-05	5.26E-07	RAIS b	6.51E-02	RAIS b	8.15E-06
Ethylene glycol monoethyl ether acetate	111-15-9	148.17	NA	7.28E-05	1.77E-06	RAIS b	4.58E-02	RAIS b	7.56E-06
Ethylene thiourea	96-45-7	102.17	NA	6.23E-09	1.52E-10	RAIS b	7.15E-02	RAIS b	1.02E-05
Furfural	98-01-1	96.08	NA	3.75E-01	3.71E-06	Properties a	8.72E-02	RAIS b	1.04E-05
Maleic hydrazide	123-33-1	112.09	NA	2.70E-01	6.59E-03	RAIS b	8.17E-02	EPA6 calc c	9.46E-06
Malononitrile	109-77-3	66.062	NA	NA	NA	NA	1.16E-01	EPA6 calc c	1.35E-05
Methyl styrene (mixed isomers)	25013-15-4	118.18	NA	NA	NA	NA	5.46E-02	RAIS b	8.08E-06
Methylhydrazine	60-34-4	46.072	NA	NA	NA	NA	1.48E-01	EPA6 calc c	1.71E-05
N,N-Diphenylamine	122-39-4	169.23	NA	3.50E-02	3.47E-07	Properties a	6.21E-02	EPA6 calc c	7.19E-06
Nitric acid, propyl ester	627-13-4	105.09	NA	NA	NA	NA	8.53E-02	EPA6 calc c	9.88E-06
N-Nitrosodi-n-butylamine	924-16-3	158.2	NA	5.47E-05	5.47E-05	EPA6 e	6.50E-02	EPA6 e	7.52E-06
N-Nitrosomorpholine	59-89-2	116.12	NA	NA	NA	NA	7.98E-02	EPA6 calc c	9.24E-06

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Mol. Wt.		Henry's Constant			Diffusivity		
Constituent of Potential Concern	CAS Registry Number	MW (g/mol)	pH	H (Published Value)	H (atm*m^3/mol)	Source	D_a (cm^2/s)	Source	D_w (cm^2/s)
N-Nitroso-N,N-dimethylamine	62-75-9	74.1	NA	3.20E-05	7.80E-07	RAIS b	1.13E-01	RAIS b	1.24E-05
o-Anisidine	90-04-0	123.15	NA	NA	NA	NA	7.68E-02	EPA6 calc c	8.89E-06
Oxalic acid	144-62-7	90.035	NA	NA	NA	NA	9.46E-02	EPA6 calc c	1.10E-05
Phthalic Anhydride	85-44-9	148.11	NA	6.28E-09	6.28E-09	EPA6 e	4.04E-02	EPA6 e	8.97E-06
p-Phthalic acid	100-21-0	166.13	NA	4.26E-07	1.04E-08	RAIS b	3.90E-02	RAIS b	7.14E-06
Pyridine	110-86-1	79.1	NA	6.86E-03	6.86E-03	EPA6 e	1.10E-01	EPA6 e	1.08E-05
Quinoline	91-22-5	129.16	NA	1.11E-05	2.70E-07	RAIS b	5.46E-02	RAIS b	8.31E-06
Quinone	106-51-4	108.1	NA	NA	NA	NA	8.37E-02	EPA6 calc c	9.69E-06
Safrole	94-59-7	162.18	NA	1.19E-05	1.19E-05	EPA6 e	4.06E-02	EPA6 e	7.16E-06
Tetrahydrofuran	109-99-9	72.1	NA	NA	NA	EPA6 e	1.31E-01	EPA6 e	1.07E-05
Other Heavy Semivolatile Compounds									
1,2,4,5-Tetrachlorobenzene	95-94-3	215.89	NA	1.18E-03	1.18E-03	EPA6 e	2.11E-02	EPA6 e	8.75E-06
1,3,5-Trinitrobenzene	99-35-4	213.11	NA	8.66E-08	8.66E-08	EPA6 e	2.84E-02	EPA6 e	6.08E-06
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	220.35	NA	NA	NA	NA	5.21E-02	EPA6 calc c	6.03E-06
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	266.25	NA	NA	NA	NA	4.59E-02	EPA6 calc c	5.32E-06
2-sec-Butyl-4,6-dinitrophenol	88-85-7	240.22	NA	1.87E-05	4.56E-07	RAIS b	4.92E-02	EPA6 calc c	5.69E-06
3,3-Dichlorobenzidine	91-94-1	253.13	NA	2.08E-08	2.08E-08	EPA6 e	2.28E-02	EPA6 e	5.48E-06
3,3'-Dimethoxybenzidine	119-90-4	244.28	NA	3.36E-10	3.36E-10	EPA6 e	2.38E-02	EPA6 e	5.60E-06
4-Bromophenylphenyl ether	101-55-3	249.11	NA	4.80E-03	1.17E-04	RAIS b	4.80E-02	EPA6 calc c	5.56E-06
Ammonium perfluorooctanoate	3825-26-1	431.1	NA	NA	NA	NA	3.33E-02	EPA6 calc c	3.86E-06
Azobenzene	103-33-3	182.22	NA	2.04E-04	4.98E-06	RAIS b	3.15E-02	RAIS b	7.45E-06
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	358.54	NA	NA	NA	NA	3.76E-02	EPA6 calc c	4.36E-06
Captan	133-06-2	300.57	NA	6.00E-01	5.94E-06	Properties a	1.81E-02	RAIS b	5.00E-06
Chlorobenzilate	510-15-6	325.2	NA	7.24E-08	7.24E-08	EPA6 e	1.65E-02	EPA6 e	4.72E-06
Dibutylphosphate	107-66-4	210.21	NA	NA	NA	NA	5.37E-02	EPA6 calc c	6.22E-06
Dimethyl aminoazobenzene	60-11-7	225.29	NA	NA	NA	NA	5.13E-02	EPA6 calc c	5.94E-06
Hexachlorobenzene	118-74-1	284.8	NA	5.35E-04	5.35E-04	EPA6 e	1.41E-02	EPA6 e	7.84E-06
Hexachlorobutadiene	87-68-3	260.76	NA	2.39E-02	2.39E-02	EPA6 e	1.73E-02	EPA6 e	7.33E-06
Hexachlorocyclopentadiene	77-47-4	272.77	NA	1.72E-02	1.72E-02	EPA6 e	1.61E-02	EPA6 e	7.21E-06

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Mol. Wt.		Henry's Constant			Diffusivity		
Constituent of Potential Concern	CAS Registry Number	MW (g/mol)	pH	H (Published Value)	H (atm*m^3/mol)	Source	D_a (cm^2/s)	Source	D_w (cm^2/s)
Hexachloroethane	67-72-1	236.74	NA	3.60E-03	3.60E-03	EPA6 e	1.77E-02	EPA6 e	8.88E-06
Hexachlorophene	70-30-4	406.92	NA	4.88E-10	4.88E-10	EPA6 e	3.46E-02	EPA6 e	4.01E-06
Hexamethylene-1,5-diisocyanate	822-06-0	168.22	NA	1.09E-05	2.66E-07	RAIS b	3.80E-02	RAIS b	7.17E-06
Mirex	2385-85-5	545.55	NA	5.32E+01	5.27E-04	Properties a	2.85E-02	EPA6 calc c	3.30E-06
Nitrofen	1836-75-5	284.1	NA	NA	NA	NA	4.40E-02	EPA6 calc c	5.09E-06
Pentachlorobenzene	608-93-5	250.34	NA	2.43E-02	2.43E-02	EPA6 e	1.86E-02	EPA6 e	7.34E-06
Pentachloronitrobenzene	82-68-8	295.36	NA	2.86E-02	2.86E-02	EPA6 e	1.87E-02	EPA6 e	5.00E-06
Pentachlorophenol*	87-86-5	266.34	2	1.41E-05	1.41E-05	EPA6 e	1.56E-02	EPA6 e	8.01E-06
Pentachlorophenol*	87-86-5	266.34	3	1.41E-05	1.41E-05	EPA6 e	1.56E-02	EPA6 e	8.01E-06
Pentachlorophenol*	87-86-5	266.34	4	1.41E-05	1.41E-05	EPA6 e	1.56E-02	EPA6 e	8.01E-06
Pentachlorophenol*	87-86-5	266.34	5	1.41E-05	1.41E-05	EPA6 e	1.56E-02	EPA6 e	8.01E-06
Pentachlorophenol*	87-86-5	266.34	6	1.41E-05	1.41E-05	EPA6 e	1.56E-02	EPA6 e	8.01E-06
Pentachlorophenol*	87-86-5	266.34	7	1.41E-05	1.41E-05	EPA6 e	1.56E-02	EPA6 e	8.01E-06
Pentachlorophenol*	87-86-5	266.34	8	1.41E-05	1.41E-05	EPA6 e	1.56E-02	EPA6 e	8.01E-06
Pentachlorophenol*	87-86-5	266.34	9	1.41E-05	1.41E-05	EPA6 e	1.56E-02	EPA6 e	8.01E-06
Pentachlorophenol*	87-86-5	266.34	10	1.41E-05	1.41E-05	EPA6 e	1.56E-02	EPA6 e	8.01E-06
Pentachlorophenol*	87-86-5	266.34	11	1.41E-05	1.41E-05	EPA6 e	1.56E-02	EPA6 e	8.01E-06
Pentachlorophenol*	87-86-5	266.34	12	1.41E-05	1.41E-05	EPA6 e	1.56E-02	EPA6 e	8.01E-06
Pentachlorophenol*	87-86-5	266.34	13	1.41E-05	1.41E-05	EPA6 e	1.56E-02	EPA6 e	8.01E-06
Pentachlorophenol*	87-86-5	266.34	14	1.41E-05	1.41E-05	EPA6 e	1.56E-02	EPA6 e	8.01E-06
Picric acid	88-89-1	229.11	NA	3.79E-13	3.79E-13	Chemfate i	5.07E-02	EPA6 calc c	5.88E-06
Pronamide	23950-58-5	256.13	NA	9.05E-06	9.05E-06	EPA6 e	4.71E-02	EPA6 e	5.45E-06
Strychnine	57-24-9	334.4	NA	4.90E-13	4.90E-13	EPA6 e	1.38E-02	EPA6 e	5.58E-06
Terphenyls	26140-60-3	230.31	NA	NA	NA	NA	5.06E-02	EPA6 calc c	5.86E-06
Tributyl phosphate	126-73-8	266.32	NA	1.50E-07	1.50E-07	Chemfate i	4.59E-02	EPA6 calc c	5.31E-06
Trifluralin	1582-09-8	335.29	NA	4.03E+00	3.99E-05	Properties a	1.49E-02	RAIS b	5.04E-06
Triphenylamine	603-34-9	245.32	NA	NA	NA	NA	4.85E-02	EPA6 calc c	5.61E-06
Herbicides and Organopesticides									
2,4,5-T	93-76-5	255.48	NA	8.79E-04	8.70E-09	Properties a	1.92E-02	RAIS b	6.70E-06

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Mol. Wt.		Henry's Constant			Diffusivity		
Constituent of Potential Concern	CAS Registry Number	MW (g/mol)	pH	H (Published Value)	H (atm*m^3/mol)	Source	D_a (cm^2/s)	Source	D_w (cm^2/s)
2,4-D and esters	94-75-7	221.04	NA	1.38E-05	1.37E-10	Properties a	2.31E-02	RAIS b	7.31E-06
4,4-DDD	72-54-8	320.05	NA	8.51E+00	8.43E-05	Properties a	1.69E-02	RAIS b	4.76E-06
4,4-DDT	50-29-3	354.5	NA	1.31E+00	1.30E-05	Properties a	1.37E-02	RAIS b	4.95E-06
Aldrin	309-00-2	364.93	NA	1.35E-07	1.35E-07	EPA6 e	1.43E-02	EPA6 e	4.40E-06
alpha-BHC	319-84-6	290	NA	6.78E-06	6.78E-06	EPA6 e	1.91E-02	EPA6 e	5.04E-06
beta-BHC	319-85-7	290.83	NA	3.46E-07	3.46E-07	EPA6 e	1.90E-02	EPA6 e	5.40E-06
Chlordane	57-74-9	409.8	NA	2.64E-05	2.64E-05	EPA6 e	1.18E-02	EPA6 e	4.37E-06
DDE	72-55-9	319.03	NA	1.24E-04	1.24E-04	EPA6 e	1.70E-02	EPA6 e	4.78E-06
Delta-BHC	319-86-8	290.85	NA	8.25E-02	8.17E-07	Properties a	4.33E-02	EPA6 calc c	5.01E-06
Dieldrin	60-57-1	380.93	NA	3.51E-09	3.51E-09	EPA6 e	1.36E-02	EPA6 e	4.29E-06
Endothall	145-73-3	186.16	NA	1.06E-08	2.59E-10	RAIS b	5.83E-02	EPA6 calc c	6.75E-06
Endrin	72-20-8	380.93	NA	1.19E-06	1.19E-06	EPA6 e	1.07E-02	EPA6 e	5.76E-06
gamma-BHC(Lindane)	58-89-9	290.83	NA	1.58E-01	1.56E-06	Properties a	4.33E-02	EPA6 calc c	5.01E-06
Heptachlor	76-44-8	373.35	NA	5.87E-06	5.87E-06	EPA6 e	1.12E-02	EPA6 e	5.69E-06
Isodrin	465-73-6	364.91	NA	NA	NA	NA	3.72E-02	EPA6 calc c	4.31E-06
Methoxychlor	72-43-5	345.65	NA	6.33E-06	6.33E-06	EPA6 e	1.30E-02	EPA6 e	5.59E-06
Silvex (2,4,5-TP)	93-72-1	269.51	NA	2.55E-01	6.23E-03	RAIS b	1.94E-02	RAIS b	5.83E-06
Toxaphene	8001-35-2	414	NA	4.56E+01	4.51E-04	Properties a	1.16E-02	RAIS b	4.34E-06

NA = Value not available
** = Varies with pH; values given for a pH of 7.0.
a = Illustrated Handbook of Physical - Chemical Properties and Environmental Fate for Organic Chemicals, 1992.
b = RAIS database: http://risk.lsd.ornl.gov/rap_hp.shtml
c = Equations derived from EPA Region 6 Guidance Appendix A-3
d = Handbook of Environmental degradation Rates. 1991.
e = EPA Region 6 Hazardous Waste Combustion Guidance
f = "A Critical Review of Henry's Law Constants for Chemicals of Environmental Interest", J. Phys. Chem. Ref. Data, Vol. 10, No. 4, 1981.
g = Handbook of Environmental Fate and Exposure Data for Organic Chemicals
h = http://www.epa.gov/region09/waste/sfund/prg/.
i = http://esc.syrres.com/efdb/Chemfate.htm
j = Hansch and Leo 1985 in Syracuse 1996

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Mol. Wt.		Henry's Constant			Diffusivity		
Constituent of Potential Concern	CAS Registry Number	MW (g/mol)	pH	H (Published Value)	H (atm*m^3/mol)	Source	D_a (cm^2/s)	Source	D_w (cm^2/s)

k = Schwarzenbach et al. 1993
l = Tomlin 1994 in Syracuse 1996
m = http://www.speclab.com/compound/c959988.htm

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Octanol / Water Partitioning Coefficients					
Constituent of Potential Concern	CAS Registry Number	Source	K_{ow} (Published Value)	K_{ow} (unitless)	Source	K_{oc} (Published Value)	K_{oc} (mL/g)	Source
Aromatic Halogenated Hydrocarbons								
2,3,4,6-Tetrachlorophenol	58-90-2	RAIS b	4.42E+00	2.63E+04	Properties a	3.82E+00	6.61E+03	Properties a
4-Chloro-3-methylphenol	59-50-7	EPA6 calc c	3.10E+00	1.26E+03	Properties a	2.78E+00	6.03E+02	Properties a
Aromatic Nonhalogenated Hydrocarbons								
2-Nitrotoluene	88-72-2	RAIS b	2.30E+00	2.00E+02	Properties a	2.63E+00	4.27E+02	Properties a
4-Nitrobiphenyl	92-93-3	EPA6 calc c	3.77E+00	5.89E+03	Chemfate i	NA	NA	NA
Benzaldehyde	100-52-7	EPA6 e	3.00E+01	3.00E+01	EPA6 e	2.01E-01	2.01E-01	EPA6 e
Benzene	71-43-2	EPA6 e	1.37E+02	1.37E+02	EPA6 e	6.20E+01	6.20E+01	EPA6 e
Benzyl alcohol	100-51-6	EPA6 e	1.26E+01	1.26E+01	EPA6 e	1.02E+01	1.02E+01	EPA6 e
Ethyl Benzene	100-41-4	EPA6 e	1.33E+03	1.33E+03	EPA6 e	2.04E+02	2.04E+02	EPA6 e
m-Xylene	108-38-3	EPA6 e	1.59E+03	1.59E+03	EPA6 e	1.96E+02	1.96E+02	EPA6 e
o-Xylene	95-47-6	EPA6 e	1.35E+03	1.35E+03	EPA6 e	2.41E+02	2.41E+02	EPA6 e
p-Xylene	106-42-3	EPA6 e	1.48E+03	1.48E+03	EPA6 e	3.11E+02	3.11E+02	EPA6 e
Styrene	100-42-5	EPA6 e	8.49E+02	8.49E+02	EPA6 e	9.12E+02	9.12E+02	EPA6 e
Toluene	108-88-3	EPA6 e	4.65E+02	4.65E+02	EPA6 e	1.40E+02	1.40E+02	EPA6 e
Non-aromatic Nonhalogenated Hydrocarbons								
1,2-Epoxybutane	106-88-7	RAIS b	1.44E+00	2.76E+01	RAIS b	7.94E+00	7.94E+00	RAIS b
1,3-Butadiene	106-99-0	EPA6 calc c	1.99E+00	9.77E+01	Organics g	NA	9.04E+01	EPA6 calc c
1,4-Dioxane	123-91-1	EPA6 e	5.40E-01	5.40E-01	EPA6 e	8.76E-01	8.76E-01	EPA6 e
1-Methylpropyl alcohol	78-92-2	EPA6 calc c	6.10E-01	4.07E+00	Properties a	NA	3.98E+00	EPA6 calc c
1-Nitropropane	108-03-2	EPA6 calc c	8.70E-01	7.41E+00	Hansch j	NA	7.17E+00	EPA6 calc c
2,2,4-Trimethlylpentane	540-84-1	EPA6 calc c	5.02E+00	1.05E+05	Properties a	NA	8.61E+04	EPA6 calc c
2-Butanone	78-93-3	EPA6 e	1.91E+00	1.91E+00	EPA6 e	2.34E+00	2.34E+00	EPA6 e
2-Butenaldehyde (2-Butenal)	4170-30-3	EPA6 calc c	NA	NA	NA	NA	NA	NA
2-Ethoxyethanol	110-80-5	RAIS b	-1.00E-01	7.94E-01	Organics g	1.32E+00	2.09E+01	Organics g
2-Heptanone	110-43-0	EPA6 calc c	1.98E+00	9.55E+01	Properties a	NA	8.84E+01	EPA6 calc c
2-Hexanone	591-78-6	EPA6 calc c	1.38E+00	2.40E+01	Properties a	2.13E+00	1.34E+02	Properties a
2-Methoxyethanol	109-86-4	RAIS b	2.52E-01	1.79E+00	RAIS b	8.53E+00	8.53E+00	RAIS b
2-Methyl-2-propanol	75-65-0	EPA6 calc c	3.50E-01	2.24E+00	Properties a	1.57E+00	3.72E+01	Properties a

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Octanol / Water Partitioning Coefficients					
Constituent of Potential Concern	CAS Registry Number	Source	K_{ow} (Published Value)	K_{ow} (unitless)	Source	K_{oc} (Published Value)	K_{oc} (mL/g)	Source
2-Methyl-2-propenenitrile	126-98-7	EPA6 e	3.47E+00	3.47E+00	EPA6 e	3.74E+00	3.74E+00	EPA6 e
2-Methylaziridine	75-55-8	EPA6 calc c	NA	NA	NA	NA	NA	NA
2-Methylpropyl alcohol	78-83-1	RAIS b	7.60E-01	5.75E+00	Properties a	NA	5.59E+00	EPA6 calc c
2-Pentanone	107-87-9	EPA6 calc c	9.10E-01	8.13E+00	Properties a	1.87E+00	7.40E+01	Properties a
2-Propanone (Acetone)	67-64-1	EPA6 e	6.00E-01	6.00E-01	EPA6 e	9.51E-01	9.51E-01	EPA6 e
2-Propene-1-ol	107-18-6	RAIS b	1.70E-01	1.48E+00	Properties a	NA	1.47E+00	EPA6 calc c
2-Propyl alcohol	67-63-0	EPA6 calc c	5.00E-02	1.12E+00	Properties a	NA	1.12E+00	EPA6 calc c
3-Heptanone	106-35-4	EPA6 calc c	NA	NA	NA	NA	NA	NA
3-Methyl-1-butanol	123-51-3	EPA6 calc c	NA	NA	NA	NA	NA	NA
3-Methyl-2-butanone	563-80-4	EPA6 calc c	NA	NA	NA	NA	NA	NA
3-Pentanone	96-22-0	EPA6 calc c	9.90E-01	9.77E+00	Properties a	1.08E+00	1.20E+01	Properties a
4-Heptanone	123-19-3	EPA6 calc c	NA	NA	NA	NA	NA	NA
4-Methyl-2-pentanone	108-10-1	EPA6 e	1.55E+01	1.55E+01	EPA6 e	1.20E+01	1.20E+01	EPA6 e
4-Methyl-3-penten-2-one	141-79-7	EPA6 calc c	NA	NA	NA	NA	NA	NA
5-Methyl-2-hexanone	110-12-3	EPA6 calc c	NA	NA	NA	NA	NA	NA
Acetaldehyde	75-07-0	EPA6 e	6.02E-01	6.02E-01	EPA6 e	9.53E-01	9.53E-01	EPA6 e
Acetamide	60-35-5	EPA6 calc c	-1.26E+00	5.50E-02	Properties a	-1.55E+00	2.82E-02	Properties a
Acetic acid	64-19-7	EPA6 calc c	-1.70E-01	6.76E-01	Properties a	0.00E+00	1.00E+00	Properties a
Acetic acid ethyl ester	141-78-6	RAIS b	7.30E-01	5.37E+00	Properties a	3.61E-01	2.30E+00	Properties a
Acetic acid n-butyl ester	123-86-4	EPA6 calc c	1.73E+00	5.37E+01	Properties a	NA	5.02E+01	EPA6 calc c
Acetonitrile	75-05-8	EPA6 e	4.57E-01	4.57E-01	EPA6 e	7.69E-01	7.69E-01	EPA6 e
Acrolein	107-02-8	EPA6 e	9.80E-01	9.80E-01	EPA6 e	1.39E+00	1.39E+00	EPA6 e
Acrylonitrile	107-13-1	EPA6 e	1.78E+00	1.78E+00	EPA6 e	2.22E+00	2.22E+00	EPA6 e
Bis(isopropyl)ether	108-20-3	EPA6 calc c	NA	NA	NA	NA	NA	NA
Butane	106-97-8	EPA6 calc c	2.89E+00	7.76E+02	k	NA	6.94E+02	EPA6 calc c
Carbon disulfide	75-15-0	EPA6 e	1.00E+02	1.00E+02	EPA6 e	5.14E+01	5.14E+01	EPA6 e
Cyanogen	460-19-5	RAIS b	8.08E-01	6.42E+00	RAIS b	5.22E+00	5.22E+00	RAIS b
Cyclohexane	110-82-7	EPA6 calc c	3.44E+00	2.75E+03	Properties a	2.68E+00	4.79E+02	Properties a
Cyclohexanone	108-94-1	RAIS b	8.10E-01	6.46E+00	Properties a	NA	6.26E+00	EPA6 calc c

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Octanol / Water Partitioning Coefficients					
Constituent of Potential Concern	CAS Registry Number	Source	K_{ow} (Published Value)	K_{ow} (unitless)	Source	K_{oc} (Published Value)	K_{oc} (mL/g)	Source
Cyclohexene	110-83-8	EPA6 calc c	2.86E+00	7.24E+02	Properties a	NA	6.48E+02	EPA6 calc c
Cyclopentane	287-92-3	EPA6 calc c	3.00E+00	1.00E+03	Properties a	NA	8.90E+02	EPA6 calc c
Ethyl alcohol	64-17-5	EPA6 calc c	-3.10E-01	4.90E-01	Organics g	NA	4.96E-01	EPA6 calc c
Ethyl ether	60-29-7	RAIS b	8.90E-01	7.76E+00	Properties a	NA	7.50E+00	EPA6 calc c
Ethyl methacrylate	97-63-2	EPA6 e	3.89E+01	3.89E+01	EPA6 e	2.46E+01	2.46E+01	EPA6 e
Formaldehyde	50-00-0	EPA6 e	2.20E+00	2.20E+00	EPA6 e	2.62E+00	2.62E+00	EPA6 e
Formamide	75-12-7	EPA6 calc c	-1.51E+00	3.09E-02	Chemfate i	NA	3.28E-02	EPA6 calc c
Formic acid	64-18-6	EPA6 e	2.90E-01	2.90E-01	EPA6 e	5.39E+00	5.39E+00	EPA6 e
Formic acid, methyl ester	107-31-3	EPA6 calc c	2.64E-01	5.45E-01	Chemfate i	NA	5.50E-01	EPA6 calc c
Glycidylaldehyde	765-34-4	EPA6 calc c	-7.30E-01	1.86E-01	RAIS b	1.00E-01	1.00E-01	RAIS b
Methyl acetate	79-20-9	RAIS b	1.80E-01	1.51E+00	Properties a	6.81E-01	4.80E+00	Properties a
Methyl alcohol	67-56-1	EPA6 e	1.95E-01	1.95E-01	EPA6 e	3.96E-01	3.96E-01	EPA6 e
Methyl isocyanate	624-83-9	EPA6 calc c	NA	NA	NA	NA	NA	NA
Methyl methacrylate	80-62-6	RAIS b	7.90E-01	6.17E+00	Properties a	1.80E+00	6.31E+01	Properties a
Methyl tert-butyl ether	1634-04-4	RAIS b	9.40E-01	8.71E+00	Properties a	NA	8.40E+00	EPA6 calc c
Methylacetylene	74-99-7	EPA6 calc c	9.40E-01	8.71E+00	Chemfate i	NA	8.40E+00	EPA6 calc c
Methylcyclohexane	108-87-2	EPA6 calc c	4.10E+00	1.26E+04	Properties a	NA	1.07E+04	EPA6 calc c
N,N-Dimethylacetamide	127-19-5	EPA6 calc c	NA	NA	NA	NA	NA	NA
n-Butyl alcohol	71-36-3	RAIS b	8.80E-01	7.59E+00	Properties a	NA	7.33E+00	EPA6 calc c
n-Heptane	142-82-5	EPA6 calc c	4.66E+00	4.57E+04	Properties a	NA	3.81E+04	EPA6 calc c
n-Hexane	110-54-3	EPA6 calc c	4.11E+00	1.29E+04	Properties a	NA	1.10E+04	EPA6 calc c
Nitromethane	75-52-5	EPA6 calc c	-3.50E-01	4.47E-01	Hansch j	NA	4.53E-01	EPA6 calc c
n-Nonane	111-84-2	EPA6 calc c	5.65E+00	4.47E+05	Properties a	NA	3.58E+05	EPA6 calc c
n-Octane	111-65-9	EPA6 calc c	4.00E+00	1.00E+04	Properties a	NA	8.55E+03	EPA6 calc c
n-Pentane	109-66-0	EPA6 calc c	3.21E+00	1.62E+03	Properties a	NA	1.43E+03	EPA6 calc c
n-Propionaldehyde	123-38-6	EPA6 calc c	5.90E-01	3.89E+00	Properties a	NA	3.80E+00	EPA6 calc c
n-Propyl alcohol	71-23-8	EPA6 calc c	2.50E-01	1.78E+00	Chemfate i	NA	1.76E+00	EPA6 calc c
n-Valeraldehyde	110-62-3	EPA6 calc c	NA	NA	NA	NA	NA	NA
Oxirane	75-21-8	EPA6 e	5.01E-01	5.01E-01	EPA6 e	8.26E-01	8.26E-01	EPA6 e

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Octanol / Water Partitioning Coefficients					
Constituent of Potential Concern	CAS Registry Number	Source	K_{ow} (Published Value)	K_{ow} (unitless)	Source	K_{oc} (Published Value)	K_{oc} (mL/g)	Source
p-Cymene	99-87-6	EPA6 calc c	4.10E+00	1.26E+04	Chemfate i	NA	1.07E+04	EPA6 calc c
Phosgene	75-44-5	EPA6 calc c	NA	NA	NA	NA	NA	NA
Propargyl alcohol	107-19-7	RAIS b	2.60E-01	1.82E+00	RAIS b	1.93E+00	1.93E+00	RAIS b
Propionic acid	79-09-4	EPA6 calc c	3.30E-01	2.14E+00	Properties a	NA	2.11E+00	EPA6 calc c
Propionitrile	107-12-0	EPA6 calc c	1.60E-01	1.45E+00	Properties a	NA	1.44E+00	EPA6 calc c
Propylene glycol monomethyl ether	107-98-2	RAIS b	-1.80E-01	6.61E-01	RAIS b	1.26E+01	1.26E+01	RAIS b
p-tert-Butyltoluene	98-51-1	EPA6 calc c	NA	NA	NA	NA	NA	NA
Triethylamine	121-44-8	RAIS b	NA	NA	NA	NA	NA	NA
Trimethylamine	75-50-3	EPA6 calc c	1.60E-01	1.45E+00	Properties a	6.02E-01	4.00E+00	Properties a
Vinyl acetate	108-05-4	EPA6 e	5.00E+00	5.00E+00	EPA6 e	4.97E+00	4.97E+00	EPA6 e
Non-aromatic Halogenated Hydrocarbons								
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	EPA6 calc c	NA	NA	NA	NA	NA	NA
1,1,1,2-Tetrachloroethane	630-20-6	EPA6 e	4.27E+02	4.27E+02	EPA6 e	1.59E+02	1.59E+02	EPA6 e
1,1,1-Trichloroethane	71-55-6	EPA6 e	2.64E+02	2.64E+02	EPA6 e	1.35E+05	1.35E+05	EPA6 e
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	EPA6 calc c	3.73E+00	5.37E+03	Properties a	2.50E+00	3.16E+02	Properties a
1,1,2,2-Tetrachloroethane	79-34-5	EPA6 e	4.40E+04	4.40E+04	EPA6 e	7.90E+01	7.90E+01	EPA6 e
1,1,2,2-Tetrachloroethene	127-18-4	EPA6 e	3.51E+02	3.51E+02	EPA6 e	2.65E+02	2.65E+02	EPA6 e
1,1,2-Trichloro-1,2,2-trifluoroethane	76-13-1	RAIS b	3.16E+00	1.45E+03	Properties a	2.41E+00	2.57E+02	Properties a
1,1,2-Trichloroethane	79-00-5	EPA6 e	1.25E+02	1.25E+02	EPA6 e	7.50E+01	7.50E+01	EPA6 e
1,1,2-Trichloroethylene	79-01-6	EPA6 e	2.71E+02	2.71E+02	EPA6 e	9.40E+01	9.40E+01	EPA6 e
1,1-Dichloroethane	75-34-3	EPA6 e	6.20E+01	6.20E+01	EPA6 e	5.30E+01	5.30E+01	EPA6 e
1,1-Dichloroethene	75-35-4	EPA6 e	1.32E+02	1.32E+02	EPA6 e	6.50E+01	6.50E+01	EPA6 e
1,2,3-Trichloropropane	96-18-4	EPA6 e	1.78E+02	1.78E+02	EPA6 e	8.05E+01	8.05E+01	EPA6 e
1,2-dibromo-3-chloropropane	96-12-8	EPA6 e	2.19E+02	2.19E+02	EPA6 e	9.47E+01	9.47E+01	EPA6 e
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	EPA6 calc c	2.82E+00	6.61E+02	Chemfate i	NA	NA	NA
1,2-Dichloroethane	107-06-2	EPA6 e	2.90E+01	2.90E+01	EPA6 e	1.96E+01	1.96E+01	EPA6 e
1,2-Dichloroethylene	540-59-0	EPA6 calc c	4.80E-01	3.02E+00	RAIS b	2.87E+00	2.87E+00	RAIS b
1,2-Dichloropropane	78-87-5	EPA6 e	1.78E+02	1.78E+02	EPA6 e	4.70E+01	4.70E+01	EPA6 e
1,3-Dichloropropene	542-75-6	EPA6 e	5.60E+01	5.60E+01	EPA6 e	2.70E+01	2.70E+01	EPA6 e

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Octanol / Water Partitioning Coefficients					
Constituent of Potential Concern	CAS Registry Number	Source	K_{ow} (Published Value)	K_{ow} (unitless)	Source	K_{oc} (Published Value)	K_{oc} (mL/g)	Source
1,4-Dichloro-2-butene	764-41-0	EPA6 calc c	8.73E-01	7.46E+00	RAIS b	1.82E+02	1.82E+02	RAIS b
1-Chloroethene	75-01-4	EPA6 e	1.40E+01	1.40E+01	EPA6 e	1.11E+02	1.11E+02	EPA6 e
2,2-Dichloropropionic acid	75-99-0	EPA6 calc c	7.78E-01	6.00E+00	RAIS b	5.82E+00	5.82E+00	RAIS b
2-Chloropropane	75-29-6	EPA6 calc c	1.90E+00	7.94E+01	Properties a	NA	7.38E+01	EPA6 calc c
3-Chloropropene (allyl chloride)	107-05-1	EPA6 calc c	9.55E-01	9.01E+00	RAIS b	4.79E+01	4.79E+01	RAIS b
Bromochloromethane	74-97-5	RAIS b	1.41E+00	2.57E+01	Properties a	NA	2.43E+01	EPA6 calc c
Bromodichloromethane	75-27-4	EPA6 e	1.06E+02	1.06E+02	EPA6 e	5.38E+01	5.38E+01	EPA6 e
Bromoethene	593-60-2	EPA6 calc c	1.07E+00	1.17E+01	RAIS b	1.51E+02	1.51E+02	RAIS b
Bromoform	75-25-2	EPA6 e	2.24E+02	2.24E+02	EPA6 e	1.26E+02	1.26E+02	EPA6 e
Bromomethane	74-83-9	EPA6 e	1.30E+01	1.30E+01	EPA6 e	9.00E+00	9.00E+00	EPA6 e
Carbon tetrachloride	56-23-5	EPA6 e	5.21E+02	5.21E+02	EPA6 e	1.52E+02	1.52E+02	EPA6 e
Chlordibromomethane	124-48-1	EPA6 e	1.50E+02	1.50E+02	EPA6 e	7.05E+01	7.05E+01	EPA6 e
Chlorodifluoromethane	75-45-6	EPA6 e	1.20E+01	1.20E+01	EPA6 e	9.83E+00	9.83E+00	EPA6 e
Chloroethane	75-00-3	EPA6 e	1.26E+03	1.26E+03	EPA6 e	3.71E+02	3.71E+02	EPA6 e
Chloroform	67-66-3	EPA6 e	8.90E+01	8.90E+01	EPA6 e	5.30E+01	5.30E+01	EPA6 e
Chloromethane	74-87-3	EPA6 e	8.00E+00	8.00E+00	EPA6 e	6.00E+00	6.00E+00	EPA6 e
Chloropentafluoroethane	76-15-3	EPA6 calc c	NA	NA	NA	NA	NA	NA
cis-1,2-Dichloroethene	156-59-2	EPA6 e	9.60E+01	9.60E+01	EPA6 e	4.98E+02	4.98E+02	EPA6 e
cis-1,3-Dichloropropene	10061-01-5	EPA6 calc c	NA	NA	NA	NA	NA	NA
Cyanogen bromide	506-68-3	EPA6 calc c	NA	NA	NA	NA	NA	NA
Cyanogen chloride	506-77-4	EPA6 calc c	2.00E-01	1.58E+00	RAIS b	1.72E+00	1.72E+00	RAIS b
Dichlorodifluoromethane	75-71-8	EPA6 e	1.44E+02	1.44E+02	EPA6 e	6.85E+00	6.85E+00	EPA6 e
Dichlorofluoromethane	75-43-4	EPA6 calc c	NA	NA	NA	NA	NA	NA
Dichloromethane	75-09-2	EPA6 e	1.80E+01	1.80E+01	EPA6 e	1.00E+01	1.00E+01	EPA6 e
Difluorodibromomethane	75-61-6	EPA6 calc c	NA	NA	NA	NA	NA	NA
Hexafluoroacetone	684-16-2	EPA6 calc c	NA	NA	NA	NA	NA	NA
Iodomethane	74-88-4	EPA6 calc c	1.69E+00	4.90E+01	Properties a	NA	4.59E+01	EPA6 calc c
Methylene bromide	74-95-3	EPA6 e	4.17E+01	4.17E+01	EPA6 e	2.60E-01	2.60E-01	EPA6 e
Pentachloroethane	76-01-7	EPA6 calc c	3.05E+00	1.12E+03	Properties a	NA	9.96E+02	EPA6 calc c

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Octanol / Water Partitioning Coefficients					
Constituent of Potential Concern	CAS Registry Number	Source	K_{ow} (Published Value)	K_{ow} (unitless)	Source	K_{oc} (Published Value)	K_{oc} (mL/g)	Source
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	EPA6 calc c	NA	NA	NA	NA	NA	NA
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	EPA6 calc c	NA	NA	NA	NA	NA	NA
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	EPA6 calc c	NA	NA	NA	NA	NA	NA
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	EPA6 calc c	NA	NA	NA	NA	NA	NA
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	EPA6 calc c	NA	NA	NA	NA	NA	NA
2',3,4,4',5-Pentachlorobiphenyl	no cas #	EPA6 calc c	NA	NA	NA	NA	NA	NA
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	EPA6 calc c	7.12E+00	1.32E+07	Chemfate i	NA	9.98E+06	EPA calc c
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	EPA6 calc c	7.41E+00	2.56E+07	Properties a	6.60E+00	3.98E+06	Properties a
3,3',4,4',5-Pentachlorobiphenyl	no cas #	EPA6 calc c	NA	NA	NA	NA	NA	NA
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	EPA6 calc c	NA	NA	NA	NA	NA	NA
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	EPA6 calc c	NA	NA	NA	NA	NA	NA
Polychlorinated biphenyls (PCBs)**	1336-36-3	RAIS b	7.31E+00	2.03E+07	RAIS b	3.09E+05	3.09E+05	RAIS b
Phthalates								
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	EPA6 e	1.60E+05	1.60E+05	EPA6 e	1.11E+09	1.11E+09	EPA6 e
Butylbenzyl phthalate	85-68-7	EPA6 e	2.59E+04	2.59E+04	EPA6 e	1.37E+04	1.37E+04	EPA6 e
Dibutyl phthalate	84-74-2	EPA6 e	5.25E+04	5.25E+04	EPA6 e	1.57E+03	1.57E+03	EPA6 e
Diethyl phthalate	84-66-2	EPA6 e	2.73E+04	2.73E+04	EPA6 e	8.20E+01	8.20E+01	EPA6 e
Dimethyl phthalate	131-11-3	EPA6 e	4.30E+01	4.30E+01	EPA6 e	2.66E+02	2.66E+02	EPA6 e
n-Dioctyl phthalate	117-84-0	EPA6 e	2.14E+09	2.14E+09	EPA6 e	9.03E+08	9.03E+08	EPA6 e
Light Polycyclic Aromatic Hydrocarbons								
2-Chloronaphthalene	91-58-7	EPA6 e	1.17E+04	1.17E+04	EPA6 e	7.14E+03	7.14E+03	EPA6 e
2-Methyl naphthalene	91-57-6	EPA6 calc c	3.86E+00	7.24E+03	Properties a	NA	6.23E+03	EPA6 calc c
5-Nitroacenaphthene	602-87-9	EPA6 calc c	NA	NA	NA	NA	NA	NA
Acenaphthene	83-32-9	EPA6 e	9.22E+03	9.22E+03	EPA6 e	4.90E+03	4.90E+03	EPA6 e
Acenaphthylene	208-96-8	RAIS b	4.07E+00	1.17E+04	Properties a	3.83E+00	6.76E+03	Properties a
Anthracene	120-12-7	EPA6 e	2.95E+04	2.95E+04	EPA6 e	2.35E+04	2.35E+04	EPA6 e
Fluorene	86-73-7	RAIS b	4.18E+00	1.51E+04	Properties a	4.15E+00	1.41E+04	Properties a
Indene	95-13-6	EPA6 calc c	NA	NA	NA	NA	NA	NA
Napthalene	91-20-3	EPA6 e	2.36E+03	2.36E+03	EPA6 e	1.19E+03	1.19E+03	EPA6 e

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Octanol / Water Partitioning Coefficients					
Constituent of Potential Concern	CAS Registry Number	Source	K_{ow} (Published Value)	K_{ow} (unitless)	Source	K_{oc} (Published Value)	K_{oc} (mL/g)	Source
Phenanthrene	85-01-8	EPA6 e	3.55E+04	3.55E+04	EPA6 e	5.01E+04	5.01E+04	EPA6 e
Pyrene	129-00-0	EPA6 e	1.00E+05	1.00E+05	EPA6 e	6.80E+04	6.80E+04	EPA6 e
Heavy Polycyclic Aromatic Hydrocarbons								
3-Methylcholanthrene	56-49-5	EPA6 calc c	7.11E+00	1.29E+07	Properties a	6.18E+00	1.51E+06	Properties a
5-Methylchrysene	3697-24-3	EPA6 calc c	NA	NA	NA	NA	NA	NA
Benzo(a)anthracene	56-55-3	EPA6 e	4.77E+05	4.77E+05	EPA6 e	2.60E+05	2.60E+05	EPA6 e
Benzo(a)pyrene	50-32-8	EPA6 e	1.35E+06	1.35E+06	EPA6 e	9.69E+05	9.69E+05	EPA6 e
Benzo(b)fluoranthene	205-99-2	EPA6 e	1.59E+06	1.59E+06	EPA6 e	8.36E+05	8.36E+05	EPA6 e
Benzo(e)pyrene	192-97-2	EPA6 calc c	7.40E+00	2.51E+07	Properties a	7.20E+00	1.58E+07	Properties a
Benzo(g,h,i)perylene	191-24-2	EPA6 calc c	7.10E+00	1.26E+07	Properties a	6.26E+00	1.82E+06	Properties a
Benzo(j)fluoranthene	205-82-3	EPA6 calc c	6.44E+00	2.75E+06	Properties a	NA	2.15E+06	EPA6 calc c
Benzo(k)fluoranthene	207-08-9	EPA6 e	1.56E+06	1.56E+06	EPA6 e	8.32E-05	8.32E-05	EPA6 e
Benzo[a,i]pyrene	191-30-0	EPA6 calc c	NA	NA	NA	NA	NA	NA
Chrysene	218-01-9	EPA6 e	5.48E+05	5.48E+05	EPA6 e	2.97E+05	2.97E+05	EPA6 e
Dibenz(a,h)anthracene	53-70-3	EPA6 e	3.53E+06	3.53E+06	EPA6 e	1.79E+06	1.79E+06	EPA6 e
Dibenz[a,h]acridine	226-36-8	EPA6 calc c	NA	NA	NA	NA	NA	NA
Dibenz[a,j]acridine	224-42-0	EPA6 calc c	NA	NA	NA	NA	NA	NA
Dibenzo(a,e)fluoranthene	5385-75-1	EPA6 calc c	NA	NA	NA	NA	NA	NA
Dibenzo(a,h)fluoranthene	no cas #	EPA6 calc c	NA	NA	NA	NA	NA	NA
Dibenzo[a,e]pyrene	192-65-4	EPA6 calc c	NA	NA	NA	NA	NA	NA
Dibenzo[a,h]pyrene	189-64-0	EPA6 calc c	NA	NA	NA	NA	NA	NA
Dibenzo[a,i]pyrene	189-55-9	EPA6 calc c	7.29E+00	1.95E+07	Chemfate i	NA	1.47E+07	EPA6 calc c
Fluoranthene	206-44-0	EPA6 e	1.21E+05	1.21E+05	EPA6 e	4.91E+04	4.91E+04	EPA6 e
Hexachloronaphthalene	1335-87-1	EPA6 calc c	7.59E+00	3.89E+07	Chemfate i	NA	2.89E+07	EPA6 calc c
Indeno(1,2,3-cd)pyrene	193-39-5	EPA6 e	8.22E+06	8.22E+06	EPA6 e	4.11E+06	4.11E+06	EPA6 e
Octachloronaphthalene	2234-13-1	EPA6 calc c	6.42E+00	2.63E+06	Chemfate i	NA	2.05E+06	EPA6 calc c
Pentachloronaphthalene	1321-64-8	EPA6 calc c	NA	NA	NA	NA	NA	NA
Tetrachloronaphthalene	1335-88-2	EPA6 calc c	NA	NA	NA	NA	NA	NA
Trichloronaphthalene	1321-65-9	EPA6 calc c	NA	NA	NA	NA	NA	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Octanol / Water Partitioning Coefficients					
Constituent of Potential Concern	CAS Registry Number	Source	K_{ow} (Published Value)	K_{ow} (unitless)	Source	K_{oc} (Published Value)	K_{oc} (mL/g)	Source
Light Substituted Benzene Compounds								
1,2,3-Trichlorobenzene	87-61-6	EPA6 e	1.11E+04	1.11E+04	EPA6 e	2.02E+04	2.02E+04	EPA6 e
1,2,4-Trichlorobenzene	120-82-1	EPA6 e	9.73E+03	9.73E+03	EPA6 e	1.66E+03	1.66E+03	EPA6 e
1,2,4-Trimethlyl benzene	25551-13-7	EPA6 calc c	3.40E+00	2.51E+03	RAIS b	5.90E+02	5.90E+02	RAIS b
1,2,4-Trimethyl benzene	95-63-6	EPA6 calc c	3.65E+00	4.47E+03	Properties a	NA	3.87E+03	EPA6 calc c
1,2-Dichlorobenzene	95-50-1	EPA6 e	2.79E+03	2.79E+03	EPA6 e	3.79E+02	3.79E+02	EPA6 e
1,3,5-Trimethyl benzene	108-67-8	EPA6 e	2.63E+03	2.63E+03	EPA6 e	1.67E+03	1.67E+03	EPA6 e
1,3-Dichlorobenzene	541-73-1	EPA6 e	3.39E+03	3.39E+03	EPA6 e	8.03E+03	8.03E+03	EPA6 e
1,3-Dinitrobenzene	99-65-0	EPA6 e	3.10E+01	3.10E+01	EPA6 e	2.06E+01	2.06E+01	EPA6 e
1,4-Dichlorobenzene	106-46-7	EPA6 e	2.58E+03	2.58E+03	EPA6 e	6.16E+02	6.16E+02	EPA6 e
1,4-Dinitrobenzene	100-25-4	EPA6 calc c	1.50E+00	3.16E+01	RAIS b	2.98E+01	2.98E+01	RAIS b
2,4,5-Trichorophenol	95-95-4	EPA6 e	7.41E+03	7.41E+03	EPA6 e	1.13E+03**	1.13E+03**	EPA6 e
2,4,6-Trichlorophenol	88-06-2	EPA6 e	5.15E+03	5.15E+03	EPA6 e	2.26E+02**	2.26E+02**	EPA6 e
2,4-Dichlorophenol	120-83-2	EPA6 e	1.09E+03	1.09E+03	EPA6 e	1.40E+02**	1.40E+02**	EPA6 e
2,4-Dimethylphenol	105-67-9	EPA6 e	2.29E+02	2.29E+02	EPA6 e	1.26E+02**	1.26E+02**	EPA6 e
2,4-Dinitrophenol	51-28-5	EPA6 e	3.30E+01	3.30E+01	EPA6 e	1.00E-02**	1.00E-02**	EPA6 e
2,4-Dinitrotoluene	121-14-2	EPA6 e	9.90E+01	9.90E+01	EPA6 e	5.10E+01	5.10E+01	EPA6 e
2,6-Dinitrotoluene	606-20-2	EPA6 e	7.70E+01	7.70E+01	EPA6 e	4.19E+01	4.19E+01	EPA6 e
2-Chlorotoluene	95-49-8	EPA6 calc c	3.54E+00	3.44E+03	RAIS b	7.56E+02	7.56E+02	RAIS b
2-Chorophenol	95-57-8	EPA6 e	1.45E+02	1.45E+02	EPA6 e	3.87E+02**	3.87E+02**	EPA6 e
2-Nitrophenol	88-75-5	EPA6 e	6.17E+01	6.17E+01	EPA6 e	3.53E+02	3.53E+02	EPA6 e
4,6-Dinitro-o-cresol	534-52-1	RAIS b	2.85E+00	7.08E+02	RAIS b	6.34E+02	6.34E+02	RAIS b
4-Chlorotoluene	106-43-4	EPA6 calc c	3.33E+00	2.14E+03	Hansch j	NA	1.88E+03	EPA6 calc c
4-Nitrophenol	100-02-7	EPA6 e	8.13E+01	8.13E+01	EPA6 e	4.37E+02	4.37E+02	EPA6 e
alpha-Methylstyrene	98-83-9	RAIS b	3.46E+00	2.91E+03	RAIS b	6.62E+02	6.62E+02	RAIS b
Aniline	62-53-3	EPA6 e	9.55E+00	9.55E+00	EPA6 e	8.23E+00	8.23E+00	EPA6 e
Benzotrichloride	98-07-7	RAIS b	2.92E+00	8.32E+02	Organics g	7.42E+02	7.42E+02	RAIS b
Benzyl Chloride	100-44-7	EPA6 e	2.30E+00	2.30E+00	EPA6 e	2.71E+00	2.71E+00	EPA6 e
Bromobenzene	108-86-1	EPA6 calc c	2.99E+00	9.77E+02	Properties a	2.65E+00	4.47E+02	Properties a

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Octanol / Water Partitioning Coefficients					
Constituent of Potential Concern	CAS Registry Number	Source	K_{ow} (Published Value)	K_{ow} (unitless)	Source	K_{oc} (Published Value)	K_{oc} (mL/g)	Source
Chlorobenzene	108-90-7	EPA6 e	6.16E+02	6.16E+02	EPA6 e	2.24E+02	2.24E+02	EPA6 e
Cumene	98-82-8	EPA6 e	4.10E+03	4.10E+03	EPA6 e	9.31E+03	9.31E+03	EPA6 e
m-Cresol	108-39-4	EPA6 e	9.10E+01	9.10E+01	EPA6 e	4.78E+01	4.78E+01	EPA6 e
n-Butyl benzene	104-51-8	EPA6 calc c	4.28E+00	1.91E+04	Properties a	3.40E+00	2.51E+03	Properties a
Nitrobenzene	98-95-3	EPA6 e	6.80E+01	6.80E+01	EPA6 e	1.19E+02	1.19E+02	EPA6 e
n-Propyl benzene	103-65-1	EPA6 calc c	3.69E+00	4.90E+03	Properties a	2.86E+00	7.24E+02	Properties a
o-Cresol	95-48-7	EPA6 e	1.05E+02	1.05E+02	EPA6 e	5.34E+01	5.34E+01	EPA6 e
o-Dinitrobenzene	528-29-0	EPA6 calc c	1.60E+00	3.98E+01	RAIS b	3.74E+01	3.74E+01	RAIS b
o-Nitroaniline	88-74-4	EPA6 e	7.08E+01	7.08E+01	EPA6 e	3.93E+02	3.93E+02	EPA6 e
o-Toluidine	95-53-4	EPA6 e	2.19E+01	2.19E+01	EPA6 e	1.57E+01	1.57E+01	EPA6 e
p-Chloroaniline	106-47-8	EPA6 e	7.40E+01	7.40E+01	EPA6 e	4.10E+01	4.10E+01	EPA6 e
p-Cresol	106-44-5	EPA6 e	8.70E+01	8.70E+01	EPA6 e	4.61E+01	4.61E+01	EPA6 e
Phenol	108-95-2	EPA6 e	3.00E+01	3.00E+01	EPA6 e	2.20E+01**	2.20E+01**	EPA6 e
p-Nitrochlorobenzene	100-00-5	EPA6 calc c	2.39E+00	2.45E+02	Chemfate i	NA	2.24E+02	EPA6 calc c
p-Toluidine	106-49-0	RAIS b	1.40E+00	2.51E+01	Properties a	NA	2.38E+01	EPA6 calc c
sec-Butyl benzene	135-98-8	EPA6 calc c	NA	NA	NA	NA	NA	NA
tert-Butyl benzene	98-06-6	EPA6 calc c	4.11E+00	1.29E+04	Properties a	NA	1.10E+04	EPA6 calc c
Toluene-2,6-diamine	823-40-5	RAIS b	1.45E+00	2.85E+01	RAIS b	2.69E+01	2.69E+01	RAIS b
Other Light Semivolatile Compounds								
1,1 Dimethyl hydrazine	57-14-7	RAIS b	5.95E-01	3.94E+00	RAIS b	3.85E+00	3.85E+00	RAIS b .
1,1-Biphenyl	92-52-4	EPA6 calc c	3.90E+00	7.94E+03	Properties a	3.40E+00	2.51E+03	Properties a
1,2 Diphenylhydrazine	122-66-7	EPA6 e	8.71E+02	8.71E+02	EPA6 e	2.78E+02	2.78E+02	EPA6 e
1,2-Dimethylhydrazine	540-73-8	EPA6 calc c	-1.37E+00	4.29E-02	Chemfate i	NA	4.52E-02	EPA6 calc c
1,3-Propane sultone	1120-71-4	EPA6 calc c	-5.23E-01	3.00E-01	Chemfate i	NA	3.06E-01	EPA6 calc c
2,4-Toluene diisocyanate	584-84-9	EPA6 calc c	NA	NA	NA	NA	NA	NA
2-Chloroacetophenone	532-27-4	RAIS b	2.59E+00	3.93E+02	RAIS b	7.60E+01	7.60E+01	RAIS b
2-Propenoic acid	79-10-7	RAIS b	4.30E-01	2.69E+00	Properties a	NA	2.65E+00	EPA6 calc c
4,4-Methylenedianiline	101-77-9	RAIS b	3.38E+00	2.40E+03	RAIS b	2.10E+03	2.10E+03	RAIS b
Acetophenone	98-86-2	EPA6 e	4.37E+01	4.37E+01	EPA6 e	2.69E+01	2.69E+01	EPA6 e

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Octanol / Water Partitioning Coefficients					
Constituent of Potential Concern	CAS Registry Number	Source	K_{ow} (Published Value)	K_{ow} (unitless)	Source	K_{oc} (Published Value)	K_{oc} (mL/g)	Source
Benzoic acid	65-85-0	RAIS b	1.86E+00	7.24E+01	RAIS b	5.50E-1**	5.50E-1**	EPA6 e
bis(2-Chloroethoxy)methane	111-91-1	NA	NA	NA	NA	NA	NA	NA
bis(2-Chloroethyl) ether	111-44-4	EPA6 e	2.00E+01	2.00E+01	EPA6 e	7.60E+01	7.60E+01	EPA6 e
Chlorocyclopentadiene	41851-50-7	EPA6 calc c	2.43E+00	2.69E+02	RAIS b	2.45E+02	2.45E+02	RAIS b
Cyclohexanol	108-93-0	EPA6 calc c	1.23E+00	1.70E+01	Properties a	1.11E+00	1.30E+01	Properties a
Dichloroisopropyl ether	108-60-1	RAIS b	2.58E+00	3.80E+02	Properties a	1.79E+00	6.10E+01	Properties a
Dichloromethyl ether	542-88-1	EPA6 calc c	-3.80E-01	4.17E-01	RAIS b	7.94E+01	7.94E+01	RAIS b
Dichloropentadiene	no cas #	EPA6 calc c	NA	NA	NA	NA	NA	NA
Dimethyl sulfate	77-78-1	RAIS b	3.18E-01	2.08E+00	RAIS b	2.06E+00	2.06E+00	RAIS b
Dimethylaniline	121-69-7	RAIS b	-8.80E-01	1.32E-01	RAIS b	1.37E-01	1.37E-01	RAIS b
di-n-propylnitrosamine	621-64-7	EPA6 e	2.40E+01	2.40E+01	EPA6 e	1.70E+01	1.70E+01	EPA6 e
Diphenyl ether	101-84-8	EPA6 calc c	4.21E+00	1.62E+04	Properties a	NA	1.38E+04	EPA6 calc c
Epichlorohydrin	106-89-8	EPA6 e	1.78E+00	1.78E+00	EPA6 e	2.22E+00	2.22E+00	EPA6 e
Ethyl carbomate (urethane)	51-79-6	EPA6 calc c	-1.50E-01	7.08E-01	Chemfate i	NA	7.12E-01	EPA6 calc c
Ethyl methane sulfanate	62-50-0	EPA6 e	1.12E+00	1.12E+00	EPA6 e	1.55E+00	1.55E+00	EPA6 e
Ethylene dibromide	106-93-4	EPA6 e	5.62E+01	5.62E+01	EPA6 e	3.28E+01	3.28E+01	EPA6 e
Ethylene glycol	107-21-1	RAIS b	-9.14E-01	1.22E-01	RAIS b	1.26E-01	1.26E-01	RAIS b
Ethylene glycol monobutyl ether	111-76-2	RAIS b	1.55E+00	3.58E+01	RAIS b	6.17E+01	6.17E+01	RAIS b
Ethylene glycol monoethyl ether acetate	111-15-9	RAIS b	6.19E-01	4.16E+00	RAIS b	4.07E+00	4.07E+00	RAIS b
Ethylene thiourea	96-45-7	RAIS b	-6.42E-01	2.28E-01	RAIS b	2.34E-01	2.34E-01	RAIS b
Furfural	98-01-1	RAIS b	9.60E-01	9.12E+00	Properties a	NA	8.79E+00	EPA6 calc c
Maleic hydrazide	123-33-1	EPA6 calc c	-7.40E-01	1.82E-01	RAIS b	1.87E-01	1.87E-01	RAIS b
Malononitrile	109-77-3	EPA6 calc c	4.00E-02	1.10E+00	RAIS b	1.29E+00	1.29E+00	RAIS b
Methyl styrene (mixed isomers)	25013-15-4	RAIS b	3.35E+00	2.24E+03	RAIS b	3.70E+02	3.70E+02	RAIS b
Methylhydrazine	60-34-4	EPA6 calc c	-1.05E+00	8.91E-02	Chemfate i	NA	9.29E-02	EPA6 calc c
N,N-Diphenylamine	122-39-4	EPA6 calc c	3.50E+00	3.16E+03	Properties a	2.54E+00	3.47E+02	Properties a
Nitric acid, propyl ester	627-13-4	EPA6 calc c	NA	NA	NA	NA	NA	NA
N-Nitrosodi-n-butylamine	924-16-3	EPA6 e	2.57E+02	2.57E+02	EPA6 e	1.07E+02	1.07E+02	EPA6 e
N-Nitrosomorpholine	59-89-2	EPA6 calc c	-4.40E-01	9.56E+00	Chemfate i	NA	9.20E+00	EPA6 calc c

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Octanol / Water Partitioning Coefficients					
Constituent of Potential Concern	CAS Registry Number	Source	K_{ow} (Published Value)	K_{ow} (unitless)	Source	K_{oc} (Published Value)	K_{oc} (mL/g)	Source
N-Nitroso-N,N-dimethylamine	62-75-9	RAIS b	-4.70E-01	3.39E-01	RAIS b	8.30E+00	8.30E+00	RAIS b
o-Anisidine	90-04-0	EPA6 calc c	1.18E+00	1.51E+01	Chemfate i	NA	1.45E+01	EPA6 calc c
Oxalic acid	144-62-7	EPA6 calc c	NA	NA	NA	NA	NA	NA
Phthalic Anhydride	85-44-9	EPA6 e	2.50E-01	2.50E-01	EPA6 e	4.80E-01	4.80E-01	EPA6 e
p-Phthalic acid	100-21-0	RAIS b	8.23E-01	6.65E+00	RAIS b	6.44E+00	6.44E+00	RAIS b
Pyridine	110-86-1	EPA6 e	4.68E+00	4.68E+00	EPA6 e	4.72E+00	4.72E+00	EPA6 e
Quinoline	91-22-5	RAIS b	2.03E+00	1.07E+02	RAIS b	9.90E+01	9.90E+01	RAIS b
Quinone	106-51-4	EPA6 calc c	2.00E-01	1.58E+00	Chemfate i	NA	1.57E+00	EPA6 calc c
Safrole	94-59-7	EPA6 e	4.57E+02	4.57E+02	EPA6 e	1.68E+02	1.68E+02	EPA6 e
Tetrahydrofuran	109-99-9	EPA6 e	2.80E+00	2.80E+00	EPA6 e	3.16E+00	3.16E+00	EPA6 e
Other Heavy Semivolatile Compounds								
1,2,4,5-Tetrachlorobenzene	95-94-3	EPA6 e	4.36E+04	4.36E+04	EPA6 e	5.89E+03	5.89E+03	EPA6 e
1,3,5-Trinitrobenzene	99-35-4	EPA6 e	1.51E+01	1.51E+01	EPA6 e	1.18E+01	1.18E+01	EPA6 e
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	EPA6 calc c	4.17E+00	1.48E+04	Chemfate i	NA	1.26E+04	EPA6 calc c
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	EPA6 calc c	-2.70E+00	2.00E-03	RAIS b	2.22E-03	2.22E-03	RAIS b
2-sec-Butyl-4,6-dinitrophenol	88-85-7	EPA6 calc c	3.14E+00	1.38E+03	RAIS b	1.22E+03	1.22E+03	RAIS b
3,3-Dichlorobenzidine	91-94-1	EPA6 e	3.76E+03	3.76E+03	EPA6 e	8.70E+02	8.70E+02	EPA6 e
3,3'-Dimethoxybenzidine	119-90-4	EPA6 e	6.46E+01	6.46E+01	EPA6 e	3.65E+01	3.65E+01	EPA6 e
4-Bromophenylphenyl ether	101-55-3	EPA6 calc c	5.00E+00	1.00E+05	RAIS b	8.23E+04	8.23E+04	RAIS b
Ammonium perfluorooctanoate	3825-26-1	EPA6 calc c	NA	NA	NA	NA	NA	NA
Azobenzene	103-33-3	RAIS b	3.82E+00	6.61E+03	RAIS b	2.69E+03	2.69E+03	RAIS b
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	EPA6 calc c	NA	NA	NA	NA	NA	NA
Captan	133-06-2	RAIS b	2.35E+00	2.24E+02	Properties a	2.30E+00	2.00E+02	Properties a
Chlorobenzilate	510-15-6	EPA6 e	2.40E+04	2.40E+04	EPA6 e	3.69E+03	3.69E+03	EPA6 e
Dibutylphosphate	107-66-4	EPA6 calc c	NA	NA	NA	NA	NA	NA
Dimethyl aminoazobenzene	60-11-7	EPA6 calc c	4.58E+00	3.80E+04	Chemfate i	NA	3.18E+04	EPA6 calc c
Hexachlorobenzene	118-74-1	EPA6 e	3.18E+05	3.18E+05	EPA6 e	8.00E+04	8.00E+04	EPA6 e
Hexachlorobutadiene	87-68-3	EPA6 e	5.38E+04	5.38E+04	EPA6 e	6.94E+03	6.94E+03	EPA6 e
Hexachlorocyclopentadiene	77-47-4	EPA6 e	8.07E-04	8.07E-04	EPA6 e	9.51E+03	9.51E+03	EPA6 e

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Octanol / Water Partitioning Coefficients					
Constituent of Potential Concern	CAS Registry Number	Source	K_{ow} (Published Value)	K_{ow} (unitless)	Source	K_{oc} (Published Value)	K_{oc} (mL/g)	Source
Hexachloroethane	67-72-1	EPA6 e	9.66E+03	9.66E+03	EPA6 e	1.82E+04	1.82E+04	EPA6 e
Hexachlorophene	70-30-4	EPA6 e	3.47E+07	3.47E+07	EPA6 e	1.08E+06	1.08E+06	EPA6 e
Hexamethylene-1,5-diisocyanate	822-06-0	RAIS b	1.27E+00	1.88E+01	RAIS b	1.79E+01	1.79E+01	RAIS b
Mirex	2385-85-5	EPA6 calc c	6.89E+00	7.76E+06	Properties a	6.00E+00	1.00E+06	Properties a
Nitrofen	1836-75-5	EPA6 calc c	5.53E+00	3.39E+05	Chemfate i	NA	2.73E+05	EPA6 calc c
Pentachlorobenzene	608-93-5	EPA6 e	1.22E+05	1.22E+05	EPA6 e	3.21E+04	3.21E+04	EPA6 e
Pentachloronitrobenzene	82-68-8	EPA6 e	4.37E+04	4.37E+04	EPA6 e	5.89E+03	5.89E+03	EPA6 e
Pentachlorophenol*	87-86-5	EPA6 e	1.20E+05	1.20E+05	EPA6 e	1.99E+04	1.99E+04	EPA6 e
Pentachlorophenol*	87-86-5	EPA6 e	1.20E+05	1.20E+05	EPA6 e	1.96E+04	1.96E+04	EPA6 e
Pentachlorophenol*	87-86-5	EPA6 e	1.20E+05	1.20E+05	EPA6 e	1.69E+04	1.69E+04	EPA6 e
Pentachlorophenol*	87-86-5	EPA6 e	1.20E+05	1.20E+05	EPA6 e	7.33E+03	7.33E+03	EPA6 e
Pentachlorophenol*	87-86-5	EPA6 e	1.20E+05	1.20E+05	EPA6 e	1.42E+03	1.42E+03	EPA6 e
Pentachlorophenol*	87-86-5	EPA6 e	1.20E+05	1.20E+05	EPA6 e	5.05E+02	5.05E+02	EPA6 e
Pentachlorophenol*	87-86-5	EPA6 e	1.20E+05	1.20E+05	EPA6 e	4.09E+02	4.09E+02	EPA6 e
Pentachlorophenol*	87-86-5	EPA6 e	1.20E+05	1.20E+05	EPA6 e	3.99E+02	3.99E+02	EPA6 e
Pentachlorophenol*	87-86-5	EPA6 e	1.20E+05	1.20E+05	EPA6 e	3.98E+02	3.98E+02	EPA6 e
Pentachlorophenol*	87-86-5	EPA6 e	1.20E+05	1.20E+05	EPA6 e	3.98E+02	3.98E+02	EPA6 e
Pentachlorophenol*	87-86-5	EPA6 e	1.20E+05	1.20E+05	EPA6 e	3.98E+02	3.98E+02	EPA6 e
Pentachlorophenol*	87-86-5	EPA6 e	1.20E+05	1.20E+05	EPA6 e	3.98E+02	3.98E+02	EPA6 e
Pentachlorophenol*	87-86-5	EPA6 e	1.20E+05	1.20E+05	EPA6 e	1.99E+04	1.99E+04	EPA6 e
Picric acid	88-89-1	EPA6 calc c	2.03E+00	1.07E+02	Chemfate i	NA	9.90E+01	EPA6 calc c
Pronamide	23950-58-5	EPA6 e	3.24E+03	3.24E+03	EPA6 e	7.74E+02	7.74E+02	EPA6 e
Strychnine	57-24-9	EPA6 e	8.51E+01	8.51E+01	EPA6 e	4.53E+02	4.53E+02	EPA6 e
Terphenyls	26140-60-3	EPA6 calc c	NA	NA	NA	NA	NA	NA
Tributyl phosphate	126-73-8	EPA6 calc c	4.00E+00	1.00E+04	Chemfate i	NA	8.55E+03	EPA6 calc c
Trifluralin	1582-09-8	RAIS b	5.34E+00	2.19E+05	Properties a	3.78E+00	6.03E+03	Properties a
Triphenylamine	603-34-9	EPA6 calc c	NA	NA	NA	NA	NA	NA
Herbicides and Organopesticides								
2,4,5-T	93-76-5	RAIS b	3.36E+00	2.29E+03	Properties a	1.72E+00	5.25E+01	Properties a

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Octanol / Water Partitioning Coefficients					
Constituent of Potential Concern	CAS Registry Number	Source	K_{ow} (Published Value)	K_{ow} (unitless)	Source	K_{oc} (Published Value)	K_{oc} (mL/g)	Source
2,4-D and esters	94-75-7	RAIS b	2.81E+00	6.46E+02	Properties a	1.30E+00	2.00E+01	Properties a
4,4-DDD	72-54-8	RAIS b	6.20E+00	1.58E+06	Properties a	5.00E+00	1.00E+05	Properties a
4,4-DDT	50-29-3	RAIS b	6.00E+00	1.00E+06	Properties a	5.38E+00	2.40E+05	Properties a
Aldrin	309-00-2	EPA6 e	1.51E+06	1.51E+06	EPA6 e	4.87E+04	4.87E+04	EPA6 e
alpha-BHC	319-84-6	EPA6 e	6.30E+03	6.30E+03	EPA6 e	1.76E+03	1.76E+03	EPA6 e
beta-BHC	319-85-7	EPA6 e	6.81E+03	6.81E+03	EPA6 e	2.14E+03	2.14E+03	EPA6 e
Chlordane	57-74-9	EPA6 e	8.66E+05	8.66E+05	EPA6 e	5.13E+04	5.13E+04	EPA6 e
DDE	72-55-9	EPA6 e	1.80E+06	1.80E+06	EPA6 e	8.64E+02	8.64E+02	EPA6 e
Delta-BHC	319-86-8	EPA6 calc c	4.14E+00	1.38E+04	Properties a	2.82E+00	6.61E+02	Properties a
Dieldrin	60-57-1	EPA6 e	1.86E+05	1.86E+05	EPA6 e	2.55E+04	2.55E+04	EPA6 e
Endothall	145-73-3	EPA6 calc c	-8.70E-01	1.35E-01	Organics g	NA	1.40E-01	EPA6 calc c
Endrin	72-20-8	EPA6 e	7.79E+04	7.79E+04	EPA6 e	1.08E+08	1.08E+08	EPA6 e
gamma-BHC(Lindane)	58-89-9	EPA6 calc c	3.72E+00	5.25E+03	Properties a	3.03E+00	1.07E+03	Properties a
Heptachlor	76-44-8	EPA6 e	1.04E+05	1.04E+05	EPA6 e	9.53E+03	9.53E+03	EPA6 e
Isodrin	465-73-6	EPA6 calc c	NA	NA	NA	NA	NA	NA
Methoxychlor	72-43-5	EPA6 e	3.36E+04	3.36E+04	EPA6 e	8.00E+04	8.00E+04	EPA6 e
Silvex (2,4,5-TP)	93-72-1	RAIS b	4.07E+00	1.18E+04	RAIS b	1.01E+04	1.01E+04	RAIS b
Toxaphene	8001-35-2	RAIS b	5.50E+00	3.16E+05	Properties a	5.00E+00	1.00E+05	Properties a

NA = Value not available
** = Varies with pH; values given for a pH of 7.0.
a = Illustrated Handbook of Physical - Chemical Properties and Environmental Fate f
b = RAIS database: http://risk.lsd.ornl.gov/rap_hp.shtml
c = Equations derived from EPA Region 6 Guidance Appendix A-3
d = Handbook of Environmental degradation Rates. 1991.
e = EPA Region 6 Hazardous Waste Combustion Guidance
f = "A Critical Review of Henry's Law Constants for Chemicals of Environmental In
g = Handbook of Environmental Fate and Exposure Data for Organic Chemicals
h = http://www.epa.gov/region09/waste/sfund/prg/.
i = http://esc.syres.com/efdb/Chemfate.htm
j = Hansch and Leo 1985 in Syracuse 1996

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Octanol / Water Partitioning Coefficients					
Constituent of Potential Concern	CAS Registry Number	Source	K_{ow} (Published Value)	K_{ow} (unitless)	Source	K_{oc} (Published Value)	K_{oc} (mL/g)	Source

k = Schwarzenbach et al. 1993
l = Tomlin 1994 in Syracuse 1996
m = http://www.speclab.com/compound/c959988.htm

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Partitioning Coefficients						
Constituent of Potential Concern	CAS Registry Number	Kd_s (cm^3/g)	Source	Kd_{sw} (L/kg)	Source	Kd_{bs} (cm^3/g)	Source	
Aromatic Halogenated Hydrocarbons								
2,3,4,6-Tetrachlorophenol	58-90-2	6.61E+01	EPA6 calc c	4.96E+02	EPA6 calc c	2.64E+02	EPA6 calc c	
4-Chloro-3-methylphenol	59-50-7	6.03E+00	EPA6 calc c	4.52E+01	EPA6 calc c	2.41E+01	EPA6 calc c	
Aromatic Nonhalogenated Hydrocarbons								
2-Nitrotoluene	88-72-2	4.27E+00	EPA6 calc c	3.20E+01	EPA6 calc c	1.71E+01	EPA6 calc c	
4-Nitrobiphenyl	92-93-3	NA	NA	NA	NA	NA	NA	
Benzaldehyde	100-52-7	2.01E-01	EPA6 e	1.51E+00	EPA6 e	8.04E-01	EPA6 e	
Benzene	71-43-2	6.20E-01	EPA6 e	4.65E+00	EPA6 e	2.48E+00	EPA6 e	
Benzyl alcohol	100-51-6	1.02E-01	EPA6 e	7.66E-01	EPA6 e	4.09E-01	EPA6 e	
Ethyl Benzene	100-41-4	2.04E+00	EPA6 e	1.53E+01	EPA6 e	8.16E+00	EPA6 e	
m-Xylene	108-38-3	1.96E+00	EPA6 e	1.47E+01	EPA6 e	7.84E+00	EPA6 e	
o-Xylene	95-47-6	2.41E+00	EPA6 e	1.81E+01	EPA6 e	9.64E+00	EPA6 e	
p-Xylene	106-42-3	3.11E+00	EPA6 e	2.33E+01	EPA6 e	1.24E+01	EPA6 e	
Styrene	100-42-5	9.12E+00	EPA6 e	6.84E+01	EPA6 e	3.65E+01	EPA6 e	
Toluene	108-88-3	1.40E+00	EPA6 e	1.05E+01	EPA6 e	5.60E+00	EPA6 e	
Non-aromatic Nonhalogenated Hydrocarbons								
1,2-Epoxybutane	106-88-7	7.94E-02	EPA6 calc c	5.96E-01	EPA6 calc c	3.18E-01	EPA6 calc c	
1,3-Butadiene	106-99-0	9.04E-01	EPA6 calc c	6.78E+00	EPA6 calc c	3.62E+00	EPA6 calc c	
1,4-Dioxane	123-91-1	8.76E-03	EPA6 e	6.57E-02	EPA6 e	3.50E-02	EPA6 e	
1-Methylpropyl alcohol	78-92-2	3.98E-02	EPA6 calc c	2.98E-01	EPA6 calc c	1.59E-01	EPA6 calc c	
1-Nitropropane	108-03-2	7.17E-02	EPA6 calc c	5.38E-01	EPA6 calc c	2.87E-01	EPA6 calc c	
2,2,4-Trimethlylpentane	540-84-1	8.61E+02	EPA6 calc c	6.46E+03	EPA6 calc c	3.44E+03	EPA6 calc c	
2-Butanone	78-93-3	2.34E-02	EPA6 e	1.76E-01	EPA6 e	9.36E-02	EPA6 e	
2-Butenaldehyde (2-Butenal)	4170-30-3	NA	NA	NA	NA	NA	NA	
2-Ethoxyethanol	110-80-5	2.09E-01	EPA6 calc c	1.57E+00	EPA6 calc c	8.36E-01	EPA6 calc c	
2-Heptanone	110-43-0	8.84E-01	EPA6 calc c	6.63E+00	EPA6 calc c	3.54E+00	EPA6 calc c	
2-Hexanone	591-78-6	1.34E+00	EPA6 calc c	1.00E+01	EPA6 calc c	5.36E+00	EPA6 calc c	
2-Methoxyethanol	109-86-4	8.53E-02	EPA6 calc c	6.40E-01	EPA6 calc c	3.41E-01	EPA6 calc c	
2-Methyl-2-propanol	75-65-0	3.72E-01	EPA6 calc c	2.79E+00	EPA6 calc c	1.49E+00	EPA6 calc c	

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Partitioning Coefficients						
Constituent of Potential Concern	CAS Registry Number	Kd_s (cm^3/g)	Source	Kd_{sw} (L/kg)	Source	Kd_{bs} (cm^3/g)	Source	
2-Methyl-2-propenenitrile	126-98-7	3.74E-02	EPA6 e	2.80E-01	EPA6 e	1.49E-01	EPA6 e	
2-Methylaziridine	75-55-8	NA	NA	NA	NA	NA	NA	
2-Methylpropyl alcohol	78-83-1	5.59E-02	EPA6 calc c	4.19E-01	EPA6 calc c	2.24E-01	EPA6 calc c	
2-Pentanone	107-87-9	7.40E-01	EPA6 calc c	5.55E+00	EPA6 calc c	2.96E+00	EPA6 calc c	
2-Propanone (Acetone)	67-64-1	9.51E-03	EPA6 e	7.13E-02	EPA6 e	3.80E-02	EPA6 e	
2-Propene-1-ol	107-18-6	1.47E-02	EPA6 calc c	1.10E-01	EPA6 calc c	5.88E-02	EPA6 calc c	
2-Propyl alcohol	67-63-0	1.12E-02	EPA6 calc c	8.40E-02	EPA6 calc c	4.48E-02	EPA6 calc c	
3-Heptanone	106-35-4	NA	NA	NA	NA	NA	NA	
3-Methyl-1-butanol	123-51-3	NA	NA	NA	NA	NA	NA	
3-Methyl-2-butanone	563-80-4	NA	NA	NA	NA	NA	NA	
3-Pentanone	96-22-0	1.20E-01	EPA6 calc c	9.02E-01	EPA6 calc c	4.81E-01	EPA6 calc c	
4-Heptanone	123-19-3	NA	NA	NA	NA	NA	NA	
4-Methyl-2-pentanone	108-10-1	1.20E-01	EPA6 e	9.00E-01	EPA6 e	4.80E-01	EPA6 e	
4-Methyl-3-penten-2-one	141-79-7	NA	NA	NA	NA	NA	NA	
5-Methyl-2-hexanone	110-12-3	NA	NA	NA	NA	NA	NA	
Acetaldehyde	75-07-0	9.53E-03	EPA6 e	7.15E-02	EPA6 e	3.81E-02	EPA6 e	
Acetamide	60-35-5	2.82E-04	EPA6 calc c	2.11E-03	EPA6 calc c	1.13E-03	EPA6 calc c	
Acetic acid	64-19-7	1.00E-02	EPA6 calc c	7.50E-02	EPA6 calc c	4.00E-02	EPA6 calc c	
Acetic acid ethyl ester	141-78-6	2.30E-02	EPA6 calc c	1.72E-01	EPA6 calc c	9.18E-02	EPA6 calc c	
Acetic acid n-butyl ester	123-86-4	5.02E-01	EPA6 calc c	3.77E+00	EPA6 calc c	2.01E+00	EPA6 calc c	
Acetonitrile	75-05-8	7.69E-03	EPA6 e	5.76E-02	EPA6 e	3.07E-02	EPA6 e	
Acrolein	107-02-8	1.39E-02	EPA6 e	1.05E-01	EPA6 e	5.57E-02	EPA6 e	
Acrylonitrile	107-13-1	2.22E-02	EPA6 e	1.66E-01	EPA6 e	8.88E-02	EPA6 e	
Bis(isopropyl)ether	108-20-3	NA	NA	NA	NA	NA	NA	
Butane	106-97-8	6.94E+00	EPA6 calc c	5.20E+01	EPA6 calc c	2.77E+01	EPA6 calc c	
Carbon disulfide	75-15-0	5.14E-01	EPA6 e	3.86E+00	EPA6 e	2.06E+00	EPA6 e	
Cyanogen	460-19-5	5.22E-02	EPA6 calc c	3.92E-01	EPA6 calc c	2.09E-01	EPA6 calc c	
Cyclohexane	110-82-7	4.79E+00	EPA6 calc c	3.59E+01	EPA6 calc c	1.91E+01	EPA6 calc c	
Cyclohexanone	108-94-1	6.26E-02	EPA6 calc c	4.69E-01	EPA6 calc c	2.50E-01	EPA6 calc c	

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Partitioning Coefficients						
Constituent of Potential Concern	CAS Registry Number	Kd_s (cm^3/g)	Source	Kd_{sw} (L/kg)	Source	Kd_{bs} (cm^3/g)	Source	
Cyclohexene	110-83-8	6.48E+00	EPA6 calc c	4.86E+01	EPA6 calc c	2.59E+01	EPA6 calc c	
Cyclopentane	287-92-3	8.90E+00	EPA6 calc c	6.67E+01	EPA6 calc c	3.56E+01	EPA6 calc c	
Ethyl alcohol	64-17-5	4.96E-03	EPA6 calc c	3.72E-02	EPA6 calc c	1.98E-02	EPA6 calc c	
Ethyl ether	60-29-7	7.50E-02	EPA6 calc c	5.63E-01	EPA6 calc c	3.00E-01	EPA6 calc c	
Ethyl methacrylate	97-63-2	2.46E-01	EPA6 e	1.85E+00	EPA6 e	9.80E-01	EPA6 e	
Formaldehyde	50-00-0	2.62E-02	EPA6 e	1.96E-01	EPA6 e	1.05E-01	EPA6 e	
Formamide	75-12-7	3.28E-04	EPA6 calc c	2.46E-03	EPA6 calc c	1.31E-03	EPA6 calc c	
Formic acid	64-18-6	5.39E-02	EPA6 e	4.04E-01	EPA6 e	2.16E-01	EPA6 e	
Formic acid, methyl ester	107-31-3	5.50E-03	EPA6 calc c	4.13E-02	EPA6 calc c	2.20E-02	EPA6 calc c	
Glycidylaldehyde	765-34-4	1.00E-03	EPA6 calc c	7.50E-03	EPA6 calc c	4.00E-03	EPA6 calc c	
Methyl acetate	79-20-9	4.80E-02	EPA6 calc c	3.60E-01	EPA6 calc c	1.92E-01	EPA6 calc c	
Methyl alcohol	67-56-1	3.96E-03	EPA6 e	2.97E-02	EPA6 e	1.58E-02	EPA6 e	
Methyl isocyanate	624-83-9	NA	NA	NA	NA	NA	NA	
Methyl methacrylate	80-62-6	6.31E-01	EPA6 calc c	4.73E+00	EPA6 calc c	2.52E+00	EPA6 calc c	
Methyl tert-butyl ether	1634-04-4	8.40E-02	EPA6 calc c	6.30E-01	EPA6 calc c	3.36E-01	EPA6 calc c	
Methylacetylene	74-99-7	8.40E-02	EPA6 calc c	6.30E-01	EPA6 calc c	3.36E-01	EPA6 calc c	
Methylcyclohexane	108-87-2	1.07E+02	EPA6 calc c	8.05E+02	EPA6 calc c	4.29E+02	EPA6 calc c	
N,N-Dimethylacetamide	127-19-5	NA	NA	NA	NA	NA	NA	
n-Butyl alcohol	71-36-3	7.33E-02	EPA6 calc c	5.50E-01	EPA6 calc c	2.93E-01	EPA6 calc c	
n-Heptane	142-82-5	3.81E+02	EPA6 calc c	2.86E+03	EPA6 calc c	1.52E+03	EPA6 calc c	
n-Hexane	110-54-3	1.10E+02	EPA6 calc c	8.23E+02	EPA6 calc c	4.39E+02	EPA6 calc c	
Nitromethane	75-52-5	4.53E-03	EPA6 calc c	3.40E-02	EPA6 calc c	1.81E-02	EPA6 calc c	
n-Nonane	111-84-2	3.58E+03	EPA6 calc c	2.69E+04	EPA6 calc c	1.43E+04	EPA6 calc c	
n-Octane	111-65-9	8.55E+01	EPA6 calc c	6.42E+02	EPA6 calc c	3.42E+02	EPA6 calc c	
n-Pentane	109-66-0	1.43E+01	EPA6 calc c	1.07E+02	EPA6 calc c	5.72E+01	EPA6 calc c	
n-Propionaldehyde	123-38-6	3.80E-02	EPA6 calc c	2.85E-01	EPA6 calc c	1.52E-01	EPA6 calc c	
n-Propyl alcohol	71-23-8	1.76E-02	EPA6 calc c	1.32E-01	EPA6 calc c	7.05E-02	EPA6 calc c	
n-Valeraldehyde	110-62-3	NA	NA	NA	NA	NA	NA	
Oxirane	75-21-8	8.26E-03	EPA6 e	6.19E-02	EPA6 e	3.30E-02	EPA6 e	

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Partitioning Coefficients						
Constituent of Potential Concern	CAS Registry Number	Kd_s (cm^3/g)	Source	Kd_{sw} (L/kg)	Source	Kd_{bs} (cm^3/g)	Source	
p-Cymene	99-87-6	1.07E+02	EPA6 calc c	8.05E+02	EPA6 calc c	4.29E+02	EPA6 calc c	
Phosgene	75-44-5	NA	NA	NA	NA	NA	NA	
Propargyl alcohol	107-19-7	NA	NA	NA	NA	NA	NA	
Propionic acid	79-09-4	2.11E-02	EPA6 calc c	1.58E-01	EPA6 calc c	8.45E-02	EPA6 calc c	
Propionitrile	107-12-0	1.44E-02	EPA6 calc c	1.08E-01	EPA6 calc c	5.75E-02	EPA6 calc c	
Propylene glycol monomethyl ether	107-98-2	1.26E-01	EPA6 calc c	9.44E-01	EPA6 calc c	5.04E-01	EPA6 calc c	
p-tert-Butyltoluene	98-51-1	NA	NA	NA	NA	NA	NA	
Triethylamine	121-44-8	NA	NA	NA	NA	NA	NA	
Trimethylamine	75-50-3	4.00E-02	EPA6 calc c	3.00E-01	EPA6 calc c	1.60E-01	EPA6 calc c	
Vinyl acetate	108-05-4	4.97E-02	EPA6 e	3.73E-01	EPA6 e	1.99E-01	EPA6 e	
Non-aromatic Halogenated Hydrocarbons								
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	NA	NA	NA	NA	NA	NA	
1,1,1,2-Tetrachloroethane	630-20-6	1.59E+00	EPA6 e	1.20E+01	EPA6 e	6.37E+00	EPA6 e	
1,1,1-Trichloroethane	71-55-6	1.35E+03	EPA6 e	1.01E+04	EPA6 e	5.40E+03	EPA6 e	
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	3.16E+00	EPA6 calc c	2.37E+01	EPA6 calc c	1.26E+01	EPA6 calc c	
1,1,2,2-Tetrachloroethane	79-34-5	7.90E-01	EPA6 e	5.93E+00	EPA6 e	3.16E+00	EPA6 e	
1,1,2,2-Tetrachloroethene	127-18-4	2.65E+00	EPA6 e	1.99E+01	EPA6 e	1.06E+01	EPA6 e	
1,1,2-Trichloro-1,2,2-trifluoroethane	76-13-1	2.57E+00	EPA6 calc c	1.93E+01	EPA6 calc c	1.03E+01	EPA6 calc c	
1,1,2-Trichloroethane	79-00-5	7.50E-01	EPA6 e	5.63E+00	EPA6 e	3.00E+00	EPA6 e	
1,1,2-Trichloroethylene	79-01-6	9.40E-01	EPA6 e	7.05E+00	EPA6 e	3.76E+00	EPA6 e	
1,1-Dichloroethane	75-34-3	5.30E-01	EPA6 e	3.98E+00	EPA6 e	2.12E+00	EPA6 e	
1,1-Dichloroethene	75-35-4	6.50E-01	EPA6 e	4.88E+00	EPA6 e	2.60E+00	EPA6 e	
1,2,3-Trichloropropane	96-18-4	8.10E-01	EPA6 e	6.04E+00	EPA6 e	3.22E+00	EPA6 e	
1,2-dibromo-3-chloropropane	96-12-8	9.47E-01	EPA6 e	7.10E+00	EPA6 e	3.79E+00	EPA6 e	
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	NA	NA	NA	NA	NA	NA	
1,2-Dichloroethane	107-06-2	1.96E-01	EPA6 e	1.47E+00	EPA6 e	7.83E-01	EPA6 e	
1,2-Dichloroethylene	540-59-0	2.87E-02	EPA6 calc c	2.16E-01	EPA6 calc c	1.15E-01	EPA6 calc c	
1,2-Dichloropropane	78-87-5	4.70E-01	EPA6 e	3.53E+00	EPA6 e	1.88E+00	EPA6 e	
1,3-Dichloropropene	542-75-6	2.70E-01	EPA6 e	2.03E+00	EPA6 e	1.08E+00	EPA6 e	

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Partitioning Coefficients						
Constituent of Potential Concern	CAS Registry Number	Kd_s (cm^3/g)	Source	Kd_{sw} (L/kg)	Source	Kd_{bs} (cm^3/g)	Source	
1,4-Dichloro-2-butene	764-41-0	1.82E+00	EPA6 calc c	1.36E+01	EPA6 calc c	7.28E+00	EPA6 calc c	
1-Chloroethene	75-01-4	1.11E-01	EPA6 e	8.32E-01	EPA6 e	4.44E-01	EPA6 e	
2,2-Dichloropropionic acid	75-99-0	5.82E-02	EPA6 calc c	4.37E-01	EPA6 calc c	2.33E-01	EPA6 calc c	
2-Chloropropane	75-29-6	7.38E-01	EPA6 calc c	5.53E+00	EPA6 calc c	2.95E+00	EPA6 calc c	
3-Chloropropene (allyl chloride)	107-05-1	4.79E-01	EPA6 calc c	3.59E+00	EPA6 calc c	1.91E+00	EPA6 calc c	
Bromochloromethane	74-97-5	2.43E-01	EPA6 calc c	1.83E+00	EPA6 calc c	9.73E-01	EPA6 calc c	
Bromodichloromethane	75-27-4	5.38E-01	EPA6 e	4.03E+00	EPA6 e	2.15E+00	EPA6 e	
Bromoethene	593-60-2	1.51E+00	EPA6 calc c	1.14E+01	EPA6 calc c	6.05E+00	EPA6 calc c	
Bromoform	75-25-2	1.26E+00	EPA6 e	9.45E+00	EPA6 e	5.04E+00	EPA6 e	
Bromomethane	74-83-9	9.00E-02	EPA6 e	6.75E-01	EPA6 e	3.60E-01	EPA6 e	
Carbon tetrachloride	56-23-5	1.52E+00	EPA6 e	1.14E+01	EPA6 e	6.08E+00	EPA6 e	
Chlordibromomethane	124-48-1	7.05E-01	EPA6 e	5.29E+00	EPA6 e	2.82E+00	EPA6 e	
Chlorodifluoromethane	75-45-6	9.83E-02	EPA6 e	9.83E-04	EPA6 e	3.93E-01	EPA6 e	
Chloroethane	75-00-3	3.71E+00	EPA6 e	2.78E+01	EPA6 e	1.48E+01	EPA6 e	
Chloroform	67-66-3	5.30E+01	EPA6 e	3.98E+00	EPA6 e	2.12E+00	EPA6 e	
Chloromethane	74-87-3	6.00E-02	EPA6 e	4.50E-01	EPA6 e	2.40E-01	EPA6 e	
Chloropentafluoroethane	76-15-3	NA	NA	NA	NA	NA	NA	
cis-1,2-Dichloroethene	156-59-2	4.98E+00	EPA6 e	3.73E+01	EPA6 e	1.99E+01	EPA6 e	
cis-1,3-Dichloropropene	10061-01-5	NA	NA	NA	NA	NA	NA	
Cyanogen bromide	506-68-3	NA	NA	NA	NA	NA	NA	
Cyanogen chloride	506-77-4	NA	NA	NA	NA	NA	NA	
Dichlorodifluoromethane	75-71-8	6.85E-01	EPA6 e	5.14E+00	EPA6 e	2.74E+00	EPA6 e	
Dichlorofluoromethane	75-43-4	NA	NA	NA	NA	NA	NA	
Dichloromethane	75-09-2	1.00E-01	EPA6 e	7.50E-01	EPA6 e	4.00E-01	EPA6 e	
Difluorodibromomethane	75-61-6	NA	NA	NA	NA	NA	NA	
Hexafluoroacetone	684-16-2	NA	NA	NA	NA	NA	NA	
Iodomethane	74-88-4	4.59E-01	EPA6 calc c	3.44E+00	EPA6 calc c	1.83E+00	EPA6 calc c	
Methylene bromide	74-95-3	2.60E-01	EPA6 e	1.95E+00	EPA6 e	1.04E+00	EPA6 e	
Pentachloroethane	76-01-7	9.96E+00	EPA6 calc c	7.47E+01	EPA6 calc c	3.98E+01	EPA6 calc c	

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Partitioning Coefficients						
Constituent of Potential Concern	CAS Registry Number	Kd_s (cm^3/g)	Source	Kd_{sw} (L/kg)	Source	Kd_{bs} (cm^3/g)	Source	
trans-1,2-Dichloroethene	156-60-5	3.80E-01	EPA6 e	2.85E+00	EPA6 e	1.52E+00	EPA6 e	
trans-1,3-Dichloropropene	10061-02-6	1.06E+00	EPA6 calc c	7.95E+00	EPA6 calc c	4.24E+00	EPA6 calc c	
Trichloroacetic acid	76-03-9	2.03E-01	EPA6 calc c	1.52E+00	EPA6 calc c	8.12E-01	EPA6 calc c	
Trichlorofluoroethane	27154-33-2	NA	NA	NA	NA	NA	NA	
Trichlorofluoromethane	75-69-4	1.34E+00	EPA6 e	1.00E+01	EPA6 e	5.34E+00	EPA6 e	
Trifluorobromomethane	75-63-8	6.74E-01	EPA6 calc c	5.05E+00	EPA6 calc c	2.70E+00	EPA6 calc c	
Dioxin and Furan Compounds								
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	35822-46-9	9.77E+05	EPA6 e	7.33E+06	EPA6 e	3.91E+06	EPA6 e	
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	5.13E+05	EPA6 e	3.85E+06	EPA6 e	2.05E+06	EPA6 e	
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	5.13E+05	EPA6 e	3.85E+06	EPA6 e	2.05E+06	EPA6 e	
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	39227-28-6	3.80E+05	EPA6 e	2.85E+06	EPA6 e	1.52E+06	EPA6 e	
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	1.10E+05	EPA6 e	8.22E+05	EPA6 e	4.39E+05	EPA6 e	
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	57653-85-7	1.10E+05	EPA6 e	8.22E+05	EPA6 e	4.39E+05	EPA6 e	
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	1.10E+05	EPA6 e	8.22E+05	EPA6 e	4.39E+05	EPA6 e	
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	19408-74-3	1.10E+05	EPA6 e	8.22E+05	EPA6 e	4.39E+05	EPA6 e	
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	1.10E+05	EPA6 e	8.22E+05	EPA6 e	4.39E+05	EPA6 e	
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	40321-76-4	2.69E+04	EPA6 e	2.02E+05	EPA6 e	1.08E+05	EPA6 e	
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	3.80E+04	EPA6 e	2.85E+05	EPA6 e	1.52E+05	EPA6 e	
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	1.10E+05	EPA6 e	8.22E+05	EPA6 e	4.39E+05	EPA6 e	
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	5.13E+04	EPA6 e	3.85E+05	EPA6 e	2.05E+05	EPA6 e	
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1746-01-6	2.69E+04	EPA6 e	2.02E+05	EPA6 e	1.08E+05	EPA6 e	
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	2.09E+04	EPA6 e	1.57E+05	EPA6 e	8.36E+04	EPA6 e	
Dibenzofuran	132-64-9	1.81E+02	EPA6 calc c	1.35E+03	EPA6 calc c	7.22E+02	EPA6 calc c	
Octachlorodibenzo(p)dioxin	3268-87-9	2.40E+05	EPA6 e	1.80E+06	EPA6 e	9.60E+05	EPA6 e	
Octachlorodibenzofuran	39001-02-0	3.72E+06	EPA6 e	2.79E+07	EPA6 e	1.49E+07	EPA6 e	
PCBs								
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	7.08E+04	EPA6 calc c	5.31E+05	EPA6 calc c	2.83E+05	EPA6 calc c	
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	8.32E+04	EPA6 calc c	6.24E+05	EPA6 calc c	3.33E+05	EPA6 calc c	
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	NA	NA	NA	NA	NA	NA	

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Partitioning Coefficients						
Constituent of Potential Concern	CAS Registry Number	Kd_s (cm^3/g)	Source	Kd_{sw} (L/kg)	Source	Kd_{bs} (cm^3/g)	Source	
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA	
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	NA	NA	NA	NA	NA	NA	
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	NA	NA	NA	NA	NA	NA	
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA	
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	NA	NA	NA	NA	NA	NA	
2',3,4,4',5-Pentachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA	
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	9.98E+04	EPA6 calc c	7.49E+05	EPA6 calc c	3.99E+05	EPA6 calc c	
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	3.98E+04	EPA6 calc c	2.99E+05	EPA6 calc c	1.59E+05	EPA6 calc c	
3,3',4,4',5-Pentachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA	
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	NA	NA	NA	NA	NA	NA	
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	NA	NA	NA	NA	NA	NA	
Polychlorinated biphenyls (PCBs)**	1336-36-3	3.09E+03	EPA6 calc c	2.32E+04	EPA6 calc c	1.24E+04	EPA6 calc c	
Phthalates								
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	1.11E+03	EPA6 e	8.33E+03	EPA6 e	4.44E+03	EPA6 e	
Butylbenzyl phthalate	85-68-7	1.37E+02	EPA6 e	1.03E+03	EPA6 e	5.50E+02	EPA6 e	
Dibutyl phthalate	84-74-2	1.57E+01	EPA6 e	1.18E+02	EPA6 e	6.27E+01	EPA6 e	
Diethyl phthalate	84-66-2	8.20E-01	EPA6 e	6.15E+00	EPA6 e	3.28E+00	EPA6 e	
Dimethyl phthalate	131-11-3	2.66E+00	EPA6 e	2.00E+01	EPA6 e	1.06E+01	EPA6 e	
n-Dioctyl phthalate	117-84-0	9.03E+06	EPA6 e	6.78E+07	EPA6 e	3.61E+07	EPA6 e	
Light Polycyclic Aromatic Hydrocarbons								
2-Chloronaphthalene	91-58-7	7.14E+01	EPA6 e	5.36E+02	EPA6 e	2.86E+02	EPA6 e	
2-Methyl naphthalene	91-57-6	6.23E+01	EPA6 calc c	4.67E+02	EPA6 calc c	2.49E+02	EPA6 calc c	
5-Nitroacenaphthene	602-87-9	NA	NA	NA	NA	NA	NA	
Acenaphthene	83-32-9	4.90E+01	EPA6 e	3.67E+02	EPA6 e	1.96E+02	EPA6 e	
Acenaphthylene	208-96-8	6.76E+01	EPA6 calc c	5.07E+02	EPA6 calc c	2.70E+02	EPA6 calc c	
Anthracene	120-12-7	2.35E+02	EPA6 e	1.76E+03	EPA6 e	9.40E+02	EPA6 e	
Fluorene	86-73-7	1.41E+02	EPA6 calc c	1.06E+03	EPA6 calc c	5.65E+02	EPA6 calc c	
Indene	95-13-6	NA	NA	NA	NA	NA	NA	
Napthalene	91-20-3	1.19E+01	EPA6 e	8.93E+01	EPA6 e	4.76E+01	EPA6 e	

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Partitioning Coefficients						
Constituent of Potential Concern	CAS Registry Number	Kd_s (cm^3/g)	Source	Kd_{sw} (L/kg)	Source	Kd_{bs} (cm^3/g)	Source	
Phenanthrene	85-01-8	5.01E+02	EPA6 e	3.76E+03	EPA6 e	2.01E+03	EPA6 e	
Pyrene	129-00-0	6.80E+02	EPA6 e	5.10E+03	EPA6 e	2.72E+03	EPA6 e	
Heavy Polycyclic Aromatic Hydrocarbons								
3-Methylcholanthrene	56-49-5	1.51E+04	EPA6 calc c	1.14E+05	EPA6 calc c	6.05E+04	EPA6 calc c	
5-Methylchrysene	3697-24-3	NA	NA	NA	NA	NA	NA	
Benzo(a)anthracene	56-55-3	2.60E+03	EPA6 e	1.95E+04	EPA6 e	1.04E+04	EPA6 e	
Benzo(a)pyrene	50-32-8	9.69E+03	EPA6 e	7.27E+04	EPA6 e	3.87E+04	EPA6 e	
Benzo(b)fluoranthene	205-99-2	8.36E+03	EPA6 e	6.27E+04	EPA6 e	3.34E+04	EPA6 e	
Benzo(e)pyrene	192-97-2	1.58E+05	EPA6 calc c	1.19E+06	EPA6 calc c	6.34E+05	EPA6 calc c	
Benzo(g,h,i)perylene	191-24-2	1.82E+04	EPA6 calc c	1.36E+05	EPA6 calc c	7.28E+04	EPA6 calc c	
Benzo(j)fluoranthene	205-82-3	2.15E+04	EPA6 calc c	1.61E+05	EPA6 calc c	8.60E+04	EPA6 calc c	
Benzo(k)fluoranthene	207-08-9	8.32E+03	EPA6 e	6.24E+04	EPA6 e	3.33E+04	EPA6 e	
Benzo[a,i]pyrene	191-30-0	NA	NA	NA	NA	NA	NA	
Chrysene	218-01-9	2.97E+03	EPA6 e	2.23E+04	EPA6 e	1.19E+04	EPA6 e	
Dibenz(a,h)anthracene	53-70-3	1.79E+04	EPA6 e	1.34E+05	EPA6 e	7.16E+04	EPA6 e	
Dibenz[a,h]acridine	226-36-8	NA	NA	NA	NA	NA	NA	
Dibenz[a,j]acridine	224-42-0	NA	NA	NA	NA	NA	NA	
Dibenzo(a,e)fluoranthene	5385-75-1	NA	NA	NA	NA	NA	NA	
Dibenzo(a,h)fluoranthene	no cas #	NA	NA	NA	NA	NA	NA	
Dibenzo[a,e]pyrene	192-65-4	NA	NA	NA	NA	NA	NA	
Dibenzo[a,h]pyrene	189-64-0	NA	NA	NA	NA	NA	NA	
Dibenzo[a,i]pyrene	189-55-9	1.47E+05	EPA6 calc c	1.10E+06	EPA6 calc c	5.87E+05	EPA6 calc c	
Fluoranthene	206-44-0	4.91E+02	EPA6 e	3.68E+03	EPA6 e	1.96E+03	EPA6 e	
Hexachloronaphthalene	1335-87-1	2.89E+05	EPA6 calc c	2.17E+06	EPA6 calc c	1.16E+06	EPA6 calc c	
Indeno(1,2,3-cd)pyrene	193-39-5	4.11E+04	EPA6 e	3.08E+05	EPA6 e	1.64E+05	EPA6 e	
Octachloronaphthalene	2234-13-1	2.05E+04	EPA6 calc c	1.54E+05	EPA6 calc c	8.19E+04	EPA6 calc c	
Pentachloronaphthalene	1321-64-8	NA	NA	NA	NA	NA	NA	
Tetrachloronaphthalene	1335-88-2	NA	NA	NA	NA	NA	NA	
Trichloronaphthalene	1321-65-9	NA	NA	NA	NA	NA	NA	

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Partitioning Coefficients						
Constituent of Potential Concern	CAS Registry Number	Kd_s (cm^3/g)	Source	Kd_{sw} (L/kg)	Source	Kd_{bs} (cm^3/g)	Source	
Light Substituted Benzene Compounds								
1,2,3-Trichlorobenzene	87-61-6	2.02E+02	EPA6 e	1.52E+03	EPA6 e	8.10E+02	EPA6 e	
1,2,4-Trichlorobenzene	120-82-1	1.66E+01	EPA6 e	1.24E+02	EPA6 e	6.64E+01	EPA6 e	
1,2,4-Trimethlyl benzene	25551-13-7	5.90E+00	EPA6 calc c	4.43E+01	EPA6 calc c	2.36E+01	EPA6 calc c	
1,2,4-Trimethyl benzene	95-63-6	3.87E+01	EPA6 calc c	2.91E+02	EPA6 calc c	1.55E+02	EPA6 calc c	
1,2-Dichlorobenzene	95-50-1	3.79E+00	EPA6 e	2.84E+01	EPA6 e	1.52E+01	EPA6 e	
1,3,5-Trimethyl benzene	108-67-8	1.67E+01	EPA6 e	1.25E+02	EPA6 e	6.69E+01	EPA6 e	
1,3-Dichlorobenzene	541-73-1	8.03E+01	EPA6 e	6.02E+02	EPA6 e	3.21E+02	EPA6 e	
1,3-Dinitrobenzene	99-65-0	2.06E-01	EPA6 e	1.55E+00	EPA6 e	8.25E-01	EPA6 e	
1,4-Dichlorobenzene	106-46-7	6.16E+00	EPA6 e	4.62E+01	EPA6 e	2.46E+01	EPA6 e	
1,4-Dinitrobenzene	100-25-4	2.98E-01	EPA6 calc c	2.24E+00	EPA6 calc c	1.19E+00	EPA6 calc c	
2,4,5-Trichorophenol	95-95-4	1.13E+01**	EPA6 e	8.45E+01**	EPA6 e	4.51E+01**	EPA6 e	
2,4,6-Trichlorophenol	88-06-2	2.26E+00**	EPA6 e	1.70E+01**	EPA6 e	9.05E+00**	EPA6 e	
2,4-Dichlorophenol	120-83-2	1.40E+00	EPA6 e	1.05E+01	EPA6 e	5.58E+00	EPA6 e	
2,4-Dimethylphenol	105-67-9	1.26E+00	EPA6 e	9.44E+00	EPA6 e	5.04E+00	EPA6 e	
2,4-Dinitrophenol	51-28-5	1.00E-04**	EPA6 e	7.50E-04**	EPA6 e	4.00E-04**	EPA6 e	
2,4-Dinitrotoluene	121-14-2	5.10E-01	EPA6 e	3.83E+00	EPA6 e	2.04E+00	EPA6 e	
2,6-Dinitrotoluene	606-20-2	4.19E-01	EPA6 e	3.14E+00	EPA6 e	1.68E+00	EPA6 e	
2-Chlorotoluene	95-49-8	7.56E+00	EPA6 calc c	5.67E+01	EPA6 calc c	3.03E+01	EPA6 calc c	
2-Chorophenol	95-57-8	3.87E+00	EPA6 e	2.90E+01	EPA6 e	1.55E+01	EPA6 e	
2-Nitrophenol	88-75-5	3.53E+00	EPA6 e	2.65E+01	EPA6 e	1.41E+01	EPA6 e	
4,6-Dinitro-o-cresol	534-52-1	6.34E+00	EPA6 calc c	4.75E+01	EPA6 calc c	2.53E+01	EPA6 calc c	
4-Chlorotoluene	106-43-4	1.88E+01	EPA6 calc c	1.41E+02	EPA6 calc c	7.51E+01	EPA6 calc c	
4-Nitrophenol	100-02-7	4.37E+00	EPA6 e	3.28E+01	EPA6 e	1.75E+01	EPA6 e	
alpha-Methylstyrene	98-83-9	6.62E+00	EPA6 calc c	4.97E+01	EPA6 calc c	2.65E+01	EPA6 calc c	
Aniline	62-53-3	8.23E-02	EPA6 e	6.17E-01	EPA6 e	3.29E-01	EPA6 e	
Benzotrichloride	98-07-7	7.42E+00	EPA6 calc c	5.57E+01	EPA6 calc c	2.97E+01	EPA6 calc c	
Benzyl Chloride	100-44-7	2.71E-02	EPA6 e	2.03E-01	EPA6 e	1.08E-01	EPA6 e	
Bromobenzene	108-86-1	4.47E+00	EPA6 calc c	3.35E+01	EPA6 calc c	1.79E+01	EPA6 calc c	

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Partitioning Coefficients						
Constituent of Potential Concern	CAS Registry Number	Kd_s (cm^3/g)	Source	Kd_{sw} (L/kg)	Source	Kd_{bs} (cm^3/g)	Source	
Chlorobenzene	108-90-7	2.24E+00	EPA6 e	1.68E+01	EPA6 e	8.96E+00	EPA6 e	
Cumene	98-82-8	9.31E+01	EPA6 e	6.98E+02	EPA6 e	3.72E+02	EPA6 e	
m-Cresol	108-39-4	4.78E-01	EPA6 e	3.58E+00	EPA6 e	1.91E+00	EPA6 e	
n-Butyl benzene	104-51-8	2.51E+01	EPA6 calc c	1.88E+02	EPA6 calc c	1.00E+02	EPA6 calc c	
Nitrobenzene	98-95-3	1.19E+00	EPA6 e	8.93E+00	EPA6 e	4.76E+04	EPA6 e	
n-Propyl benzene	103-65-1	7.24E+00	EPA6 calc c	5.43E+01	EPA6 calc c	2.90E+01	EPA6 calc c	
o-Cresol	95-48-7	5.34E-01	EPA6 e	4.00E+00	EPA6 e	2.14E+00	EPA6 e	
o-Dinitrobenzene	528-29-0	3.74E-01	EPA6 calc c	2.81E+00	EPA6 calc c	1.50E+00	EPA6 calc c	
o-Nitroaniline	88-74-4	3.93E+00	EPA6 e	2.95E+01	EPA6 e	1.57E+01	EPA6 e	
o-Toluidine	95-53-4	1.57E-01	EPA6 e	1.18E+00	EPA6 e	6.28E-01	EPA6 e	
p-Chloroaniline	106-47-8	4.06E-01	EPA6 e	3.05E+00	EPA6 e	1.63E+00	EPA6 e	
p-Cresol	106-44-5	4.61E-01	EPA6 e	3.46E+00	EPA6 e	1.84E+00	EPA6 e	
Phenol	108-95-2	2.20E-01	EPA6 e	1.65E+00	EPA6 e	8.79E-01	EPA6 e	
p-Nitrochlorobenzene	100-00-5	2.24E+00	EPA6 calc c	1.68E+01	EPA6 calc c	8.95E+00	EPA6 calc c	
p-Toluidine	106-49-0	2.38E-01	EPA6 calc c	1.78E+00	EPA6 calc c	9.52E-01	EPA6 calc c	
sec-Butyl benzene	135-98-8	NA	NA	NA	NA	NA	NA	
tert-Butyl benzene	98-06-6	1.10E+02	EPA6 calc c	8.23E+02	EPA6 calc c	4.39E+02	EPA6 calc c	
Toluene-2,6-diamine	823-40-5	2.69E-01	EPA6 calc c	2.02E+00	EPA6 calc c	1.08E+00	EPA6 calc c	
Other Light Semivolatile Compounds								
1,1 Dimethyl hydrazine	57-14-7	3.85E-02	EPA6 calc c	2.89E-01	EPA6 calc c	1.54E-01	EPA6 calc c	
1,1-Biphenyl	92-52-4	2.51E+01	EPA6 calc c	1.88E+02	EPA6 calc c	1.00E+02	EPA6 calc c	
1,2 Diphenylhydrazine	122-66-7	2.78E+00	EPA6 e	2.09E+01	EPA6 e	1.11E+01	EPA6 e	
1,2-Dimethylhydrazine	540-73-8	4.52E-04	EPA6 calc c	3.39E-03	EPA6 calc c	1.81E-03	EPA6 calc c	
1,3-Propane sultone	1120-71-4	3.06E-03	EPA6 calc c	2.30E-02	EPA6 calc c	1.23E-02	EPA6 calc c	
2,4-Toluene diisocyanate	584-84-9	NA	NA	NA	NA	NA	NA	
2-Chloroacetophenone	532-27-4	7.60E-01	EPA6 calc c	5.70E+00	EPA6 calc c	3.04E+00	EPA6 calc c	
2-Propenoic acid	79-10-7	2.65E-02	EPA6 calc c	1.99E-01	EPA6 calc c	1.06E-01	EPA6 calc c	
4,4-Methylenedianiline	101-77-9	2.10E+01	EPA6 calc c	1.58E+02	EPA6 calc c	8.40E+01	EPA6 calc c	
Acetophenone	98-86-2	2.69E-01	EPA6 e	2.02E+00	EPA6 e	1.08E+00	EPA6 e	

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Partitioning Coefficients						
Constituent of Potential Concern	CAS Registry Number	Kd_s (cm^3/g)	Source	Kd_{sw} (L/kg)	Source	Kd_{bs} (cm^3/g)	Source	
Benzoic acid	65-85-0	5.50E-03	NA	4.13E-02	NA	2.20E-02	NA	
bis(2-Chloroethoxy)methane	111-91-1	NA	NA	NA	NA	NA	NA	
bis(2-Chloroethyl) ether	111-44-4	7.60E-01	EPA6 e	5.70E+00	EPA6 e	3.04E+00	EPA6 e	
Chlorocyclopentadiene	41851-50-7	2.45E+00	EPA6 calc c	1.84E+01	EPA6 calc c	9.80E+00	EPA6 calc c	
Cyclohexanol	108-93-0	1.30E-01	EPA6 calc c	9.75E-01	EPA6 calc c	5.20E-01	EPA6 calc c	
Dichloroisopropyl ether	108-60-1	6.10E-01	EPA6 calc c	4.57E+00	EPA6 calc c	2.44E+00	EPA6 calc c	
Dichloromethyl ether	542-88-1	7.94E-01	EPA6 calc c	5.96E+00	EPA6 calc c	3.18E+00	EPA6 calc c	
Dichloropentadiene	no cas #	NA	NA	NA	NA	NA	NA	
Dimethyl sulfate	77-78-1	2.06E-02	EPA6 calc c	1.54E-01	EPA6 calc c	8.23E-02	EPA6 calc c	
Dimethylaniline	121-69-7	1.37E-03	EPA6 calc c	1.02E-02	EPA6 calc c	5.46E-03	EPA6 calc c	
di-n-propylnitrosamine	621-64-7	1.70E-01	EPA6 e	1.28E+00	EPA6 e	6.80E-01	EPA6 e	
Diphenyl ether	101-84-8	1.38E+02	EPA6 calc c	1.03E+03	EPA6 calc c	5.50E+02	EPA6 calc c	
Epichlorohydrin	106-89-8	2.22E-02	EPA6 e	1.66E-01	EPA6 e	8.88E-02	EPA6 e	
Ethyl carbomate (urethane)	51-79-6	7.12E-03	EPA6 calc c	5.34E-02	EPA6 calc c	2.85E-02	EPA6 calc c	
Ethyl methane sulfanate	62-50-0	1.55E-02	EPA6 e	1.16E-01	EPA6 e	6.19E-02	EPA6 e	
Ethylene dibromide	106-93-4	3.28E-01	EPA6 e	2.46E+00	EPA6 e	1.31E+00	EPA6 e	
Ethylene glycol	107-21-1	1.26E-03	EPA6 calc c	9.47E-03	EPA6 calc c	5.05E-03	EPA6 calc c	
Ethylene glycol monobutyl ether	111-76-2	6.17E-01	EPA6 calc c	4.62E+00	EPA6 calc c	2.47E+00	EPA6 calc c	
Ethylene glycol monoethyl ether acetate	111-15-9	4.07E-02	EPA6 calc c	3.05E-01	EPA6 calc c	1.63E-01	EPA6 calc c	
Ethylene thiourea	96-45-7	2.34E-03	EPA6 calc c	1.76E-02	EPA6 calc c	9.37E-03	EPA6 calc c	
Furfural	98-01-1	8.79E-02	EPA6 calc c	6.59E-01	EPA6 calc c	3.52E-01	EPA6 calc c	
Maleic hydrazide	123-33-1	1.87E-03	EPA6 calc c	1.41E-02	EPA6 calc c	7.50E-03	EPA6 calc c	
Malononitrile	109-77-3	1.29E-02	EPA6 calc c	9.66E-02	EPA6 calc c	5.15E-02	EPA6 calc c	
Methyl styrene (mixed isomers)	25013-15-4	3.70E+00	EPA6 calc c	2.78E+01	EPA6 calc c	1.48E+01	EPA6 calc c	
Methylhydrazine	60-34-4	9.29E-04	EPA6 calc c	6.97E-03	EPA6 calc c	3.72E-03	EPA6 calc c	
N,N-Diphenylamine	122-39-4	3.47E+00	EPA6 calc c	2.60E+01	EPA6 calc c	1.39E+01	EPA6 calc c	
Nitric acid, propyl ester	627-13-4	NA	NA	NA	NA	NA	NA	
N-Nitrosodi-n-butylamine	924-16-3	1.07E+00	EPA6 e	8.05E+00	EPA6 e	4.29E+00	EPA6 e	
N-Nitrosomorpholine	59-89-2	9.20E-02	EPA6 calc c	6.90E-01	EPA6 calc c	3.68E-01	EPA6 calc c	

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Partitioning Coefficients						
Constituent of Potential Concern	CAS Registry Number	Kd_s (cm^3/g)	Source	Kd_{sw} (L/kg)	Source	Kd_{bs} (cm^3/g)	Source	
N-Nitroso-N,N-dimethylamine	62-75-9	8.30E-02	EPA6 calc c	6.22E-01	EPA6 calc c	3.32E-01	EPA6 calc c	
o-Anisidine	90-04-0	1.45E-01	EPA6 calc c	1.08E+00	EPA6 calc c	5.78E-01	EPA6 calc c	
Oxalic acid	144-62-7	NA	NA	NA	NA	NA	NA	
Phthalic Anhydride	85-44-9	4.80E-03	EPA6 e	3.60E-02	EPA6 e	1.92E-02	EPA6 e	
p-Phthalic acid	100-21-0	6.44E-02	EPA6 calc c	4.83E-01	EPA6 calc c	2.58E-01	EPA6 calc c	
Pyridine	110-86-1	4.72E-02	EPA6 e	3.54E-01	EPA6 e	1.89E-01	EPA6 e	
Quinoline	91-22-5	9.90E-01	EPA6 calc c	7.43E+00	EPA6 calc c	3.96E+00	EPA6 calc c	
Quinone	106-51-4	1.57E-02	EPA6 calc c	1.18E-01	EPA6 calc c	6.29E-02	EPA6 calc c	
Safrole	94-59-7	1.68E+00	EPA6 e	1.26E+01	EPA6 e	6.73E+00	EPA6 e	
Tetrahydrofuran	109-99-9	3.16E-02	EPA6 e	2.37E-01	EPA6 e	1.26E-01	EPA6 e	
Other Heavy Semivolatile Compounds								
1,2,4,5-Tetrachlorobenzene	95-94-3	5.89E+01	EPA6 e	4.42E+02	EPA6 e	2.36E+02	EPA6 e	
1,3,5-Trinitrobenzene	99-35-4	1.18E-01	EPA6 e	8.84E-01	EPA6 e	4.72E-01	EPA6 e	
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	1.26E+02	EPA6 calc c	9.43E+02	EPA6 calc c	5.03E+02	EPA6 calc c	
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	2.22E-05	EPA6 calc c	1.66E-04	EPA6 calc c	8.88E-05	EPA6 calc c	
2-sec-Butyl-4,6-dinitrophenol	88-85-7	1.22E+01	EPA6 calc c	9.16E+01	EPA6 calc c	4.89E+01	EPA6 calc c	
3,3-Dichlorobenzidine	91-94-1	8.70E+00	EPA6 e	6.52E+01	EPA6 e	3.48E+01	EPA6 e	
3,3'-Dimethoxybenzidine	119-90-4	3.65E-01	EPA6 e	2.74E+00	EPA6 e	1.46E+00	EPA6 e	
4-Bromophenylphenyl ether	101-55-3	8.23E+02	EPA6 calc c	6.17E+03	EPA6 calc c	3.29E+03	EPA6 calc c	
Ammonium perfluorooctanoate	3825-26-1	NA	NA	NA	NA	NA	NA	
Azobenzene	103-33-3	2.69E+01	EPA6 calc c	2.02E+02	EPA6 calc c	1.08E+02	EPA6 calc c	
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	NA	NA	NA	NA	NA	NA	
Captan	133-06-2	2.00E+00	EPA6 calc c	1.50E+01	EPA6 calc c	7.98E+00	EPA6 calc c	
Chlorobenzilate	510-15-6	3.69E+01	EPA6 e	2.77E+02	EPA6 e	1.48E+02	EPA6 e	
Dibutylphosphate	107-66-4	NA	NA	NA	NA	NA	NA	
Dimethyl aminoazobenzene	60-11-7	3.18E+02	EPA6 calc c	2.38E+03	EPA6 calc c	1.27E+03	EPA6 calc c	
Hexachlorobenzene	118-74-1	8.00E+02	EPA6 e	6.00E+03	EPA6 e	3.20E+03	EPA6 e	
Hexachlorobutadiene	87-68-3	6.94E+01	EPA6 e	5.20E+02	EPA6 e	2.77E+02	EPA6 e	
Hexachlorocyclopentadiene	77-47-4	9.51E+01	EPA6 e	7.13E+02	EPA6 e	3.80E+02	EPA6 e	

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Partitioning Coefficients						
Constituent of Potential Concern	CAS Registry Number	Kd_s (cm^3/g)	Source	Kd_{sw} (L/kg)	Source	Kd_{bs} (cm^3/g)	Source	
Hexachloroethane	67-72-1	1.82E+01	EPA6 e	1.36E+01	EPA6 e	7.27E+01	EPA6 e	
Hexachlorophene	70-30-4	1.08E+04	EPA6 e	8.08E+04	EPA6 e	4.31E+04	EPA6 e	
Hexamethylene-1,5-diisocyanate	822-06-0	1.79E-01	EPA6 calc c	1.34E+00	EPA6 calc c	7.15E-01	EPA6 calc c	
Mirex	2385-85-5	1.00E+04	EPA6 calc c	7.50E+04	EPA6 calc c	4.00E+04	EPA6 calc c	
Nitrofen	1836-75-5	2.73E+03	EPA6 calc c	2.05E+04	EPA6 calc c	1.09E+04	EPA6 calc c	
Pentachlorobenzene	608-93-5	3.21E+02	EPA6 e	2.41E+03	EPA6 e	1.29E+03	EPA6 e	
Pentachloronitrobenzene	82-68-8	5.89E+01	EPA6 e	4.42E+02	EPA6 e	2.36E+02	EPA6 e	
Pentachlorophenol*	87-86-5	1.99E+02	EPA6 e	1.49E+03	EPA6 e	7.97E+02	EPA6 e	
Pentachlorophenol*	87-86-5	1.96E+02	EPA6 e	1.47E+03	EPA6 e	7.84E+02	EPA6 e	
Pentachlorophenol*	87-86-5	1.69E+02	EPA6 e	1.27E+03	EPA6 e	6.78E+02	EPA6 e	
Pentachlorophenol*	87-86-5	7.33E+01	EPA6 e	5.50E+02	EPA6 e	2.93E+02	EPA6 e	
Pentachlorophenol*	87-86-5	1.42E+01	EPA6 e	1.06E+02	EPA6 e	5.67E+01	EPA6 e	
Pentachlorophenol*	87-86-5	5.05E+00	EPA6 e	3.79E+01	EPA6 e	2.02E+01	EPA6 e	
Pentachlorophenol*	87-86-5	4.09E+00	EPA6 e	3.07E+01	EPA6 e	1.64E+01	EPA6 e	
Pentachlorophenol*	87-86-5	3.99E+00	EPA6 e	2.99E+01	EPA6 e	1.60E+01	EPA6 e	
Pentachlorophenol*	87-86-5	3.98E+00	EPA6 e	2.99E+01	EPA6 e	1.59E+01	EPA6 e	
Pentachlorophenol*	87-86-5	3.98E+00	EPA6 e	2.99E+01	EPA6 e	1.59E+01	EPA6 e	
Pentachlorophenol*	87-86-5	3.98E+00	EPA6 e	2.99E+01	EPA6 e	1.59E+01	EPA6 e	
Pentachlorophenol*	87-86-5	3.98E+00	EPA6 e	2.99E+01	EPA6 e	1.59E+01	EPA6 e	
Pentachlorophenol*	87-86-5	2.00E+02	EPA6 e	1.50E+03	EPA6 e	7.98E+02	EPA6 e	
Picric acid	88-89-1	9.90E-01	EPA6 calc c	7.43E+00	EPA6 calc c	3.96E+00	EPA6 calc c	
Pronamide	23950-58-5	7.74E+00	EPA6 e	5.81E+01	EPA6 e	3.10E+01	EPA6 e	
Strychnine	57-24-9	4.53E-01	EPA6 e	3.40E+00	EPA6 e	1.81E+00	EPA6 e	
Terphenyls	26140-60-3	NA	NA	NA	NA	NA	NA	
Tributyl phosphate	126-73-8	8.55E+01	EPA6 calc c	6.42E+02	EPA6 calc c	3.42E+02	EPA6 calc c	
Trifluralin	1582-09-8	6.03E+01	EPA6 calc c	4.52E+02	EPA6 calc c	2.41E+02	EPA6 calc c	
Triphenylamine	603-34-9	NA	NA	NA	NA	NA	NA	
Herbicides and Organopesticides								
2,4,5-T	93-76-5	5.25E-01	EPA6 calc c	3.94E+00	EPA6 calc c	2.10E+00	EPA6 calc c	

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Partitioning Coefficients						
Constituent of Potential Concern	CAS Registry Number	Kd_s (cm^3/g)	Source	Kd_{sw} (L/kg)	Source	Kd_{bs} (cm^3/g)	Source	
2,4-D and esters	94-75-7	2.00E-01	EPA6 calc c	1.50E+00	EPA6 calc c	7.98E-01	EPA6 calc c	
4,4-DDD	72-54-8	1.00E+03	EPA6 calc c	7.50E+03	EPA6 calc c	4.00E+03	EPA6 calc c	
4,4-DDT	50-29-3	2.40E+03	EPA6 calc c	1.80E+04	EPA6 calc c	9.60E+03	EPA6 calc c	
Aldrin	309-00-2	4.87E+02	EPA6 e	3.65E+03	EPA6 e	1.95E+03	EPA6 e	
alpha-BHC	319-84-6	1.76E+01	EPA6 e	1.32E+02	EPA6 e	7.05E+01	EPA6 e	
beta-BHC	319-85-7	2.14E+01	EPA6 e	1.60E+02	EPA6 e	8.56E+01	EPA6 e	
Chlordane	57-74-9	5.13E+02	EPA6 e	3.85E+03	EPA6 e	2.05E+03	EPA6 e	
DDE	72-55-9	8.64E+06	EPA6 e	6.48E+03	EPA6 e	3.46E+03	EPA6 e	
Delta-BHC	319-86-8	6.61E+00	EPA6 calc c	4.96E+01	EPA6 calc c	2.64E+01	EPA6 calc c	
Dieldrin	60-57-1	2.55E+02	EPA6 e	1.91E+03	EPA6 e	1.02E+03	EPA6 e	
Endothall	145-73-3	1.40E-03	EPA6 calc c	1.05E-02	EPA6 calc c	5.59E-03	EPA6 calc c	
Endrin	72-20-8	1.08E+02	EPA6 e	8.11E+02	EPA6 e	4.32E+02	EPA6 e	
gamma-BHC(Lindane)	58-89-9	1.07E+01	EPA6 calc c	8.04E+01	EPA6 calc c	4.29E+01	EPA6 calc c	
Heptachlor	76-44-8	9.53E+01	EPA6 e	7.15E+02	EPA6 e	3.81E+02	EPA6 e	
Isodrin	465-73-6	NA	NA	NA	NA	NA	NA	
Methoxychlor	72-43-5	8.00E+02	EPA6 e	6.00E+03	EPA6 e	3.20E+03	EPA6 e	
Silvex (2,4,5-TP)	93-72-1	1.01E+02	EPA6 calc c	7.56E+02	EPA6 calc c	4.03E+02	EPA6 calc c	
Toxaphene	8001-35-2	1.00E+03	EPA6 calc c	7.50E+03	EPA6 calc c	4.00E+03	EPA6 calc c	

NA = Value not available
** = Varies with pH; values given for a pH of 7.0.
a = Illustrated Handbook of Physical - Chemical Properties and Environmental Fate f
b = RAIS database: http://risk.lsd.ornl.gov/rap_hp.shtml
c = Equations derived from EPA Region 6 Guidance Appendix A-3
d = Handbook of Environmental degradation Rates. 1991.
e = EPA Region 6 Hazardous Waste Combustion Guidance
f = "A Critical Review of Henry's Law Constants for Chemicals of Environmental In
g = Handbook of Environmental Fate and Exposure Data for Organic Chemicals
h = http://www.epa.gov/region09/waste/sfund/prg/.
i = http://esc.syrres.com/efdb/Chemfate.htm
j = Hansch and Leo 1985 in Syracuse 1996

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Partitioning Coefficients						
Constituent of Potential Concern	CAS Registry Number	Kd_s (cm^3/g)	Source	Kd_{sw} (L/kg)	Source	Kd_{bs} (cm^3/g)	Source	

k = Schwarzenbach et al. 1993
l = Tomlin 1994 in Syracuse 1996
m = http://www.speclab.com/compound/c959988.htm

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Soil Loss		Half Life		Vapor Phase Fraction	
Constituent of Potential Concern	CAS Registry Number	ksg $(year)^{-1}$	Source	$t_{1/2}$ (hours)	Source	F_v (unitless)	Source
Aromatic Halogenated Hydrocarbons							
2,3,4,6-Tetrachlorophenol	58-90-2	1.41E+00	EPA6 calc c	4.32E+03	Degradation d	NA	NA
4-Chloro-3-methylphenol	59-50-7	NA	NA	NA	NA	NA	NA
Aromatic Nonhalogenated Hydrocarbons							
2-Nitrotoluene	88-72-2	NA	NA	NA	NA	NA	NA
4-Nitrobiphenyl	92-93-3	9.03E+00	EPA6 calc c	6.72E+02	Degradation d	NA	NA
Benzaldehyde	100-52-7	NA	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Benzene	71-43-2	3.89E+00	EPA6 e	3.84E+02	Degradation d	1.00E+00	EPA6 e
Benzyl alcohol	100-51-6	NA	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Ethyl Benzene	100-41-4	2.53E+01	EPA6 e	2.40E+02	Degradation d	1.00E+00	EPA6 e
m-Xylene	108-38-3	9.03E+00	EPA6 e	6.72E+02	Degradation d	1.00E+00	EPA6 e
o-Xylene	95-47-6	9.03E+00	EPA6 e	6.72E+02	Degradation d	1.00E+00	EPA6 e
p-Xylene	106-42-3	9.03E+00	EPA6 e	6.72E+02	Degradation d	1.00E+00	EPA6 e
Styrene	100-42-5	9.03E+00	EPA6 e	6.72E+02	Degradation d	1.00E+00	EPA6 e
Toluene	108-88-3	1.15E+01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Non-aromatic Nonhalogenated Hydrocarbons							
1,2-Epoxybutane	106-88-7	1.96E+01	EPA6 calc c	3.10E+02	Degradation d	NA	NA
1,3-Butadiene	106-99-0	9.03E+00	EPA6 calc c	6.72E+02	Degradation d	NA	NA
1,4-Dioxane	123-91-1	1.41E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
1-Methylpropyl alcohol	78-92-2	3.61E+01	EPA6 calc c	1.68E+02	Degradation d	NA	NA
1-Nitropropane	108-03-2	NA	NA	NA	NA	NA	NA
2,2,4-Trimethlylpentane	540-84-1	NA	NA	NA	NA	NA	NA
2-Butanone	78-93-3	3.61E+01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
2-Butenaldehyde (2-Butenal)	4170-30-3	NA	NA	NA	NA	NA	NA
2-Ethoxyethanol	110-80-5	9.03E+00	EPA6 calc c	6.72E+02	Degradation d	NA	NA
2-Heptanone	110-43-0	NA	NA	NA	NA	NA	NA
2-Hexanone	591-78-6	NA	NA	NA	NA	NA	NA
2-Methoxyethanol	109-86-4	9.03E+00	EPA6 calc c	6.72E+02	Degradation d	NA	NA
2-Methyl-2-propanol	75-65-0	1.26E+00	EPA6 calc c	4.80E+03	Degradation d	NA	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Soil Loss		Half Life		Vapor Phase Fraction	
Constituent of Potential Concern	CAS Registry Number	ksg $(year)^{-1}$	Source	$t_{1/2}$ (hours)	Source	F_v (unitless)	Source
2-Methyl-2-propenenitrile	126-98-7	NA	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
2-Methylaziridine	75-55-8	6.98E+00	EPA6 calc c	8.70E+02	Degradation d	NA	NA
2-Methylpropyl alcohol	78-83-1	3.51E+01	EPA6 calc c	1.73E+02	Degradation d	NA	NA
2-Pentanone	107-87-9	NA	NA	NA	NA	NA	NA
2-Propanone (Acetone)	67-64-1	3.61E+01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
2-Propene-1-ol	107-18-6	3.61E+01	EPA6 calc c	1.68E+02	Degradation d	NA	NA
2-Propyl alcohol	67-63-0	3.61E+01	EPA6 calc c	1.68E+02	Degradation d	NA	NA
3-Heptanone	106-35-4	NA	NA	NA	NA	NA	NA
3-Methyl-1-butanol	123-51-3	NA	NA	NA	NA	NA	NA
3-Methyl-2-butanone	563-80-4	NA	NA	NA	NA	NA	NA
3-Pentanone	96-22-0	NA	NA	NA	NA	NA	NA
4-Heptanone	123-19-3	NA	NA	NA	NA	NA	NA
4-Methyl-2-pentanone	108-10-1	3.61E+01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
4-Methyl-3-penten-2-one	141-79-7	NA	NA	NA	NA	NA	NA
5-Methyl-2-hexanone	110-12-3	NA	NA	NA	NA	NA	NA
Acetaldehyde	75-07-0	NA	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Acetamide	60-35-5	3.61E+01	EPA6 calc c	1.68E+02	Degradation d	NA	NA
Acetic acid	64-19-7	NA	NA	NA	NA	NA	NA
Acetic acid ethyl ester	141-78-6	3.61E+01	EPA6 calc c	1.68E+02	Degradation d	NA	NA
Acetic acid n-butyl ester	123-86-4	NA	NA	NA	NA	NA	NA
Acetonitrile	75-05-8	9.03E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Acrolein	107-02-8	9.03E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Acrylonitrile	107-13-1	1.10E+01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Bis(isopropyl)ether	108-20-3	NA	NA	NA	NA	NA	NA
Butane	106-97-8	NA	NA	NA	NA	NA	NA
Carbon disulfide	75-15-0	NA	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Cyanogen	460-19-5	NA	NA	NA	NA	NA	NA
Cyclohexane	110-82-7	1.41E+00	EPA6 calc c	4.32E+03	Degradation d	NA	NA
Cyclohexanone	108-94-1	NA	NA	NA	NA	NA	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Soil Loss		Half Life		Vapor Phase Fraction	
Constituent of Potential Concern	CAS Registry Number	ksg $(year)^{-1}$	Source	$t_{1/2}$ (hours)	Source	F_v (unitless)	Source
Cyclohexene	110-83-8	NA	NA	NA	NA	NA	NA
Cyclopentane	287-92-3	NA	NA	NA	NA	NA	NA
Ethyl alcohol	64-17-5	2.53E+02	EPA6 calc c	2.40E+01	Degradation d	NA	NA
Ethyl ether	60-29-7	NA	NA	NA	NA	NA	NA
Ethyl methacrylate	97-63-2	NA	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Formaldehyde	50-00-0	3.61E+01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Formamide	75-12-7	NA	NA	NA	NA	NA	NA
Formic acid	64-18-6	3.61E+01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Formic acid, methyl ester	107-31-3	NA	NA	NA	NA	NA	NA
Glycidylaldehyde	765-34-4	9.03E+00	EPA6 calc c	6.72E+02	Degradation d	NA	NA
Methyl acetate	79-20-9	NA	NA	NA	NA	NA	NA
Methyl alcohol	67-56-1	3.61E+01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Methyl isocyanate	624-83-9	1.86E+04	EPA6 calc c	3.26E-01	Degradation d	NA	NA
Methyl methacrylate	80-62-6	9.03E+00	EPA6 calc c	6.72E+02	Degradation d	NA	NA
Methyl tert-butyl ether	1634-04-4	1.41E+00	EPA6 calc c	4.32E+03	Degradation d	NA	NA
Methylacetylene	74-99-7	NA	NA	NA	NA	NA	NA
Methylcyclohexane	108-87-2	NA	NA	NA	NA	NA	NA
N,N-Dimethylacetamide	127-19-5	NA	NA	NA	NA	NA	NA
n-Butyl alcohol	71-36-3	3.61E+01	EPA6 calc c	1.68E+02	Degradation d	NA	NA
n-Heptane	142-82-5	NA	NA	NA	NA	NA	NA
n-Hexane	110-54-3	NA	NA	NA	NA	NA	NA
Nitromethane	75-52-5	NA	NA	NA	NA	NA	NA
n-Nonane	111-84-2	NA	NA	NA	NA	NA	NA
n-Octane	111-65-9	NA	NA	NA	NA	NA	NA
n-Pentane	109-66-0	NA	NA	NA	NA	NA	NA
n-Propionaldehyde	123-38-6	3.61E+01	EPA6 calc c	1.68E+02	Degradation d	NA	NA
n-Propyl alcohol	71-23-8	NA	NA	NA	NA	NA	NA
n-Valeraldehyde	110-62-3	NA	NA	NA	NA	NA	NA
Oxirane	75-21-8	2.13E+01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Soil Loss		Half Life		Vapor Phase Fraction	
Constituent of Potential Concern	CAS Registry Number	ksg $(year)^{-1}$	Source	$t_{1/2}$ (hours)	Source	F_v (unitless)	Source
p-Cymene	99-87-6	NA	NA	NA	NA	NA	NA
Phosgene	75-44-5	6.07E+03	EPA6 calc c	1.00E+00	Degradation d	NA	NA
Propargyl alcohol	107-19-7	NA	NA	NA	NA	NA	NA
Propionic acid	79-09-4	NA	NA	NA	NA	NA	NA
Propionitrile	107-12-0	NA	NA	NA	NA	NA	NA
Propylene glycol monomethyl ether	107-98-2	9.03E+00	EPA6 calc c	6.72E+02	Degradation d	NA	NA
p-tert-Butyltoluene	98-51-1	NA	NA	NA	NA	NA	NA
Triethylamine	121-44-8	NA	NA	NA	NA	NA	NA
Trimethylamine	75-50-3	NA	NA	NA	NA	NA	NA
Vinyl acetate	108-05-4	NA	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Non-aromatic Halogenated Hydrocarbons							
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	NA	NA	NA	NA	NA	NA
1,1,1,2-Tetrachloroethane	630-20-6	5.75E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
1,1,1-Trichloroethane	71-55-6	9.26E-01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	NA	NA	NA	NA	NA	NA
1,1,2,2-Tetrachloroethane	79-34-5	5.75E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
1,1,2,2-Tetrachloroethene	127-18-4	7.03E-01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
1,1,2-Trichloro-1,2,2-trifluoroethane	76-13-1	7.03E-01	EPA6 calc c	8.64E+03	Degradation d	NA	NA
1,1,2-Trichloroethane	79-00-5	6.93E-01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
1,1,2-Trichloroethylene	79-01-6	7.03E-01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
1,1-Dichloroethane	75-34-3	1.64E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
1,1-Dichloroethene	75-35-4	1.41E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
1,2,3-Trichloropropane	96-18-4	7.03E-01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
1,2-dibromo-3-chloropropane	96-12-8	1.41E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	NA	NA	NA	NA	NA	NA
1,2-Dichloroethane	107-06-2	1.41E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
1,2-Dichloroethylene	540-59-0	1.41E+00	EPA6 calc c	4.32E+03	Degradation d	NA	NA
1,2-Dichloropropane	78-87-5	1.96E-01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
1,3-Dichloropropene	542-75-6	2.24E+01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Soil Loss		Half Life		Vapor Phase Fraction	
Constituent of Potential Concern	CAS Registry Number	ksg $(year)^{-1}$	Source	$t_{1/2}$ (hours)	Source	F_v (unitless)	Source
1,4-Dichloro-2-butene	764-41-0	NA	NA	NA	NA	NA	NA
1-Chloroethene	75-01-4	1.41E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
2,2-Dichloropropionic acid	75-99-0	4.22E+00	EPA6 calc c	1.44E+03	Degradation d	NA	NA
2-Chloropropane	75-29-6	NA	NA	NA	NA	NA	NA
3-Chloropropene (allyl chloride)	107-05-1	1.81E+01	EPA6 calc c	3.35E+02	Degradation d	NA	NA
Bromochloromethane	74-97-5	NA	NA	NA	NA	NA	NA
Bromodichloromethane	75-27-4	NA	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Bromoethene	593-60-2	1.41E+00	EPA6 calc c	4.32E+03	Degradation d	NA	NA
Bromoform	75-25-2	1.41E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Bromomethane	74-83-9	9.03E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Carbon tetrachloride	56-23-5	7.03E-01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Chlordibromomethane	124-48-1	1.41E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Chlorodifluoromethane	75-45-6	NA	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Chloroethane	75-00-3	6.72E+02	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Chloroform	67-66-3	1.41E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Chloromethane	74-87-3	9.03E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Chloropentafluoroethane	76-15-3	NA	NA	NA	NA	NA	NA
cis-1,2-Dichloroethene	156-59-2	1.41E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
cis-1,3-Dichloropropene	10061-01-5	NA	NA	NA	NA	NA	NA
Cyanogen bromide	506-68-3	NA	NA	NA	NA	NA	NA
Cyanogen chloride	506-77-4	NA	NA	NA	NA	NA	NA
Dichlorodifluoromethane	75-71-8	1.41E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Dichlorofluoromethane	75-43-4	NA	NA	NA	NA	NA	NA
Dichloromethane	75-09-2	9.03E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Difluorodibromomethane	75-61-6	NA	NA	NA	NA	NA	NA
Hexafluoroacetone	684-16-2	NA	NA	NA	NA	NA	NA
Iodomethane	74-88-4	9.03E+00	EPA6 calc c	6.72E+02	Degradation d	NA	NA
Methylene bromide	74-95-3	9.03E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Pentachloroethane	76-01-7	NA	NA	NA	NA	NA	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Soil Loss		Half Life		Vapor Phase Fraction	
Constituent of Potential Concern	CAS Registry Number	ksg $(year)^{-1}$	Source	$t_{1/2}$ (hours)	Source	F_v (unitless)	Source
trans-1,2-Dichloroethene	156-60-5	1.41E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
trans-1,3-Dichloropropene	10061-02-6	NA	NA	NA	NA	NA	NA
Trichloroacetic acid	76-03-9	NA	NA	NA	NA	NA	NA
Trichlorofluoroethane	27154-33-2	NA	NA	NA	NA	NA	NA
Trichlorofluoromethane	75-69-4	7.03E-01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Trifluorobromomethane	75-63-8	NA	NA	NA	NA	NA	NA
Dioxin and Furan Compounds							
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	35822-46-9	1.09E-01	EPA6 e	NA	EPA6 e	1.62E-02	EPA6 e
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	3.57E-01	EPA6 e	NA	EPA6 e	3.47E-02	EPA6 e
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	3.57E-01	EPA6 e	NA	EPA6 e	2.01E-02	EPA6 e
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	39227-28-6	1.09E-01	EPA6 e	NA	EPA6 e	5.96E-02	EPA6 e
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	NA	EPA6 e	NA	EPA6 e	4.86E-02	EPA6 e
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	57653-85-7	1.09E-01	EPA6 e	NA	EPA6 e	2.89E-02	EPA6 e
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	NA	EPA6 e	NA	EPA6 e	5.15E-02	EPA6 e
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	19408-74-3	1.09E-01	EPA6 e	NA	EPA6 e	1.53E-02	EPA6 e
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	NA	EPA6 e	NA	EPA6 e	5.76E+00	EPA6 e
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	40321-76-4	NA	EPA6 e	NA	EPA6 e	2.19E-01	EPA6 e
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	3.57E-01	EPA6 e	NA	EPA6 e	3.64E-01	EPA6 e
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	NA	EPA6 e	NA	EPA6 e	5.47E-02	EPA6 e
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	3.57E-01	EPA6 e	NA	EPA6 e	2.63E-01	EPA6 e
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1746-01-6	4.29E-01	EPA6 e	NA	EPA6 e	9.78E-01	EPA6 e
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	3.57E-01	EPA6 e	NA	EPA6 e	7.68E-01	EPA6 e
Dibenzofuran	132-64-9	9.03E+00	EPA6 calc c	6.72E+02	Degradation d	NA	NA
Octachlorodibenzo(p)dioxin	3268-87-9	1.09E-01	EPA6 e	NA	EPA6 e	9.93E-01	EPA6 e
Octachlorodibenzofuran	39001-02-0	1.10E-01	EPA6 e	NA	EPA6 e	1.67E-03	EPA6 e
PCBs							
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	NA	NA	NA	NA	NA	NA
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	NA	NA	NA	NA	NA	NA
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	NA	NA	NA	NA	NA	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Soil Loss		Half Life		Vapor Phase Fraction	
Constituent of Potential Concern	CAS Registry Number	ksg $(year)^{-1}$	Source	$t_{1/2}$ (hours)	Source	F_v (unitless)	Source
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	NA	NA	NA	NA	NA	NA
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	NA	NA	NA	NA	NA	NA
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	NA	NA	NA	NA	NA	NA
2',3,4,4',5-Pentachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	NA	NA	NA	NA	NA	NA
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	NA	NA	NA	NA	NA	NA
3,3',4,4',5-Pentachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	NA	NA	NA	NA	NA	NA
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	NA	NA	NA	NA	NA	NA
Polychlorinated biphenyls (PCBs)**	1336-36-3	NA	NA	NA	NA	NA	NA
Phthalates							
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	1.10E+01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Butylbenzyl phthalate	85-68-7	3.61E+01	EPA6 e	NA	EPA6 e	9.64E-01	EPA6 e
Dibutyl phthalate	84-74-2	1.11E+01	EPA6 e	NA	EPA6 e	9.89E-01	EPA6 e
Diethyl phthalate	84-66-2	4.52E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Dimethyl phthalate	131-11-3	3.61E+01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
n-Dioctyl phthalate	117-84-0	9.03E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Light Polycyclic Aromatic Hydrocarbons							
2-Chloronaphthalene	91-58-7	NA	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
2-Methyl naphthalene	91-57-6	NA	NA	NA	NA	NA	NA
5-Nitroacenaphthene	602-87-9	NA	NA	NA	NA	NA	NA
Acenaphthene	83-32-9	2.48E+00	EPA6 e	NA	NA	1.00E+00	EPA6 e
Acenaphthylene	208-96-8	4.22E+00	EPA6 calc c	1.44E+03	Degradation d	NA	NA
Anthracene	120-12-7	5.50E-01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Fluorene	86-73-7	4.22E+00	EPA6 calc c	1.44E+03	Degradation d	NA	NA
Indene	95-13-6	NA	NA	NA	NA	NA	NA
Napthalene	91-20-3	5.27E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Soil Loss		Half Life		Vapor Phase Fraction	
Constituent of Potential Concern	CAS Registry Number	ksg $(year)^{-1}$	Source	$t_{1/2}$ (hours)	Source	F_v (unitless)	Source
Phenanthrene	85-01-8	1.26E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Pyrene	129-00-0	1.33E-01	EPA6 e	NA	EPA6 e	1.96E-01	EPA6 e
Heavy Polycyclic Aromatic Hydrocarbons							
3-Methylcholanthrene	56-49-5	1.81E-01	EPA6 calc c	3.36E+04	Degradation d	NA	NA
5-Methylchrysene	3697-24-3	NA	NA	NA	NA	NA	NA
Benzo(a)anthracene	56-55-3	3.72E-01	EPA6 e	NA	EPA6 e	8.81E-01	EPA6 e
Benzo(a)pyrene	50-32-8	4.77E-01	EPA6 e	NA	EPA6 e	2.65E-01	EPA6 e
Benzo(b)fluoranthene	205-99-2	4.15E-01	EPA6 e	NA	EPA6 e	8.22E-01	EPA6 e
Benzo(e)pyrene	192-97-2	NA	NA	NA	NA	NA	NA
Benzo(g,h,i)perylene	191-24-2	3.89E-01	EPA6 calc c	1.56E+04	Degradation d	NA	NA
Benzo(j)fluoranthene	205-82-3	NA	NA	NA	NA	NA	NA
Benzo(k)fluoranthene	207-08-9	1.18E-01	EPA6 e	NA	EPA6 e	1.49E-01	EPA6 e
Benzo[a,i]pyrene	191-30-0	NA	NA	NA	NA	NA	NA
Chrysene	218-01-9	2.53E-01	EPA6 e	NA	EPA6 e	7.61E-01	EPA6 e
Dibenz(a,h)anthracene	53-70-3	2.69E-01	EPA6 e	NA	EPA6 e	1.10E-02	EPA6 e
Dibenz[a,h]acridine	226-36-8	NA	NA	NA	NA	NA	NA
Dibenz[a,j]acridine	224-42-0	NA	NA	NA	NA	NA	NA
Dibenzo(a,e)fluoranthene	5385-75-1	NA	NA	NA	NA	NA	NA
Dibenzo(a,h)fluoranthene	no cas #	NA	NA	NA	NA	NA	NA
Dibenzo[a,e]pyrene	192-65-4	NA	NA	NA	NA	NA	NA
Dibenzo[a,h]pyrene	189-64-0	NA	NA	NA	NA	NA	NA
Dibenzo[a,i]pyrene	189-55-9	7.01E-01	EPA6 calc c	8.66E+03	Degradation d	NA	NA
Fluoranthene	206-44-0	5.75E-01	EPA6 e	NA	EPA6 e	9.92E-01	EPA6 e
Hexachloronaphthalene	1335-87-1	6.93E-01	EPA6 calc c	8.76E+03	Degradation d	NA	NA
Indeno(1,2,3-cd)pyrene	193-39-5	3.47E-01	EPA6 e	NA	EPA6 e	7.00E-03	EPA6 e
Octachloronaphthalene	2234-13-1	6.93E-01	EPA6 calc c	8.76E+03	Degradation d	NA	NA
Pentachloronaphthalene	1321-64-8	NA	NA	NA	NA	NA	NA
Tetrachloronaphthalene	1335-88-2	NA	NA	NA	NA	NA	NA
Trichloronaphthalene	1321-65-9	NA	NA	NA	NA	NA	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Soil Loss		Half Life		Vapor Phase Fraction	
Constituent of Potential Concern	CAS Registry Number	ksg $(year)^{-1}$	Source	$t_{1/2}$ (hours)	Source	F_v (unitless)	Source
Light Substituted Benzene Compounds							
1,2,3-Trichlorobenzene	87-61-6	1.41E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
1,2,4-Trichlorobenzene	120-82-1	1.41E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
1,2,4-Trimethlyl benzene	25551-13-7	NA	NA	NA	NA	NA	NA
1,2,4-Trimethyl benzene	95-63-6	9.03E+00	EPA6 calc c	6.72E+02	Degradation d	NA	NA
1,2-Dichlorobenzene	95-50-1	1.41E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
1,3,5-Trimethyl benzene	108-67-8	3.16E+01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
1,3-Dichlorobenzene	541-73-1	1.41E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
1,3-Dinitrobenzene	99-65-0	1.41E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
1,4-Dichlorobenzene	106-46-7	1.41E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
1,4-Dinitrobenzene	100-25-4	NA	NA	NA	NA	NA	NA
2,4,5-Trichorophenol	95-95-4	3.67E-01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
2,4,6-Trichlorophenol	88-06-2	3.61E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
2,4-Dichlorophenol	120-83-2	3.61E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
2,4-Dimethylphenol	105-67-9	3.61E+01	EPA6 e	NA	EPA6 e	9.97E-01	EPA6 e
2,4-Dinitrophenol	51-28-5	9.62E-01	EPA6 e	NA	EPA6 e	9.99E-01	EPA6 e
2,4-Dinitrotoluene	121-14-2	1.41E+00	EPA6 e	NA	EPA6 e	9.99E-01	EPA6 e
2,6-Dinitrotoluene	606-20-2	1.41E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
2-Chlorotoluene	95-49-8	NA	NA	NA	NA	NA	NA
2-Chorophenol	95-57-8	NA	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
2-Nitrophenol	88-75-5	9.03E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
4,6-Dinitro-o-cresol	534-52-1	1.20E+01	EPA6 calc c	5.04E+02	Degradation d	NA	NA
4-Chlorotoluene	106-43-4	NA	NA	NA	NA	NA	NA
4-Nitrophenol	100-02-7	2.09E+02	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
alpha-Methylstyrene	98-83-9	NA	NA	NA	NA	NA	NA
Aniline	62-53-3	3.20E+01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Benzotrichloride	98-07-7	1.21E+05	EPA6 calc c	5.00E-02	Degradation d	NA	NA
Benzyl Chloride	100-44-7	2.09E+01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Bromobenzene	108-86-1	NA	NA	NA	NA	NA	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Soil Loss		Half Life		Vapor Phase Fraction	
Constituent of Potential Concern	CAS Registry Number	ksg (year)$^{-1}$	Source	$t_{1/2}$ (hours)	Source	F_v (unitless)	Source
Chlorobenzene	108-90-7	1.69E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Cumene	98-82-8	3.16E+01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
m-Cresol	108-39-4	8.72E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
n-Butyl benzene	104-51-8	NA	NA	NA	NA	NA	NA
Nitrobenzene	98-95-3	1.28E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
n-Propyl benzene	103-65-1	NA	NA	NA	NA	NA	NA
o-Cresol	95-48-7	3.61E+01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
o-Dinitrobenzene	528-29-0	NA	NA	NA	NA	NA	NA
o-Nitroaniline	88-74-4	NA	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
o-Toluidine	95-53-4	3.61E+01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
p-Chloroaniline	106-47-8	NA	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
p-Cresol	106-44-5	3.79E+02	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Phenol	108-95-2	2.53E+01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
p-Nitrochlorobenzene	100-00-5	NA	NA	NA	NA	NA	NA
p-Toluidine	106-49-0	NA	NA	NA	NA	NA	NA
sec-Butyl benzene	135-98-8	NA	NA	NA	NA	NA	NA
tert-Butyl benzene	98-06-6	NA	NA	NA	NA	NA	NA
Toluene-2,6-diamine	823-40-5	NA	NA	NA	NA	NA	NA
Other Light Semivolatile Compounds							
1,1 Dimethyl hydrazine	57-14-7	1.15E+01	EPA6 calc c	5.28E+02	Degradation d	NA	NA
1,1-Biphenyl	92-52-4	3.61E+01	EPA6 calc c	1.68E+02	Degradation d	NA	NA
1,2 Diphenylhydrazine	122-66-7	1.41E+00	EPA6 e	NA	EPA6 e	9.99E-01	EPA6 e
1,2-Dimethylhydrazine	540-73-8	9.03E+00	EPA6 calc c	6.72E+02	Degradation d	NA	NA
1,3-Propane sultone	1120-71-4	9.03E+00	EPA6 calc c	6.72E+02	Degradation d	NA	NA
2,4-Toluene diisocyanate	584-84-9	NA	NA	NA	NA	NA	NA
2-Chloroacetophenone	532-27-4	9.03E+00	EPA6 calc c	6.72E+02	Degradation d	NA	NA
2-Propenoic acid	79-10-7	3.61E+01	EPA6 calc c	1.68E+02	Degradation d	NA	NA
4,4-Methylenedianiline	101-77-9	3.61E+01	EPA6 calc c	1.68E+02	Degradation d	NA	NA
Acetophenone	98-86-2	NA	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Soil Loss		Half Life		Vapor Phase Fraction	
Constituent of Potential Concern	CAS Registry Number	ksg $(year)^{-1}$	Source	$t_{1/2}$ (hours)	Source	F_v (unitless)	Source
Benzoic acid	65-85-0	1.26E+02	NA	NA	NA	1.00E+00	NA
bis(2-Chloroethoxy)methane	111-91-1	NA	NA	NA	NA	NA	NA
bis(2-Chloroethyl) ether	111-44-4	1.41E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Chlorocyclopentadiene	41851-50-7	NA	NA	NA	NA	NA	NA
Cyclohexanol	108-93-0	NA	NA	NA	NA	NA	NA
Dichloroisopropyl ether	108-60-1	1.41E+00	EPA6 calc c	4.32E+03	Degradation d	NA	NA
Dichloromethyl ether	542-88-1	NA	NA	NA	NA	NA	NA
Dichloropentadiene	no cas #	NA	NA	NA	NA	NA	NA
Dimethyl sulfate	77-78-1	5.06E+02	EPA6 calc c	1.20E+01	Degradation d	NA	NA
Dimethylaniline	121-69-7	1.41E+00	EPA6 calc c	4.32E+03	Degradation d	NA	NA
di-n-propylnitrosamine	621-64-7	1.41E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Diphenyl ether	101-84-8	NA	NA	NA	NA	NA	NA
Epichlorohydrin	106-89-8	9.03E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Ethyl carbomate (urethane)	51-79-6	3.61E+01	EPA6 calc c	1.68E+02	Degradation d	NA	NA
Ethyl methane sulfanate	62-50-0	7.88E+01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Ethylene dibromide	106-93-4	1.41E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Ethylene glycol	107-21-1	2.11E+01	EPA6 calc c	2.88E+02	Degradation d	NA	NA
Ethylene glycol monobutyl ether	111-76-2	9.03E+00	EPA6 calc c	6.72E+02	Degradation d	NA	NA
Ethylene glycol monoethyl ether acetate	111-15-9	NA	NA	NA	NA	NA	NA
Ethylene thiourea	96-45-7	9.03E+00	EPA6 calc c	6.72E+02	Degradation d	NA	NA
Furfural	98-01-1	NA	NA	NA	NA	NA	NA
Maleic hydrazide	123-33-1	NA	NA	NA	NA	NA	NA
Malononitrile	109-77-3	NA	NA	NA	NA	NA	NA
Methyl styrene (mixed isomers)	25013-15-4	NA	NA	NA	NA	NA	NA
Methylhydrazine	60-34-4	1.05E+01	EPA6 calc c	5.76E+02	Degradation d	NA	NA
N,N-Diphenylamine	122-39-4	9.03E+00	EPA6 calc c	6.72E+02	Degradation d	NA	NA
Nitric acid, propyl ester	627-13-4	NA	NA	NA	NA	NA	NA
N-Nitrosodi-n-butylamine	924-16-3	7.44E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
N-Nitrosomorpholine	59-89-2	1.41E+00	EPA6 calc c	4.32E+03	Degradation d	NA	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Soil Loss		Half Life		Vapor Phase Fraction	
Constituent of Potential Concern	CAS Registry Number	ksg $(year)^{-1}$	Source	$t_{1/2}$ (hours)	Source	F_v (unitless)	Source
N-Nitroso-N,N-dimethylamine	62-75-9	1.41E+00	EPA6 calc c	4.32E+03	Degradation d	NA	NA
o-Anisidine	90-04-0	1.41E+00	EPA6 calc c	4.32E+03	Degradation d	NA	NA
Oxalic acid	144-62-7	NA	NA	NA	NA	NA	NA
Phthalic Anhydride	85-44-9	1.35E+04	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
p-Phthalic acid	100-21-0	3.61E+01	EPA6 calc c	1.68E+02	Degradation d	NA	NA
Pyridine	110-86-1	3.61E+01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Quinoline	91-22-5	2.53E+01	EPA6 calc c	2.40E+02	Degradation d	NA	NA
Quinone	106-51-4	5.06E+01	EPA6 calc c	1.20E+02	Degradation d	NA	NA
Safrole	94-59-7	9.03E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Tetrahydrofuran	109-99-9	4.43E+01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Other Heavy Semivolatile Compounds							
1,2,4,5-Tetrachlorobenzene	95-94-3	1.41E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
1,3,5-Trinitrobenzene	99-35-4	NA	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	NA	NA	NA	NA	NA	NA
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	NA	NA	NA	NA	NA	NA
2-sec-Butyl-4,6-dinitrophenol	88-85-7	2.06E+00	EPA6 calc c	2.95E+03	Degradation d	NA	NA
3,3-Dichlorobenzidine	91-94-1	1.41E+00	EPA6 e	NA	EPA6 e	8.47E-01	EPA6 e
3,3'-Dimethoxybenzidine	119-90-4	1.41E+00	EPA6 e	NA	EPA6 e	8.77E-01	EPA6 e
4-Bromophenylphenyl ether	101-55-3	NA	NA	NA	NA	NA	NA
Ammonium perfluorooctanoate	3825-26-1	NA	NA	NA	NA	NA	NA
Azobenzene	103-33-3	NA	NA	NA	NA	NA	NA
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	NA	NA	NA	NA	NA	NA
Captan	133-06-2	4.22E+00	EPA6 calc c	1.44E+03	Degradation d	NA	NA
Chlorobenzilate	510-15-6	7.23E+00	EPA6 e	NA	EPA6 e	8.62E-01	EPA6 e
Dibutylphosphate	107-66-4	NA	NA	NA	NA	NA	NA
Dimethyl aminoazobenzene	60-11-7	9.03E+00	EPA6 calc c	6.72E+02	Degradation d	NA	NA
Hexachlorobenzene	118-74-1	1.21E-01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Hexachlorobutadiene	87-68-3	1.41E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Hexachlorocyclopentadiene	77-47-4	9.03E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Soil Loss		Half Life		Vapor Phase Fraction	
Constituent of Potential Concern	CAS Registry Number	ksg $(year)^{-1}$	Source	$t_{1/2}$ (hours)	Source	F_v (unitless)	Source
Hexachloroethane	67-72-1	1.41E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Hexachlorophene	70-30-4	7.71E-01	EPA6 e	NA	EPA6 e	1.40E-04	EPA6 e
Hexamethylene-1,5-diisocyanate	822-06-0	NA	NA	NA	NA	NA	NA
Mirex	2385-85-5	NA	NA	NA	NA	NA	NA
Nitrofen	1836-75-5	NA	NA	NA	NA	NA	NA
Pentachlorobenzene	608-93-5	7.33E-01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Pentachloronitrobenzene	82-68-8	3.62E-01	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Pentachlorophenol*	87-86-5	1.42E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Pentachlorophenol*	87-86-5	1.42E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Pentachlorophenol*	87-86-5	1.42E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Pentachlorophenol*	87-86-5	1.42E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Pentachlorophenol*	87-86-5	1.42E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Pentachlorophenol*	87-86-5	1.42E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Pentachlorophenol*	87-86-5	1.42E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Pentachlorophenol*	87-86-5	1.42E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Pentachlorophenol*	87-86-5	1.42E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Pentachlorophenol*	87-86-5	1.42E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Pentachlorophenol*	87-86-5	1.42E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Picric acid	88-89-1	1.41E+00	EPA6 calc c	4.32E+03	Degradation d	NA	NA
Pronamide	23950-58-5	NA	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Strychnine	57-24-9	9.03E+00	EPA6 e	NA	EPA6 e	8.60E-02	EPA6 e
Terphenyls	26140-60-3	NA	NA	NA	NA	NA	NA
Tributyl phosphate	126-73-8	NA	NA	NA	NA	NA	NA
Trifluralin	1582-09-8	NA	NA	NA	NA	NA	NA
Triphenylamine	603-34-9	NA	NA	NA	NA	NA	NA
Herbicides and Organopesticides							
2,4,5-T	93-76-5	1.26E+01	EPA6 calc c	4.80E+02	Degradation d	NA	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Soil Loss		Half Life		Vapor Phase Fraction	
Constituent of Potential Concern	CAS Registry Number	ksg $(year)^{-1}$	Source	$t_{1/2}$ (hours)	Source	F_v (unitless)	Source
2,4-D and esters	94-75-7	5.06E+00	EPA6 calc c	1.20E+03	Degradation d	NA	NA
4,4-DDD	72-54-8	4.34E-02	EPA6 calc c	1.40E+05	Degradation d	NA	NA
4,4-DDT	50-29-3	4.34E-02	EPA6 calc c	1.40E+05	Degradation d	NA	NA
Aldrin	309-00-2	4.28E-01	EPA6 e	NA	EPA6 e	2.27E-01	EPA6 e
alpha-BHC	319-84-6	1.87E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
beta-BHC	319-85-7	2.04E+00	EPA6 e	NA	EPA6 e	9.99E-01	EPA6 e
Chlordane	57-74-9	1.83E-01	EPA6 e	NA	EPA6 e	9.97E-01	EPA6 e
DDE	72-55-9	4.34E-02	EPA6 e	NA	EPA6 e	9.81E-01	EPA6 e
Delta-BHC	319-86-8	2.53E+00	EPA6 calc c	2.40E+03	Degradation d	NA	NA
Dieldrin	60-57-1	2.34E+00	EPA6 e	NA	EPA6 e	8.30E-02	EPA6 e
Endothall	145-73-3	NA	NA	NA	NA	NA	NA
Endrin	72-20-8	3.61E+04	EPA6 e	NA	EPA6 e	NA	EPA6 e
gamma-BHC(Lindane)	58-89-9	1.05E+00	EPA6 calc c	5.77E+03	Degradation d	NA	NA
Heptachlor	76-44-8	1.41E+00	EPA6 e	NA	EPA6 e	1.00E+00	EPA6 e
Isodrin	465-73-6	NA	NA	NA	NA	NA	NA
Methoxychlor	72-43-5	6.93E-01	EPA6 e	NA	EPA6 e	9.01E-01	EPA6 e
Silvex (2,4,5-TP)	93-72-1	NA	NA	NA	NA	NA	NA
Toxaphene	8001-35-2	NA	NA	NA	NA	NA	NA

NA = Value not available
** = Varies with pH; values given for a pH of 7.0.
a = Illustrated Handbook of Physical - Chemical Properties and Environmental Fate f
b = RAIS database: http://risk.lsd.ornl.gov/rap_hp.shtml
c = Equations derived from EPA Region 6 Guidance Appendix A-3
d = Handbook of Environmental degradation Rates. 1991.
e = EPA Region 6 Hazardous Waste Combustion Guidance
f = "A Critical Review of Henry's Law Constants for Chemicals of Environmental In
g = Handbook of Environmental Fate and Exposure Data for Organic Chemicals
h = http://www.epa.gov/region09/waste/sfund/prg/.
i = http://esc.syrres.com/efdb/Chemfate.htm
j = Hansch and Leo 1985 in Syracuse 1996

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Soil Loss		Half Life		Vapor Phase Fraction	
Constituent of Potential Concern	CAS Registry Number	ksg $(year)^{-1}$	Source	$t_{1/2}$ (hours)	Source	F_v (unitless)	Source

k = Schwarzenbach et al. 1993
l = Tomlin 1994 in Syracuse 1996
m = http://www.speclab.com/compound/c959988.htm

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	RCF ((ug/g DW plant)/ (ug/mL soil water))	Source	$Br_{root\ veg}$ ((ug/g DW plant) /(ug/g soil))	Source	Br_{ag} ((ug/g DW plant) /(ug/g soil))	Source
Aromatic Halogenated Hydrocarbons							
2,3,4,6-Tetrachlorophenol	58-90-2	7.73E+01	EPA6 calc c	1.17E+00	EPA6 calc c	1.08E-01	EPA6 calc c
4-Chloro-3-methylphenol	59-50-7	8.18E+00	EPA6 calc c	1.36E+00	EPA6 calc c	6.25E-01	EPA6 calc c
Aromatic Nonhalogenated Hydrocarbons							
2-Nitrotoluene	88-72-2	2.60E+00	EPA6 calc c	6.10E-01	EPA6 calc c	1.81E+00	EPA6 calc c
4-Nitrobiphenyl	92-93-3	2.50E+01	EPA6 calc c	NA	NA	2.56E-01	EPA6 calc c
Benzaldehyde	100-52-7	9.50E+00	EPA6 e	4.72E+01	EPA6 e	5.42E+00	EPA6 e
Benzene	71-43-2	1.66E+01	EPA6 e	2.67E+01	EPA6 e	2.25E+00	EPA6 e
Benzyl alcohol	100-51-6	7.94E+00	EPA6 e	7.77E+01	EPA6 e	8.95E+00	EPA6 e
Ethyl Benzene	100-41-4	6.52E+01	EPA6 e	3.20E+01	EPA6 e	6.07E-01	EPA6 e
m-Xylene	108-38-3	7.41E+01	EPA6 e	3.78E+01	EPA6 e	5.47E-01	EPA6 e
o-Xylene	95-47-6	6.61E+01	EPA6 e	2.74E+01	EPA6 e	6.01E-01	EPA6 e
p-Xylene	106-42-3	7.05E+01	EPA6 e	2.27E+01	EPA6 e	5.70E-01	EPA6 e
Styrene	100-42-5	4.81E+01	EPA6 e	5.28E+00	EPA6 e	7.85E-01	EPA6 e
Toluene	108-88-3	3.26E+01	EPA6 e	2.33E+01	EPA6 e	1.11E+00	EPA6 e
Non-aromatic Nonhalogenated Hydrocarbons							
1,2-Epoxybutane	106-88-7	1.21E+00	EPA6 calc c	1.52E+01	EPA6 calc c	5.69E+00	EPA6 calc c
1,3-Butadiene	106-99-0	1.85E+00	EPA6 calc c	2.04E+00	EPA6 calc c	2.74E+00	EPA6 calc c
1,4-Dioxane	123-91-1	6.45E+00	EPA6 e	7.37E+02	EPA6 e	5.53E+01	EPA6 e
1-Methylpropyl alcohol	78-92-2	9.09E-01	EPA6 calc c	2.28E+01	EPA6 calc c	1.72E+01	EPA6 calc c
1-Nitropropane	108-03-2	9.61E-01	EPA6 calc c	1.34E+01	EPA6 calc c	1.22E+01	EPA6 calc c
2,2,4-Trimethlylpentane	540-84-1	2.22E+02	EPA6 calc c	2.58E-01	EPA6 calc c	4.86E-02	EPA6 calc c
2-Butanone	78-93-3	6.69E+00	EPA6 e	2.86E+02	EPA6 e	2.67E+01	EPA6 e
2-Butenaldehyde (2-Butenal)	4170-30-3	NA	NA	NA	NA	NA	NA
2-Ethoxyethanol	110-80-5	8.45E-01	EPA6 calc c	4.05E+00	EPA6 calc c	4.42E+01	EPA6 calc c
2-Heptanone	110-43-0	1.83E+00	EPA6 calc c	2.07E+00	EPA6 calc c	2.78E+00	EPA6 calc c
2-Hexanone	591-78-6	1.17E+00	EPA6 calc c	8.72E-01	EPA6 calc c	6.17E+00	EPA6 calc c
2-Methoxyethanol	109-86-4	8.67E-01	EPA6 calc c	1.02E+01	EPA6 calc c	2.77E+01	EPA6 calc c
2-Methyl-2-propanol	75-65-0	8.76E-01	EPA6 calc c	2.36E+00	EPA6 calc c	2.43E+01	EPA6 calc c

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	RCF ((ug/g DW plant)/ (ug/mL soil water))	Source	$Br_{root\ veg}$ ((ug/g DW plant) /(ug/g soil))	Source	Br_{ag} ((ug/g DW plant) /(ug/g soil))	Source
2-Methyl-2-propenenitrile	126-98-7	6.91E+00	EPA6 e	1.85E+02	EPA6 e	1.89E+01	EPA6 e
2-Methylaziridine	75-55-8	NA	NA	NA	NA	NA	NA
2-Methylpropyl alcohol	78-83-1	9.36E-01	EPA6 calc c	1.68E+01	EPA6 calc c	1.41E+01	EPA6 calc c
2-Pentanone	107-87-9	9.72E-01	EPA6 calc c	1.31E+00	EPA6 calc c	1.15E+01	EPA6 calc c
2-Propanone (Acetone)	67-64-1	6.46E+00	EPA6 e	6.80E+02	EPA6 e	5.20E+01	EPA6 e
2-Propene-1-ol	107-18-6	8.61E-01	EPA6 calc c	5.86E+01	EPA6 calc c	3.09E+01	EPA6 calc c
2-Propyl alcohol	67-63-0	8.53E-01	EPA6 calc c	7.61E+01	EPA6 calc c	3.62E+01	EPA6 calc c
3-Heptanone	106-35-4	NA	NA	NA	NA	NA	NA
3-Methyl-1-butanol	123-51-3	NA	NA	NA	NA	NA	NA
3-Methyl-2-butanone	563-80-4	NA	NA	NA	NA	NA	NA
3-Pentanone	96-22-0	9.95E-01	EPA6 calc c	8.27E+00	EPA6 calc c	1.04E+01	EPA6 calc c
4-Heptanone	123-19-3	NA	NA	NA	NA	NA	NA
4-Methyl-2-pentanone	108-10-1	8.22E+00	EPA6 e	6.85E+01	EPA6 e	7.95E+00	EPA6 e
4-Methyl-3-penten-2-one	141-79-7	NA	NA	NA	NA	NA	NA
5-Methyl-2-hexanone	110-12-3	NA	NA	NA	NA	NA	NA
Acetaldehyde	75-07-0	6.46E+00	EPA6 e	6.78E+02	EPA6 e	5.19E+01	EPA6 e
Acetamide	60-35-5	8.23E-01	EPA6 calc c	2.92E+03	EPA6 calc c	2.07E+02	EPA6 calc c
Acetic acid	64-19-7	8.42E-01	EPA6 calc c	8.42E+01	EPA6 calc c	4.86E+01	EPA6 calc c
Acetic acid ethyl ester	141-78-6	9.30E-01	EPA6 calc c	4.05E+01	EPA6 calc c	1.47E+01	EPA6 calc c
Acetic acid n-butyl ester	123-86-4	1.47E+00	EPA6 calc c	2.93E+00	EPA6 calc c	3.87E+00	EPA6 calc c
Acetonitrile	75-05-8	6.43E+00	EPA6 e	8.37E+02	EPA6 e	6.09E+01	EPA6 e
Acrolein	107-02-8	6.54E+00	EPA6 e	4.69E+02	EPA6 e	3.92E+01	EPA6 e
Acrylonitrile	107-13-1	6.67E+00	EPA6 e	3.00E+02	EPA6 e	2.77E+01	EPA6 e
Bis(isopropyl)ether	108-20-3	NA	NA	NA	NA	NA	NA
Butane	106-97-8	5.89E+00	EPA6 calc c	8.50E-01	EPA6 calc c	8.27E-01	EPA6 calc c
Carbon disulfide	75-15-0	1.44E+01	EPA6 e	2.79E+01	EPA6 e	2.70E+00	EPA6 e
Cyanogen	460-19-5	9.46E-01	EPA6 calc c	1.81E+01	EPA6 calc c	1.32E+01	EPA6 calc c
Cyclohexane	110-82-7	1.43E+01	EPA6 calc c	2.98E+00	EPA6 calc c	3.98E-01	EPA6 calc c
Cyclohexanone	108-94-1	9.47E-01	EPA6 calc c	1.51E+01	EPA6 calc c	1.32E+01	EPA6 calc c

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	RCF ((ug/g DW plant)/ (ug/mL soil water))	Source	$Br_{root\ veg}$ ((ug/g DW plant) /(ug/g soil))	Source	Br_{ag} ((ug/g DW plant) /(ug/g soil))	Source
Cyclohexene	110-83-8	5.63E+00	EPA6 calc c	8.69E-01	EPA6 calc c	8.61E-01	EPA6 calc c
Cyclopentane	287-92-3	6.99E+00	EPA6 calc c	7.85E-01	EPA6 calc c	7.14E-01	EPA6 calc c
Ethyl alcohol	64-17-5	8.37E-01	EPA6 calc c	1.69E+02	EPA6 calc c	5.85E+01	EPA6 calc c
Ethyl ether	60-29-7	9.66E-01	EPA6 calc c	1.29E+01	EPA6 calc c	1.18E+01	EPA6 calc c
Ethyl methacrylate	97-63-2	1.02E+01	EPA6 e	4.14E+01	EPA6 e	4.67E+00	EPA6 e
Formaldehyde	50-00-0	6.73E+00	EPA6 e	2.57E+02	EPA6 e	2.46E+01	EPA6 e
Formamide	75-12-7	8.22E-01	EPA6 calc c	2.51E+03	EPA6 calc c	2.89E+02	EPA6 calc c
Formic acid	64-18-6	6.40E+00	EPA6 e	1.19E+02	EPA6 e	7.92E+01	EPA6 e
Formic acid, methyl ester	107-31-3	8.39E-01	EPA6 calc c	1.52E+02	EPA6 calc c	5.50E+01	EPA6 calc c
Glycidylaldehyde	765-34-4	8.28E-01	EPA6 calc c	8.28E+02	EPA6 calc c	1.02E+02	EPA6 calc c
Methyl acetate	79-20-9	8.62E-01	EPA6 calc c	1.80E+01	EPA6 calc c	3.05E+01	EPA6 calc c
Methyl alcohol	67-56-1	6.37E+00	EPA6 e	1.61E+03	EPA6 e	9.96E+01	EPA6 e
Methyl isocyanate	624-83-9	NA	NA	NA	NA	NA	NA
Methyl methacrylate	80-62-6	9.43E-01	EPA6 calc c	1.49E+00	EPA6 calc c	1.35E+01	EPA6 calc c
Methyl tert-butyl ether	1634-04-4	9.80E-01	EPA6 calc c	1.17E+01	EPA6 calc c	1.11E+01	EPA6 calc c
Methylacetylene	74-99-7	9.80E-01	EPA6 calc c	1.17E+01	EPA6 calc c	1.11E+01	EPA6 calc c
Methylcyclohexane	108-87-2	4.42E+01	EPA6 calc c	4.12E-01	EPA6 calc c	1.65E-01	EPA6 calc c
N,N-Dimethylacetamide	127-19-5	NA	NA	NA	NA	NA	NA
n-Butyl alcohol	71-36-3	9.64E-01	EPA6 calc c	1.31E+01	EPA6 calc c	1.20E+01	EPA6 calc c
n-Heptane	142-82-5	1.18E+02	EPA6 calc c	3.09E-01	EPA6 calc c	7.84E-02	EPA6 calc c
n-Hexane	110-54-3	4.49E+01	EPA6 calc c	4.10E-01	EPA6 calc c	1.63E-01	EPA6 calc c
Nitromethane	75-52-5	8.36E-01	EPA6 calc c	1.85E+02	EPA6 calc c	6.17E+01	EPA6 calc c
n-Nonane	111-84-2	6.78E+02	EPA6 calc c	1.89E-01	EPA6 calc c	2.10E-02	EPA6 calc c
n-Octane	111-65-9	3.71E+01	EPA6 calc c	4.34E-01	EPA6 calc c	1.89E-01	EPA6 calc c
n-Pentane	109-66-0	9.77E+00	EPA6 calc c	6.83E-01	EPA6 calc c	5.40E-01	EPA6 calc c
n-Propionaldehyde	123-38-6	9.06E-01	EPA6 calc c	2.38E+01	EPA6 calc c	1.77E+01	EPA6 calc c
n-Propyl alcohol	71-23-8	NA	NA	NA	NA	NA	NA
n-Valeraldehyde	110-62-3	NA	NA	NA	NA	NA	NA
Oxirane	75-21-8	6.44E+00	EPA6 e	7.80E+02	EPA6 e	5.77E+01	EPA6 e

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	RCF ((ug/g DW plant)/ (ug/mL soil water))	Source	$Br_{root\ veg}$ ((ug/g DW plant) /(ug/g soil))	Source	Br_{ag} ((ug/g DW plant) /(ug/g soil))	Source
p-Cymene	99-87-6	4.42E+01	EPA6 calc c	4.12E-01	EPA6 calc c	1.65E-01	EPA6 calc c
Phosgene	75-44-5	NA	NA	NA	NA	NA	NA
Propargyl alcohol	107-19-7	NA	NA	NA	NA	NA	NA
Propionic acid	79-09-4	8.74E-01	EPA6 calc c	4.14E+01	EPA6 calc c	2.50E+01	EPA6 calc c
Propionitrile	107-12-0	8.60E-01	EPA6 calc c	5.99E+01	EPA6 calc c	3.13E+01	EPA6 calc c
Propylene glycol monomethyl ether	107-98-2	8.42E-01	EPA6 calc c	6.69E+00	EPA6 calc c	4.92E+01	EPA6 calc c
p-tert-Butyltoluene	98-51-1	NA	NA	NA	NA	NA	NA
Triethylamine	121-44-8	NA	NA	NA	NA	NA	NA
Trimethylamine	75-50-3	8.60E-01	EPA6 calc c	2.15E+01	EPA6 calc c	3.13E+01	EPA6 calc c
Vinyl acetate	108-05-4	7.11E+00	EPA6 e	1.43E+02	EPA6 e	1.53E+01	EPA6 e
Non-aromatic Halogenated Hydrocarbons							
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	NA	NA	NA	NA	NA	NA
1,1,1,2-Tetrachloroethane	630-20-6	3.09E+01	EPA6 e	1.94E+01	EPA6 e	1.17E+00	EPA6 e
1,1,1-Trichloroethane	71-55-6	2.33E+01	EPA6 e	1.73E-02	EPA6 e	1.54E+00	EPA6 e
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	2.33E+01	EPA6 calc c	7.37E+00	EPA6 calc c	2.70E-01	EPA6 calc c
1,1,2,2-Tetrachloroethane	79-34-5	8.80E+02	EPA6 e	1.11E+03	EPA6 e	8.02E-02	EPA6 e
1,1,2,2-Tetrachloroethene	127-18-4	2.75E+01	EPA6 e	1.04E-03	EPA6 e	1.31E+00	EPA6 e
1,1,2-Trichloro-1,2,2-trifluoroethane	76-13-1	9.01E+00	EPA6 calc c	3.50E+00	EPA6 calc c	5.77E-01	EPA6 calc c
1,1,2-Trichloroethane	79-00-5	1.59E+01	EPA6 e	2.12E+01	EPA6 e	2.38E+00	EPA6 e
1,1,2-Trichloroethylene	79-01-6	2.37E+01	EPA6 e	2.12E+01	EPA6 e	1.52E+00	EPA6 e
1,1-Dichloroethane	75-34-3	1.19E+01	EPA6 e	2.24E+01	EPA6 e	3.56E+00	EPA6 e
1,1-Dichloroethene	75-35-4	1.63E+01	EPA6 e	2.50E+01	EPA6 e	2.30E+00	EPA6 e
1,2,3-Trichloropropane	96-18-4	1.89E+01	EPA6 e	2.34E+01	EPA6 e	1.94E+00	EPA6 e
1,2-dibromo-3-chloropropane	96-12-8	2.10E+01	EPA6 e	2.22E+01	EPA6 e	1.72E+00	EPA6 e
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	NA	NA	NA	NA	NA	NA
1,2-Dichloroethane	107-06-2	9.41E+00	EPA6 e	4.81E+01	EPA6 e	5.53E+00	EPA6 e
1,2-Dichloroethylene	540-59-0	8.91E-01	EPA6 calc c	3.10E+01	EPA6 calc c	2.04E+01	EPA6 calc c
1,2-Dichloropropane	78-87-5	1.89E+01	EPA6 e	4.01E+01	EPA6 e	1.94E+00	EPA6 e
1,3-Dichloropropene	542-75-6	1.15E+01	EPA6 e	4.25E+01	EPA6 e	3.78E+00	EPA6 e

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	RCF ((ug/g DW plant)/ (ug/mL soil water))	Source	$Br_{root\ veg}$ ((ug/g DW plant) /(ug/g soil))	Source	Br_{ag} ((ug/g DW plant) /(ug/g soil))	Source
1,4-Dichloro-2-butene	764-41-0	9.62E-01	EPA6 calc c	5.29E-01	EPA6 calc c	1.21E+01	EPA6 calc c
1-Chloroethene	75-01-4	8.08E+00	EPA6 e	7.29E+01	EPA6 e	8.42E+00	EPA6 e
2,2-Dichloropropionic acid	75-99-0	9.40E-01	EPA6 calc c	1.61E+01	EPA6 calc c	1.38E+01	EPA6 calc c
2-Chloropropane	75-29-6	1.70E+00	EPA6 calc c	2.30E+00	EPA6 calc c	3.09E+00	EPA6 calc c
3-Chloropropene (allyl chloride)	107-05-1	9.84E-01	EPA6 calc c	2.06E+00	EPA6 calc c	1.09E+01	EPA6 calc c
Bromochloromethane	74-97-5	1.19E+00	EPA6 calc c	4.88E+00	EPA6 calc c	5.93E+00	EPA6 calc c
Bromodichloromethane	75-27-4	1.47E+01	EPA6 e	2.74E+01	EPA6 e	2.61E+00	EPA6 e
Bromoethene	593-60-2	1.02E+00	EPA6 calc c	6.74E-01	EPA6 calc c	9.37E+00	EPA6 calc c
Bromoform	75-25-2	2.13E+01	EPA6 e	1.69E+01	EPA6 e	1.70E+00	EPA6 e
Bromomethane	74-83-9	7.98E+00	EPA6 e	8.87E+01	EPA6 e	8.79E+00	EPA6 e
Carbon tetrachloride	56-23-5	3.50E+01	EPA6 e	2.30E+01	EPA6 e	1.04E+00	EPA6 e
Chlordibromomethane	124-48-1	1.73E+01	EPA6 e	2.45E+01	EPA6 e	2.14E+00	EPA6 e
Chlorodifluoromethane	75-45-6	7.88E+00	EPA6 e	8.01E+01	EPA6 e	9.21E+00	EPA6 e
Chloroethane	75-00-3	6.30E+01	EPA6 e	1.70E+01	EPA6 e	6.25E-01	EPA6 e
Chloroform	67-66-3	1.37E+01	EPA6 e	2.58E+01	EPA6 e	2.89E+00	EPA6 e
Chloromethane	74-87-3	7.46E+00	EPA6 e	1.24E+02	EPA6 e	1.16E+01	EPA6 e
Chloropentafluoroethane	76-15-3	NA	NA	NA	NA	NA	NA
cis-1,2-Dichloroethene	156-59-2	1.41E+01	EPA6 e	2.38E+00	EPA6 e	2.77E+00	EPA6 e
cis-1,3-Dichloropropene	10061-01-5	NA	NA	NA	NA	NA	NA
Cyanogen bromide	506-68-3	NA	NA	NA	NA	NA	NA
Cyanogen chloride	506-77-4	NA	NA	NA	NA	NA	NA
Dichlorodifluoromethane	75-71-8	1.70E+01	EPA6 e	2.48E+01	EPA6 e	2.19E+00	EPA6 e
Dichlorofluoromethane	75-43-4	NA	NA	NA	NA	NA	NA
Dichloromethane	75-09-2	8.46E+00	EPA6 e	8.46E+01	EPA6 e	7.29E+00	EPA6 e
Difluorodibromomethane	75-61-6	NA	NA	NA	NA	NA	NA
Hexafluoroacetone	684-16-2	NA	NA	NA	NA	NA	NA
Iodomethane	74-88-4	1.42E+00	EPA6 calc c	3.11E+00	EPA6 calc c	4.08E+00	EPA6 calc c
Methylene bromide	74-95-3	1.04E+01	EPA6 e	4.01E+01	EPA6 e	4.48E+00	EPA6 e
Pentachloroethane	76-01-7	7.56E+00	EPA6 calc c	7.59E-01	EPA6 calc c	6.68E-01	EPA6 calc c

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	RCF ((ug/g DW plant)/ (ug/mL soil water))	Source	$Br_{root\ veg}$ ((ug/g DW plant) /(ug/g soil))	Source	Br_{ag} ((ug/g DW plant) /(ug/g soil))	Source
trans-1,2-Dichloroethene	156-60-5	1.41E+01	EPA6 e	3.71E+01	EPA6 e	2.77E+00	EPA6 e
trans-1,3-Dichloropropene	10061-02-6	1.98E+00	EPA6 calc c	1.87E+00	EPA6 calc c	2.50E+00	EPA6 calc c
Trichloroacetic acid	76-03-9	1.14E+00	EPA6 calc c	5.61E+00	EPA6 calc c	6.60E+00	EPA6 calc c
Trichlorofluoroethane	27154-33-2	NA	NA	NA	NA	NA	NA
Trichlorofluoromethane	75-69-4	2.70E+01	EPA6 e	2.02E+00	EPA6 e	1.33E+00	EPA6 e
Trifluorobromomethane	75-63-8	1.64E+00	EPA6 calc c	2.43E+00	EPA6 calc c	3.26E+00	EPA6 calc c
Dioxin and Furan Compounds							
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	35822-46-9	4.79E+05	EPA6 e	4.90E-01	EPA6 e	7.05E-04	EPA6 e
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	2.91E+05	EPA6 e	5.68E-01	EPA6 e	1.02E-03	EPA6 e
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	2.91E+05	EPA6 e	5.68E-01	EPA6 e	1.02E-03	EPA6 e
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	39227-28-6	2.31E+05	EPA6 e	6.09E-01	EPA6 e	1.22E-03	EPA6 e
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	8.88E+04	EPA6 e	8.10E-01	EPA6 e	2.50E-03	EPA6 e
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	57653-85-7	8.88E+04	EPA6 e	8.10E-01	EPA6 e	2.50E-03	EPA6 e
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	8.88E+04	EPA6 e	8.10E-01	EPA6 e	2.50E-03	EPA6 e
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	19408-74-3	8.88E+04	EPA6 e	8.10E-01	EPA6 e	2.50E-03	EPA6 e
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	8.88E+04	EPA6 e	8.10E-01	EPA6 e	2.50E-03	EPA6 e
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	40321-76-4	3.01E+04	EPA6 e	1.12E+00	EPA6 e	5.62E-03	EPA6 e
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	3.93E+04	EPA6 e	1.03E+00	EPA6 e	4.61E-03	EPA6 e
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	8.88E+04	EPA6 e	8.10E-01	EPA6 e	2.50E-03	EPA6 e
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	4.95E+04	EPA6 e	9.65E-01	EPA6 e	3.87E-03	EPA6 e
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1746-01-6	3.01E+04	EPA6 e	1.12E+00	EPA6 e	5.62E-03	EPA6 e
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	2.48E+04	EPA6 e	1.19E+00	EPA6 e	6.51E-03	EPA6 e
Dibenzofuran	132-64-9	6.60E+01	EPA6 calc c	3.66E-01	EPA6 calc c	1.22E-01	EPA6 calc c
Octachlorodibenzo(p)dioxin	3268-87-9	1.62E+05	EPA6 e	6.77E-01	EPA6 e	1.59E-03	EPA6 e
Octachlorodibenzofuran	39001-02-0	1.34E+06	EPA6 e	3.60E-01	EPA6 e	3.26E-04	EPA6 e
PCBs							
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	8.54E+03	EPA6 calc c	1.21E-01	EPA6 calc c	3.13E-03	EPA6 calc c
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	9.17E+03	EPA6 calc c	1.10E-01	EPA6 calc c	2.97E-03	EPA6 calc c
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	NA	NA	NA	NA	NA	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	RCF ((ug/g DW plant)/ (ug/mL soil water))	Source	$Br_{root\ veg}$ ((ug/g DW plant) /(ug/g soil))	Source	Br_{ag} ((ug/g DW plant) /(ug/g soil))	Source
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	NA	NA	NA	NA	NA	NA
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	NA	NA	NA	NA	NA	NA
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	NA	NA	NA	NA	NA	NA
2',3,4,4',5-Pentachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	9.17E+03	EPA6 calc c	9.19E-02	EPA6 calc c	2.97E-03	EPA6 calc c
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	1.53E+04	EPA6 calc c	3.84E-01	EPA6 calc c	2.02E-03	EPA6 calc c
3,3',4,4',5-Pentachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	NA	NA	NA	NA	NA	NA
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	NA	NA	NA	NA	NA	NA
Polychlorinated biphenyls (PCBs)**	1336-36-3	1.28E+04	EPA6 calc c	4.15E+00	EPA6 calc c	2.31E-03	EPA6 calc c
Phthalates							
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	2.37E+03	EPA6 e	2.13E+00	EPA6 e	3.80E-02	EPA6 e
Butylbenzyl phthalate	85-68-7	5.87E+02	EPA6 e	4.27E+00	EPA6 e	1.09E-01	EPA6 e
Dibutyl phthalate	84-74-2	1.01E+03	EPA6 e	6.43E+01	EPA6 e	7.24E-02	EPA6 e
Diethyl phthalate	84-66-2	6.12E+02	EPA6 e	7.46E+02	EPA6 e	1.06E-01	EPA6 e
Dimethyl phthalate	131-11-3	1.05E+01	EPA6 e	3.95E+00	EPA6 e	4.40E+00	EPA6 e
n-Dioctyl phthalate	117-84-0	3.55E+06	EPA6 e	3.93E-01	EPA6 e	1.57E-04	EPA6 e
Light Polycyclic Aromatic Hydrocarbons							
2-Chloronaphthalene	91-58-7	3.23E+02	EPA6 e	4.51E+00	EPA6 e	1.72E-01	EPA6 e
2-Methyl naphthalene	91-57-6	2.91E+01	EPA6 calc c	4.68E-01	EPA6 calc c	2.27E-01	EPA6 calc c
5-Nitroacenaphthene	602-87-9	NA	NA	NA	NA	NA	NA
Acenaphthene	83-32-9	2.69E+02	EPA6 e	5.48E+00	EPA6 e	1.98E-01	EPA6 e
Acenaphthylene	208-96-8	4.19E+01	EPA6 calc c	6.20E-01	EPA6 calc c	1.72E-01	EPA6 calc c
Anthracene	120-12-7	6.49E+02	EPA6 e	2.76E+00	EPA6 e	1.01E-01	EPA6 e
Fluorene	86-73-7	5.08E+01	EPA6 calc c	3.59E-01	EPA6 calc c	1.49E-01	EPA6 calc c
Indene	95-13-6	NA	NA	NA	NA	NA	NA
Napthalene	91-20-3	9.81E+01	EPA6 e	8.23E+00	EPA6 e	4.35E-01	EPA6 e

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	RCF ((ug/g DW plant)/ (ug/mL soil water))	Source	$Br_{root\ veg}$ ((ug/g DW plant) /(ug/g soil))	Source	Br_{ag} ((ug/g DW plant) /(ug/g soil))	Source
Phenanthrene	85-01-8	7.47E+02	EPA6 e	1.49E+00	EPA6 e	9.08E-02	EPA6 e
Pyrene	129-00-0	1.66E+03	EPA6 e	2.44E+00	EPA6 e	4.98E-02	EPA6 e
Heavy Polycyclic Aromatic Hydrocarbons							
3-Methylcholanthrene	56-49-5	9.01E+03	EPA6 calc c	5.95E-01	EPA6 calc c	3.01E-03	EPA6 calc c
5-Methylchrysene	3697-24-3	NA	NA	NA	NA	NA	NA
Benzo(a)anthracene	56-55-3	5.48E+03	EPA6 e	2.11E+00	EPA6 e	2.02E-02	EPA6 e
Benzo(a)pyrene	50-32-8	1.22E+04	EPA6 e	1.26E+00	EPA6 e	1.11E-02	EPA6 e
Benzo(b)fluoranthene	205-99-2	1.39E+04	EPA6 e	1.66E+00	EPA6 e	1.01E-02	EPA6 e
Benzo(e)pyrene	192-97-2	1.51E+04	EPA6 calc c	9.51E-02	EPA6 calc c	2.05E-03	EPA6 calc c
Benzo(g,h,i)perylene	191-24-2	8.85E+03	EPA6 calc c	4.86E-01	EPA6 calc c	3.05E-03	EPA6 calc c
Benzo(j)fluoranthene	205-82-3	2.75E+03	EPA6 calc c	1.28E-01	EPA6 calc c	7.34E-03	EPA6 calc c
Benzo(k)fluoranthene	207-08-9	1.38E+04	EPA6 e	1.66E+00	EPA6 e	1.01E-02	EPA6 e
Benzo[a,i]pyrene	191-30-0	NA	NA	NA	NA	NA	NA
Chrysene	218-01-9	6.10E+03	EPA6 e	2.05E+00	EPA6 e	1.87E-02	EPA6 e
Dibenz(a,h)anthracene	53-70-3	2.56E+04	EPA6 e	1.43E-04	EPA6 e	6.36E-03	EPA6 e
Dibenz[a,h]acridine	226-36-8	NA	NA	NA	NA	NA	NA
Dibenz[a,j]acridine	224-42-0	NA	NA	NA	NA	NA	NA
Dibenzo(a,e)fluoranthene	5385-75-1	NA	NA	NA	NA	NA	NA
Dibenzo(a,h)fluoranthene	no cas #	NA	NA	NA	NA	NA	NA
Dibenzo[a,e]pyrene	192-65-4	NA	NA	NA	NA	NA	NA
Dibenzo[a,h]pyrene	189-64-0	NA	NA	NA	NA	NA	NA
Dibenzo[a,i]pyrene	189-55-9	1.24E+04	EPA6 calc c	8.45E-02	EPA6 calc c	2.37E-03	EPA6 calc c
Fluoranthene	206-44-0	1.92E+03	EPA6 e	3.90E+00	EPA6 e	4.46E-02	EPA6 e
Hexachloronaphthalene	1335-87-1	2.11E+04	EPA6 calc c	7.30E-02	EPA6 calc c	1.59E-03	EPA6 calc c
Indeno(1,2,3-cd)pyrene	193-39-5	4.91E+04	EPA6 e	1.19E+00	EPA6 e	3.90E-03	EPA6 e
Octachloronaphthalene	2234-13-1	2.65E+03	EPA6 calc c	1.30E-01	EPA6 calc c	7.54E-03	EPA6 calc c
Pentachloronaphthalene	1321-64-8	NA	NA	NA	NA	NA	NA
Tetrachloronaphthalene	1335-88-2	NA	NA	NA	NA	NA	NA
Trichloronaphthalene	1321-65-9	NA	NA	NA	NA	NA	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Octanol / Water Partitioning Coefficients					
Constituent of Potential Concern	CAS Registry Number	Source	K_{ow} (Published Value)	K_{ow} (unitless)	Source	K_{oc} (Published Value)	K_{oc} (mL/g)	Source
trans-1,2-Dichloroethene	156-60-5	EPA6 e	9.60E+01	9.60E+01	EPA6 e	3.80E+01	3.80E+01	EPA6 e
trans-1,3-Dichloropropene	10061-02-6	EPA6 calc c	2.06E+00	1.15E+02	m	NA	1.06E+02	EPA6 calc c
Trichloroacetic acid	76-03-9	EPA6 calc c	1.33E+00	2.14E+01	Properties a	NA	2.03E+01	EPA6 calc c
Trichlorofluoroethane	27154-33-2	EPA6 calc c	NA	NA	NA	NA	NA	NA
Trichlorofluoromethane	75-69-4	EPA6 e	3.40E+02	3.40E+02	EPA6 e	NA	3.08E+02	EPA6 e
Trifluorobromomethane	75-63-8	EPA6 calc c	1.86E+00	7.24E+01	Hansch j	NA	6.74E+01	EPA6 calc c
Dioxin and Furan Compounds								
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	35822-46-9	EPA6 e	1.58E+08	1.58E+08	EPA6 e	9.77E+07	9.77E+07	EPA6 e
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	EPA6 e	8.32E+07	8.32E+07	EPA6 e	5.13E+07	5.13E+07	EPA6 e
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	EPA6 e	8.32E+07	8.32E+07	EPA6 e	5.13E+07	5.13E+07	EPA6 e
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	39227-28-6	EPA6 e	6.17E+07	6.17E+07	EPA6 e	3.80E+07	3.80E+07	EPA6 e
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	EPA6 e	1.78E+07	1.78E+07	EPA6 e	1.10E+07	1.10E+07	EPA6 e
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	57653-85-7	EPA6 e	1.78E+07	1.78E+07	EPA6 e	1.10E+07	1.10E+07	EPA6 e
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	EPA6 e	1.78E+07	1.78E+07	EPA6 e	1.10E+07	1.10E+07	EPA6 e
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	19408-74-3	EPA6 e	1.78E+07	1.78E+07	EPA6 e	1.10E+07	1.10E+07	EPA6 e
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	EPA6 e	1.78E+07	1.78E+07	EPA6 e	1.10E+07	1.10E+07	EPA6 e
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	40321-76-4	EPA6 e	4.37E+06	4.37E+06	EPA6 e	2.69E+06	2.69E+06	EPA6 e
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	EPA6 e	6.17E+06	6.17E+06	EPA6 e	3.80E+06	3.80E+06	EPA6 e
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	EPA6 e	1.78E+07	1.78E+07	EPA6 e	1.10E+07	1.10E+07	EPA6 e
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	EPA6 e	8.32E+06	8.32E+06	EPA6 e	5.13E+06	5.13E+06	EPA6 e
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1746-01-6	EPA6 e	4.37E+06	4.37E+06	EPA6 e	2.69E+04	2.69E+04	EPA6 e
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	EPA6 e	3.39E+06	3.39E+06	EPA6 e	2.09E+06	2.09E+06	EPA6 e
Dibenzofuran	132-64-9	EPA6 calc c	4.33E+00	2.14E+04	Properties a	NA	1.81E+04	EPA6 calc c
Octachlorodibenzo(p)dioxin	3268-87-9	EPA6 e	3.89E+07	3.89E+07	EPA6 e	2.40E+07	2.40E+07	EPA6 e
Octachlorodibenzofuran	39001-02-0	EPA6 e	6.03E+08	6.03E+08	EPA6 e	3.72E+08	3.72E+08	EPA6 e
PCBs								
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	EPA6 calc c	7.08E+00	1.20E+07	Properties a	6.85E+00	7.08E+06	Properties a
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	EPA6 calc c	7.12E+00	1.32E+07	Properties a	6.92E+00	8.32E+06	Properties a
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	EPA6 calc c	NA	NA	NA	NA	NA	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	RCF ((ug/g DW plant)/ (ug/mL soil water))	Source	$Br_{root\ veg}$ ((ug/g DW plant) /(ug/g soil))	Source	Br_{ag} ((ug/g DW plant) /(ug/g soil))	Source
Light Substituted Benzene Compounds							
1,2,3-Trichlorobenzene	87-61-6	3.09E+02	EPA6 e	1.53E+00	EPA6 e	1.78E-01	EPA6 e
1,2,4-Trichlorobenzene	120-82-1	2.80E+02	EPA6 e	1.69E+01	EPA6 e	1.92E-01	EPA6 e
1,2,4-Trimethlyl benzene	25551-13-7	1.34E+01	EPA6 calc c	2.26E+00	EPA6 calc c	4.20E-01	EPA6 calc c
1,2,4-Trimethyl benzene	95-63-6	2.03E+01	EPA6 calc c	5.25E-01	EPA6 calc c	3.01E-01	EPA6 calc c
1,2-Dichlorobenzene	95-50-1	1.11E+02	EPA6 e	2.92E+01	EPA6 e	3.95E-01	EPA6 e
1,3,5-Trimethyl benzene	108-67-8	1.06E+02	EPA6 e	6.35E+00	EPA6 e	4.09E-01	EPA6 e
1,3-Dichlorobenzene	541-73-1	1.28E+02	EPA6 e	1.59E+00	EPA6 e	3.53E-01	EPA6 e
1,3-Dinitrobenzene	99-65-0	9.58E+00	EPA6 e	4.64E+01	EPA6 e	5.32E+00	EPA6 e
1,4-Dichlorobenzene	106-46-7	1.05E+02	EPA6 e	1.70E+01	EPA6 e	4.13E-01	EPA6 e
1,4-Dinitrobenzene	100-25-4	1.25E+00	EPA6 calc c	4.19E+00	EPA6 calc c	5.26E+00	EPA6 calc c
2,4,5-Trichorophenol	95-95-4	2.28E+02	EPA6 e	2.02E+01	EPA6 e	2.24E-01	EPA6 e
2,4,6-Trichlorophenol	88-06-2	1.74E+01	EPA6 e	7.69E+01	EPA6 e	2.77E-01	EPA6 e
2,4-Dichlorophenol	120-83-2	5.68E+01	EPA6 e	4.07E+01	EPA6 e	6.82E-01	EPA6 e
2,4-Dimethylphenol	105-67-9	2.16E+01	EPA6 e	1.71E+01	EPA6 e	1.68E+00	EPA6 e
2,4-Dinitrophenol	51-28-5	9.74E+00	EPA6 e	9.74E+04	EPA6 e	5.13E+00	EPA6 e
2,4-Dinitrotoluene	121-14-2	1.43E+01	EPA6 e	2.80E+01	EPA6 e	2.72E+00	EPA6 e
2,6-Dinitrotoluene	606-20-2	1.29E+01	EPA6 e	3.08E+01	EPA6 e	3.15E+00	EPA6 e
2-Chlorotoluene	95-49-8	1.68E+01	EPA6 calc c	2.22E+00	EPA6 calc c	3.50E-01	EPA6 calc c
2-Chorophenol	95-57-8	1.70E+01	EPA6 e	4.40E+00	EPA6 e	2.18E+00	EPA6 e
2-Nitrophenol	88-75-5	1.19E+01	EPA6 e	3.36E+00	EPA6 e	3.57E+00	EPA6 e
4,6-Dinitro-o-cresol	534-52-1	5.55E+00	EPA6 calc c	8.75E-01	EPA6 calc c	8.72E-01	EPA6 calc c
4-Chlorotoluene	106-43-4	1.19E+01	EPA6 calc c	6.33E-01	EPA6 calc c	4.61E-01	EPA6 calc c
4-Nitrophenol	100-02-7	1.32E+01	EPA6 e	3.01E+00	EPA6 e	3.05E+00	EPA6 e
alpha-Methylstyrene	98-83-9	1.48E+01	EPA6 calc c	2.24E+00	EPA6 calc c	3.86E-01	EPA6 calc c
Aniline	62-53-3	7.63E+00	EPA6 e	9.27E+01	EPA6 e	1.05E+01	EPA6 e
Benzotrichloride	98-07-7	6.17E+00	EPA6 calc c	8.31E-01	EPA6 calc c	7.95E-01	EPA6 calc c
Benzyl Chloride	100-44-7	6.75E+00	EPA6 e	2.49E+02	EPA6 e	2.39E+01	EPA6 e
Bromobenzene	108-86-1	6.88E+00	EPA6 calc c	1.54E+00	EPA6 calc c	7.24E-01	EPA6 calc c

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	RCF ((ug/g DW plant)/ (ug/mL soil water))	Source	$Br_{root\ veg}$ ((ug/g DW plant) /(ug/g soil))	Source	Br_{ag} ((ug/g DW plant) /(ug/g soil))	Source
Chlorobenzene	108-90-7	3.90E+01	EPA6 e	1.74E+01	EPA6 e	9.45E-01	EPA6 e
Cumene	98-82-8	1.47E+02	EPA6 e	1.58E+00	EPA6 e	3.16E-01	EPA6 e
m-Cresol	108-39-4	1.38E+01	EPA6 e	2.89E+01	EPA6 e	2.86E+00	EPA6 e
n-Butyl benzene	104-51-8	6.05E+01	EPA6 calc c	2.41E+00	EPA6 calc c	1.30E-01	EPA6 calc c
Nitrobenzene	98-95-3	1.23E+01	EPA6 e	1.03E+01	EPA6 e	3.38E+00	EPA6 e
n-Propyl benzene	103-65-1	2.18E+01	EPA6 calc c	3.01E+00	EPA6 calc c	2.85E-01	EPA6 calc c
o-Cresol	95-48-7	1.47E+01	EPA6 e	2.75E+01	EPA6 e	2.63E+00	EPA6 e
o-Dinitrobenzene	528-29-0	1.34E+00	EPA6 calc c	3.57E+00	EPA6 calc c	4.60E+00	EPA6 calc c
o-Nitroaniline	88-74-4	1.25E+01	EPA6 e	3.18E+00	EPA6 e	3.30E+00	EPA6 e
o-Toluidine	95-53-4	8.81E+00	EPA6 e	5.61E+01	EPA6 e	6.51E+00	EPA6 e
p-Chloroaniline	106-47-8	1.27E+01	EPA6 e	3.12E+01	EPA6 e	3.22E+00	EPA6 e
p-Cresol	106-44-5	1.35E+01	EPA6 e	2.94E+01	EPA6 e	2.93E+00	EPA6 e
Phenol	108-95-2	9.50E+00	EPA6 e	4.32E+01	EPA6 e	5.42E+00	EPA6 e
p-Nitrochlorobenzene	100-00-5	2.91E+00	EPA6 calc c	1.30E+00	EPA6 calc c	1.61E+00	EPA6 calc c
p-Toluidine	106-49-0	1.18E+00	EPA6 calc c	4.97E+00	EPA6 calc c	6.01E+00	EPA6 calc c
sec-Butyl benzene	135-98-8	NA	NA	NA	NA	NA	NA
tert-Butyl benzene	98-06-6	4.49E+01	EPA6 calc c	4.10E-01	EPA6 calc c	1.63E-01	EPA6 calc c
Toluene-2,6-diamine	823-40-5	1.22E+00	EPA6 calc c	4.52E+00	EPA6 calc c	5.59E+00	EPA6 calc c
Other Light Semivolatile Compounds							
1,1 Dimethyl hydrazine	57-14-7	9.07E-01	EPA6 calc c	2.36E+01	EPA6 calc c	1.75E+01	EPA6 calc c
1,1-Biphenyl	92-52-4	3.12E+01	EPA6 calc c	1.24E+00	EPA6 calc c	2.16E-01	EPA6 calc c
1,2 Diphenylhydrazine	122-66-7	4.90E+01	EPA6 e	1.76E+01	EPA6 e	7.74E-01	EPA6 e
1,2-Dimethylhydrazine	540-73-8	8.23E-01	EPA6 calc c	1.82E+03	EPA6 calc c	2.39E+02	EPA6 calc c
1,3-Propane sultone	1120-71-4	8.32E-01	EPA6 calc c	2.72E+02	EPA6 calc c	7.77E+01	EPA6 calc c
2,4-Toluene diisocyanate	584-84-9	NA	NA	NA	NA	NA	NA
2-Chloroacetophenone	532-27-4	3.83E+00	EPA6 calc c	5.03E+00	EPA6 calc c	1.23E+00	EPA6 calc c
2-Propenoic acid	79-10-7	8.85E-01	EPA6 calc c	3.34E+01	EPA6 calc c	2.19E+01	EPA6 calc c
4,4-Methylenedianiline	101-77-9	1.29E+01	EPA6 calc c	6.14E-01	EPA6 calc c	4.31E-01	EPA6 calc c
Acetophenone	98-86-2	1.06E+01	EPA6 e	3.92E+01	EPA6 e	4.37E+00	EPA6 e

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	RCF ((ug/g DW plant)/ (ug/mL soil water))	Source	$Br_{root\ veg}$ ((ug/g DW plant) /(ug/g soil))	Source	Br_{ag} ((ug/g DW plant) /(ug/g soil))	Source
Benzoic acid	65-85-0	1.28E+01	NA	2.33E+03	NA	3.17E+00	NA
bis(2-Chloroethoxy)methane	111-91-1	NA	NA	NA	NA	NA	NA
bis(2-Chloroethyl) ether	111-44-4	8.64E+00	EPA6 e	1.14E+01	EPA6 e	6.85E+00	EPA6 e
Chlorocyclopentadiene	41851-50-7	3.06E+00	EPA6 calc c	1.25E+00	EPA6 calc c	1.53E+00	EPA6 calc c
Cyclohexanol	108-93-0	1.09E+00	EPA6 calc c	8.36E+00	EPA6 calc c	7.53E+00	EPA6 calc c
Dichloroisopropyl ether	108-60-1	3.75E+00	EPA6 calc c	6.15E+00	EPA6 calc c	1.25E+00	EPA6 calc c
Dichloromethyl ether	542-88-1	8.35E-01	EPA6 calc c	1.05E+00	EPA6 calc c	6.42E+01	EPA6 calc c
Dichloropentadiene	no cas #	NA	NA	NA	NA	NA	NA
Dimethyl sulfate	77-78-1	8.73E-01	EPA6 calc c	4.25E+01	EPA6 calc c	2.54E+01	EPA6 calc c
Dimethylaniline	121-69-7	8.26E-01	EPA6 calc c	6.05E+02	EPA6 calc c	1.25E+02	EPA6 calc c
di-n-propylnitrosamine	621-64-7	8.99E+00	EPA6 e	5.29E+01	EPA6 e	6.17E+00	EPA6 e
Diphenyl ether	101-84-8	5.35E+01	EPA6 calc c	3.89E-01	EPA6 calc c	1.43E-01	EPA6 calc c
Epichlorohydrin	106-89-8	6.67E+00	EPA6 e	3.00E+02	EPA6 e	2.77E+01	EPA6 e
Ethyl carbomate (urethane)	51-79-6	8.43E-01	EPA6 calc c	1.18E+02	EPA6 calc c	4.73E+01	EPA6 calc c
Ethyl methane sulfanate	62-50-0	6.56E+00	EPA6 e	4.24E+02	EPA6 e	3.63E+01	EPA6 e
Ethylene dibromide	106-93-4	1.15E+01	EPA6 e	3.50E+01	EPA6 e	3.77E+00	EPA6 e
Ethylene glycol	107-21-1	8.26E-01	EPA6 calc c	6.54E+02	EPA6 calc c	1.31E+02	EPA6 calc c
Ethylene glycol monobutyl ether	111-76-2	1.30E+00	EPA6 calc c	2.10E+00	EPA6 calc c	4.89E+00	EPA6 calc c
Ethylene glycol monoethyl ether acetate	111-15-9	9.11E-01	EPA6 calc c	2.24E+01	EPA6 calc c	1.70E+01	EPA6 calc c
Ethylene thiourea	96-45-7	8.30E-01	EPA6 calc c	3.54E+02	EPA6 calc c	9.10E+01	EPA6 calc c
Furfural	98-01-1	9.86E-01	EPA6 calc c	1.12E+01	EPA6 calc c	1.08E+01	EPA6 calc c
Maleic hydrazide	123-33-1	8.28E-01	EPA6 calc c	4.42E+02	EPA6 calc c	1.04E+02	EPA6 calc c
Malononitrile	109-77-3	8.52E-01	EPA6 calc c	6.62E+01	EPA6 calc c	3.67E+01	EPA6 calc c
Methyl styrene (mixed isomers)	25013-15-4	1.23E+01	EPA6 calc c	3.32E+00	EPA6 calc c	4.48E-01	EPA6 calc c
Methylhydrazine	60-34-4	8.25E-01	EPA6 calc c	8.88E+02	EPA6 calc c	1.57E+02	EPA6 calc c
N,N-Diphenylamine	122-39-4	1.58E+01	EPA6 calc c	4.55E+00	EPA6 calc c	3.67E-01	EPA6 calc c
Nitric acid, propyl ester	627-13-4	NA	NA	NA	NA	NA	NA
N-Nitrosodi-n-butylamine	924-16-3	2.30E+01	EPA6 e	2.14E+01	EPA6 e	1.57E+00	EPA6 e
N-Nitrosomorpholine	59-89-2	9.92E-01	EPA6 calc c	1.08E+01	EPA6 calc c	1.05E+01	EPA6 calc c

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	RCF ((ug/g DW plant)/ (ug/mL soil water))	Source	$Br_{root\ veg}$ ((ug/g DW plant) /(ug/g soil))	Source	Br_{ag} ((ug/g DW plant) /(ug/g soil))	Source
N-Nitroso-N,N-dimethylamine	62-75-9	8.33E-01	EPA6 calc c	1.00E+01	EPA6 calc c	7.24E+01	EPA6 calc c
o-Anisidine	90-04-0	1.06E+00	EPA6 calc c	7.36E+00	EPA6 calc c	8.05E+00	EPA6 calc c
Oxalic acid	144-62-7	NA	NA	NA	NA	NA	NA
Phthalic Anhydride	85-44-9	6.39E+00	EPA6 e	1.33E+03	EPA6 e	8.63E+01	EPA6 e
p-Phthalic acid	100-21-0	9.50E-01	EPA6 calc c	1.47E+01	EPA6 calc c	1.30E+01	EPA6 calc c
Pyridine	110-86-1	7.07E+00	EPA6 e	1.50E+02	EPA6 e	1.59E+01	EPA6 e
Quinoline	91-22-5	1.92E+00	EPA6 calc c	1.94E+00	EPA6 calc c	2.60E+00	EPA6 calc c
Quinone	106-51-4	8.63E-01	EPA6 calc c	5.49E+01	EPA6 calc c	2.97E+01	EPA6 calc c
Safrole	94-59-7	3.23E+01	EPA6 e	1.92E+01	EPA6 e	1.12E+00	EPA6 e
Tetrahydrofuran	109-99-9	6.82E+00	EPA6 e	2.16E+02	EPA6 e	2.14E+01	EPA6 e
Other Heavy Semivolatile Compounds							
1,2,4,5-Tetrachlorobenzene	95-94-3	8.75E+02	EPA6 e	1.49E+01	EPA6 e	8.06E-02	EPA6 e
1,3,5-Trinitrobenzene	99-35-4	8.19E+00	EPA6 e	6.95E+01	EPA6 e	8.05E+00	EPA6 e
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	4.99E+01	EPA6 calc c	3.97E-01	EPA6 calc c	1.51E-01	EPA6 calc c
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	8.20E-01	EPA6 calc c	3.70E+04	EPA6 calc c	1.41E+03	EPA6 calc c
2-sec-Butyl-4,6-dinitrophenol	88-85-7	8.72E+00	EPA6 calc c	7.14E-01	EPA6 calc c	5.93E-01	EPA6 calc c
3,3-Dichlorobenzidine	91-94-1	1.38E+02	EPA6 e	1.58E+01	EPA6 e	3.32E-01	EPA6 e
3,3'-Dimethoxybenzidine	119-90-4	1.21E+01	EPA6 e	3.30E+01	EPA6 e	3.48E+00	EPA6 e
4-Bromophenylphenyl ether	101-55-3	2.15E+02	EPA6 calc c	2.61E-01	EPA6 calc c	4.99E-02	EPA6 calc c
Ammonium perfluorooctanoate	3825-26-1	NA	NA	NA	NA	NA	NA
Azobenzene	103-33-3	2.72E+01	EPA6 calc c	1.01E+00	EPA6 calc c	2.40E-01	EPA6 calc c
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	NA	NA	NA	NA	NA	NA
Captan	133-06-2	2.77E+00	EPA6 calc c	1.39E+00	EPA6 calc c	1.70E+00	EPA6 calc c
Chlorobenzilate	510-15-6	5.54E+02	EPA6 e	1.50E+01	EPA6 e	1.14E-01	EPA6 e
Dibutylphosphate	107-66-4	NA	NA	NA	NA	NA	NA
Dimethyl aminoazobenzene	60-11-7	1.02E+02	EPA6 calc c	3.22E-01	EPA6 calc c	8.72E-02	EPA6 calc c
Hexachlorobenzene	118-74-1	4.02E+03	EPA6 e	5.02E+00	EPA6 e	2.56E-02	EPA6 e
Hexachlorobutadiene	87-68-3	1.03E+03	EPA6 e	1.48E+01	EPA6 e	7.14E-02	EPA6 e
Hexachlorocyclopentadiene	77-47-4	1.40E+03	EPA6 e	1.47E+01	EPA6 e	5.65E-02	EPA6 e

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	RCF ((ug/g DW plant)/ (ug/mL soil water))	Source	$Br_{root\ veg}$ ((ug/g DW plant) /(ug/g soil))	Source	Br_{ag} ((ug/g DW plant) /(ug/g soil))	Source
Hexachloroethane	67-72-1	2.78E+02	EPA6 e	1.53E+01	EPA6 e	1.93E-01	EPA6 e
Hexachlorophene	70-30-4	1.49E+05	EPA6 e	1.38E+00	EPA6 e	1.70E-03	EPA6 e
Hexamethylene-1,5-diisocyanate	822-06-0	1.11E+00	EPA6 calc c	6.21E+00	EPA6 calc c	7.11E+00	EPA6 calc c
Mirex	2385-85-5	6.10E+03	EPA6 calc c	6.10E-01	EPA6 calc c	4.03E-03	EPA6 calc c
Nitrofen	1836-75-5	5.48E+02	EPA6 calc c	2.01E-01	EPA6 calc c	2.46E-02	EPA6 calc c
Pentachlorobenzene	608-93-5	1.93E+03	EPA6 e	5.99E+00	EPA6 e	4.44E-02	EPA6 e
Pentachloronitrobenzene	82-68-8	8.75E+02	EPA6 e	1.49E+01	EPA6 e	8.06E-02	EPA6 e
Pentachlorophenol*	87-86-5	1.90E+03	EPA6 e	3.77E+02	EPA6 e	4.48E-02	EPA6 e
Pentachlorophenol*	87-86-5	1.90E+03	EPA6 e	3.77E+02	EPA6 e	4.48E-02	EPA6 e
Pentachlorophenol*	87-86-5	1.90E+03	EPA6 e	3.77E+02	EPA6 e	4.48E-02	EPA6 e
Pentachlorophenol*	87-86-5	1.90E+03	EPA6 e	3.77E+02	EPA6 e	4.48E-02	EPA6 e
Pentachlorophenol*	87-86-5	1.90E+03	EPA6 e	3.77E+02	EPA6 e	4.48E-02	EPA6 e
Pentachlorophenol*	87-86-5	1.90E+03	EPA6 e	3.77E+02	EPA6 e	4.48E-02	EPA6 e
Pentachlorophenol*	87-86-5	1.90E+03	EPA6 e	3.77E+02	EPA6 e	4.48E-02	EPA6 e
Pentachlorophenol*	87-86-5	1.90E+03	EPA6 e	3.77E+02	EPA6 e	4.48E-02	EPA6 e
Pentachlorophenol*	87-86-5	1.90E+03	EPA6 e	3.77E+02	EPA6 e	4.48E-02	EPA6 e
Pentachlorophenol*	87-86-5	1.90E+03	EPA6 e	3.77E+02	EPA6 e	4.48E-02	EPA6 e
Pentachlorophenol*	87-86-5	1.90E+03	EPA6 e	3.77E+02	EPA6 e	4.48E-02	EPA6 e
Pentachlorophenol*	87-86-5	1.90E+03	EPA6 e	3.77E+02	EPA6 e	4.48E-02	EPA6 e
Picric acid	88-89-1	1.92E+00	EPA6 calc c	1.94E+00	EPA6 calc c	2.60E+00	EPA6 calc c
Pronamide	23950-58-5	1.23E+02	EPA6 e	1.59E+01	EPA6 e	3.62E-01	EPA6 e
Strychnine	57-24-9	1.34E+01	EPA6 e	2.96E+01	EPA6 e	2.97E+00	EPA6 e
Terphenyls	26140-60-3	NA	NA	NA	NA	NA	NA
Tributyl phosphate	126-73-8	3.71E+01	EPA6 calc c	4.34E-01	EPA6 calc c	1.89E-01	EPA6 calc c
Trifluralin	1582-09-8	3.91E+02	EPA6 calc c	6.50E+00	EPA6 calc c	3.17E-02	EPA6 calc c
Triphenylamine	603-34-9	NA	NA	NA	NA	NA	NA
Herbicides and Organopesticides							
2,4,5-T	93-76-5	1.25E+01	EPA6 calc c	2.38E+01	EPA6 calc c	4.43E-01	EPA6 calc c

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	RCF ((ug/g DW plant)/ (ug/mL soil water))	Source	$Br_{root\ veg}$ ((ug/g DW plant) /(ug/g soil))	Source	Br_{ag} ((ug/g DW plant) /(ug/g soil))	Source
2,4-D and esters	94-75-7	5.22E+00	EPA6 calc c	2.62E+01	EPA6 calc c	9.20E-01	EPA6 calc c
4,4-DDD	72-54-8	1.80E+03	EPA6 calc c	1.80E+00	EPA6 calc c	1.01E-02	EPA6 calc c
4,4-DDT	50-29-3	1.26E+03	EPA6 calc c	5.25E-01	EPA6 calc c	1.32E-02	EPA6 calc c
Aldrin	309-00-2	1.33E+04	EPA6 e	2.73E+01	EPA6 e	1.04E-02	EPA6 e
alpha-BHC	319-84-6	2.02E+02	EPA6 e	1.15E+01	EPA6 e	2.47E-01	EPA6 e
beta-BHC	319-85-7	2.14E+02	EPA6 e	1.00E+01	EPA6 e	2.36E-01	EPA6 e
Chlordane	57-74-9	8.67E+03	EPA6 e	1.69E+01	EPA6 e	1.43E-02	EPA6 e
DDE	72-55-9	1.53E+04	EPA6 e	1.77E+01	EPA6 e	9.37E-03	EPA6 e
Delta-BHC	319-86-8	4.74E+01	EPA6 calc c	7.17E+00	EPA6 calc c	1.57E-01	EPA6 calc c
Dieldrin	60-57-1	2.66E+03	EPA6 e	1.04E+01	EPA6 e	3.49E-02	EPA6 e
Endothall	145-73-3	8.26E-01	EPA6 calc c	5.92E+02	EPA6 calc c	1.23E+02	EPA6 calc c
Endrin	72-20-8	1.36E+03	EPA6 e	1.26E+01	EPA6 e	5.76E-02	EPA6 e
gamma-BHC(Lindane)	58-89-9	2.29E+01	EPA6 calc c	2.14E+00	EPA6 calc c	2.74E-01	EPA6 calc c
Heptachlor	76-44-8	1.70E+03	EPA6 e	1.78E+01	EPA6 e	4.89E-02	EPA6 e
Isodrin	465-73-6	NA	NA	NA	NA	NA	NA
Methoxychlor	72-43-5	7.16E+02	EPA6 e	8.95E-01	EPA6 e	9.38E-02	EPA6 e
Silvex (2,4,5-TP)	93-72-1	4.21E+01	EPA6 calc c	4.18E-01	EPA6 calc c	1.71E-01	EPA6 calc c
Toxaphene	8001-35-2	5.20E+02	EPA6 calc c	5.20E-01	EPA6 calc c	2.56E-02	EPA6 calc c

NA = Value not available
** = Varies with pH; values given for a pH of 7.0.
a = Illustrated Handbook of Physical - Chemical Properties and Environmental Fate f
b = RAIS database: http://risk.lsd.ornl.gov/rap_hp.shtml
c = Equations derived from EPA Region 6 Guidance Appendix A-3
d = Handbook of Environmental degradation Rates. 1991.
e = EPA Region 6 Hazardous Waste Combustion Guidance
f = "A Critical Review of Henry's Law Constants for Chemicals of Environmental In
g = Handbook of Environmental Fate and Exposure Data for Organic Chemicals
h = http://www.epa.gov/region09/waste/sfund/prg/.
i = http://esc.syrres.com/efdb/Chemfate.htm
j = Hansch and Leo 1985 in Syracuse 1996

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	RCF ((ug/g DW plant)/ (ug/mL soil water))	Source	$Br_{root\ veg}$ ((ug/g DW plant) /(ug/g soil))	Source	Br_{ag} ((ug/g DW plant) /(ug/g soil))	Source

k = Schwarzenbach et al. 1993
l = Tomlin 1994 in Syracuse 1996
m = http://www.speclab.com/compound/c959988.htm

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	Br_{forage} ((ug/g DW plant) /(ug/g soil))	Source	Br_{grain} ((ug/g DW plant) /(ug/g soil))	Bv_{ag} ((ug/g DW plant) /(ug/g air))	Source	Bv_{forage} ((ug/g DW plant) /(ug/g air))
Aromatic Halogenated Hydrocarbons							
2,3,4,6-Tetrachlorophenol	58-90-2	1.08E-01	EPA6 calc c	NA	8.13E+02	EPA6 calc c	8.13E+02
4-Chloro-3-methylphenol	59-50-7	6.25E-01	EPA6 calc c	NA	NA	NA	NA
Aromatic Nonhalogenated Hydrocarbons							
2-Nitrotoluene	88-72-2	1.81E+00	EPA6 calc c	NA	2.74E-01	EPA6 calc c	2.74E-01
4-Nitrobiphenyl	92-93-3	2.56E-01	EPA6 calc c	NA	NA	NA	NA
Benzaldehyde	100-52-7	5.42E+00	EPA6 e	NA	5.00E-02	EPA6 e	5.00E-02
Benzene	71-43-2	2.25E+00	EPA6 e	NA	1.92E-03	EPA6 e	1.92E-03
Benzyl alcohol	100-51-6	8.95E+00	EPA6 e	NA	2.19E+00	EPA6 e	2.19E+00
Ethyl Benzene	100-41-4	6.07E-01	EPA6 e	NA	1.53E-02	EPA6 e	1.53E-02
m-Xylene	108-38-3	5.47E-01	EPA6 e	NA	1.55E+01	EPA6 e	1.55E+01
o-Xylene	95-47-6	6.01E-01	EPA6 e	NA	1.79E+01	EPA6 e	1.79E+01
p-Xylene	106-42-3	5.70E-01	EPA6 e	NA	1.93E+01	EPA6 e	1.93E+01
Styrene	100-42-5	7.85E-01	EPA6 e	NA	2.21E-02	EPA6 e	2.21E-02
Toluene	108-88-3	1.11E+00	EPA6 e	NA	6.33E-03	EPA6 e	6.33E-03
Non-aromatic Nonhalogenated Hydrocarbons							
1,2-Epoxybutane	106-88-7	5.69E+00	EPA6 calc c	NA	4.17E-03	EPA6 calc c	4.17E-03
1,3-Butadiene	106-99-0	2.74E+00	EPA6 calc c	NA	9.99E-05	EPA6 calc c	8.01E-05
1,4-Dioxane	123-91-1	5.53E+01	EPA6 e	NA	5.93E-03	EPA6 e	5.93E-03
1-Methylpropyl alcohol	78-92-2	1.72E+01	EPA6 calc c	NA	2.45E-02	EPA6 calc c	2.45E-02
1-Nitropropane	108-03-2	1.22E+01	EPA6 calc c	NA	NA	NA	NA
2,2,4-Trimethylpentane	540-84-1	4.86E-02	EPA6 calc c	NA	4.12E-03	EPA6 calc c	4.12E-03
2-Butanone	78-93-3	2.67E+01	EPA6 e	NA	3.08E-03	EPA6 e	3.08E-03
2-Butenaldehyde (2-Butenal)	4170-30-3	NA	NA	NA	NA	NA	NA
2-Ethoxyethanol	110-80-5	4.42E+01	EPA6 calc c	NA	8.55E-07	EPA6 calc c	8.55E-07
2-Heptanone	110-43-0	2.78E+00	EPA6 calc c	NA	4.98E-02	EPA6 calc c	4.98E-02
2-Hexanone	591-78-6	6.17E+00	EPA6 calc c	NA	1.72E-02	EPA6 calc c	1.72E-02
2-Methoxyethanol	109-86-4	2.77E+01	EPA6 calc c	NA	1.26E-01	EPA6 calc c	1.26E-01
2-Methyl-2-propanol	75-65-0	2.43E+01	EPA6 calc c	NA	1.10E-02	EPA6 calc c	1.10E-02

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	Br_{forage} ((ug/g DW plant) /(ug/g soil))	Source	Br_{grain} ((ug/g DW plant) /(ug/g soil))	Bv_{ag} ((ug/g DW plant) /(ug/g air))	Source	Bv_{forage} ((ug/g DW plant) /(ug/g air))
2-Methyl-2-propenenitrile	126-98-7	1.89E+01	EPA6 e	NA	8.81E-04	EPA6 e	8.81E-04
2-Methylaziridine	75-55-8	NA	NA	NA	NA	NA	NA
2-Methylpropyl alcohol	78-83-1	1.41E+01	EPA6 calc c	NA	3.01E-02	EPA6 calc c	3.01E-02
2-Pentanone	107-87-9	1.15E+01	EPA6 calc c	NA	8.09E-03	EPA6 calc c	8.09E-03
2-Propanone (Acetone)	67-64-1	5.20E+01	EPA6 e	NA	1.13E-03	EPA6 e	1.13E-03
2-Propene-1-ol	107-18-6	3.09E+01	EPA6 calc c	NA	1.70E-02	EPA6 calc c	1.70E-02
2-Propyl alcohol	67-63-0	3.62E+01	EPA6 calc c	NA	7.80E-03	EPA6 calc c	7.80E-03
3-Heptanone	106-35-4	NA	NA	NA	NA	NA	NA
3-Methyl-1-butanol	123-51-3	NA	NA	NA	NA	NA	NA
3-Methyl-2-butanone	563-80-4	NA	NA	NA	NA	NA	NA
3-Pentanone	96-22-0	1.04E+01	EPA6 calc c	NA	7.89E-03	EPA6 calc c	7.89E-03
4-Heptanone	123-19-3	NA	NA	NA	NA	NA	NA
4-Methyl-2-pentanone	108-10-1	7.95E+00	EPA6 e	NA	8.26E-03	EPA6 e	8.26E-03
4-Methyl-3-penten-2-one	141-79-7	NA	NA	NA	NA	NA	NA
5-Methyl-2-hexanone	110-12-3	NA	NA	NA	NA	NA	NA
Acetaldehyde	75-07-0	5.19E+01	EPA6 e	NA	NA	EPA6 e	NA
Acetamide	60-35-5	2.07E+02	EPA6 calc c	NA	7.30E-01	EPA6 calc c	7.30E-01
Acetic acid	64-19-7	4.86E+01	EPA6 calc c	NA	1.22E-01	EPA6 calc c	1.22E-01
Acetic acid ethyl ester	141-78-6	1.47E+01	EPA6 calc c	NA	2.49E-03	EPA6 calc c	2.49E-03
Acetic acid n-butyl ester	123-86-4	3.87E+00	EPA6 calc c	NA	1.18E-02	EPA6 calc c	1.18E-02
Acetonitrile	75-05-8	6.09E+01	EPA6 e	NA	6.41E-10	EPA6 e	6.41E-10
Acrolein	107-02-8	3.92E+01	EPA6 e	NA	5.86E-04	EPA6 e	5.86E-04
Acrylonitrile	107-13-1	2.77E+01	EPA6 e	NA	1.04E-03	EPA6 e	1.04E-03
Bis(isopropyl)ether	108-20-3	NA	NA	NA	NA	NA	NA
Butane	106-97-8	8.27E-01	EPA6 calc c	NA	7.06E-05	EPA6 calc c	1.18E-02
Carbon disulfide	75-15-0	2.70E+00	EPA6 e	NA	5.92E-04	EPA6 e	5.92E-04
Cyanogen	460-19-5	1.32E+01	EPA6 calc c	NA	8.19E-05	EPA6 calc c	1.18E-02
Cyclohexane	110-82-7	3.98E-01	EPA6 calc c	NA	1.33E-03	EPA6 calc c	1.33E-03
Cyclohexanone	108-94-1	1.32E+01	EPA6 calc c	NA	3.39E-02	EPA6 calc c	3.39E-02

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	Br_{forage} ((ug/g DW plant) /(ug/g soil))	Source	Br_{grain} ((ug/g DW plant) /(ug/g soil))	Bv_{ag} ((ug/g DW plant) /(ug/g air))	Source	Bv_{forage} ((ug/g DW plant) /(ug/g air))
Cyclohexene	110-83-8	8.61E-01	EPA6 calc c	NA	1.57E-03	EPA6 calc c	1.57E-03
Cyclopentane	287-92-3	7.14E-01	EPA6 calc c	NA	4.72E-04	EPA6 calc c	4.72E-04
Ethyl alcohol	64-17-5	5.85E+01	EPA6 calc c	NA	4.17E-03	EPA6 calc c	1.91E-02
Ethyl ether	60-29-7	1.18E+01	EPA6 calc c	NA	3.86E-04	EPA6 calc c	3.86E-04
Ethyl methacrylate	97-63-2	4.67E+00	EPA6 e	NA	2.00E-02	EPA6 e	2.00E-02
Formaldehyde	50-00-0	2.46E+01	EPA6 e	NA	4.65E-04	EPA6 e	4.65E-04
Formamide	75-12-7	2.89E+02	EPA6 calc c	NA	NA	NA	1.18E-02
Formic acid	64-18-6	7.92E+01	EPA6 e	NA	6.02E-03	EPA6 e	6.02E-03
Formic acid, methyl ester	107-31-3	5.50E+01	EPA6 calc c	NA	NA	NA	NA
Glycidylaldehyde	765-34-4	1.02E+02	EPA6 calc c	NA	1.83E-02	EPA6 calc c	1.18E-02
Methyl acetate	79-20-9	3.05E+01	EPA6 calc c	NA	1.00E-03	EPA6 calc c	1.00E-03
Methyl alcohol	67-56-1	9.96E+01	EPA6 e	NA	6.82E-05	EPA6 e	6.82E-05
Methyl isocyanate	624-83-9	NA	NA	NA	NA	NA	NA
Methyl methacrylate	80-62-6	1.35E+01	EPA6 calc c	NA	1.22E-03	EPA6 calc c	1.22E-03
Methyl tert-butyl ether	1634-04-4	1.11E+01	EPA6 calc c	NA	9.55E-04	EPA6 calc c	9.55E-04
Methylacetylene	74-99-7	1.11E+01	EPA6 calc c	NA	NA	NA	NA
Methylcyclohexane	108-87-2	1.65E-01	EPA6 calc c	NA	2.99E-03	EPA6 calc c	2.99E-03
N,N-Dimethylacetamide	127-19-5	NA	NA	NA	NA	NA	NA
n-Butyl alcohol	71-36-3	1.20E+01	EPA6 calc c	NA	5.70E-02	EPA6 calc c	5.70E-02
n-Heptane	142-82-5	7.84E-02	EPA6 calc c	NA	2.26E-03	EPA6 calc c	2.26E-03
n-Hexane	110-54-3	1.63E-01	EPA6 calc c	NA	7.35E-04	EPA6 calc c	7.35E-04
Nitromethane	75-52-5	6.17E+01	EPA6 calc c	NA	NA	NA	NA
n-Nonane	111-84-2	2.10E-02	EPA6 calc c	NA	1.18E-02	EPA6 calc c	1.18E-02
n-Octane	111-65-9	1.89E-01	EPA6 calc c	NA	3.15E-04	EPA6 calc c	3.15E-04
n-Pentane	109-66-0	5.40E-01	EPA6 calc c	NA	1.17E-04	EPA6 calc c	1.17E-04
n-Propionaldehyde	123-38-6	1.77E+01	EPA6 calc c	NA	3.21E-03	EPA6 calc c	3.21E-03
n-Propyl alcohol	71-23-8	NA	NA	NA	NA	NA	NA
n-Valeraldehyde	110-62-3	NA	NA	NA	NA	NA	NA
Oxirane	75-21-8	5.77E+01	EPA6 e	NA	1.60E-04	EPA6 e	1.60E-04

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	Br_{forage} ((ug/g DW plant) /(ug/g soil))	Source	Br_{grain} ((ug/g DW plant) /(ug/g soil))	Bv_{ag} ((ug/g DW plant) /(ug/g air))	Source	Bv_{forage} ((ug/g DW plant) /(ug/g air))
p-Cymene	99-87-6	1.65E-01	EPA6 calc c	NA	1.65E-01	EPA6 calc c	2.83E-01
Phosgene	75-44-5	NA	NA	NA	NA	NA	NA
Propargyl alcohol	107-19-7	NA	NA	NA	NA	NA	NA
Propionic acid	79-09-4	2.50E+01	EPA6 calc c	NA	2.83E-01	EPA6 calc c	2.83E-01
Propionitrile	107-12-0	3.13E+01	EPA6 calc c	NA	2.23E-03	EPA6 calc c	2.23E-03
Propylene glycol monomethyl ether	107-98-2	4.92E+01	EPA6 calc c	NA	3.47E-02	EPA6 calc c	3.47E-02
p-tert-Butyltoluene	98-51-1	NA	NA	NA	NA	NA	NA
Triethylamine	121-44-8	NA	NA	NA	NA	NA	NA
Trimethylamine	75-50-3	3.13E+01	EPA6 calc c	NA	7.95E-04	EPA6 calc c	1.26E-03
Vinyl acetate	108-05-4	1.53E+01	EPA6 e	NA	5.65E-04	EPA6 e	5.65E-04
Non-aromatic Halogenated Hydrocarbons							
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	NA	NA	NA	NA	NA	NA
1,1,1,2-Tetrachloroethane	630-20-6	1.17E+00	EPA6 e	NA	1.45E-02	EPA6 e	1.45E-02
1,1,1-Trichloroethane	71-55-6	1.54E+00	EPA6 e	NA	1.14E-03	EPA6 e	1.14E-03
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	2.70E-01	EPA6 calc c	NA	5.38E-03	EPA6 calc c	5.38E-03
1,1,2,2-Tetrachloroethane	79-34-5	8.02E-02	EPA6 e	NA	1.33E+01	EPA6 e	1.33E+01
1,1,2,2-Tetrachloroethene	127-18-4	1.31E+00	EPA6 e	NA	1.66E-03	EPA6 e	1.66E-03
1,1,2-Trichloro-1,2,2-trifluoroethane	76-13-1	5.77E-01	EPA6 calc c	NA	2.46E-04	EPA6 calc c	2.46E-04
1,1,2-Trichloroethane	79-00-5	2.38E+00	EPA6 e	NA	9.53E-03	EPA6 e	9.53E-03
1,1,2-Trichloroethylene	79-01-6	1.52E+00	EPA6 e	NA	2.07E-03	EPA6 e	2.07E-03
1,1-Dichloroethane	75-34-3	3.56E+00	EPA6 e	NA	7.88E-04	EPA6 e	7.88E-04
1,1-Dichloroethene	75-35-4	2.30E+00	EPA6 e	NA	3.98E-04	EPA6 e	3.98E-04
1,2,3-Trichloropropane	96-18-4	1.94E+00	EPA6 e	NA	3.66E-02	EPA6 e	3.66E-02
1,2-dibromo-3-chloropropane	96-12-8	1.72E+00	EPA6 e	NA	8.81E-02	EPA6 e	8.81E-02
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	NA	NA	NA	NA	NA	NA
1,2-Dichloroethane	107-06-2	5.53E+00	EPA6 e	NA	1.58E-03	EPA6 e	1.58E-03
1,2-Dichloroethylene	540-59-0	2.04E+01	EPA6 calc c	NA	2.41E-05	EPA6 calc c	2.41E-05
1,2-Dichloropropane	78-87-5	1.94E+00	EPA6 e	NA	4.96E-03	EPA6 e	4.96E-03
1,3-Dichloropropene	542-75-6	3.78E+00	EPA6 e	NA	1.38E-03	EPA6 e	1.38E-03

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	Br_{forage} ((ug/g DW plant) /(ug/g soil))	Source	Br_{grain} ((ug/g DW plant) /(ug/g soil))	Bv_{ag} ((ug/g DW plant) /(ug/g air))	Source	Bv_{forage} ((ug/g DW plant) /(ug/g air))
1,4-Dichloro-2-butene	764-41-0	1.21E+01	EPA6 calc c	NA	1.84E-03	EPA6 calc c	1.84E-03
1-Chloroethene	75-01-4	8.42E+00	EPA6 e	NA	2.95E-06	EPA6 e	2.95E-06
2,2-Dichloropropionic acid	75-99-0	1.38E+01	EPA6 calc c	NA	5.87E+00	EPA6 calc c	5.87E+00
2-Chloropropane	75-29-6	3.09E+00	EPA6 calc c	NA	3.62E-04	EPA6 calc c	3.62E-04
3-Chloropropene (allyl chloride)	107-05-1	1.09E+01	EPA6 calc c	NA	5.35E-05	EPA6 calc c	5.35E-05
Bromochloromethane	74-97-5	5.93E+00	EPA6 calc c	NA	6.87E-08	EPA6 calc c	1.68E-09
Bromodichloromethane	75-27-4	2.61E+00	EPA6 e	NA	2.53E-03	EPA6 e	2.53E-03
Bromoethene	593-60-2	9.37E+00	EPA6 calc c	NA	1.14E-04	EPA6 calc c	1.14E-04
Bromoform	75-25-2	1.70E+00	EPA6 e	NA	2.89E-02	EPA6 e	2.89E-02
Bromomethane	74-83-9	8.79E+00	EPA6 e	NA	6.07E-05	EPA6 e	6.07E-05
Carbon tetrachloride	56-23-5	1.04E+00	EPA6 e	NA	1.52E-03	EPA6 e	1.52E-03
Chlordibromomethane	124-48-1	2.14E+00	EPA6 e	NA	9.59E-03	EPA6 e	9.59E-03
Chlorodifluoromethane	75-45-6	9.21E+00	EPA6 e	NA	4.69E-06	EPA6 e	4.69E-06
Chloroethane	75-00-3	6.25E-01	EPA6 e	NA	6.05E-05	EPA6 e	6.05E-05
Chloroform	67-66-3	2.89E+00	EPA6 e	NA	1.65E-03	EPA6 e	1.65E-03
Chloromethane	74-87-3	1.16E+01	EPA6 e	NA	1.13E-05	EPA6 e	1.13E-05
Chloropentafluoroethane	76-15-3	NA	NA	NA	NA	NA	NA
cis-1,2-Dichloroethene	156-59-2	2.77E+00	EPA6 e	NA	1.60E-03	EPA6 e	1.60E-03
cis-1,3-Dichloropropene	10061-01-5	NA	NA	NA	NA	NA	NA
Cyanogen bromide	506-68-3	NA	NA	NA	NA	NA	NA
Cyanogen chloride	506-77-4	NA	NA	NA	NA	NA	NA
Dichlorodifluoromethane	75-71-8	2.19E+00	EPA6 e	NA	4.33E-06	EPA6 e	4.33E-06
Dichlorofluoromethane	75-43-4	NA	NA	NA	NA	NA	NA
Dichloromethane	75-09-2	7.29E+00	EPA6 e	NA	5.11E-04	EPA6 e	5.11E-04
Difluorodibromomethane	75-61-6	NA	NA	NA	NA	NA	NA
Hexafluoroacetone	684-16-2	NA	NA	NA	NA	NA	NA
Iodomethane	74-88-4	4.08E+00	EPA6 calc c	NA	6.44E-04	EPA6 calc c	6.44E-04
Methylene bromide	74-95-3	4.48E+00	EPA6 e	NA	1.13E-04	EPA6 e	1.13E-04
Pentachloroethane	76-01-7	6.68E-01	EPA6 calc c	NA	4.04E-02	EPA6 calc c	4.04E-02

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	Br_{forage} ((ug/g DW plant) /(ug/g soil))	Source	Br_{grain} ((ug/g DW plant) /(ug/g soil))	Bv_{ag} ((ug/g DW plant) /(ug/g air))	Source	Bv_{forage} ((ug/g DW plant) /(ug/g air))
trans-1,2-Dichloroethene	156-60-5	2.77E+00	EPA6 e	NA	9.70E-04	EPA6 e	9.70E-04
trans-1,3-Dichloropropene	10061-02-6	2.50E+00	EPA6 calc c	NA	NA	NA	NA
Trichloroacetic acid	76-03-9	6.60E+00	EPA6 calc c	NA	8.14E+00	EPA6 calc c	8.14E+00
Trichlorofluoroethane	27154-33-2	NA	NA	NA	NA	NA	NA
Trichlorofluoromethane	75-69-4	1.33E+00	EPA6 e	NA	2.02E-04	EPA6 e	2.02E-04
Trifluorobromomethane	75-63-8	3.26E+00	EPA6 calc c	NA	NA	NA	NA
Dioxin and Furan Compounds							
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	35822-46-9	7.05E-04	EPA6 e	NA	3.50E+05	EPA6 e	3.50E+05
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	1.02E-03	EPA6 e	NA	4.40E+05	EPA6 e	4.40E+05
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	1.02E-03	EPA6 e	NA	4.40E+05	EPA6 e	4.40E+05
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	39227-28-6	1.22E-03	EPA6 e	NA	4.50E+05	EPA6 e	4.50E+05
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	2.50E-03	EPA6 e	NA	1.50E+05	EPA6 e	1.50E+05
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	57653-85-7	2.50E-03	EPA6 e	NA	4.50E+05	EPA6 e	4.50E+05
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	2.50E-03	EPA6 e	NA	1.50E+05	EPA6 e	1.50E+05
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	19408-74-3	2.50E-03	EPA6 e	NA	4.50E+05	EPA6 e	4.50E+05
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	2.50E-03	EPA6 e	NA	1.50E+05	EPA6 e	1.50E+05
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	40321-76-4	5.62E-03	EPA6 e	NA	1.20E+05	EPA6 e	1.20E+05
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	4.61E-03	EPA6 e	NA	4.30E+04	EPA6 e	4.60E+04
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	2.50E-03	EPA6 e	NA	1.50E+05	EPA6 e	1.50E+05
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	3.87E-03	EPA6 e	NA	4.60E+04	EPA6 e	4.60E+04
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1746-01-6	5.62E-03	EPA6 e	NA	6.10E+04	EPA6 e	6.10E+04
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	6.51E-03	EPA6 e	NA	8.10E+04	EPA6 e	8.10E+04
Dibenzofuran	132-64-9	1.22E-01	EPA6 calc c	NA	1.76E+02	EPA6 calc c	1.76E+02
Octachlorodibenzo(p)dioxin	3268-87-9	1.59E-03	EPA6 e	NA	8.60E+06	EPA6 e	8.60E+06
Octachlorodibenzofuran	39001-02-0	3.26E-04	EPA6 e	NA	1.30E+06	EPA6 e	1.30E+06
PCBs							
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	3.13E-03	EPA6 calc c	NA	1.02E+04	EPA6 calc c	1.02E+04
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	2.97E-03	EPA6 calc c	NA	7.13E+03	EPA6 calc c	7.13E+03
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	NA	NA	NA	NA	NA	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	Br_{forage} ((ug/g DW plant) /(ug/g soil))	Source	Br_{grain} ((ug/g DW plant) /(ug/g soil))	Bv_{ag} ((ug/g DW plant) /(ug/g air))	Source	Bv_{forage} ((ug/g DW plant) /(ug/g air))
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	NA	NA	NA	NA	NA	NA
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	NA	NA	NA	NA	NA	NA
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	NA	NA	NA	NA	NA	NA
2',3,4,4',5-Pentachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	2.97E-03	EPA6 calc c	NA	NA	NA	NA
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	2.02E-03	EPA6 calc c	NA	7.34E+04	EPA6 calc c	7.34E+04
3,3',4,4',5-Pentachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	NA	NA	NA	NA	NA	NA
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	NA	NA	NA	NA	NA	NA
Polychlorinated biphenyls (PCBs)**	1336-36-3	2.31E-03	EPA6 calc c	NA	3.94E+03	EPA6 calc c	3.94E+03
Phthalates							
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	3.80E-02	EPA6 e	NA	1.77E+06	EPA6 e	1.77E+06
Butylbenzyl phthalate	85-68-7	1.09E-01	EPA6 e	NA	1.46E+03	EPA6 e	1.46E+03
Dibutyl phthalate	84-74-2	7.24E-02	EPA6 e	NA	4.16E+03	EPA6 e	4.16E+03
Diethyl phthalate	84-66-2	1.06E-01	EPA6 e	NA	5.42E+03	EPA6 e	5.42E+03
Dimethyl phthalate	131-11-3	4.40E+00	EPA6 e	NA	3.05E+01	EPA6 e	3.05E+01
n-Dioctyl phthalate	117-84-0	1.57E-04	EPA6 e	NA	6.28E+08	EPA6 e	6.28E+08
Light Polycyclic Aromatic Hydrocarbons							
2-Chloronaphthalene	91-58-7	1.72E-01	EPA6 e	NA	8.46E+00	EPA6 e	8.46E+00
2-Methyl naphthalene	91-57-6	2.27E-01	EPA6 calc c	NA	1.80E+00	EPA6 calc c	1.80E+00
5-Nitroacenaphthene	602-87-9	NA	NA	NA	NA	NA	NA
Acenaphthene	83-32-9	1.98E-01	EPA6 e	NA	4.66E+00	EPA6 e	4.66E+00
Acenaphthylene	208-96-8	1.72E-01	EPA6 calc c	NA	1.03E+01	EPA6 calc c	1.03E+01
Anthracene	120-12-7	1.01E-01	EPA6 e	NA	2.90E+01	EPA6 e	2.90E+01
Fluorene	86-73-7	1.49E-01	EPA6 calc c	NA	1.88E+01	EPA6 calc c	1.88E+01
Indene	95-13-6	NA	NA	NA	NA	NA	NA
Napthalene	91-20-3	4.35E-01	EPA6 e	NA	4.52E-01	EPA6 e	4.52E-01

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	Br_{forage} ((ug/g DW plant) /(ug/g soil))	Source	Br_{grain} ((ug/g DW plant) /(ug/g soil))	Bv_{ag} ((ug/g DW plant) /(ug/g air))	Source	Bv_{forage} ((ug/g DW plant) /(ug/g air))
Phenanthrene	85-01-8	9.08E-02	EPA6 e	NA	2.08E-02	EPA6 e	2.08E-02
Pyrene	129-00-0	4.98E-02	EPA6 e	NA	1.04E+06	EPA6 e	1.04E+06
Heavy Polycyclic Aromatic Hydrocarbons							
3-Methylcholanthrene	56-49-5	3.01E-03	EPA6 calc c	NA	1.68E+07	EPA6 calc c	1.68E+07
5-Methylchrysene	3697-24-3	NA	NA	NA	NA	NA	NA
Benzo(a)anthracene	56-55-3	2.02E-02	EPA6 e	NA	1.72E+04	EPA6 e	1.72E+04
Benzo(a)pyrene	50-32-8	1.11E-02	EPA6 e	NA	2.25E+05	EPA6 e	2.25E+05
Benzo(b)fluoranthene	205-99-2	1.01E-02	EPA6 e	NA	3.65E+04	EPA6 e	3.65E+04
Benzo(e)pyrene	192-97-2	2.05E-03	EPA6 calc c	NA	9.22E+06	EPA6 calc c	9.22E+06
Benzo(g,h,i)perylene	191-24-2	3.05E-03	EPA6 calc c	NA	2.91E+05	EPA6 calc c	2.91E+05
Benzo(j)fluoranthene	205-82-3	7.34E-03	EPA6 calc c	NA	NA	NA	NA
Benzo(k)fluoranthene	207-08-9	1.01E-02	EPA6 e	NA	5.40E+05	EPA6 e	5.40E+05
Benzo[a,i]pyrene	191-30-0	NA	NA	NA	NA	NA	NA
Chrysene	218-01-9	1.87E-02	EPA6 e	NA	5.97E+04	EPA6 e	5.97E+04
Dibenz(a,h)anthracene	53-70-3	6.36E-03	EPA6 e	NA	4.68E+07	EPA6 e	4.68E+07
Dibenz[a,h]acridine	226-36-8	NA	NA	NA	NA	NA	NA
Dibenz[a,j]acridine	224-42-0	NA	NA	NA	NA	NA	NA
Dibenzo(a,e)fluoranthene	5385-75-1	NA	NA	NA	NA	NA	NA
Dibenzo(a,h)fluoranthene	no cas #	NA	NA	NA	NA	NA	NA
Dibenzo[a,e]pyrene	192-65-4	NA	NA	NA	NA	NA	NA
Dibenzo[a,h]pyrene	189-64-0	NA	NA	NA	NA	NA	NA
Dibenzo[a,i]pyrene	189-55-9	2.37E-03	EPA6 calc c	NA	NA	NA	NA
Fluoranthene	206-44-0	4.46E-02	EPA6 e	NA	1.56E+03	EPA6 e	1.56E+03
Hexachloronaphthalene	1335-87-1	1.59E-03	EPA6 calc c	NA	NA	NA	NA
Indeno(1,2,3-cd)pyrene	193-39-5	3.90E-03	EPA6 e	NA	2.67E+08	EPA6 e	2.67E+08
Octachloronaphthalene	2234-13-1	7.54E-03	EPA6 calc c	NA	NA	NA	NA
Pentachloronaphthalene	1321-64-8	NA	NA	NA	NA	NA	NA
Tetrachloronaphthalene	1335-88-2	NA	NA	NA	NA	NA	NA
Trichloronaphthalene	1321-65-9	NA	NA	NA	NA	NA	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	Br_{forage} ((ug/g DW plant) /(ug/g soil))	Source	Br_{grain} ((ug/g DW plant) /(ug/g soil))	Bv_{ag} ((ug/g DW plant) /(ug/g air))	Source	Bv_{forage} ((ug/g DW plant) /(ug/g air))
Light Substituted Benzene Compounds							
1,2,3-Trichlorobenzene	87-61-6	1.78E-01	EPA6 e	NA	4.01E-01	EPA6 e	4.01E-01
1,2,4-Trichlorobenzene	120-82-1	1.92E-01	EPA6 e	NA	3.78E-01	EPA6 e	3.78E-01
1,2,4-Trimethlyl benzene	25551-13-7	4.20E-01	EPA6 calc c	NA	NA	NA	NA
1,2,4-Trimethyl benzene	95-63-6	3.01E-01	EPA6 calc c	NA	7.52E-02	EPA6 calc c	7.52E-02
1,2-Dichlorobenzene	95-50-1	3.95E-01	EPA6 e	NA	1.24E-01	EPA6 e	1.24E-01
1,3,5-Trimethyl benzene	108-67-8	4.09E-01	EPA6 e	NA	3.14E-02	EPA6 e	3.14E-02
1,3-Dichlorobenzene	541-73-1	3.53E-01	EPA6 e	NA	8.02E-02	EPA6 e	8.02E-02
1,3-Dinitrobenzene	99-65-0	5.32E+00	EPA6 e	NA	1.74E+01	EPA6 e	1.74E+01
1,4-Dichlorobenzene	106-46-7	4.13E-01	EPA6 e	NA	8.60E-02	EPA6 e	8.60E-02
1,4-Dinitrobenzene	100-25-4	5.26E+00	EPA6 calc c	NA	NA	NA	NA
2,4,5-Trichorophenol	95-95-4	2.24E-01	EPA6 e	NA	1.31E+02	EPA6 e	1.31E+02
2,4,6-Trichlorophenol	88-06-2	2.77E-01	EPA6 e	NA	1.23E+02	EPA6 e	1.23E+02
2,4-Dichlorophenol	120-83-2	6.82E-01	EPA6 e	NA	4.01E+02	EPA6 e	4.01E+02
2,4-Dimethylphenol	105-67-9	1.68E+00	EPA6 e	NA	4.27E+03	EPA6 e	4.27E+03
2,4-Dinitrophenol	51-28-5	5.13E+00	EPA6 e	NA	4.80E+02	EPA6 e	4.80E+02
2,4-Dinitrotoluene	121-14-2	2.72E+00	EPA6 e	NA	5.10E+01	EPA6 e	5.10E+01
2,6-Dinitrotoluene	606-20-2	3.15E+00	EPA6 e	NA	4.41E+01	EPA6 e	4.41E+01
2-Chlorotoluene	95-49-8	3.50E-01	EPA6 calc c	NA	5.15E-01	EPA6 calc c	5.15E-01
2-Chorophenol	95-57-8	2.18E+00	EPA6 e	NA	6.76E-01	EPA6 e	6.76E-01
2-Nitrophenol	88-75-5	3.57E+00	EPA6 e	NA	3.08E-01	EPA6 e	3.08E-01
4,6-Dinitro-o-cresol	534-52-1	8.72E-01	EPA6 calc c	NA	4.33E+01	EPA6 calc c	4.33E+01
4-Chlorotoluene	106-43-4	4.61E-01	EPA6 calc c	NA	NA	NA	NA
4-Nitrophenol	100-02-7	3.05E+00	EPA6 e	NA	8.26E+02	EPA6 e	8.26E+02
alpha-Methylstyrene	98-83-9	3.86E-01	EPA6 calc c	NA	4.63E-02	EPA6 calc c	4.63E-02
Aniline	62-53-3	1.05E+01	EPA6 e	NA	2.72E-01	EPA6 e	2.72E-01
Benzotrichloride	98-07-7	7.95E-01	EPA6 calc c	NA	7.36E-02	EPA6 calc c	7.36E-02
Benzyl Chloride	100-44-7	2.39E+01	EPA6 e	NA	3.28E-04	EPA6 e	3.28E-04
Bromobenzene	108-86-1	7.24E-01	EPA6 calc c	NA	4.10E-02	EPA6 calc c	4.10E-02

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	Br_{forage} ((ug/g DW plant) /(ug/g soil))	Source	Br_{grain} ((ug/g DW plant) /(ug/g soil))	Bv_{ag} ((ug/g DW plant) /(ug/g air))	Source	Bv_{forage} ((ug/g DW plant) /(ug/g air))
Chlorobenzene	108-90-7	9.45E-01	EPA6 e	NA	1.19E-02	EPA6 e	1.19E-02
Cumene	98-82-8	3.16E-01	EPA6 e	NA	3.06E-02	EPA6 e	3.06E-02
m-Cresol	108-39-4	2.86E+00	EPA6 e	NA	7.64E+00	EPA6 e	7.64E+00
n-Butyl benzene	104-51-8	1.30E-01	EPA6 calc c	NA	1.57E-01	EPA6 calc c	1.57E-01
Nitrobenzene	98-95-3	3.38E+00	EPA6 e	NA	2.43E-01	EPA6 e	2.43E-01
n-Propyl benzene	103-65-1	2.85E-01	EPA6 calc c	NA	4.34E-02	EPA6 calc c	4.34E-02
o-Cresol	95-48-7	2.63E+00	EPA6 e	NA	4.89E+00	EPA6 e	4.89E+00
o-Dinitrobenzene	528-29-0	4.60E+00	EPA6 calc c	NA	NA	NA	NA
o-Nitroaniline	88-74-4	3.30E+00	EPA6 e	NA	4.47E+00	EPA6 e	4.47E+00
o-Toluidine	95-53-4	6.51E+00	EPA6 e	NA	6.16E-01	EPA6 e	6.16E-01
p-Chloroaniline	106-47-8	3.22E+00	EPA6 e	NA	4.66E+00	EPA6 e	4.66E+00
p-Cresol	106-44-5	2.93E+00	EPA6 e	NA	8.13E+00	EPA6 e	8.13E+00
Phenol	108-95-2	5.42E+00	EPA6 e	NA	3.52E+00	EPA6 e	3.52E+00
p-Nitrochlorobenzene	100-00-5	1.61E+00	EPA6 calc c	NA	NA	NA	NA
p-Toluidine	106-49-0	6.01E+00	EPA6 calc c	NA	2.67E-01	EPA6 calc c	2.67E-01
sec-Butyl benzene	135-98-8	NA	NA	NA	NA	NA	NA
tert-Butyl benzene	98-06-6	1.63E-01	EPA6 calc c	NA	1.12E-01	EPA6 calc c	1.12E-01
Toluene-2,6-diamine	823-40-5	5.59E+00	EPA6 calc c	NA	1.60E+05	EPA6 calc c	1.60E+05
Other Light Semivolatile Compounds							
1,1 Dimethyl hydrazine	57-14-7	1.75E+01	EPA6 calc c	NA	1.47E-01	EPA6 calc c	1.47E-01
1,1-Biphenyl	92-52-4	2.16E-01	EPA6 calc c	NA	2.88E+00	EPA6 calc c	2.88E+00
1,2 Diphenylhydrazine	122-66-7	7.74E-01	EPA6 e	NA	5.89E+02	EPA6 e	5.89E+02
1,2-Dimethylhydrazine	540-73-8	2.39E+02	EPA6 calc c	NA	NA	NA	NA
1,3-Propane sultone	1120-71-4	7.77E+01	EPA6 calc c	NA	NA	NA	NA
2,4-Toluene diisocyanate	584-84-9	NA	NA	NA	NA	NA	NA
2-Chloroacetophenone	532-27-4	1.23E+00	EPA6 calc c	NA	3.73E+01	EPA6 calc c	3.73E+01
2-Propenoic acid	79-10-7	2.19E+01	EPA6 calc c	NA	3.87E-01	EPA6 calc c	2.83E-01
4,4-Methylenedianiline	101-77-9	4.31E-01	EPA6 calc c	NA	7.93E+03	EPA6 calc c	7.93E+03
Acetophenone	98-86-2	4.37E+00	EPA6 e	NA	3.04E-01	EPA6 e	3.04E-01

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	Br_{forage} ((ug/g DW plant) /(ug/g soil))	Source	Br_{grain} ((ug/g DW plant) /(ug/g soil))	Bv_{ug} ((ug/g DW plant) /(ug/g air))	Source	Bv_{forage} ((ug/g DW plant) /(ug/g air))
Benzoic acid	65-85-0	3.17E+00	NA	NA	1.69E+01	NA	1.69E+01
bis(2-Chloroethoxy)methane	111-91-1	NA	NA	NA	NA	NA	NA
bis(2-Chloroethyl) ether	111-44-4	6.85E+00	EPA6 e	NA	6.37E-02	EPA6 e	6.37E-02
Chlorocyclopentadiene	41851-50-7	1.53E+00	EPA6 calc c	NA	9.69E-04	EPA6 calc c	9.69E-04
Cyclohexanol	108-93-0	7.53E+00	EPA6 calc c	NA	1.99E-01	EPA6 calc c	1.99E-01
Dichloroisopropyl ether	108-60-1	1.25E+00	EPA6 calc c	NA	2.78E-01	EPA6 calc c	2.78E-01
Dichloromethyl ether	542-88-1	6.42E+01	EPA6 calc c	NA	2.45E-04	EPA6 calc c	2.45E-04
Dichloropentadiene	no cas #	NA	NA	NA	NA	NA	NA
Dimethyl sulfate	77-78-1	2.54E+01	EPA6 calc c	NA	3.05E-02	EPA6 calc c	3.05E-02
Dimethylaniline	121-69-7	1.25E+02	EPA6 calc c	NA	4.67E-04	EPA6 calc c	4.67E-04
di-n-propylnitrosamine	621-64-7	6.17E+00	EPA6 e	NA	3.04E+01	EPA6 e	3.04E+01
Diphenyl ether	101-84-8	1.43E-01	EPA6 calc c	NA	6.87E+00	EPA6 calc c	6.87E+00
Epichlorohydrin	106-89-8	2.77E+01	EPA6 e	NA	3.35E-03	EPA6 e	3.35E-03
Ethyl carbomate (urethane)	51-79-6	4.73E+01	EPA6 calc c	NA	NA	NA	NA
Ethyl methane sulfanate	62-50-0	3.63E+01	EPA6 e	NA	7.11E-01	EPA6 e	7.11E-01
Ethylene dibromide	106-93-4	3.77E+00	EPA6 e	NA	9.13E-03	EPA6 e	9.13E-03
Ethylene glycol	107-21-1	1.31E+02	EPA6 calc c	NA	3.31E+00	EPA6 calc c	3.31E+00
Ethylene glycol monobutyl ether	111-76-2	4.89E+00	EPA6 calc c	NA	4.82E+00	EPA6 calc c	4.82E+00
Ethylene glycol monoethyl ether acetate	111-15-9	1.70E+01	EPA6 calc c	NA	1.44E-01	EPA6 calc c	1.44E-01
Ethylene thiourea	96-45-7	9.10E+01	EPA6 calc c	NA	7.65E+01	EPA6 calc c	7.65E+01
Furfural	98-01-1	1.08E+01	EPA6 calc c	NA	1.59E-01	EPA6 calc c	1.59E-01
Maleic hydrazide	123-33-1	1.04E+02	EPA6 calc c	NA	1.39E-06	EPA6 calc c	1.39E-06
Malononitrile	109-77-3	3.67E+01	EPA6 calc c	NA	NA	NA	NA
Methyl styrene (mixed isomers)	25013-15-4	4.48E-01	EPA6 calc c	NA	NA	NA	NA
Methylhydrazine	60-34-4	1.57E+02	EPA6 calc c	NA	NA	NA	NA
N,N-Diphenylamine	122-39-4	3.67E-01	EPA6 calc c	NA	8.64E+02	EPA6 calc c	8.64E+02
Nitric acid, propyl ester	627-13-4	NA	NA	NA	NA	NA	NA
N-Nitrosodi-n-butylamine	924-16-3	1.57E+00	EPA6 e	NA	3.77E-01	EPA6 e	3.77E-01
N-Nitrosomorpholine	59-89-2	1.05E+01	EPA6 calc c	NA	NA	NA	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	Br_{forage} ((ug/g DW plant) /(ug/g soil))	Source	Br_{grain} ((ug/g DW plant) /(ug/g soil))	Bv_{ag} ((ug/g DW plant) /(ug/g air))	Source	Bv_{forage} ((ug/g DW plant) /(ug/g air))
N-Nitroso-N,N-dimethylamine	62-75-9	7.24E+01	EPA6 calc c	NA	2.27E-02	EPA6 calc c	2.27E-02
o-Anisidine	90-04-0	8.05E+00	EPA6 calc c	NA	NA	NA	NA
Oxalic acid	144-62-7	NA	NA	NA	NA	NA	NA
Phthalic Anhydride	85-44-9	8.63E+01	EPA6 e	NA	2.03E+00	EPA6 e	2.03E+00
p-Phthalic acid	100-21-0	1.30E+01	EPA6 calc c	NA	4.05E+01	EPA6 calc c	4.05E+01
Pyridine	110-86-1	1.59E+01	EPA6 e	NA	4.22E-05	EPA6 e	4.22E-05
Quinoline	91-22-5	2.60E+00	EPA6 calc c	NA	3.01E+01	EPA6 calc c	3.01E+01
Quinone	106-51-4	2.97E+01	EPA6 calc c	NA	NA	NA	NA
Safrole	94-59-7	1.12E+00	EPA6 e	NA	3.20E+00	EPA6 e	3.20E+00
Tetrahydrofuran	109-99-9	2.14E+01	EPA6 e	NA	NA	EPA6 e	NA
Other Heavy Semivolatile Compounds							
1,2,4,5-Tetrachlorobenzene	95-94-3	8.06E-02	EPA6 e	NA	4.14E+00	EPA6 e	4.14E+00
1,3,5-Trinitrobenzene	99-35-4	8.05E+00	EPA6 e	NA	1.17E+01	EPA6 e	1.17E+01
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	1.51E-01	EPA6 calc c	NA	NA	NA	NA
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	1.41E+03	EPA6 calc c	NA	NA	NA	NA
2-sec-Butyl-4,6-dinitrophenol	88-85-7	5.93E-01	EPA6 calc c	NA	2.71E+02	EPA6 calc c	2.71E+02
3,3-Dichlorobenzidine	91-94-1	3.32E-01	EPA6 e	NA	1.73E+04	EPA6 e	1.73E+04
3,3'-Dimethoxybenzidine	119-90-4	3.48E+00	EPA6 e	NA	1.41E+04	EPA6 e	1.41E+04
4-Bromophenylphenyl ether	101-55-3	4.99E-02	EPA6 calc c	NA	1.01E+02	EPA6 calc c	1.01E+02
Ammonium perfluorooctanoate	3825-26-1	NA	NA	NA	NA	NA	NA
Azobenzene	103-33-3	2.40E-01	EPA6 calc c	NA	1.32E+02	EPA6 calc c	1.32E+02
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	NA	NA	NA	NA	NA	NA
Captan	133-06-2	1.70E+00	EPA6 calc c	NA	3.00E+00	EPA6 calc c	3.00E+00
Chlorobenzilate	510-15-6	1.14E-01	EPA6 e	NA	3.57E+04	EPA6 e	3.57E+04
Dibutylphosphate	107-66-4	NA	NA	NA	NA	NA	NA
Dimethyl aminoazobenzene	60-11-7	8.72E-02	EPA6 calc c	NA	NA	NA	NA
Hexachlorobenzene	118-74-1	2.56E-02	EPA6 e	NA	7.57E+01	EPA6 e	7.57E+01
Hexachlorobutadiene	87-68-3	7.14E-02	EPA6 e	NA	2.55E-01	EPA6 e	2.55E-01
Hexachlorocyclopentadiene	77-47-4	5.65E-02	EPA6 e	NA	5.47E-01	EPA6 e	5.47E-01

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	Br_{forage} ((ug/g DW plant) /(ug/g soil))	Source	Br_{grain} ((ug/g DW plant) /(ug/g soil))	Bv_{ag} ((ug/g DW plant) /(ug/g air))	Source	Bv_{forage} ((ug/g DW plant) /(ug/g air))
Hexachloroethane	67-72-1	1.93E-01	EPA6 e	NA	2.72E-01	EPA6 e	2.72E-01
Hexachlorophene	70-30-4	1.70E-03	EPA6 e	NA	1.23E+10	EPA6 e	1.23E+10
Hexamethylene-1,5-diisocyanate	822-06-0	7.11E+00	EPA6 calc c	NA	4.78E+00	EPA6 calc c	4.78E+00
Mirex	2385-85-5	4.03E-03	EPA6 calc c	NA	2.32E+03	EPA6 calc c	2.32E+03
Nitrofen	1836-75-5	2.46E-02	EPA6 calc c	NA	NA	NA	NA
Pentachlorobenzene	608-93-5	4.44E-02	EPA6 e	NA	6.04E-01	EPA6 e	6.04E-01
Pentachloronitrobenzene	82-68-8	8.06E-02	EPA6 e	NA	1.71E-01	EPA6 e	1.71E-01
Pentachlorophenol*	87-86-5	4.48E-02	EPA6 e	NA	1.02E+03	EPA6 e	1.02E+03
Pentachlorophenol*	87-86-5	4.48E-02	EPA6 e	NA	1.02E+03	EPA6 e	1.02E+03
Pentachlorophenol*	87-86-5	4.48E-02	EPA6 e	NA	1.02E+03	EPA6 e	1.02E+03
Pentachlorophenol*	87-86-5	4.48E-02	EPA6 e	NA	1.02E+03	EPA6 e	1.02E+03
Pentachlorophenol*	87-86-5	4.48E-02	EPA6 e	NA	1.02E+03	EPA6 e	1.02E+03
Pentachlorophenol*	87-86-5	4.48E-02	EPA6 e	NA	1.02E+03	EPA6 e	1.02E+03
Pentachlorophenol*	87-86-5	4.48E-02	EPA6 e	NA	1.02E+03	EPA6 e	1.02E+03
Pentachlorophenol*	87-86-5	4.48E-02	EPA6 e	NA	1.02E+03	EPA6 e	1.02E+03
Pentachlorophenol*	87-86-5	4.48E-02	EPA6 e	NA	1.02E+03	EPA6 e	1.02E+03
Pentachlorophenol*	87-86-5	4.48E-02	EPA6 e	NA	1.02E+03	EPA6 e	1.02E+03
Pentachlorophenol*	87-86-5	4.48E-02	EPA6 e	NA	1.02E+03	EPA6 e	1.02E+03
Pentachlorophenol*	87-86-5	4.48E-02	EPA6 e	NA	1.02E+03	EPA6 e	1.02E+03
Picric acid	88-89-1	2.60E+00	EPA6 calc c	NA	2.15E+07	EPA6 calc c	2.15E+07
Pronamide	23950-58-5	3.62E-01	EPA6 e	NA	3.38E+01	EPA6 e	3.38E+01
Strychnine	57-24-9	2.97E+00	EPA6 e	NA	1.29E+07	EPA6 e	1.29E+07
Terphenyls	26140-60-3	NA	NA	NA	NA	NA	NA
Tributyl phosphate	126-73-8	1.89E-01	EPA6 calc c	NA	6.80E+03	EPA6 calc c	6.80E+03
Trifluralin	1582-09-8	3.17E-02	EPA6 calc c	NA	6.84E+02	EPA6 calc c	6.84E+02
Triphenylamine	603-34-9	NA	NA	NA	NA	NA	NA
Herbicides and Organopesticides							
2,4,5-T	93-76-5	4.43E-01	EPA6 calc c	NA	2.44E+04	EPA6 calc c	2.44E+04

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	Br_{forage} ((ug/g DW plant) /(ug/g soil))	Source	Br_{grain} ((ug/g DW plant) /(ug/g soil))	Bv_{ag} ((ug/g DW plant) /(ug/g air))	Source	Bv_{forage} ((ug/g DW plant) /(ug/g air))
2,4-D and esters	94-75-7	9.20E-01	EPA6 calc c	NA	4.03E+05	EPA6 calc c	4.03E+05
4,4-DDD	72-54-8	1.01E-02	EPA6 calc c	NA	2.67E+03	EPA6 calc c	2.67E+03
4,4-DDT	50-29-3	1.32E-02	EPA6 calc c	NA	1.06E+04	EPA6 calc c	1.06E+04
Aldrin	309-00-2	1.04E-02	EPA6 e	NA	1.58E+06	EPA6 e	1.58E+06
alpha-BHC	319-84-6	2.47E-01	EPA6 e	NA	9.17E+01	EPA6 e	9.17E+01
beta-BHC	319-85-7	2.36E-01	EPA6 e	NA	1.95E+03	EPA6 e	1.95E+03
Chlordane	57-74-9	1.43E-02	EPA6 e	NA	4.46E+03	EPA6 e	4.46E+03
DDE	72-55-9	9.37E-03	EPA6 e	NA	2.08E+03	EPA6 e	2.08E+03
Delta-BHC	319-86-8	1.57E-01	EPA6 calc c	NA	1.76E+03	EPA6 calc c	1.76E+03
Dieldrin	60-57-1	3.49E-02	EPA6 e	NA	6.50E+06	EPA6 e	6.50E+06
Endothall	145-73-3	1.23E+02	EPA6 calc c	NA	2.56E+01	EPA6 calc c	2.56E+01
Endrin	72-20-8	5.76E-02	EPA6 e	NA	7.62E+03	EPA6 e	7.62E+03
gamma-BHC(Lindane)	58-89-9	2.74E-01	EPA6 calc c	NA	3.28E+02	EPA6 calc c	3.28E+02
Heptachlor	76-44-8	4.89E-02	EPA6 e	NA	2.09E+03	EPA6 e	2.09E+03
Isodrin	465-73-6	NA	NA	NA	NA	NA	NA
Methoxychlor	72-43-5	9.38E-02	EPA6 e	NA	5.83E+02	EPA6 e	5.83E+02
Silvex (2,4,5-TP)	93-72-1	1.71E-01	EPA6 calc c	NA	1.96E-01	EPA6 calc c	1.96E-01
Toxaphene	8001-35-2	2.56E-02	EPA6 calc c	NA	8.94E+01	EPA6 calc c	8.94E+01

NA = Value not available
** = Varies with pH; values given for a pH of 7.0.
a = Illustrated Handbook of Physical - Chemical Properties and Environmental Fate f
b = RAIS database: http://risk.lsd.ornl.gov/rap_hp.shtml
c = Equations derived from EPA Region 6 Guidance Appendix A-3
d = Handbook of Environmental degradation Rates. 1991.
e = EPA Region 6 Hazardous Waste Combustion Guidance
f = "A Critical Review of Henry's Law Constants for Chemicals of Environmental In
g = Handbook of Environmental Fate and Exposure Data for Organic Chemicals
h = http://www.epa.gov/region09/waste/sfund/prg/.
i = http://esc.syrres.com/efdb/Chemfate.htm
j = Hansch and Leo 1985 in Syracuse 1996

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Plant Uptake Factors					
Constituent of Potential Concern	CAS Registry Number	Br_{forage} ((ug/g DW plant) /(ug/g soil))	Source	Br_{grain} ((ug/g DW plant) /(ug/g soil))	Bv_{ag} ((ug/g DW plant) /(ug/g air))	Source	Bv_{forage} ((ug/g DW plant) /(ug/g air))

k = Schwarzenbach et al. 1993

l = Tomlin 1994 in Syracuse 1996

m = http://www.speclab.com/compound/c959988.htm

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Bioconcentration / Bioaccumulation Factors					
Constituent of Potential Concern	CAS Registry Number	Source	Ba_{milk} (day/kg FW)	Source	Ba_{beef} (day/kg FW)	Source	Ba_{pork} (day/kg FW)	Source
Aromatic Halogenated Hydrocarbons								
2,3,4,6-Tetrachlorophenol	58-90-2	EPA6 calc c	2.09E-04	EPA6 calc c	6.61E-04	EPA6 calc c	7.93E-04	EPA6 calc c
4-Chloro-3-methylphenol	59-50-7	EPA6 calc c	1.00E-05	EPA6 calc c	3.16E-05	EPA6 calc c	3.79E-05	EPA6 calc c
Aromatic Nonhalogenated Hydrocarbons								
2-Nitrotoluene	88-72-2	EPA6 calc c	1.58E-06	EPA6 calc c	5.01E-06	EPA6 calc c	6.01E-06	EPA6 calc c
4-Nitrobiphenyl	92-93-3	NA	4.68E-05	EPA6 calc c	1.48E-04	EPA6 calc c	1.77E-04	EPA6 calc c
Benzaldehyde	100-52-7	EPA6 e	2.38E-07	EPA6 e	7.54E-07	EPA6 e	9.12E-07	EPA6 e
Benzene	71-43-2	EPA6 e	1.09E-06	EPA6 e	3.44E-06	EPA6 e	4.17E-06	EPA6 e
Benzyl alcohol	100-51-6	EPA6 e	1.00E-07	EPA6 e	3.16E-07	EPA6 e	3.83E-07	EPA6 e
Ethyl Benzene	100-41-4	EPA6 e	1.05E-05	EPA6 e	3.33E-05	EPA6 e	4.03E-05	EPA6 e
m-Xylene	108-38-3	EPA6 e	1.26E-05	EPA6 e	3.99E-05	EPA6 e	4.83E-05	EPA6 e
o-Xylene	95-47-6	EPA6 e	1.07E-05	EPA6 e	3.39E-05	EPA6 e	4.10E-05	EPA6 e
p-Xylene	106-42-3	EPA6 e	1.18E-05	EPA6 e	3.72E-05	EPA6 e	4.50E-05	EPA6 e
Styrene	100-42-5	EPA6 e	6.74E-06	EPA6 e	2.13E-05	EPA6 e	2.58E-05	EPA6 e
Toluene	108-88-3	EPA6 e	3.69E-06	EPA6 e	1.17E-05	EPA6 e	1.41E-05	EPA6 e
Non-aromatic Nonhalogenated Hydrocarbons								
1,2-Epoxybutane	106-88-7	EPA6 calc c	2.19E-07	EPA6 calc c	6.94E-07	EPA6 calc c	8.33E-07	EPA6 calc c
1,3-Butadiene	106-99-0	EPA6 calc c	7.76E-07	EPA6 calc c	2.45E-06	EPA6 calc c	2.95E-06	EPA6 calc c
1,4-Dioxane	123-91-1	EPA6 e	4.29E-09	EPA6 e	1.36E-08	EPA6 e	1.64E-08	EPA6 e
1-Methylpropyl alcohol	78-92-2	EPA6 calc c	3.24E-08	EPA6 calc c	1.02E-07	EPA6 calc c	1.23E-07	EPA6 calc c
1-Nitropropane	108-03-2	NA	5.89E-08	EPA6 calc c	1.86E-07	EPA6 calc c	2.23E-07	EPA6 calc c
2,2,4-Trimethlylpentane	540-84-1	EPA6 calc c	8.32E-04	EPA6 calc c	2.63E-03	EPA6 calc c	3.16E-03	EPA6 calc c
2-Butanone	78-93-3	EPA6 e	1.51E-08	EPA6 e	4.79E-08	EPA6 e	5.79E-08	EPA6 e
2-Butenaldehyde (2-Butenal)	4170-30-3	NA	NA	NA	NA	NA	NA	NA
2-Ethoxyethanol	110-80-5	EPA6 calc c	6.31E-09	EPA6 calc c	2.00E-08	EPA6 calc c	2.39E-08	EPA6 calc c
2-Heptanone	110-43-0	EPA6 calc c	7.59E-07	EPA6 calc c	2.40E-06	EPA6 calc c	2.88E-06	EPA6 calc c
2-Hexanone	591-78-6	EPA6 calc c	1.91E-07	EPA6 calc c	6.03E-07	EPA6 calc c	7.23E-07	EPA6 calc c
2-Methoxyethanol	109-86-4	EPA6 calc c	1.42E-08	EPA6 calc c	4.49E-08	EPA6 calc c	5.39E-08	EPA6 calc c
2-Methyl-2-propanol	75-65-0	EPA6 calc c	1.78E-08	EPA6 calc c	5.62E-08	EPA6 calc c	6.75E-08	EPA6 calc c

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Bioconcentration / Bioaccumulation Factors					
Constituent of Potential Concern	CAS Registry Number	Source	Ba_{milk} (day/kg FW)	Source	Ba_{beef} (day/kg FW)	Source	Ba_{pork} (day/kg FW)	Source
2-Methyl-2-propenenitrile	126-98-7	EPA6 e	2.76E-08	EPA6 e	8.72E-08	EPA6 e	1.06E-07	EPA6 e
2-Methylaziridine	75-55-8	NA	NA	NA	NA	NA	NA	NA
2-Methylpropyl alcohol	78-83-1	EPA6 calc c	4.57E-08	EPA6 calc c	1.45E-07	EPA6 calc c	1.73E-07	EPA6 calc c
2-Pentanone	107-87-9	EPA6 calc c	6.46E-08	EPA6 calc c	2.04E-07	EPA6 calc c	2.45E-07	EPA6 calc c
2-Propanone (Acetone)	67-64-1	EPA6 e	4.77E-09	EPA6 e	1.15E-08	EPA6 e	1.82E-08	EPA6 e
2-Propene-1-ol	107-18-6	EPA6 calc c	1.17E-08	EPA6 calc c	3.72E-08	EPA6 calc c	4.46E-08	EPA6 calc c
2-Propyl alcohol	67-63-0	EPA6 calc c	8.91E-09	EPA6 calc c	2.82E-08	EPA6 calc c	3.38E-08	EPA6 calc c
3-Heptanone	106-35-4	NA	NA	NA	NA	NA	NA	NA
3-Methyl-1-butanol	123-51-3	NA	NA	NA	NA	NA	NA	NA
3-Methyl-2-butanone	563-80-4	NA	NA	NA	NA	NA	NA	NA
3-Pentanone	96-22-0	EPA6 calc c	7.76E-08	EPA6 calc c	2.45E-07	EPA6 calc c	2.95E-07	EPA6 calc c
4-Heptanone	123-19-3	NA	NA	NA	NA	NA	NA	NA
4-Methyl-2-pentanone	108-10-1	EPA6 e	1.23E-07	EPA6 e	3.89E-07	EPA6 e	4.71E-07	EPA6 e
4-Methyl-3-penten-2-one	141-79-7	NA	NA	NA	NA	NA	NA	NA
5-Methyl-2-hexanone	110-12-3	NA	NA	NA	NA	NA	NA	NA
Acetaldehyde	75-07-0	EPA6 e	4.78E+09	EPA6 e	1.51E-08	EPA6 e	1.83E-08	EPA6 e
Acetamide	60-35-5	EPA6 calc c	4.37E-10	EPA6 calc c	1.38E-09	EPA6 calc c	1.66E-09	EPA6 calc c
Acetic acid	64-19-7	EPA6 calc c	5.37E-09	EPA6 calc c	1.70E-08	EPA6 calc c	2.04E-08	EPA6 calc c
Acetic acid ethyl ester	141-78-6	EPA6 calc c	4.27E-08	EPA6 calc c	1.35E-07	EPA6 calc c	1.62E-07	EPA6 calc c
Acetic acid n-butyl ester	123-86-4	EPA6 calc c	4.27E-07	EPA6 calc c	1.35E-06	EPA6 calc c	1.62E-06	EPA6 calc c
Acetonitrile	75-05-8	EPA6 e	3.63E-09	EPA6 e	1.15E-08	EPA6 e	1.39E-08	EPA6 e
Acrolein	107-02-8	EPA6 e	7.78E-09	EPA6 e	2.46E-08	EPA6 e	2.98E-08	EPA6 e
Acrylonitrile	107-13-1	EPA6 e	1.41E-08	EPA6 e	4.47E-08	EPA6 e	5.41E-08	EPA6 e
Bis(isopropyl)ether	108-20-3	NA	NA	NA	NA	NA	NA	NA
Butane	106-97-8	EPA6 calc c	6.17E-06	EPA6 calc c	1.95E-05	EPA6 calc c	2.34E-05	EPA6 calc c
Carbon disulfide	75-15-0	EPA6 e	7.94E-07	EPA6 e	2.51E-06	EPA6 e	3.04E-06	EPA6 e
Cyanogen	460-19-5	EPA6 calc c	5.10E-08	EPA6 calc c	1.61E-07	EPA6 calc c	1.94E-07	EPA6 calc c
Cyclohexane	110-82-7	EPA6 calc c	2.19E-05	EPA6 calc c	6.92E-05	EPA6 calc c	8.30E-05	EPA6 calc c
Cyclohexanone	108-94-1	EPA6 calc c	5.13E-08	EPA6 calc c	1.62E-07	EPA6 calc c	1.95E-07	EPA6 calc c

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Bioconcentration / Bioaccumulation Factors					
Constituent of Potential Concern	CAS Registry Number	Source	Ba_{milk} (day/kg FW)	Source	Ba_{beef} (day/kg FW)	Source	Ba_{pork} (day/kg FW)	Source
Cyclohexene	110-83-8	EPA6 calc c	5.75E-06	EPA6 calc c	1.82E-05	EPA6 calc c	2.18E-05	EPA6 calc c
Cyclopentane	287-92-3	EPA6 calc c	7.94E-06	EPA6 calc c	2.51E-05	EPA6 calc c	3.01E-05	EPA6 calc c
Ethyl alcohol	64-17-5	EPA6 calc c	3.89E-09	EPA6 calc c	1.23E-08	EPA6 calc c	1.48E-08	EPA6 calc c
Ethyl ether	60-29-7	EPA6 calc c	6.17E-08	EPA6 calc c	1.95E-07	EPA6 calc c	2.34E-07	EPA6 calc c
Ethyl methacrylate	97-63-2	EPA6 e	3.09E-07	EPA6 e	9.77E-07	EPA6 e	1.18E-06	EPA6 e
Formaldehyde	50-00-0	EPA6 e	1.75E-08	EPA6 e	5.53E-08	EPA6 e	6.69E-08	EPA6 e
Formamide	75-12-7	EPA6 calc c	2.45E-10	EPA6 calc c	7.76E-10	EPA6 calc c	9.31E-10	EPA6 calc c
Formic acid	64-18-6	EPA6 e	2.30E-09	EPA6 e	7.28E-09	EPA6 e	8.82E-09	EPA6 e
Formic acid, methyl ester	107-31-3	NA	4.33E-09	EPA6 calc c	1.37E-08	EPA6 calc c	1.64E-08	EPA6 calc c
Glycidylaldehyde	765-34-4	EPA6 calc c	1.48E-09	EPA6 calc c	4.68E-09	EPA6 calc c	5.61E-09	EPA6 calc c
Methyl acetate	79-20-9	EPA6 calc c	1.20E-08	EPA6 calc c	3.80E-08	EPA6 calc c	4.56E-08	EPA6 calc c
Methyl alcohol	67-56-1	EPA6 e	1.30E-09	EPA6 e	4.30E-09	EPA6 e	5.21E-09	EPA6 e
Methyl isocyanate	624-83-9	NA	NA	NA	NA	NA	NA	NA
Methyl methacrylate	80-62-6	EPA6 calc c	4.90E-08	EPA6 calc c	1.55E-07	EPA6 calc c	1.86E-07	EPA6 calc c
Methyl tert-butyl ether	1634-04-4	EPA6 calc c	6.92E-08	EPA6 calc c	2.19E-07	EPA6 calc c	2.63E-07	EPA6 calc c
Methylacetylene	74-99-7	NA	6.92E-08	EPA6 calc c	2.19E-07	EPA6 calc c	2.63E-07	EPA6 calc c
Methylcyclohexane	108-87-2	EPA6 calc c	1.00E-04	EPA6 calc c	3.16E-04	EPA6 calc c	3.79E-04	EPA6 calc c
N,N-Dimethylacetamide	127-19-5	NA	NA	NA	NA	NA	NA	NA
n-Butyl alcohol	71-36-3	EPA6 calc c	6.03E-08	EPA6 calc c	1.91E-07	EPA6 calc c	2.29E-07	EPA6 calc c
n-Heptane	142-82-5	EPA6 calc c	3.63E-04	EPA6 calc c	1.15E-03	EPA6 calc c	1.38E-03	EPA6 calc c
n-Hexane	110-54-3	EPA6 calc c	1.02E-04	EPA6 calc c	3.24E-04	EPA6 calc c	3.88E-04	EPA6 calc c
Nitromethane	75-52-5	NA	3.55E-09	EPA6 calc c	1.12E-08	EPA6 calc c	1.35E-08	EPA6 calc c
n-Nonane	111-84-2	EPA6 calc c	3.55E-03	EPA6 calc c	1.12E-02	EPA6 calc c	1.35E-02	EPA6 calc c
n-Octane	111-65-9	EPA6 calc c	7.94E-05	EPA6 calc c	2.51E-04	EPA6 calc c	3.01E-04	EPA6 calc c
n-Pentane	109-66-0	EPA6 calc c	1.29E-05	EPA6 calc c	4.07E-05	EPA6 calc c	4.89E-05	EPA6 calc c
n-Propionaldehyde	123-38-6	EPA6 calc c	3.09E-08	EPA6 calc c	9.77E-08	EPA6 calc c	1.17E-07	EPA6 calc c
n-Propyl alcohol	71-23-8	NA	NA	NA	NA	NA	NA	NA
n-Valeraldehyde	110-62-3	NA	NA	NA	NA	NA	NA	NA
Oxirane	75-21-8	EPA6 e	3.98E-09	EPA6 e	1.26E-08	EPA6 e	1.52E-08	EPA6 e

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Bioconcentration / Bioaccumulation Factors					
Constituent of Potential Concern	CAS Registry Number	Source	Ba_{milk} (day/kg FW)	Source	Ba_{beef} (day/kg FW)	Source	Ba_{pork} (day/kg FW)	Source
p-Cymene	99-87-6	EPA6 calc c	1.00E-04	EPA6 calc c	3.16E-04	EPA6 calc c	3.79E-04	EPA6 calc c
Phosgene	75-44-5	NA	NA	NA	NA	NA	NA	NA
Propargyl alcohol	107-19-7	NA	NA	NA	NA	NA	NA	NA
Propionic acid	79-09-4	EPA6 calc c	1.70E-08	EPA6 calc c	5.37E-08	EPA6 calc c	6.44E-08	EPA6 calc c
Propionitrile	107-12-0	EPA6 calc c	1.15E-08	EPA6 calc c	3.63E-08	EPA6 calc c	4.36E-08	EPA6 calc c
Propylene glycol monomethyl ether	107-98-2	EPA6 calc c	5.25E-09	EPA6 calc c	1.66E-08	EPA6 calc c	1.99E-08	EPA6 calc c
p-tert-Butyltoluene	98-51-1	NA	NA	NA	NA	NA	NA	NA
Triethylamine	121-44-8	NA	NA	NA	NA	NA	NA	NA
Trimethylamine	75-50-3	EPA6 calc c	1.15E-08	EPA6 calc c	3.63E-08	EPA6 calc c	4.36E-08	EPA6 calc c
Vinyl acetate	108-05-4	EPA6 e	3.97E-08	EPA6 e	1.26E-07	EPA6 e	1.52E-07	EPA6 e
Non-aromatic Halogenated Hydrocarbons								
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	NA	NA	NA	NA	NA	NA	NA
1,1,1,2-Tetrachloroethane	630-20-6	EPA6 e	3.39E-06	EPA6 e	1.07E-05	EPA6 e	1.30E-05	EPA6 e
1,1,1-Trichloroethane	71-55-6	EPA6 e	2.10E-06	EPA6 e	6.63E-06	EPA6 e	8.03E-06	EPA6 e
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	EPA6 calc c	4.27E-05	EPA6 calc c	1.35E-04	EPA6 calc c	1.62E-04	EPA6 calc c
1,1,2,2-Tetrachloroethane	79-34-5	EPA6 e	3.50E-04	EPA6 e	1.11E-03	EPA6 e	1.34E-03	EPA6 e
1,1,2,2-Tetrachloroethene	127-18-4	EPA6 e	2.79E-06	EPA6 e	8.82E-06	EPA6 e	1.07E-05	EPA6 e
1,1,2-Trichloro-1,2,2-trifluoroethane	76-13-1	EPA6 calc c	1.15E-05	EPA6 calc c	3.63E-05	EPA6 calc c	4.36E-05	EPA6 calc c
1,1,2-Trichloroethane	79-00-5	EPA6 e	9.93E-07	EPA6 e	3.14E-06	EPA6 e	3.80E-06	EPA6 e
1,1,2-Trichloroethylene	79-01-6	EPA6 e	2.15E-06	EPA6 e	6.81E-06	EPA6 e	8.24E-06	EPA6 e
1,1-Dichloroethane	75-34-3	EPA6 e	4.93E-07	EPA6 e	1.56E-06	EPA6 e	1.89E-06	EPA6 e
1,1-Dichloroethene	75-35-4	EPA6 e	1.05E-06	EPA6 e	3.32E-06	EPA6 e	4.01E-06	EPA6 e
1,2,3-Trichloropropane	96-18-4	EPA6 e	1.41E-06	EPA6 e	4.47E-06	EPA6 e	5.41E-06	EPA6 e
1,2-dibromo-3-chloropropane	96-12-8	EPA6 e	1.74E-06	EPA6 e	5.50E-06	EPA6 e	6.65E-06	EPA6 e
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	NA	NA	NA	NA	NA	NA	NA
1,2-Dichloroethane	107-06-2	EPA6 e	2.30E-07	EPA6 e	7.28E-07	EPA6 e	8.82E-07	EPA6 e
1,2-Dichloroethylene	540-59-0	EPA6 calc c	2.40E-08	EPA6 calc c	7.59E-08	EPA6 calc c	9.10E-08	EPA6 calc c
1,2-Dichloropropane	78-87-5	EPA6 e	1.41E-06	EPA6 e	4.47E-06	EPA6 e	5.41E-06	EPA6 e
1,3-Dichloropropene	542-75-6	EPA6 e	4.45E-07	EPA6 e	1.41E-06	EPA6 e	1.70E-06	EPA6 e

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Bioconcentration / Bioaccumulation Factors					
Constituent of Potential Concern	CAS Registry Number	Source	Ba_{milk} (day/kg FW)	Source	Ba_{beef} (day/kg FW)	Source	Ba_{pork} (day/kg FW)	Source
1,4-Dichloro-2-butene	764-41-0	EPA6 calc c	5.92E-08	EPA6 calc c	1.87E-07	EPA6 calc c	2.25E-07	EPA6 calc c
1-Chloroethene	75-01-4	EPA6 e	1.11E-07	EPA6 e	3.52E-07	EPA6 e	4.26E-07	EPA6 e
2,2-Dichloropropionic acid	75-99-0	EPA6 calc c	4.76E-08	EPA6 calc c	1.51E-07	EPA6 calc c	1.81E-07	EPA6 calc c
2-Chloropropane	75-29-6	EPA6 calc c	6.31E-07	EPA6 calc c	2.00E-06	EPA6 calc c	2.39E-06	EPA6 calc c
3-Chloropropene (allyl chloride)	107-05-1	EPA6 calc c	7.15E-08	EPA6 calc c	2.26E-07	EPA6 calc c	2.71E-07	EPA6 calc c
Bromochloromethane	74-97-5	EPA6 calc c	2.04E-07	EPA6 calc c	6.46E-07	EPA6 calc c	7.75E-07	EPA6 calc c
Bromodichloromethane	75-27-4	EPA6 e	8.42E-07	EPA6 e	2.66E-06	EPA6 e	3.22E-06	EPA6 e
Bromoethene	593-60-2	EPA6 calc c	9.26E-08	EPA6 calc c	2.93E-07	EPA6 calc c	3.51E-07	EPA6 calc c
Bromoform	75-25-2	EPA6 e	1.78E-06	EPA6 e	5.63E-06	EPA6 e	6.81E-06	EPA6 e
Bromomethane	74-83-9	EPA6 e	1.03E-07	EPA6 e	3.27E-07	EPA6 e	3.95E-07	EPA6 e
Carbon tetrachloride	56-23-5	EPA6 e	4.14E-06	EPA6 e	1.30E-05	EPA6 e	1.58E-05	EPA6 e
Chlordibromomethane	124-48-1	EPA6 e	1.19E-06	EPA6 e	3.77E-06	EPA6 e	4.56E-06	EPA6 e
Chlorodifluoromethane	75-45-6	EPA6 e	9.53E-08	EPA6 e	3.01E-07	EPA6 e	3.65E-07	EPA6 e
Chloroethane	75-00-3	EPA6 e	1.00E-05	EPA6 e	3.16E-05	EPA6 e	3.83E-05	EPA6 e
Chloroform	67-66-3	EPA6 e	7.07E-07	EPA6 e	2.23E-06	EPA6 e	2.71E-06	EPA6 e
Chloromethane	74-87-3	EPA6 e	6.36E-08	EPA6 e	2.01E-07	EPA6 e	2.43E-07	EPA6 e
Chloropentafluoroethane	76-15-3	NA	NA	NA	NA	NA	NA	NA
cis-1,2-Dichloroethene	156-59-2	EPA6 e	7.63E-07	EPA6 e	2.41E-06	EPA6 e	2.92E-06	EPA6 e
cis-1,3-Dichloropropene	10061-01-5	NA	NA	NA	NA	NA	NA	NA
Cyanogen bromide	506-68-3	NA	NA	NA	NA	NA	NA	NA
Cyanogen chloride	506-77-4	NA	NA	NA	NA	NA	NA	NA
Dichlorodifluoromethane	75-71-8	EPA6 e	1.15E-06	EPA6 e	3.63E-06	EPA6 e	4.40E-06	EPA6 e
Dichlorofluoromethane	75-43-4	NA	NA	NA	NA	NA	NA	NA
Dichloromethane	75-09-2	EPA6 e	1.43E-07	EPA6 e	4.52E-07	EPA6 e	5.47E-07	EPA6 e
Difluorodibromomethane	75-61-6	NA	NA	NA	NA	NA	NA	NA
Hexafluoroacetone	684-16-2	NA	NA	NA	NA	NA	NA	NA
Iodomethane	74-88-4	EPA6 calc c	3.89E-07	EPA6 calc c	1.23E-06	EPA6 calc c	1.48E-06	EPA6 calc c
Methylene bromide	74-95-3	EPA6 e	3.31E-07	EPA6 e	1.05E-06	EPA6 e	1.27E-06	EPA6 e
Pentachloroethane	76-01-7	EPA6 calc c	8.91E-06	EPA6 calc c	2.82E-05	EPA6 calc c	3.38E-05	EPA6 calc c

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Bioconcentration / Bioaccumulation Factors					
Constituent of Potential Concern	CAS Registry Number	Source	Ba_{milk} (day/kg FW)	Source	Ba_{beef} (day/kg FW)	Source	Ba_{pork} (day/kg FW)	Source
trans-1,2-Dichloroethene	156-60-5	EPA6 e	7.63E-07	EPA6 e	2.41E-06	EPA6 e	2.92E-06	EPA6 e
trans-1,3-Dichloropropene	10061-02-6	NA	9.12E-07	EPA6 calc c	2.88E-06	EPA6 calc c	3.46E-06	EPA6 calc c
Trichloroacetic acid	76-03-9	EPA6 calc c	1.70E-07	EPA6 calc c	5.37E-07	EPA6 calc c	6.44E-07	EPA6 calc c
Trichlorofluoroethane	27154-33-2	NA	NA	NA	NA	NA	NA	NA
Trichlorofluoromethane	75-69-4	EPA6 e	2.70E-06	EPA6 e	8.54E-06	EPA6 e	1.03E-05	EPA6 e
Trifluorobromomethane	75-63-8	NA	5.75E-07	EPA6 calc c	1.82E-06	EPA6 calc c	2.18E-06	EPA6 calc c
Dioxin and Furan Compounds								
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	35822-46-9	EPA6 e	1.00E-03	EPA6 e	5.40E-03	EPA6 e	6.57E-03	EPA6 e
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	EPA6 e	1.00E-03	EPA6 e	5.43E-03	EPA6 e	6.57E-03	EPA6 e
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	EPA6 e	3.00E-03	EPA6 e	1.63E-02	EPA6 e	1.97E-02	EPA6 e
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	39227-28-6	EPA6 e	6.00E-03	EPA6 e	3.26E-02	EPA6 e	3.94E-02	EPA6 e
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	EPA6 e	7.00E-03	EPA6 e	3.80E-02	EPA6 e	4.60E-02	EPA6 e
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	57653-85-7	EPA6 e	5.00E-03	EPA6 e	2.71E-02	EPA6 e	3.29E-02	EPA6 e
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	EPA6 e	6.00E-03	EPA6 e	3.26E-02	EPA6 e	3.94E-02	EPA6 e
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	19408-74-3	EPA6 e	5.00E-03	EPA6 e	2.71E-02	EPA6 e	3.29E-02	EPA6 e
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	EPA6 e	6.00E-03	EPA6 e	3.26E-02	EPA6 e	3.94E-02	EPA6 e
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	40321-76-4	EPA6 e	1.00E-02	EPA6 e	5.43E-02	EPA6 e	6.57E-02	EPA6 e
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	EPA6 e	2.00E-03	EPA6 e	1.09E-02	EPA6 e	1.31E-02	EPA6 e
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	EPA6 e	5.00E-03	EPA6 e	2.71E-02	EPA6 e	3.29E-02	EPA6 e
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	EPA6 e	9.00E-03	EPA6 e	4.89E-02	EPA6 e	5.91E-02	EPA6 e
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1746-01-6	EPA6 e	1.00E-02	EPA6 e	5.43E-02	EPA6 e	6.57E-02	EPA6 e
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	EPA6 e	3.00E-03	EPA6 e	1.63E-02	EPA6 e	1.97E-02	EPA6 e
Dibenzofuran	132-64-9	EPA6 calc c	1.70E-04	EPA6 calc c	5.37E-04	EPA6 calc c	6.44E-04	EPA6 calc c
Octachlorodibenzo(p)dioxin	3268-87-9	EPA6 e	1.00E-03	EPA6 e	5.43E-03	EPA6 e	6.57E-03	EPA6 e
Octachlorodibenzofuran	39001-02-0	EPA6 e	1.00E-03	EPA6 e	5.43E-03	EPA6 e	6.57E-03	EPA6 e
PCBs								
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	EPA6 calc c	9.55E-02	EPA6 calc c	3.02E-01	EPA6 calc c	3.62E-01	EPA6 calc c
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	EPA6 calc c	1.05E-01	EPA6 calc c	3.31E-01	EPA6 calc c	3.97E-01	EPA6 calc c
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	NA	NA	NA	NA	NA	NA	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Bioconcentration / Bioaccumulation Factors					
Constituent of Potential Concern	CAS Registry Number	Source	Ba_{milk} (day/kg FW)	Source	Ba_{beef} (day/kg FW)	Source	Ba_{pork} (day/kg FW)	Source
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA	NA
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	NA	NA	NA	NA	NA	NA	NA
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	NA	NA	NA	NA	NA	NA	NA
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA	NA
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	NA	NA	NA	NA	NA	NA	NA
2',3,4,4',5-Pentachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA	NA
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	NA	1.05E-01	EPA6 calc c	3.31E-01	EPA6 calc c	3.97E-01	EPA6 calc c
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	EPA6 calc c	2.03E-01	EPA6 calc c	6.43E-01	EPA6 calc c	7.71E-01	EPA6 calc c
3,3',4,4',5-Pentachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA	NA
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	NA	NA	NA	NA	NA	NA	NA
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	NA	NA	NA	NA	NA	NA	NA
Polychlorinated biphenyls (PCBs)**	1336-36-3	EPA6 calc c	1.62E-01	EPA6 calc c	5.11E-01	EPA6 calc c	6.13E-01	EPA6 calc c
Phthalates								
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	EPA6 e	1.27E-03	EPA6 e	4.03E-03	EPA6 e	4.88E-03	EPA6 e
Butylbenzyl phthalate	85-68-7	EPA6 e	2.06E-04	EPA6 e	6.50E-04	EPA6 e	7.87E-04	EPA6 e
Dibutyl phthalate	84-74-2	EPA6 e	4.17E-04	EPA6 e	1.32E-03	EPA6 e	1.60E-03	EPA6 e
Diethyl phthalate	84-66-2	EPA6 e	2.17E-04	EPA6 e	6.87E-04	EPA6 e	8.31E-04	EPA6 e
Dimethyl phthalate	131-11-3	EPA6 e	3.42E-07	EPA6 e	1.08E-06	EPA6 e	1.31E-06	EPA6 e
n-Dioctyl phthalate	117-84-0	EPA6 e	1.70E+01	EPA6 e	5.37E+01	EPA6 e	6.50E+01	EPA6 e
Light Polycyclic Aromatic Hydrocarbons								
2-Chloronaphthalene	91-58-7	EPA6 e	9.33E-05	EPA6 e	2.95E-04	EPA6 e	3.57E-04	EPA6 e
2-Methyl naphthalene	91-57-6	EPA6 calc c	5.75E-05	EPA6 calc c	1.82E-04	EPA6 calc c	2.18E-04	EPA6 calc c
5-Nitroacenaphthene	602-87-9	NA	NA	NA	NA	NA	NA	NA
Acenaphthene	83-32-9	EPA6 e	7.32E-05	EPA6 e	2.31E-04	EPA6 e	2.80E-04	EPA6 e
Acenaphthylene	208-96-8	EPA6 calc c	9.33E-05	EPA6 calc c	2.95E-04	EPA6 calc c	3.54E-04	EPA6 calc c
Anthracene	120-12-7	EPA6 e	2.34E-04	EPA6 e	7.41E-04	EPA6 e	8.98E-04	EPA6 e
Fluorene	86-73-7	EPA6 calc c	1.20E-04	EPA6 calc c	3.80E-04	EPA6 calc c	4.56E-04	EPA6 calc c
Indene	95-13-6	NA	NA	NA	NA	NA	NA	NA
Napthalene	91-20-3	EPA6 e	1.87E-05	EPA6 e	5.92E-05	EPA6 e	7.16E-05	EPA6 e

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Bioconcentration / Bioaccumulation Factors					
Constituent of Potential Concern	CAS Registry Number	Source	Ba_{milk} (day/kg FW)	Source	Ba_{beef} (day/kg FW)	Source	Ba_{pork} (day/kg FW)	Source
Phenanthrene	85-01-8	EPA6 e	2.82E-04	EPA6 e	8.92E-04	EPA6 e	1.08E-03	EPA6 e
Pyrene	129-00-0	EPA6 e	7.98E-04	EPA6 e	2.52E-03	EPA6 e	3.06E-03	EPA6 e
Heavy Polycyclic Aromatic Hydrocarbons								
3-Methylcholanthrene	56-49-5	EPA6 calc c	1.02E-01	EPA6 calc c	3.24E-01	EPA6 calc c	3.88E-01	EPA6 calc c
5-Methylchrysene	3697-24-3	NA	NA	NA	NA	NA	NA	NA
Benzo(a)anthracene	56-55-3	EPA6 e	3.79E-03	EPA6 e	1.20E-02	EPA6 e	1.45E-02	EPA6 e
Benzo(a)pyrene	50-32-8	EPA6 e	1.07E-02	EPA6 e	3.38E-02	EPA6 e	4.10E-02	EPA6 e
Benzo(b)fluoranthene	205-99-2	EPA6 e	1.27E-02	EPA6 e	4.00E-02	EPA6 e	4.84E-02	EPA6 e
Benzo(e)pyrene	192-97-2	EPA6 calc c	2.00E-01	EPA6 calc c	6.31E-01	EPA6 calc c	7.57E-01	EPA6 calc c
Benzo(g,h,i)perylene	191-24-2	EPA6 calc c	1.00E-01	EPA6 calc c	3.16E-01	EPA6 calc c	3.79E-01	EPA6 calc c
Benzo(j)fluoranthene	205-82-3	EPA6 calc c	2.19E-02	EPA6 calc c	6.92E-02	EPA6 calc c	8.30E-02	EPA6 calc c
Benzo(k)fluoranthene	207-08-9	EPA6 e	1.26E-02	EPA6 e	3.98E-02	EPA6 e	4.82E-02	EPA6 e
Benzo[a,i]pyrene	191-30-0	NA	NA	NA	NA	NA	NA	NA
Chrysene	218-01-9	EPA6 e	4.36E-03	EPA6 e	1.38E-02	EPA6 e	1.67E-02	EPA6 e
Dibenz(a,h)anthracene	53-70-3	EPA6 e	2.80E-02	EPA6 e	8.86E-02	EPA6 e	1.07E-01	EPA6 e
Dibenz[a,h]acridine	226-36-8	NA	NA	NA	NA	NA	NA	NA
Dibenz[a,j]acridine	224-42-0	NA	NA	NA	NA	NA	NA	NA
Dibenzo(a,e)fluoranthene	5385-75-1	NA	NA	NA	NA	NA	NA	NA
Dibenzo(a,h)fluoranthene	no cas #	NA	NA	NA	NA	NA	NA	NA
Dibenzo[a,e]pyrene	192-65-4	NA	NA	NA	NA	NA	NA	NA
Dibenzo[a,h]pyrene	189-64-0	NA	NA	NA	NA	NA	NA	NA
Dibenzo[a,i]pyrene	189-55-9	NA	1.55E-01	EPA6 calc c	4.90E-01	EPA6 calc c	5.88E-01	EPA6 calc c
Fluoranthene	206-44-0	EPA6 e	9.65E-04	EPA6 e	3.05E-03	EPA6 e	3.69E-03	EPA6 e
Hexachloronaphthalene	1335-87-1	NA	3.09E-01	EPA6 calc c	9.77E-01	EPA6 calc c	1.17E+00	EPA6 calc c
Indeno(1,2,3-cd)pyrene	193-39-5	EPA6 e	6.53E-02	EPA6 e	2.07E-01	EPA6 e	2.50E-01	EPA6 e
Octachloronaphthalene	2234-13-1	NA	2.09E-02	EPA6 calc c	6.61E-02	EPA6 calc c	7.93E-02	EPA6 calc c
Pentachloronaphthalene	1321-64-8	NA	NA	NA	NA	NA	NA	NA
Tetrachloronaphthalene	1335-88-2	NA	NA	NA	NA	NA	NA	NA
Trichloronaphthalene	1321-65-9	NA	NA	NA	NA	NA	NA	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Bioconcentration / Bioaccumulation Factors					
Constituent of Potential Concern	CAS Registry Number	Source	Ba_{milk} (day/kg FW)	Source	Ba_{beef} (day/kg FW)	Source	Ba_{pork} (day/kg FW)	Source
Light Substituted Benzene Compounds								
1,2,3-Trichlorobenzene	87-61-6	EPA6 e	8.82E-05	EPA6 e	2.79E-04	EPA6 e	3.38E-04	EPA6 e
1,2,4-Trichlorobenzene	120-82-1	EPA6 e	7.73E-05	EPA6 e	2.45E-04	EPA6 e	2.96E-04	EPA6 e
1,2,4-Trimethlyl benzene	25551-13-7	NA	2.00E-05	EPA6 calc c	6.31E-05	EPA6 calc c	7.57E-05	EPA6 calc c
1,2,4-Trimethyl benzene	95-63-6	EPA6 calc c	3.55E-05	EPA6 calc c	1.12E-04	EPA6 calc c	1.35E-04	EPA6 calc c
1,2-Dichlorobenzene	95-50-1	EPA6 e	2.21E-05	EPA6 e	7.00E-05	EPA6 e	8.48E-05	EPA6 e
1,3,5-Trimethyl benzene	108-67-8	EPA6 e	2.09E-05	EPA6 e	6.61E-05	EPA6 e	8.00E-05	EPA6 e
1,3-Dichlorobenzene	541-73-1	EPA6 e	2.69E-05	EPA6 e	8.52E-05	EPA6 e	1.03E-04	EPA6 e
1,3-Dinitrobenzene	99-65-0	EPA6 e	2.46E-07	EPA6 e	7.79E-07	EPA6 e	9.43E-07	EPA6 e
1,4-Dichlorobenzene	106-46-7	EPA6 e	2.05E-05	EPA6 e	6.49E-05	EPA6 e	7.86E-05	EPA6 e
1,4-Dinitrobenzene	100-25-4	NA	2.51E-07	EPA6 calc c	7.94E-07	EPA6 calc c	9.53E-07	EPA6 calc c
2,4,5-Trichorophenol	95-95-4	EPA6 e	5.89E-05	EPA6 e	1.86E-04	EPA6 e	2.25E-04	EPA6 e
2,4,6-Trichlorophenol	88-06-2	EPA6 e	4.09E-05	EPA6 e	1.29E-04	EPA6 e	1.57E-04	EPA6 e
2,4-Dichlorophenol	120-83-2	EPA6 e	8.62E-06	EPA6 e	2.73E-05	EPA6 e	3.30E-05	EPA6 e
2,4-Dimethylphenol	105-67-9	EPA6 e	1.82E-06	EPA6 e	5.75E-06	EPA6 e	6.96E-06	EPA6 e
2,4-Dinitrophenol	51-28-5	EPA6 e	2.62E-07	EPA6 e	8.29E-07	EPA6 e	1.00E-06	EPA6 e
2,4-Dinitrotoluene	121-14-2	EPA6 e	7.86E-07	EPA6 e	2.49E-06	EPA6 e	3.01E-06	EPA6 e
2,6-Dinitrotoluene	606-20-2	EPA6 e	6.12E-07	EPA6 e	1.93E-06	EPA6 e	2.34E-06	EPA6 e
2-Chlorotoluene	95-49-8	EPA6 calc c	2.73E-05	EPA6 calc c	8.64E-05	EPA6 calc c	1.04E-04	EPA6 calc c
2-Chorophenol	95-57-8	EPA6 e	1.15E-06	EPA6 e	3.64E-06	EPA6 e	4.41E-06	EPA6 e
2-Nitrophenol	88-75-5	EPA6 e	4.90E-07	EPA6 e	1.55E-06	EPA6 e	1.88E-06	EPA6 e
4,6-Dinitro-o-cresol	534-52-1	EPA6 calc c	5.62E-06	EPA6 calc c	1.78E-05	EPA6 calc c	2.13E-05	EPA6 calc c
4-Chlorotoluene	106-43-4	NA	1.70E-05	EPA6 calc c	5.37E-05	EPA6 calc c	6.44E-05	EPA6 calc c
4-Nitrophenol	100-02-7	EPA6 e	6.46E-07	EPA6 e	2.04E-06	EPA6 e	2.47E-06	EPA6 e
alpha-Methylstyrene	98-83-9	EPA6 calc c	2.31E-05	EPA6 calc c	7.30E-05	EPA6 calc c	8.76E-05	EPA6 calc c
Aniline	62-53-3	EPA6 e	7.59E-08	EPA6 e	2.40E-07	EPA6 e	2.90E-07	EPA6 e
Benzotrichloride	98-07-7	EPA6 calc c	6.61E-06	EPA6 calc c	2.09E-05	EPA6 calc c	2.51E-05	EPA6 calc c
Benzyl Chloride	100-44-7	EPA6 e	1.83E-08	EPA6 e	5.78E-08	EPA6 e	6.99E-08	EPA6 e
Bromobenzene	108-86-1	EPA6 calc c	7.76E-06	EPA6 calc c	2.45E-05	EPA6 calc c	2.95E-05	EPA6 calc c

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Bioconcentration / Bioaccumulation Factors					
Constituent of Potential Concern	CAS Registry Number	Source	Ba_{milk} (day/kg FW)	Source	Ba_{beef} (day/kg FW)	Source	Ba_{pork} (day/kg FW)	Source
Chlorobenzene	108-90-7	EPA6 e	4.89E-06	EPA6 e	1.55E-05	EPA6 e	1.87E-05	EPA6 e
Cumene	98-82-8	EPA6 e	3.26E-05	EPA6 e	1.03E-04	EPA6 e	1.25E-04	EPA6 e
m-Cresol	108-39-4	EPA6 e	7.23E-07	EPA6 e	2.29E-06	EPA6 e	2.77E-06	EPA6 e
n-Butyl benzene	104-51-8	EPA6 calc c	1.51E-04	EPA6 calc c	4.79E-04	EPA6 calc c	5.74E-04	EPA6 calc c
Nitrobenzene	98-95-3	EPA6 e	5.40E-07	EPA6 e	1.71E-06	EPA6 e	2.07E-06	EPA6 e
n-Propyl benzene	103-65-1	EPA6 calc c	3.89E-05	EPA6 calc c	1.23E-04	EPA6 calc c	1.48E-04	EPA6 calc c
o-Cresol	95-48-7	EPA6 e	8.34E-07	EPA6 e	2.64E-06	EPA6 e	3.19E-06	EPA6 e
o-Dinitrobenzene	528-29-0	NA	3.16E-07	EPA6 calc c	1.00E-06	EPA6 calc c	1.20E-06	EPA6 calc c
o-Nitroaniline	88-74-4	EPA6 e	5.62E-07	EPA6 e	1.78E-06	EPA6 e	2.15E-06	EPA6 e
o-Toluidine	95-53-4	EPA6 e	1.74E-07	EPA6 e	5.50E-07	EPA6 e	6.65E-07	EPA6 e
p-Chloroaniline	106-47-8	EPA6 e	5.88E-07	EPA6 e	1.86E-06	EPA6 e	2.25E-06	EPA6 e
p-Cresol	106-44-5	EPA6 e	6.91E-07	EPA6 e	2.19E-06	EPA6 e	2.65E-06	EPA6 e
Phenol	108-95-2	EPA6 e	2.38E-07	EPA6 e	7.54E-07	EPA6 e	9.12E-07	EPA6 e
p-Nitrochlorobenzene	100-00-5	NA	1.95E-06	EPA6 calc c	6.17E-06	EPA6 calc c	7.40E-06	EPA6 calc c
p-Toluidine	106-49-0	EPA6 calc c	2.00E-07	EPA6 calc c	6.31E-07	EPA6 calc c	7.57E-07	EPA6 calc c
sec-Butyl benzene	135-98-8	NA	NA	NA	NA	NA	NA	NA
tert-Butyl benzene	98-06-6	EPA6 calc c	1.02E-04	EPA6 calc c	3.24E-04	EPA6 calc c	3.88E-04	EPA6 calc c
Toluene-2,6-diamine	823-40-5	EPA6 calc c	2.26E-07	EPA6 calc c	7.16E-07	EPA6 calc c	8.59E-07	EPA6 calc c
Other Light Semivolatile Compounds								
1,1 Dimethyl hydrazine	57-14-7	EPA6 calc c	3.13E-08	EPA6 calc c	9.89E-08	EPA6 calc c	1.19E-07	EPA6 calc c
1,1-Biphenyl	92-52-4	EPA6 calc c	6.31E-05	EPA6 calc c	2.00E-04	EPA6 calc c	2.39E-04	EPA6 calc c
1,2 Diphenylhydrazine	122-66-7	EPA6 e	6.92E-06	EPA6 e	2.19E-05	EPA6 e	2.65E-05	EPA6 e
1,2-Dimethylhydrazine	540-73-8	NA	3.40E-10	EPA6 calc c	1.08E-09	EPA6 calc c	1.29E-09	EPA6 calc c
1,3-Propane sultone	1120-71-4	NA	2.38E-09	EPA6 calc c	7.53E-09	EPA6 calc c	9.04E-09	EPA6 calc c
2,4-Toluene diisocyanate	584-84-9	NA	NA	NA	NA	NA	NA	NA
2-Chloroacetophenone	532-27-4	EPA6 calc c	3.12E-06	EPA6 calc c	9.88E-06	EPA6 calc c	1.19E-05	EPA6 calc c
2-Propenoic acid	79-10-7	EPA6 calc c	2.14E-08	EPA6 calc c	6.76E-08	EPA6 calc c	8.11E-08	EPA6 calc c
4,4-Methylenedianiline	101-77-9	EPA6 calc c	1.90E-05	EPA6 calc c	6.02E-05	EPA6 calc c	7.22E-05	EPA6 calc c
Acetophenone	98-86-2	EPA6 e	3.47E-07	EPA6 e	1.10E-06	EPA6 e	1.33E-06	EPA6 e

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Bioconcentration / Bioaccumulation Factors					
Constituent of Potential Concern	CAS Registry Number	Source	Ba_{milk} (day/kg FW)	Source	Ba_{beef} (day/kg FW)	Source	Ba_{pork} (day/kg FW)	Source
Benzoic acid	65-85-0	NA	6.04E-07	NA	1.91E-06	NA	2.31E-06	NA
bis(2-Chloroethoxy)methane	111-91-1	NA	NA	NA	NA	NA	NA	NA
bis(2-Chloroethyl) ether	111-44-4	EPA6 e	1.59E-07	EPA6 e	5.02E-07	EPA6 e	6.08E-07	EPA6 e
Chlorocyclopentadiene	41851-50-7	EPA6 calc c	2.14E-06	EPA6 calc c	6.76E-06	EPA6 calc c	8.11E-06	EPA6 calc c
Cyclohexanol	108-93-0	EPA6 calc c	1.35E-07	EPA6 calc c	4.27E-07	EPA6 calc c	5.12E-07	EPA6 calc c
Dichloroisopropyl ether	108-60-1	EPA6 calc c	3.02E-06	EPA6 calc c	9.55E-06	EPA6 calc c	1.15E-05	EPA6 calc c
Dichloromethyl ether	542-88-1	EPA6 calc c	3.31E-09	EPA6 calc c	1.05E-08	EPA6 calc c	1.26E-08	EPA6 calc c
Dichloropentadiene	no cas #	NA	NA	NA	NA	NA	NA	NA
Dimethyl sulfate	77-78-1	EPA6 calc c	1.65E-08	EPA6 calc c	5.23E-08	EPA6 calc c	6.27E-08	EPA6 calc c
Dimethylaniline	121-69-7	EPA6 calc c	1.05E-09	EPA6 calc c	3.31E-09	EPA6 calc c	3.97E-09	EPA6 calc c
di-n-propylnitrosamine	621-64-7	EPA6 e	1.91E-07	EPA6 e	6.03E-07	EPA6 e	7.30E-07	EPA6 e
Diphenyl ether	101-84-8	EPA6 calc c	1.29E-04	EPA6 calc c	4.07E-04	EPA6 calc c	4.89E-04	EPA6 calc c
Epichlorohydrin	106-89-8	EPA6 e	1.41E-08	EPA6 e	4.47E-08	EPA6 e	5.41E-08	EPA6 e
Ethyl carbomate (urethane)	51-79-6	NA	5.62E-09	EPA6 calc c	1.78E-08	EPA6 calc c	2.13E-08	EPA6 calc c
Ethyl methane sulfanate	62-50-0	EPA6 e	8.90E-09	EPA6 e	2.81E-08	EPA6 e	3.41E-08	EPA6 e
Ethylene dibromide	106-93-4	EPA6 e	4.47E-07	EPA6 e	1.41E-06	EPA6 e	1.71E-06	EPA6 e
Ethylene glycol	107-21-1	EPA6 calc c	9.67E-10	EPA6 calc c	3.06E-09	EPA6 calc c	3.67E-09	EPA6 calc c
Ethylene glycol monobutyl ether	111-76-2	EPA6 calc c	2.85E-07	EPA6 calc c	9.00E-07	EPA6 calc c	1.08E-06	EPA6 calc c
Ethylene glycol monoethyl ether acetate	111-15-9	EPA6 calc c	3.31E-08	EPA6 calc c	1.05E-07	EPA6 calc c	1.25E-07	EPA6 calc c
Ethylene thiourea	96-45-7	EPA6 calc c	1.81E-09	EPA6 calc c	5.73E-09	EPA6 calc c	6.88E-09	EPA6 calc c
Furfural	98-01-1	EPA6 calc c	7.24E-08	EPA6 calc c	2.29E-07	EPA6 calc c	2.75E-07	EPA6 calc c
Maleic hydrazide	123-33-1	EPA6 calc c	1.45E-09	EPA6 calc c	4.57E-09	EPA6 calc c	5.49E-09	EPA6 calc c
Malononitrile	109-77-3	NA	8.71E-09	EPA6 calc c	2.75E-08	EPA6 calc c	3.31E-08	EPA6 calc c
Methyl styrene (mixed isomers)	25013-15-4	NA	1.78E-05	EPA6 calc c	5.62E-05	EPA6 calc c	6.75E-05	EPA6 calc c
Methylhydrazine	60-34-4	NA	7.08E-10	EPA6 calc c	2.24E-09	EPA6 calc c	2.69E-09	EPA6 calc c
N,N-Diphenylamine	122-39-4	EPA6 calc c	2.51E-05	EPA6 calc c	7.94E-05	EPA6 calc c	9.53E-05	EPA6 calc c
Nitric acid, propyl ester	627-13-4	NA	NA	NA	NA	NA	NA	NA
N-Nitrosodi-n-butylamine	924-16-3	EPA6 e	2.04E-06	EPA6 e	6.46E-06	EPA6 e	7.82E-06	EPA6 e
N-Nitrosomorpholine	59-89-2	NA	7.59E-08	EPA6 calc c	2.40E-07	EPA6 calc c	2.88E-07	EPA6 calc c

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Bioconcentration / Bioaccumulation Factors					
Constituent of Potential Concern	CAS Registry Number	Source	Ba_{milk} (day/kg FW)	Source	Ba_{beef} (day/kg FW)	Source	Ba_{pork} (day/kg FW)	Source
N-Nitroso-N,N-dimethylamine	62-75-9	EPA6 calc c	2.69E-09	EPA6 calc c	8.51E-09	EPA6 calc c	1.02E-08	EPA6 calc c
o-Anisidine	90-04-0	NA	1.20E-07	EPA6 calc c	3.80E-07	EPA6 calc c	4.56E-07	EPA6 calc c
Oxalic acid	144-62-7	NA	NA	NA	NA	NA	NA	NA
Phthalic Anhydride	85-44-9	EPA6 e	1.99E-09	EPA6 e	6.28E-09	EPA6 e	7.60E-09	EPA6 e
p-Phthalic acid	100-21-0	EPA6 calc c	5.28E-08	EPA6 calc c	1.67E-07	EPA6 calc c	2.00E-07	EPA6 calc c
Pyridine	110-86-1	EPA6 e	3.72E-08	EPA6 e	1.18E-07	EPA6 e	1.42E-07	EPA6 e
Quinoline	91-22-5	EPA6 calc c	8.51E-07	EPA6 calc c	2.69E-06	EPA6 calc c	3.23E-06	EPA6 calc c
Quinone	106-51-4	NA	1.26E-08	EPA6 calc c	3.98E-08	EPA6 calc c	4.78E-08	EPA6 calc c
Safrole	94-59-7	EPA6 e	3.63E-06	EPA6 e	1.15E-05	EPA6 e	1.39E-05	EPA6 e
Tetrahydrofuran	109-99-9	EPA6 e	2.22E-08	EPA6 e	7.03E-08	EPA6 e	8.51E-08	EPA6 e
Other Heavy Semivolatile Compounds								
1,2,4,5-Tetrachlorobenzene	95-94-3	EPA6 e	3.47E-04	EPA6 e	1.10E-03	EPA6 e	1.33E-03	EPA6 e
1,3,5-Trinitrobenzene	99-35-4	EPA6 e	1.20E-07	EPA6 e	3.80E-07	EPA6 e	4.60E-07	EPA6 e
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	NA	1.17E-04	EPA6 calc c	3.72E-04	EPA6 calc c	4.46E-04	EPA6 calc c
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	NA	1.58E-11	EPA6 calc c	5.01E-11	EPA6 calc c	6.01E-11	EPA6 calc c
2-sec-Butyl-4,6-dinitrophenol	88-85-7	EPA6 calc c	1.10E-05	EPA6 calc c	3.47E-05	EPA6 calc c	4.16E-05	EPA6 calc c
3,3-Dichlorobenzidine	91-94-1	EPA6 e	2.99E-05	EPA6 e	9.44E-05	EPA6 e	1.14E-04	EPA6 e
3,3'-Dimethoxybenzidine	119-90-4	EPA6 e	5.13E-07	EPA6 e	1.62E-06	EPA6 e	1.96E-06	EPA6 e
4-Bromophenylphenyl ether	101-55-3	EPA6 calc c	7.94E-04	EPA6 calc c	2.51E-03	EPA6 calc c	3.01E-03	EPA6 calc c
Ammonium perfluorooctanoate	3825-26-1	NA	NA	NA	NA	NA	NA	NA
Azobenzene	103-33-3	EPA6 calc c	5.25E-05	EPA6 calc c	1.66E-04	EPA6 calc c	1.99E-04	EPA6 calc c
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	NA	NA	NA	NA	NA	NA	NA
Captan	133-06-2	EPA6 calc c	1.78E-06	EPA6 calc c	5.62E-06	EPA6 calc c	6.75E-06	EPA6 calc c
Chlorobenzilate	510-15-6	EPA6 e	1.91E-04	EPA6 e	6.03E-04	EPA6 e	7.29E-04	EPA6 e
Dibutylphosphate	107-66-4	NA	NA	NA	NA	NA	NA	NA
Dimethyl aminoazobenzene	60-11-7	NA	3.02E-04	EPA6 calc c	9.55E-04	EPA6 calc c	1.15E-03	EPA6 calc c
Hexachlorobenzene	118-74-1	EPA6 e	2.53E-03	EPA6 e	7.99E-03	EPA6 e	9.68E-03	EPA6 e
Hexachlorobutadiene	87-68-3	EPA6 e	4.28E-04	EPA6 e	1.35E-03	EPA6 e	1.64E-03	EPA6 e
Hexachlorocyclopentadiene	77-47-4	EPA6 e	6.41E-04	EPA6 e	2.03E-03	EPA6 e	2.45E-03	EPA6 e

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Bioconcentration / Bioaccumulation Factors					
Constituent of Potential Concern	CAS Registry Number	Source	Ba_{milk} (day/kg FW)	Source	Ba_{beef} (day/kg FW)	Source	Ba_{pork} (day/kg FW)	Source
Hexachloroethane	67-72-1	EPA6 e	7.67E-05	EPA6 e	2.43E-04	EPA6 e	2.94E-04	EPA6 e
Hexachlorophene	70-30-4	EPA6 e	2.75E-01	EPA6 e	8.71E-01	EPA6 e	1.05E+00	EPA6 e
Hexamethylene-1,5-diisocyanate	822-06-0	EPA6 calc c	1.49E-07	EPA6 calc c	4.71E-07	EPA6 calc c	5.66E-07	EPA6 calc c
Mirex	2385-85-5	EPA6 calc c	6.17E-02	EPA6 calc c	1.95E-01	EPA6 calc c	2.34E-01	EPA6 calc c
Nitrofen	1836-75-5	NA	2.69E-03	EPA6 calc c	8.51E-03	EPA6 calc c	1.02E-02	EPA6 calc c
Pentachlorobenzene	608-93-5	EPA6 e	9.72E-04	EPA6 e	3.07E-03	EPA6 e	3.72E-03	EPA6 e
Pentachloronitrobenzene	82-68-8	EPA6 e	3.47E-04	EPA6 e	1.10E-03	EPA6 e	1.33E-03	EPA6 e
Pentachlorophenol*	87-86-5	EPA6 e	9.55E-04	EPA6 e	3.02E-03	EPA6 e	3.66E-03	EPA6 e
Pentachlorophenol*	87-86-5	EPA6 e	9.55E-04	EPA6 e	3.02E-03	EPA6 e	3.66E-03	EPA6 e
Pentachlorophenol*	87-86-5	EPA6 e	9.55E-04	EPA6 e	3.02E-03	EPA6 e	3.66E-03	EPA6 e
Pentachlorophenol*	87-86-5	EPA6 e	9.55E-04	EPA6 e	3.02E-03	EPA6 e	3.66E-03	EPA6 e
Pentachlorophenol*	87-86-5	EPA6 e	9.55E-04	EPA6 e	3.02E-03	EPA6 e	3.66E-03	EPA6 e
Pentachlorophenol*	87-86-5	EPA6 e	9.55E-04	EPA6 e	3.02E-03	EPA6 e	3.66E-03	EPA6 e
Pentachlorophenol*	87-86-5	EPA6 e	9.55E-04	EPA6 e	3.02E-03	EPA6 e	3.66E-03	EPA6 e
Pentachlorophenol*	87-86-5	EPA6 e	9.55E-04	EPA6 e	3.02E-03	EPA6 e	3.66E-03	EPA6 e
Pentachlorophenol*	87-86-5	EPA6 e	9.55E-04	EPA6 e	3.02E-03	EPA6 e	3.66E-03	EPA6 e
Pentachlorophenol*	87-86-5	EPA6 e	9.55E-04	EPA6 e	3.02E-03	EPA6 e	3.66E-03	EPA6 e
Pentachlorophenol*	87-86-5	EPA6 e	9.55E-04	EPA6 e	3.02E-03	EPA6 e	3.66E-03	EPA6 e
Pentachlorophenol*	87-86-5	EPA6 e	9.55E-04	EPA6 e	3.02E-03	EPA6 e	3.66E-03	EPA6 e
Picric acid	88-89-1	EPA6 calc c	8.51E-07	EPA6 calc c	2.69E-06	EPA6 calc c	3.23E-06	EPA6 calc c
Pronamide	23950-58-5	EPA6 e	2.57E-05	EPA6 e	8.13E-05	EPA6 e	9.84E-05	EPA6 e
Strychnine	57-24-9	EPA6 e	6.76E-07	EPA6 e	2.14E-06	EPA6 e	2.59E-06	EPA6 e
Terphenyls	26140-60-3	NA	NA	NA	NA	NA	NA	NA
Tributyl phosphate	126-73-8	EPA6 calc c	7.94E-05	EPA6 calc c	2.51E-04	EPA6 calc c	3.01E-04	EPA6 calc c
Trifluralin	1582-09-8	EPA6 calc c	1.74E-03	EPA6 calc c	5.50E-03	EPA6 calc c	6.59E-03	EPA6 calc c
Triphenylamine	603-34-9	NA	NA	NA	NA	NA	NA	NA
Herbicides and Organopesticides								
2,4,5-T	93-76-5	EPA6 calc c	1.82E-05	EPA6 calc c	5.75E-05	EPA6 calc c	6.91E-05	EPA6 calc c

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Bioconcentration / Bioaccumulation Factors					
Constituent of Potential Concern	CAS Registry Number	Source	Ba_{milk} (day/kg FW)	Source	Ba_{beef} (day/kg FW)	Source	Ba_{pork} (day/kg FW)	Source
2,4-D and esters	94-75-7	EPA6 calc c	5.13E-06	EPA6 calc c	1.62E-05	EPA6 calc c	1.95E-05	EPA6 calc c
4,4-DDD	72-54-8	EPA6 calc c	1.26E-02	EPA6 calc c	3.98E-02	EPA6 calc c	4.78E-02	EPA6 calc c
4,4-DDT	50-29-3	EPA6 calc c	7.94E-03	EPA6 calc c	2.51E-02	EPA6 calc c	3.01E-02	EPA6 calc c
Aldrin	309-00-2	EPA6 e	1.20E-02	EPA6 e	3.79E-02	EPA6 e	4.59E-02	EPA6 e
alpha-BHC	319-84-6	EPA6 e	5.00E-05	EPA6 e	1.58E-04	EPA6 e	1.92E-04	EPA6 e
beta-BHC	319-85-7	EPA6 e	5.41E-05	EPA6 e	1.71E-04	EPA6 e	2.07E-04	EPA6 e
Chlordane	57-74-9	EPA6 e	6.88E-03	EPA6 e	2.17E-02	EPA6 e	2.63E-02	EPA6 e
DDE	72-55-9	EPA6 e	1.43E-02	EPA6 e	4.53E-02	EPA6 e	5.49E-02	EPA6 e
Delta-BHC	319-86-8	EPA6 calc c	1.10E-04	EPA6 calc c	3.47E-04	EPA6 calc c	4.16E-04	EPA6 calc c
Dieldrin	60-57-1	EPA6 e	1.48E-03	EPA6 e	4.67E-03	EPA6 e	5.65E-03	EPA6 e
Endothall	145-73-3	EPA6 calc c	1.07E-09	EPA6 calc c	3.39E-09	EPA6 calc c	4.07E-09	EPA6 calc c
Endrin	72-20-8	EPA6 e	6.19E-04	EPA6 e	1.96E-03	EPA6 e	2.37E-03	EPA6 e
gamma-BHC(Lindane)	58-89-9	EPA6 calc c	4.17E-05	EPA6 calc c	1.32E-04	EPA6 calc c	1.58E-04	EPA6 calc c
Heptachlor	76-44-8	EPA6 e	8.22E-04	EPA6 e	2.60E-03	EPA6 e	3.15E-03	EPA6 e
Isodrin	465-73-6	NA	NA	NA	NA	NA	NA	NA
Methoxychlor	72-43-5	EPA6 e	2.67E-04	EPA6 e	8.43E-04	EPA6 e	1.02E-03	EPA6 e
Silvex (2,4,5-TP)	93-72-1	EPA6 calc c	9.38E-05	EPA6 calc c	2.97E-04	EPA6 calc c	3.56E-04	EPA6 calc c
Toxaphene	8001-35-2	EPA6 calc c	2.51E-03	EPA6 calc c	7.94E-03	EPA6 calc c	9.53E-03	EPA6 calc c

NA = Value not available
** = Varies with pH; values given for a pH of 7.0.
a = Illustrated Handbook of Physical - Chemical Properties and Environmental Fate i
b = RAIS database: http://risk.lsd.ornl.gov/rap_hp.shtml
c = Equations derived from EPA Region 6 Guidance Appendix A-3
d = Handbook of Environmental degradation Rates. 1991.
e = EPA Region 6 Hazardous Waste Combustion Guidance
f = "A Critical Review of Henry's Law Constants for Chemicals of Environmental Im
g = Handbook of Environmental Fate and Exposure Data for Organic Chemicals
h = http://www.epa.gov/region09/waste/sfund/prg/.
i = http://esc.syrres.com/efdb/Chemfate.htm
j = Hansch and Leo 1985 in Syracuse 1996

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

			Bioconcentration / Bioaccumulation Factors					
Constituent of Potential Concern	CAS Registry Number	Source	Ba_{milk} (day/kg FW)	Source	Ba_{beef} (day/kg FW)	Source	Ba_{pork} (day/kg FW)	Source

k = Schwarzenbach et al. 1993
l = Tomlin 1994 in Syracuse 1996
m = http://www.speclab.com/compound/c959988.htm

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors						
Constituent of Potential Concern	CAS Registry Number	Ba_{egg} (day/kg FW)	Source	$Ba_{chicken}$ (day/kg FW)	Source	BCF_{fish} (L/kg FW tissue)	Source	BAF_{fish} (L/kg FW)
Aromatic Halogenated Hydrocarbons								
2,3,4,6-Tetrachlorophenol	58-90-2	2.09E-01	EPA6 calc c	5.22E-04	EPA6 calc c	NA	NA	NA
4-Chloro-3-methylphenol	59-50-7	1.00E-02	EPA6 calc c	2.50E-05	EPA6 calc c	1.34E+02	EPA6 calc c	NA
Aromatic Nonhalogenated Hydrocarbons								
2-Nitrotoluene	88-72-2	1.58E-03	EPA6 calc c	3.96E-06	EPA6 calc c	3.30E+01	EPA6 calc c	NA
4-Nitrobiphenyl	92-93-3	4.68E-02	EPA6 calc c	1.17E-04	EPA6 calc c	4.32E+02	EPA6 calc c	NA
Benzaldehyde	100-52-7	2.38E-04	EPA6 e	5.95E-07	EPA6 e	7.81E+00	EPA6 e	NA
Benzene	71-43-2	1.09E-03	EPA6 e	2.72E-06	EPA6 e	2.48E+01	EPA6 e	NA
Benzyl alcohol	100-51-6	1.00E-04	EPA6 e	2.50E-07	EPA6 e	4.04E+00	EPA6 e	NA
Ethyl Benzene	100-41-4	1.05E-02	EPA6 e	2.63E-05	EPA6 e	1.39E+02	EPA6 e	NA
m-Xylene	108-38-3	1.26E-02	EPA6 e	3.15E-05	EPA6 e	1.60E+02	EPA6 e	NA
o-Xylene	95-47-6	1.07E-02	EPA6 e	2.68E-05	EPA6 e	1.41E+02	EPA6 e	NA
p-Xylene	106-42-3	1.18E-02	EPA6 e	2.93E-05	EPA6 e	1.51E+02	EPA6 e	NA
Styrene	100-42-5	6.74E-03	EPA6 e	1.68E-05	EPA6 e	9.91E+01	EPA6 e	NA
Toluene	108-88-3	3.69E-03	EPA6 e	9.22E-06	EPA6 e	6.27E+01	EPA6 e	NA
Non-aromatic Nonhalogenated Hydrocarbons								
1,2-Epoxybutane	106-88-7	2.19E-04	EPA6 calc c	5.48E-07	EPA6 calc c	7.34E+00	EPA6 calc c	NA
1,3-Butadiene	106-99-0	7.76E-04	EPA6 calc c	1.94E-06	EPA6 calc c	1.92E+01	EPA6 calc c	NA
1,4-Dioxane	123-91-1	4.29E-06	EPA6 e	1.07E-08	EPA6 e	3.69E-01	EPA6 e	NA
1-Methylpropyl alcohol	78-92-2	3.24E-05	EPA6 calc c	8.08E-08	EPA6 calc c	1.71E+00	EPA6 calc c	NA
1-Nitropropane	108-03-2	5.89E-05	EPA6 calc c	1.47E-07	EPA6 calc c	2.70E+00	EPA6 calc c	NA
2,2,4-Trimethlylpentane	540-84-1	8.32E-01	EPA6 calc c	2.08E-03	EPA6 calc c	NA	EPA6 calc c	NA
2-Butanone	78-93-3	1.51E-05	EPA6 e	3.78E-08	EPA6 e	9.61E-01	EPA6 e	NA
2-Butenaldehyde (2-Butenal)	4170-30-3	NA	NA	NA	NA	NA	NA	NA
2-Ethoxyethanol	110-80-5	6.31E-06	EPA6 calc c	1.58E-08	EPA6 calc c	4.94E-01	EPA6 calc c	NA
2-Heptanone	110-43-0	7.59E-04	EPA6 calc c	1.89E-06	EPA6 calc c	1.88E+01	EPA6 calc c	NA
2-Hexanone	591-78-6	1.91E-04	EPA6 calc c	4.76E-07	EPA6 calc c	6.59E+00	EPA6 calc c	NA
2-Methoxyethanol	109-86-4	1.42E-05	EPA6 calc c	3.54E-08	EPA6 calc c	9.16E-01	EPA6 calc c	NA
2-Methyl-2-propanol	75-65-0	1.78E-05	EPA6 calc c	4.44E-08	EPA6 calc c	1.09E+00	EPA6 calc c	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors						
Constituent of Potential Concern	CAS Registry Number	Ba_{egg} (day/kg FW)	Source	$Ba_{chicken}$ (day/kg FW)	Source	BCF_{fish} (L/kg FW tissue)	Source	BAF_{fish} (L/kg FW)
2-Methyl-2-propenenitrile	126-98-7	2.76E-05	EPA6 e	6.88E-08	EPA6 e	1.52E+00	EPA6 e	NA
2-Methylaziridine	75-55-8	NA	NA	NA	NA	NA	NA	NA
2-Methylpropyl alcohol	78-83-1	4.57E-05	EPA6 calc c	1.14E-07	EPA6 calc c	2.23E+00	EPA6 calc c	NA
2-Pentanone	107-87-9	6.46E-05	EPA6 calc c	1.61E-07	EPA6 calc c	2.89E+00	EPA6 calc c	NA
2-Propanone (Acetone)	67-64-1	4.77E-06	EPA6 e	1.19E-08	EPA6 e	4.00E-01	EPA6 e	NA
2-Propene-1-ol	107-18-6	1.17E-05	EPA6 calc c	2.93E-08	EPA6 calc c	7.93E-01	EPA6 calc c	NA
2-Propyl alcohol	67-63-0	8.91E-06	EPA6 calc c	2.23E-08	EPA6 calc c	6.43E-01	EPA6 calc c	NA
3-Heptanone	106-35-4	NA	NA	NA	NA	NA	NA	NA
3-Methyl-1-butanol	123-51-3	NA	NA	NA	NA	NA	NA	NA
3-Methyl-2-butanone	563-80-4	NA	NA	NA	NA	NA	NA	NA
3-Pentanone	96-22-0	7.76E-05	EPA6 calc c	1.94E-07	EPA6 calc c	3.33E+00	EPA6 calc c	NA
4-Heptanone	123-19-3	NA	NA	NA	NA	NA	NA	NA
4-Methyl-2-pentanone	108-10-1	1.23E-04	EPA6 e	3.07E-07	EPA6 e	4.73E+00	EPA6 e	NA
4-Methyl-3-penten-2-one	141-79-7	NA	NA	NA	NA	NA	NA	NA
5-Methyl-2-hexanone	110-12-3	NA	NA	NA	NA	NA	NA	NA
Acetaldehyde	75-07-0	4.78E-06	EPA6 e	1.19E-08	EPA6 e	4.00E-01	EPA6 e	NA
Acetamide	60-35-5	4.37E-07	EPA6 calc c	1.09E-09	EPA6 calc c	6.49E-02	EPA6 calc c	NA
Acetic acid	64-19-7	5.37E-06	EPA6 calc c	1.34E-08	EPA6 calc c	4.37E-01	EPA6 calc c	NA
Acetic acid ethyl ester	141-78-6	4.27E-05	EPA6 calc c	1.06E-07	EPA6 calc c	2.11E+00	EPA6 calc c	NA
Acetic acid n-butyl ester	123-86-4	4.27E-04	EPA6 calc c	1.06E-06	EPA6 calc c	1.22E+01	EPA6 calc c	NA
Acetonitrile	75-05-8	3.63E-06	EPA6 e	9.06E-09	EPA6 e	3.25E-01	EPA6 e	NA
Acrolein	107-02-8	7.78E-06	EPA6 e	1.94E-08	EPA6 e	5.80E-01	EPA6 e	NA
Acrylonitrile	107-13-1	1.41E-05	EPA6 e	3.53E-08	EPA6 e	4.80E+01	EPA6 e	NA
Bis(isopropyl)ether	108-20-3	NA	NA	NA	NA	NA	NA	NA
Butane	106-97-8	6.17E-03	EPA6 calc c	1.54E-05	EPA6 calc c	9.26E+01	EPA6 calc c	NA
Carbon disulfide	75-15-0	7.94E-04	EPA6 e	1.98E-06	EPA6 e	1.95E+01	EPA6 e	NA
Cyanogen	460-19-5	5.10E-05	EPA6 calc c	1.27E-07	EPA6 calc c	2.42E+00	EPA6 calc c	NA
Cyclohexane	110-82-7	2.19E-02	EPA6 calc c	5.46E-05	EPA6 calc c	2.42E+02	EPA6 calc c	NA
Cyclohexanone	108-94-1	5.13E-05	EPA6 calc c	1.28E-07	EPA6 calc c	2.43E+00	EPA6 calc c	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors						
Constituent of Potential Concern	CAS Registry Number	Ba_{egg} (day/kg FW)	Source	$Ba_{chicken}$ (day/kg FW)	Source	BCF_{fish} (L/kg FW tissue)	Source	BAF_{fish} (L/kg FW)
Cyclohexene	110-83-8	5.75E-03	EPA6 calc c	1.44E-05	EPA6 calc c	8.78E+01	EPA6 calc c	NA
Cyclopentane	287-92-3	7.94E-03	EPA6 calc c	1.98E-05	EPA6 calc c	1.12E+02	EPA6 calc c	NA
Ethyl alcohol	64-17-5	3.89E-06	EPA6 calc c	9.71E-09	EPA6 calc c	3.42E-01	EPA6 calc c	NA
Ethyl ether	60-29-7	6.17E-05	EPA6 calc c	1.54E-07	EPA6 calc c	2.80E+00	EPA6 calc c	NA
Ethyl methacrylate	97-63-2	3.09E-04	EPA6 e	7.71E-07	EPA6 e	9.51E+00	EPA6 e	NA
Formaldehyde	50-00-0	1.75E-05	EPA6 e	4.36E-08	EPA6 e	1.07E+00	EPA6 e	NA
Formamide	75-12-7	2.45E-07	EPA6 calc c	6.13E-10	EPA6 calc c	4.19E-02	EPA6 calc c	NA
Formic acid	64-18-6	2.30E-06	EPA6 e	5.75E-09	EPA6 e	2.30E-01	EPA6 e	NA
Formic acid, methyl ester	107-31-3	4.33E-06	EPA6 calc c	1.08E-08	EPA6 calc c	3.71E-01	EPA6 calc c	NA
Glycidylaldehyde	765-34-4	1.48E-06	EPA6 calc c	3.69E-09	EPA6 calc c	1.64E-01	EPA6 calc c	NA
Methyl acetate	79-20-9	1.20E-05	EPA6 calc c	3.00E-08	EPA6 calc c	8.07E-01	EPA6 calc c	NA
Methyl alcohol	67-56-1	1.55E-06	EPA6 e	3.39E-09	EPA6 e	1.70E-01	EPA6 e	NA
Methyl isocyanate	624-83-9	NA	NA	NA	NA	NA	NA	NA
Methyl methacrylate	80-62-6	4.90E-05	EPA6 calc c	1.22E-07	EPA6 calc c	2.35E+00	EPA6 calc c	NA
Methyl tert-butyl ether	1634-04-4	6.92E-05	EPA6 calc c	1.73E-07	EPA6 calc c	3.05E+00	EPA6 calc c	NA
Methylacetylene	74-99-7	6.92E-05	EPA6 calc c	1.73E-07	EPA6 calc c	3.05E+00	EPA6 calc c	NA
Methylcyclohexane	108-87-2	1.00E-01	EPA6 calc c	2.50E-04	EPA6 calc c	NA	NA	NA
N,N-Dimethylacetamide	127-19-5	NA	NA	NA	NA	NA	NA	NA
n-Butyl alcohol	71-36-3	6.03E-05	EPA6 calc c	1.50E-07	EPA6 calc c	2.75E+00	EPA6 calc c	NA
n-Heptane	142-82-5	3.63E-01	EPA6 calc c	9.06E-04	EPA6 calc c	NA	NA	NA
n-Hexane	110-54-3	1.02E-01	EPA6 calc c	2.55E-04	EPA6 calc c	NA	NA	NA
Nitromethane	75-52-5	3.55E-06	EPA6 calc c	8.86E-09	EPA6 calc c	NA	NA	NA
n-Nonane	111-84-2	3.55E+00	EPA6 calc c	8.86E-03	EPA6 calc c	NA	NA	NA
n-Octane	111-65-9	7.94E-02	EPA6 calc c	1.98E-04	EPA6 calc c	NA	NA	NA
n-Pentane	109-66-0	1.29E-02	EPA6 calc c	3.22E-05	EPA6 calc c	1.62E+02	EPA6 calc c	NA
n-Propionaldehyde	123-38-6	3.09E-05	EPA6 calc c	7.72E-08	EPA6 calc c	1.65E+00	EPA6 calc c	NA
n-Propyl alcohol	71-23-8	NA	NA	NA	NA	NA	NA	NA
n-Valeraldehyde	110-62-3	NA	NA	NA	NA	NA	NA	NA
Oxirane	75-21-8	3.98E-06	EPA6 e	9.94E-09	EPA6 e	3.48E-01	EPA6 e	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors						
Constituent of Potential Concern	CAS Registry Number	Ba_{egg} (day/kg FW)	Source	$Ba_{chicken}$ (day/kg FW)	Source	BCF_{fish} (L/kg FW tissue)	Source	BAF_{fish} (L/kg FW)
p-Cymene	99-87-6	1.00E-01	EPA6 calc c	2.50E-04	EPA6 calc c	NA	NA	NA
Phosgene	75-44-5	NA	NA	NA	NA	NA	NA	NA
Propargyl alcohol	107-19-7	NA	NA	NA	NA	NA	NA	NA
Propionic acid	79-09-4	1.70E-05	EPA6 calc c	4.24E-08	EPA6 calc c	1.05E+00	EPA6 calc c	NA
Propionitrile	107-12-0	1.15E-05	EPA6 calc c	2.87E-08	EPA6 calc c	7.79E-01	EPA6 calc c	NA
Propylene glycol monomethyl ether	107-98-2	5.25E-06	EPA6 calc c	1.31E-08	EPA6 calc c	4.30E-01	EPA6 calc c	NA
p-tert-Butyltoluene	98-51-1	NA	NA	NA	NA	NA	NA	NA
Triethylamine	121-44-8	NA	NA	NA	NA	NA	NA	NA
Trimethylamine	75-50-3	1.15E-05	EPA6 calc c	2.87E-08	EPA6 calc c	7.79E-01	EPA6 calc c	NA
Vinyl acetate	108-05-4	3.97E-05	EPA6 e	9.92E-08	EPA6 e	2.00E+00	EPA6 e	NA
Non-aromatic Halogenated Hydrocarbons								
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	NA	NA	NA	NA	NA	NA	NA
1,1,1,2-Tetrachloroethane	630-20-6	3.39E-03	EPA6 e	8.46E-06	EPA6 e	5.87E+01	EPA6 e	NA
1,1,1-Trichloroethane	71-55-6	2.10E-03	EPA6 e	5.24E-06	EPA6 e	4.08E+01	EPA6 e	NA
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	4.27E-02	EPA6 calc c	1.06E-04	EPA6 calc c	4.03E+02	EPA6 calc c	NA
1,1,2,2-Tetrachloroethane	79-34-5	3.50E-01	EPA6 e	8.73E-04	EPA6 e	NA	EPA6 e	4.33E+03
1,1,2,2-Tetrachloroethene	127-18-4	2.79E-03	EPA6 e	6.96E-06	EPA6 e	5.06E+01	EPA6 e	NA
1,1,2-Trichloro-1,2,2-trifluoroethane	76-13-1	1.15E-02	EPA6 calc c	2.87E-05	EPA6 calc c	1.48E+02	EPA6 calc c	NA
1,1,2-Trichloroethane	79-00-5	9.93E-04	EPA6 e	2.48E-06	EPA6 e	2.31E+01	EPA6 e	NA
1,1,2-Trichloroethylene	79-01-6	2.15E-03	EPA6 e	5.37E-06	EPA6 e	4.16E+01	EPA6 e	NA
1,1-Dichloroethane	75-34-3	4.92E-04	EPA6 e	1.23E-06	EPA6 e	1.36E+01	EPA6 e	NA
1,1-Dichloroethene	75-35-4	1.05E-03	EPA6 e	2.62E-06	EPA6 e	2.41E+01	EPA6 e	NA
1,2,3-Trichloropropane	96-18-4	1.41E-03	EPA6 e	3.53E-06	EPA6 e	3.02E+01	EPA6 e	NA
1,2-dibromo-3-chloropropane	96-12-8	1.74E-03	EPA6 e	4.34E-06	EPA6 e	3.54E+01	EPA6 e	NA
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	NA	NA	NA	NA	NA	NA	NA
1,2-Dichloroethane	107-06-2	2.30E-04	EPA6 e	5.75E-07	EPA6 e	7.61E+00	EPA6 e	NA
1,2-Dichloroethylene	540-59-0	2.40E-05	EPA6 calc c	5.99E-08	EPA6 calc c	1.36E+00	EPA6 calc c	NA
1,2-Dichloropropane	78-87-5	1.41E-03	EPA6 e	3.53E-06	EPA6 e	3.02E+01	EPA6 e	NA
1,3-Dichloropropene	542-75-6	4.45E-04	EPA6 e	1.11E-06	EPA6 e	1.25E+01	EPA6 e	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors						
Constituent of Potential Concern	CAS Registry Number	Ba_{egg} (day/kg FW)	Source	$Ba_{chicken}$ (day/kg FW)	Source	BCF_{fish} (L/kg FW tissue)	Source	BAF_{fish} (L/kg FW)
1,4-Dichloro-2-butene	764-41-0	5.92E-05	EPA6 calc c	1.48E-07	EPA6 calc c	2.71E+00	EPA6 calc c	NA
1-Chloroethene	75-01-4	1.11E-04	EPA6 e	2.78E-07	EPA6 e	4.37E+00	EPA6 e	NA
2,2-Dichloropropionic acid	75-99-0	4.76E-05	EPA6 calc c	1.19E-07	EPA6 calc c	2.30E+00	EPA6 calc c	NA
2-Chloropropane	75-29-6	6.31E-04	EPA6 calc c	1.58E-06	EPA6 calc c	1.64E+01	EPA6 calc c	NA
3-Chloropropene (allyl chloride)	107-05-1	7.15E-05	EPA6 calc c	1.79E-07	EPA6 calc c	3.13E+00	EPA6 calc c	NA
Bromochloromethane	74-97-5	2.04E-04	EPA6 calc c	5.10E-07	EPA6 calc c	6.94E+00	EPA6 calc c	NA
Bromodichloromethane	75-27-4	8.42E-04	EPA6 e	2.10E-06	EPA6 e	2.04E+01	EPA6 e	NA
Bromoethene	593-60-2	9.26E-05	EPA6 calc c	2.31E-07	EPA6 calc c	3.81E+00	EPA6 calc c	NA
Bromoform	75-25-2	1.78E-03	EPA6 e	4.44E-06	EPA6 e	3.60E+01	EPA6 e	NA
Bromomethane	74-83-9	1.03E-04	EPA6 e	2.58E-07	EPA6 e	4.14E+00	EPA6 e	NA
Carbon tetrachloride	56-23-5	4.14E-03	EPA6 e	1.05E-05	EPA6 e	3.00E+01	EPA6 e	NA
Chlordibromomethane	124-48-1	1.19E-03	EPA6 e	2.97E-06	EPA6 e	2.65E+01	EPA6 e	NA
Chlorodifluoromethane	75-45-6	9.53E-05	EPA6 e	2.38E-07	EPA6 e	3.89E+00	EPA6 e	NA
Chloroethane	75-00-3	1.00E-02	EPA6 e	2.50E-05	EPA6 e	1.34E+02	EPA6 e	NA
Chloroform	67-66-3	7.07E-04	EPA6 e	1.76E-06	EPA6 e	3.59E+00	EPA6 e	NA
Chloromethane	74-87-3	6.35E-05	EPA6 e	1.59E-07	EPA6 e	2.86E+00	EPA6 e	NA
Chloropentafluoroethane	76-15-3	NA	NA	NA	NA	NA	NA	NA
cis-1,2-Dichloroethene	156-59-2	7.63E-04	EPA6 e	1.90E-06	EPA6 e	1.89E+01	EPA6 e	NA
cis-1,3-Dichloropropene	10061-01-5	NA	NA	NA	NA	NA	NA	NA
Cyanogen bromide	506-68-3	NA	NA	NA	NA	NA	NA	NA
Cyanogen chloride	506-77-4	NA	NA	NA	NA	NA	NA	NA
Dichlorodifluoromethane	75-71-8	1.15E-03	EPA6 e	2.87E-06	EPA6 e	2.58E+01	EPA6 e	NA
Dichlorofluoromethane	75-43-4	NA	NA	NA	NA	NA	NA	NA
Dichloromethane	75-09-2	1.43E-04	EPA6 e	3.57E-07	EPA6 e	5.30E+00	EPA6 e	NA
Difluorodibromomethane	75-61-6	NA	NA	NA	NA	NA	NA	NA
Hexafluoroacetone	684-16-2	NA	NA	NA	NA	NA	NA	NA
Iodomethane	74-88-4	3.89E-04	EPA6 calc c	9.71E-07	EPA6 calc c	1.13E+01	EPA6 calc c	NA
Methylene bromide	74-95-3	3.31E-04	EPA6 e	8.27E-07	EPA6 e	1.00E+01	EPA6 e	NA
Pentachloroethane	76-01-7	8.91E-03	EPA6 calc c	2.23E-05	EPA6 calc c	1.22E+02	EPA6 calc c	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors						
Constituent of Potential Concern	CAS Registry Number	Ba_{egg} (day/kg FW)	Source	$Ba_{chicken}$ (day/kg FW)	Source	BCF_{fish} (L/kg FW tissue)	Source	BAF_{fish} (L/kg FW)
trans-1,2-Dichloroethene	156-60-5	7.63E-04	EPA6 e	1.90E-06	EPA6 e	1.89E+01	EPA6 e	NA
trans-1,3-Dichloropropene	10061-02-6	9.12E-04	EPA6 calc c	2.28E-06	EPA6 calc c	2.17E+01	EPA6 calc c	NA
Trichloroacetic acid	76-03-9	1.70E-04	EPA6 calc c	4.24E-07	EPA6 calc c	6.04E+00	EPA6 calc c	NA
Trichlorofluoroethane	27154-33-2	NA	NA	NA	NA	NA	NA	NA
Trichlorofluoromethane	75-69-4	2.70E-03	EPA6 e	6.74E-06	EPA6 e	4.94E+01	EPA6 e	NA
Trifluorobromomethane	75-63-8	5.75E-04	EPA6 calc c	1.44E-06	EPA6 calc c	1.53E+01	EPA6 calc c	NA
Dioxin and Furan Compounds								
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	35822-46-9	2.55E-02	EPA6 e	8.58E-03	EPA6 e	NA	EPA6 e	NA
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	2.09E-02	EPA6 e	7.04E-03	EPA6 e	NA	EPA6 e	NA
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	2.42E-02	EPA6 e	1.06E-02	EPA6 e	NA	EPA6 e	NA
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	39227-28-6	4.53E-02	EPA6 e	4.03E-02	EPA6 e	NA	EPA6 e	NA
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	4.51E-02	EPA6 e	3.48E-02	EPA6 e	NA	EPA6 e	NA
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	57653-85-7	3.70E-02	EPA6 e	2.57E-02	EPA6 e	NA	EPA6 e	NA
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	4.53E-02	EPA6 e	3.56E-02	EPA6 e	NA	EPA6 e	NA
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	19408-74-3	2.33E-02	EPA6 e	1.39E-02	EPA6 e	NA	EPA6 e	NA
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	NA	EPA6 e	NA	EPA6 e	NA	EPA6 e	NA
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	40321-76-4	4.71E-02	EPA6 e	5.50E-02	EPA6 e	NA	EPA6 e	NA
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	NA	EPA6 e	NA	EPA6 e	NA	EPA6 e	NA
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	2.11E-02	EPA6 e	1.74E-02	EPA6 e	NA	EPA6 e	NA
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	5.61E-02	EPA6 e	7.32E-02	EPA6 e	NA	EPA6 e	NA
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1746-01-6	5.42E-02	EPA6 e	7.30E-02	EPA6 e	NA	EPA6 e	NA
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	3.61E-02	EPA6 e	5.63E-02	EPA6 e	NA	EPA6 e	NA
Dibenzofuran	132-64-9	1.70E-01	EPA6 calc c	4.24E-04	EPA6 calc c	NA	NA	NA
Octachlorodibenzo(p)dioxin	3268-87-9	9.90E-03	EPA6 e	1.10E-03	EPA6 e	NA	EPA6 e	NA
Octachlorodibenzofuran	39001-02-0	7.92E-03	EPA6 e	4.40E-04	EPA6 e	NA	EPA6 e	NA
PCBs								
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	9.55E+01	EPA6 calc c	2.38E-01	EPA6 calc c	NA	NA	NA
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	1.05E+02	EPA6 calc c	2.61E-01	EPA6 calc c	NA	NA	NA
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	NA	NA	NA	NA	NA	NA	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors						
Constituent of Potential Concern	CAS Registry Number	Ba_{egg} (day/kg FW)	Source	$Ba_{chicken}$ (day/kg FW)	Source	BCF_{fish} (L/kg FW tissue)	Source	BAF_{fish} (L/kg FW)
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA	NA
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	NA	NA	NA	NA	NA	NA	NA
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	NA	NA	NA	NA	NA	NA	NA
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA	NA
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	NA	NA	NA	NA	NA	NA	NA
2',3,4,4',5-Pentachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA	NA
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	1.05E+02	EPA6 calc c	2.61E-01	EPA6 calc c	NA	NA	NA
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	2.03E+02	EPA6 calc c	5.07E-01	EPA6 calc c	NA	NA	NA
3,3',4,4',5-Pentachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA	NA
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	NA	NA	NA	NA	NA	NA	NA
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	NA	NA	NA	NA	NA	NA	NA
Polychlorinated biphenyls (PCBs)**	1336-36-3	1.62E+02	EPA6 calc c	4.04E-01	EPA6 calc c	NA	NA	2.10E+04
Phthalates								
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	1.27E+00	EPA6 e	3.18E-03	EPA6 e	NA	EPA6 e	3.60E+02
Butylbenzyl phthalate	85-68-7	2.06E-01	EPA6 e	5.13E-04	EPA6 e	NA	EPA6 e	2.35E+03
Dibutyl phthalate	84-74-2	4.17E-01	EPA6 e	1.04E-03	EPA6 e	NA	EPA6 e	5.58E+03
Diethyl phthalate	84-66-2	2.17E-01	EPA6 e	5.42E-04	EPA6 e	NA	EPA6 e	2.45E+03
Dimethyl phthalate	131-11-3	3.42E-04	EPA6 e	8.53E-07	EPA6 e	1.03E+01	EPA6 e	NA
n-Dioctyl phthalate	117-84-0	1.70E+04	EPA6 e	4.24E+01	EPA6 e	NA	EPA6 e	3.88E+03
Light Polycyclic Aromatic Hydrocarbons								
2-Chloronaphthalene	91-58-7	9.33E-02	EPA6 e	2.33E-04	EPA6 e	NA	EPA6 e	9.60E+02
2-Methyl naphthalene	91-57-6	5.75E-02	EPA6 calc c	1.44E-04	EPA6 calc c	5.05E+02	EPA6 calc c	NA
5-Nitroacenaphthene	602-87-9	NA	NA	NA	NA	NA	NA	NA
Acenaphthene	83-32-9	7.32E-02	EPA6 e	1.83E-04	EPA6 e	6.07E+02	EPA6 e	NA
Acenaphthylene	208-96-8	9.33E-02	EPA6 calc c	2.33E-04	EPA6 calc c	NA	NA	NA
Anthracene	120-12-7	2.34E-01	EPA6 e	5.85E-04	EPA6 e	NA	EPA6 e	2.60E+03
Fluorene	86-73-7	1.20E-01	EPA6 calc c	3.00E-04	EPA6 calc c	NA	NA	NA
Indene	95-13-6	NA	NA	NA	NA	NA	NA	NA
Napthalene	91-20-3	1.87E-02	EPA6 e	4.67E-05	EPA6 e	2.15E+02	EPA6 e	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors						
Constituent of Potential Concern	CAS Registry Number	Ba_{egg} (day/kg FW)	Source	$Ba_{chicken}$ (day/kg FW)	Source	BCF_{fish} (L/kg FW tissue)	Source	BAF_{fish} (L/kg FW)
Phenanthrene	85-01-8	2.82E-01	EPA6 e	7.04E-04	EPA6 e	NA	EPA6 e	3.30E+03
Pyrene	129-00-0	7.98E-01	EPA6 e	1.99E-03	EPA6 e	NA	EPA6 e	1.19E+04
Heavy Polycyclic Aromatic Hydrocarbons								
3-Methylcholanthrene	56-49-5	1.02E+02	EPA6 calc c	2.55E-01	EPA6 calc c	NA	NA	NA
5-Methylchrysene	3697-24-3	NA	NA	NA	NA	NA	NA	NA
Benzo(a)anthracene	56-55-3	3.79E+00	EPA6 e	9.46E-03	EPA6 e	NA	EPA6 e	5.10E+03
Benzo(a)pyrene	50-32-8	1.07E+01	EPA6 e	2.67E-02	EPA6 e	NA	EPA6 e	9.95E+03
Benzo(b)fluoranthene	205-99-2	1.27E+01	EPA6 e	3.16E-02	EPA6 e	NA	EPA6 e	9.95E+03
Benzo(e)pyrene	192-97-2	2.00E+02	EPA6 calc c	4.98E-01	EPA6 calc c	NA	NA	NA
Benzo(g,h,i)perylene	191-24-2	1.00E+02	EPA6 calc c	2.50E-01	EPA6 calc c	NA	NA	NA
Benzo(j)fluoranthene	205-82-3	2.19E+01	EPA6 calc c	5.46E-02	EPA6 calc c	NA	NA	NA
Benzo(k)fluoranthene	207-08-9	1.26E+01	EPA6 e	3.14E-02	EPA6 e	NA	EPA6 e	9.95E+03
Benzo[a,i]pyrene	191-30-0	NA	NA	NA	NA	NA	NA	NA
Chrysene	218-01-9	4.35E+00	EPA6 e	1.09E-02	EPA6 e	NA	EPA6 e	6.03E+03
Dibenz(a,h)anthracene	53-70-3	2.80E+01	EPA6 e	7.00E-02	EPA6 e	NA	EPA6 e	1.28E+04
Dibenz[a,h]acridine	226-36-8	NA	NA	NA	NA	NA	NA	NA
Dibenz[a,j]acridine	224-42-0	NA	NA	NA	NA	NA	NA	NA
Dibenzo(a,e)fluoranthene	5385-75-1	NA	NA	NA	NA	NA	NA	NA
Dibenzo(a,h)fluoranthene	no cas #	NA	NA	NA	NA	NA	NA	NA
Dibenzo[a,e]pyrene	192-65-4	NA	NA	NA	NA	NA	NA	NA
Dibenzo[a,h]pyrene	189-64-0	NA	NA	NA	NA	NA	NA	NA
Dibenzo[a,i]pyrene	189-55-9	1.55E+02	EPA6 calc c	3.87E-01	EPA6 calc c	NA	NA	NA
Fluoranthene	206-44-0	9.65E-01	EPA6 e	2.41E-03	EPA6 e	NA	EPA6 e	1.57E+04
Hexachloronaphthalene	1335-87-1	3.09E+02	EPA6 calc c	7.72E-01	EPA6 calc c	NA	NA	NA
Indeno(1,2,3-cd)pyrene	193-39-5	6.53E+01	EPA6 e	1.63E-01	EPA6 e	NA	EPA6 e	1.31E+04
Octachloronaphthalene	2234-13-1	2.09E+01	EPA6 calc c	5.22E-02	EPA6 calc c	NA	NA	NA
Pentachloronaphthalene	1321-64-8	NA	NA	NA	NA	NA	NA	NA
Tetrachloronaphthalene	1335-88-2	NA	NA	NA	NA	NA	NA	NA
Trichloronaphthalene	1321-65-9	NA	NA	NA	NA	NA	NA	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors						
Constituent of Potential Concern	CAS Registry Number	Ba_{egg} (day/kg FW)	Source	$Ba_{chicken}$ (day/kg FW)	Source	BCF_{fish} (L/kg FW tissue)	Source	BAF_{fish} (L/kg FW)
Light Substituted Benzene Compounds								
1,2,3-Trichlorobenzene	87-61-6	8.82E-02	EPA6 e	2.20E-04	EPA6 e	NA	EPA6 e	8.76E+02
1,2,4-Trichlorobenzene	120-82-1	7.73E-02	EPA6 e	1.93E-04	EPA6 e	6.33E+02	EPA6 e	NA
1,2,4-Trimethlyl benzene	25551-13-7	2.00E-02	EPA6 calc c	4.98E-05	EPA6 calc c	2.26E+02	EPA6 calc c	NA
1,2,4-Trimethyl benzene	95-63-6	3.55E-02	EPA6 calc c	8.86E-05	EPA6 calc c	3.50E+02	EPA6 calc c	NA
1,2-Dichlorobenzene	95-50-1	2.21E-02	EPA6 e	5.53E-05	EPA6 e	2.45E+02	EPA6 e	NA
1,3,5-Trimethyl benzene	108-67-8	2.09E-02	EPA6 e	5.22E-05	EPA6 e	2.34E+02	EPA6 e	NA
1,3-Dichlorobenzene	541-73-1	2.69E-02	EPA6 e	6.72E-05	EPA6 e	2.84E+02	EPA6 e	NA
1,3-Dinitrobenzene	99-65-0	2.46E-04	EPA6 e	6.15E-07	EPA6 e	7.40E+01	EPA6 e	NA
1,4-Dichlorobenzene	106-46-7	2.05E-02	EPA6 e	5.12E-05	EPA6 e	2.31E+02	EPA6 e	NA
1,4-Dinitrobenzene	100-25-4	2.51E-04	EPA6 calc c	6.27E-07	EPA6 calc c	8.13E+00	EPA6 calc c	NA
2,4,5-Trichorophenol	95-95-4	5.89E-02	EPA6 e	1.47E-04	EPA6 e	5.14E+02	EPA6 e	NA
2,4,6-Trichlorophenol	88-06-2	4.09E-02	EPA6 e	1.02E-04	EPA6 e	3.90E+02	EPA6 e	NA
2,4-Dichlorophenol	120-83-2	8.62E-03	EPA6 e	2.15E-05	EPA6 e	1.19E+02	EPA6 e	NA
2,4-Dimethylphenol	105-67-9	1.82E-03	EPA6 e	4.54E-06	EPA6 e	3.66E+01	EPA6 e	NA
2,4-Dinitrophenol	51-28-5	2.62E-04	EPA6 e	6.54E-07	EPA6 e	8.40E+00	EPA6 e	NA
2,4-Dinitrotoluene	121-14-2	7.86E-04	EPA6 e	1.96E-06	EPA6 e	5.92E+00	EPA6 e	NA
2,6-Dinitrotoluene	606-20-2	6.12E-04	EPA6 e	1.53E-06	EPA6 e	5.92E+00	EPA6 e	NA
2-Chlorotoluene	95-49-8	2.73E-02	EPA6 calc c	6.82E-05	EPA6 calc c	2.87E+02	EPA6 calc c	NA
2-Chorophenol	95-57-8	1.15E-03	EPA6 e	2.88E-06	EPA6 e	2.59E+01	EPA6 e	NA
2-Nitrophenol	88-75-5	4.90E-04	EPA6 e	1.22E-06	EPA6 e	1.35E+01	EPA6 e	NA
4,6-Dinitro-o-cresol	534-52-1	5.62E-03	EPA6 calc c	1.40E-05	EPA6 calc c	8.63E+01	EPA6 calc c	NA
4-Chlorotoluene	106-43-4	1.70E-02	EPA6 calc c	4.24E-05	EPA6 calc c	2.00E+02	EPA6 calc c	NA
4-Nitrophenol	100-02-7	6.46E-04	EPA6 e	1.61E-06	EPA6 e	1.67E+01	EPA6 e	NA
alpha-Methylstyrene	98-83-9	2.31E-02	EPA6 calc c	5.77E-05	EPA6 calc c	2.53E+02	EPA6 calc c	NA
Aniline	62-53-3	7.59E-05	EPA6 e	1.89E-07	EPA6 e	3.27E+00	EPA6 e	NA
Benzotrichloride	98-07-7	6.61E-03	EPA6 calc c	1.65E-05	EPA6 calc c	9.75E+01	EPA6 calc c	NA
Benzyl Chloride	100-44-7	1.83E-05	EPA6 e	4.56E-08	EPA6 e	1.11E+00	EPA6 e	NA
Bromobenzene	108-86-1	7.76E-03	EPA6 calc c	1.94E-05	EPA6 calc c	1.10E+02	EPA6 calc c	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors						
Constituent of Potential Concern	CAS Registry Number	Ba_{egg} (day/kg FW)	Source	$Ba_{chicken}$ (day/kg FW)	Source	BCF_{fish} (L/kg FW tissue)	Source	BAF_{fish} (L/kg FW)
Chlorobenzene	108-90-7	4.89E-03	EPA6 e	1.22E-05	EPA6 e	7.76E+01	EPA6 e	NA
Cumene	98-82-8	3.26E-02	EPA6 e	8.13E-05	EPA6 e	3.28E+02	EPA6 e	NA
m-Cresol	108-39-4	7.23E-04	EPA6 e	1.86E-06	EPA6 e	1.81E+01	EPA6 e	NA
n-Butyl benzene	104-51-8	1.51E-01	EPA6 calc c	3.78E-04	EPA6 calc c	NA	NA	NA
Nitrobenzene	98-95-3	5.40E-04	EPA6 e	1.35E-06	EPA6 e	5.92E+00	EPA6 e	NA
n-Propyl benzene	103-65-1	3.89E-02	EPA6 calc c	9.71E-05	EPA6 calc c	3.75E+02	EPA6 calc c	NA
o-Cresol	95-48-7	8.34E-04	EPA6 e	2.08E-06	EPA6 e	2.02E+01	EPA6 e	NA
o-Dinitrobenzene	528-29-0	3.16E-04	EPA6 calc c	7.89E-07	EPA6 calc c	9.68E+00	EPA6 calc c	NA
o-Nitroaniline	88-74-4	5.62E-04	EPA6 e	1.40E-06	EPA6 e	1.50E+01	EPA6 e	NA
o-Toluidine	95-53-4	1.74E-04	EPA6 e	4.34E-07	EPA6 e	6.14E+00	EPA6 e	NA
p-Chloroaniline	106-47-8	5.88E-04	EPA6 e	1.47E-06	EPA6 e	1.55E+01	EPA6 e	NA
p-Cresol	106-44-5	6.91E-04	EPA6 e	1.73E-06	EPA6 e	1.75E+01	EPA6 e	NA
Phenol	108-95-2	2.38E-04	EPA6 e	5.95E-07	EPA6 e	7.81E+00	EPA6 e	NA
p-Nitrochlorobenzene	100-00-5	1.95E-03	EPA6 calc c	4.87E-06	EPA6 calc c	3.86E+01	EPA6 calc c	NA
p-Toluidine	106-49-0	2.00E-04	EPA6 calc c	4.98E-07	EPA6 calc c	6.82E+00	EPA6 calc c	NA
sec-Butyl benzene	135-98-8	NA	NA	NA	NA	NA	NA	NA
tert-Butyl benzene	98-06-6	1.02E-01	EPA6 calc c	2.55E-04	EPA6 calc c	NA	NA	NA
Toluene-2,6-diamine	823-40-5	2.26E-04	EPA6 calc c	5.65E-07	EPA6 calc c	7.51E+00	EPA6 calc c	NA
Other Light Semivolatile Compounds								
1,1 Dimethyl hydrazine	57-14-7	3.13E-05	EPA6 calc c	7.81E-08	EPA6 calc c	1.67E+00	EPA6 calc c	NA
1,1-Biphenyl	92-52-4	6.31E-02	EPA6 calc c	1.58E-04	EPA6 calc c	5.42E+02	EPA6 calc c	NA
1,2 Diphenylhydrazine	122-66-7	6.92E-03	EPA6 e	1.73E-05	EPA6 e	1.01E+02	EPA6 e	NA
1,2-Dimethylhydrazine	540-73-8	3.40E-07	EPA6 calc c	8.50E-10	EPA6 calc c	5.37E-02	EPA6 calc c	NA
1,3-Propane sultone	1120-71-4	2.38E-06	EPA6 calc c	5.95E-09	EPA6 calc c	2.36E-01	EPA6 calc c	NA
2,4-Toluene diisocyanate	584-84-9	NA	NA	NA	NA	NA	NA	NA
2-Chloroacetophenone	532-27-4	3.12E-03	EPA6 calc c	7.80E-06	EPA6 calc c	5.52E+01	EPA6 calc c	NA
2-Propenoic acid	79-10-7	2.14E-05	EPA6 calc c	5.34E-08	EPA6 calc c	1.25E+00	EPA6 calc c	NA
4,4-Methylenedianiline	101-77-9	1.90E-02	EPA6 calc c	4.75E-05	EPA6 calc c	2.18E+02	EPA6 calc c	NA
Acetophenone	98-86-2	3.47E-04	EPA6 e	8.66E-07	EPA6 e	1.04E+01	EPA6 e	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors						
Constituent of Potential Concern	CAS Registry Number	Ba_{egg} (day/kg FW)	Source	$Ba_{chicken}$ (day/kg FW)	Source	BCF_{fish} (L/kg FW tissue)	Source	BAF_{fish} (L/kg FW)
Benzoic acid	65-85-0	6.04E-04	NA	1.51E-06	NA	1.58E+01	EPA6 calc c	NA
bis(2-Chloroethoxy)methane	111-91-1	NA	NA	NA	NA	NA	NA	NA
bis(2-Chloroethyl) ether	111-44-4	1.59E-04	EPA6 e	3.97E-07	EPA6 e	5.74E+00	EPA6 e	NA
Chlorocyclopentadiene	41851-50-7	2.14E-03	EPA6 calc c	5.34E-06	EPA6 calc c	4.14E+01	EPA6 calc c	NA
Cyclohexanol	108-93-0	1.35E-04	EPA6 calc c	3.37E-07	EPA6 calc c	5.07E+00	EPA6 calc c	NA
Dichloroisopropyl ether	108-60-1	3.02E-03	EPA6 calc c	7.54E-06	EPA6 calc c	5.38E+01	EPA6 calc c	NA
Dichloromethyl ether	542-88-1	3.31E-06	EPA6 calc c	8.27E-09	EPA6 calc c	3.03E-01	EPA6 calc c	1.10E+00
Dichloropentadiene	no cas #	NA	NA	NA	NA	NA	NA	NA
Dimethyl sulfate	77-78-1	1.65E-05	EPA6 calc c	4.13E-08	EPA6 calc c	1.03E+00	EPA6 calc c	NA
Dimethylaniline	121-69-7	1.05E-06	EPA6 calc c	2.61E-09	EPA6 calc c	1.26E-01	EPA6 calc c	NA
di-n-propylnitrosamine	621-64-7	1.91E-04	EPA6 e	4.76E-07	EPA6 e	6.59E+00	EPA6 e	NA
Diphenyl ether	101-84-8	1.29E-01	EPA6 calc c	3.22E-04	EPA6 calc c	NA	NA	NA
Epichlorohydrin	106-89-8	1.41E-05	EPA6 e	3.53E-08	EPA6 e	9.13E-01	EPA6 e	NA
Ethyl carbomate (urethane)	51-79-6	5.62E-06	EPA6 calc c	1.40E-08	EPA6 calc c	4.53E-01	EPA6 calc c	NA
Ethyl methane sulfanate	62-50-0	8.90E-06	EPA6 e	2.22E-08	EPA6 e	6.42E-01	EPA6 e	NA
Ethylene dibromide	106-93-4	4.47E-04	EPA6 e	1.12E-06	EPA6 e	1.26E+01	EPA6 e	NA
Ethylene glycol	107-21-1	9.67E-07	EPA6 calc c	2.41E-09	EPA6 calc c	1.19E-01	EPA6 calc c	NA
Ethylene glycol monobutyl ether	111-76-2	2.85E-04	EPA6 calc c	7.11E-07	EPA6 calc c	8.94E+00	EPA6 calc c	NA
Ethylene glycol monoethyl ether acetate	111-15-9	3.31E-05	EPA6 calc c	8.25E-08	EPA6 calc c	1.74E+00	EPA6 calc c	NA
Ethylene thiourea	96-45-7	1.81E-06	EPA6 calc c	4.53E-09	EPA6 calc c	1.92E-01	EPA6 calc c	NA
Furfural	98-01-1	7.24E-05	EPA6 calc c	1.81E-07	EPA6 calc c	3.16E+00	EPA6 calc c	NA
Maleic hydrazide	123-33-1	1.45E-06	EPA6 calc c	3.61E-09	EPA6 calc c	1.61E-01	EPA6 calc c	NA
Malononitrile	109-77-3	8.71E-06	EPA6 calc c	2.17E-08	EPA6 calc c	6.32E-01	EPA6 calc c	NA
Methyl styrene (mixed isomers)	25013-15-4	1.78E-02	EPA6 calc c	4.44E-05	EPA6 calc c	2.07E+02	EPA6 calc c	NA
Methylhydrazine	60-34-4	7.08E-07	EPA6 calc c	1.77E-09	EPA6 calc c	9.38E-02	EPA6 calc c	NA
N,N-Diphenylamine	122-39-4	2.51E-02	EPA6 calc c	6.27E-05	EPA6 calc c	2.69E+02	EPA6 calc c	NA
Nitric acid, propyl ester	627-13-4	NA	NA	NA	NA	NA	NA	NA
N-Nitrosodi-n-butylamine	924-16-3	2.04E-03	EPA6 e	5.10E-06	EPA6 e	4.00E+01	EPA6 e	NA
N-Nitrosomorpholine	59-89-2	7.59E-05	EPA6 calc c	1.90E-07	EPA6 calc c	3.27E+00	EPA6 calc c	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors						
Constituent of Potential Concern	CAS Registry Number	Ba_{egg} (day/kg FW)	Source	$Ba_{chicken}$ (day/kg FW)	Source	BCF_{fish} (L/kg FW tissue)	Source	BAF_{fish} (L/kg FW)
N-Nitroso-N,N-dimethylamine	62-75-9	2.69E-06	EPA6 calc c	6.72E-09	EPA6 calc c	2.59E-01	EPA6 calc c	NA
o-Anisidine	90-04-0	1.20E-04	EPA6 calc c	3.00E-07	EPA6 calc c	4.64E+00	EPA6 calc c	NA
Oxalic acid	144-62-7	NA	NA	NA	NA	NA	NA	NA
Phthalic Anhydride	85-44-9	1.99E-06	EPA6 e	4.96E-09	EPA6 e	2.05E-01	EPA6 e	NA
p-Phthalic acid	100-21-0	5.28E-05	EPA6 calc c	1.32E-07	EPA6 calc c	2.48E+00	EPA6 calc c	NA
Pyridine	110-86-1	3.72E-05	EPA6 e	9.28E-08	EPA6 e	1.90E+00	EPA6 e	NA
Quinoline	91-22-5	8.51E-04	EPA6 calc c	2.12E-06	EPA6 calc c	2.05E+01	EPA6 calc c	NA
Quinone	106-51-4	1.26E-05	EPA6 calc c	3.14E-08	EPA6 calc c	8.36E-01	EPA6 calc c	NA
Safrole	94-59-7	3.63E-03	EPA6 e	9.06E-06	EPA6 e	6.19E+01	EPA6 e	NA
Tetrahydrofuran	109-99-9	2.22E-05	EPA6 e	5.55E-08	EPA6 e	1.29E+00	EPA6 e	NA
Other Heavy Semivolatile Compounds								
1,2,4,5-Tetrachlorobenzene	95-94-3	3.47E-01	EPA6 e	8.66E-04	EPA6 e	NA	EPA6 e	4.30E+03
1,3,5-Trinitrobenzene	99-35-4	1.20E-04	EPA6 e	3.00E-07	EPA6 e	4.64E+00	EPA6 e	NA
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	1.17E-01	EPA6 calc c	2.93E-04	EPA6 calc c	NA	NA	NA
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	1.58E-08	EPA6 calc c	3.96E-11	EPA6 calc c	5.22E-03	EPA6 calc c	NA
2-sec-Butyl-4,6-dinitrophenol	88-85-7	1.10E-02	EPA6 calc c	2.74E-05	EPA6 calc c	1.43E+02	EPA6 calc c	NA
3,3-Dichlorobenzidine	91-94-1	2.99E-02	EPA6 e	7.45E-05	EPA6 e	3.07E+02	EPA6 e	NA
3,3'-Dimethoxybenzidine	119-90-4	5.13E-04	EPA6 e	1.28E-06	EPA6 e	1.40E+01	EPA6 e	NA
4-Bromophenylphenyl ether	101-55-3	7.94E-01	EPA6 calc c	1.98E-03	EPA6 calc c	NA	NA	NA
Ammonium perfluorooctanoate	3825-26-1	NA	NA	NA	NA	NA	NA	NA
Azobenzene	103-33-3	5.25E-02	EPA6 calc c	1.31E-04	EPA6 calc c	4.71E+02	EPA6 calc c	NA
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	NA	NA	NA	NA	NA	NA	NA
Captan	133-06-2	1.78E-03	EPA6 calc c	4.44E-06	EPA6 calc c	3.60E+01	EPA6 calc c	NA
Chlorobenzilate	510-15-6	1.91E-01	EPA6 e	4.76E-04	EPA6 e	NA	EPA6 e	2.03E+03
Dibutylphosphate	107-66-4	NA	NA	NA	NA	NA	NA	NA
Dimethyl aminoazobenzene	60-11-7	3.02E-01	EPA6 calc c	7.54E-04	EPA6 calc c	NA	NA	NA
Hexachlorobenzene	118-74-1	2.53E+00	EPA6 e	6.31E-03	EPA6 e	NA	EPA6 e	5.52E+04
Hexachlorobutadiene	87-68-3	4.27E-01	EPA6 e	1.07E-03	EPA6 e	NA	EPA6 e	5.69E+03
Hexachlorocyclopentadiene	77-47-4	6.41E-01	EPA6 e	1.60E-03	EPA6 e	NA	EPA6 e	5.25E+02

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors						
Constituent of Potential Concern	CAS Registry Number	Ba_{egg} (day/kg FW)	Source	$Ba_{chicken}$ (day/kg FW)	Source	BCF_{fish} (L/kg FW tissue)	Source	BAF_{fish} (L/kg FW)
Hexachloroethane	67-72-1	7.67E-02	EPA6 e	1.92E-04	EPA6 e	6.29E+02	EPA6 e	NA
Hexachlorophene	70-30-4	2.75E+02	EPA6 e	6.88E-01	EPA6 e	NA	EPA6 e	4.66E+03
Hexamethylene-1,5-diisocyanate	822-06-0	1.49E-04	EPA6 calc c	3.72E-07	EPA6 calc c	5.47E+00	EPA6 calc c	NA
Mirex	2385-85-5	6.17E+01	EPA6 calc c	1.54E-01	EPA6 calc c	NA	NA	NA
Nitrofen	1836-75-5	2.69E+00	EPA6 calc c	6.72E-03	EPA6 calc c	NA	NA	NA
Pentachlorobenzene	608-93-5	9.72E-01	EPA6 e	2.43E-03	EPA6 e	NA	EPA6 e	3.61E+04
Pentachloronitrobenzene	82-68-8	3.47E-01	EPA6 e	8.66E-04	EPA6 e	NA	EPA6 e	4.65E+02
Pentachlorophenol*	87-86-5	9.55E-01	EPA6 e	2.39E-03	EPA6 e	NA	EPA6 e	3.97E+02
Pentachlorophenol*	87-86-5	9.55E-01	EPA6 e	2.39E-03	EPA6 e	NA	EPA6 e	3.97E+02
Pentachlorophenol*	87-86-5	9.55E-01	EPA6 e	2.39E-03	EPA6 e	NA	EPA6 e	3.97E+02
Pentachlorophenol*	87-86-5	9.55E-01	EPA6 e	2.39E-03	EPA6 e	NA	EPA6 e	3.97E+02
Pentachlorophenol*	87-86-5	9.55E-01	EPA6 e	2.39E-03	EPA6 e	NA	EPA6 e	3.97E+02
Pentachlorophenol*	87-86-5	9.55E-01	EPA6 e	2.39E-03	EPA6 e	NA	EPA6 e	3.97E+02
Pentachlorophenol*	87-86-5	9.55E-01	EPA6 e	2.39E-03	EPA6 e	NA	EPA6 e	3.97E+02
Pentachlorophenol*	87-86-5	9.55E-01	EPA6 e	2.39E-03	EPA6 e	NA	EPA6 e	3.97E+02
Pentachlorophenol*	87-86-5	9.55E-01	EPA6 e	2.39E-03	EPA6 e	NA	EPA6 e	3.97E+02
Pentachlorophenol*	87-86-5	9.55E-01	EPA6 e	2.39E-03	EPA6 e	NA	EPA6 e	3.97E+02
Pentachlorophenol*	87-86-5	9.55E-01	EPA6 e	2.39E-03	EPA6 e	NA	EPA6 e	3.97E+02
Pentachlorophenol*	87-86-5	9.55E-01	EPA6 e	2.39E-03	EPA6 e	NA	EPA6 e	3.97E+02
Picric acid	88-89-1	8.51E-04	EPA6 calc c	2.12E-06	EPA6 calc c	2.05E+01	EPA6 calc c	NA
Pronamide	23950-58-5	2.57E-02	EPA6 e	6.42E-05	EPA6 e	2.74E+02	EPA6 e	NA
Strychnine	57-24-9	6.76E-04	EPA6 e	1.69E-06	EPA6 e	1.72E+01	EPA6 e	NA
Terphenyls	26140-60-3	NA	NA	NA	NA	NA	NA	NA
Tributyl phosphate	126-73-8	7.94E-02	EPA6 calc c	1.98E-04	EPA6 calc c	NA	NA	NA
Trifluralin	1582-09-8	1.74E+00	EPA6 calc c	4.34E-03	EPA6 calc c	NA	NA	NA
Triphenylamine	603-34-9	NA	NA	NA	NA	NA	NA	NA
Herbicides and Organopesticides								
2,4,5-T	93-76-5	1.82E-02	EPA6 calc c	4.54E-05	EPA6 calc c	2.11E+02	EPA6 calc c	NA

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors						
Constituent of Potential Concern	CAS Registry Number	Ba_{egg} (day/kg FW)	Source	$Ba_{chicken}$ (day/kg FW)	Source	BCF_{fish} (L/kg FW tissue)	Source	BAF_{fish} (L/kg FW)
2,4-D and esters	94-75-7	5.13E-03	EPA6 calc c	1.28E-05	EPA6 calc c	8.05E+01	EPA6 calc c	NA
4,4-DDD	72-54-8	1.26E+01	EPA6 calc c	3.14E-02	EPA6 calc c	NA	NA	NA
4,4-DDT	50-29-3	7.94E+00	EPA6 calc c	1.98E-02	EPA6 calc c	NA	NA	NA
Aldrin	309-00-2	1.20E+01	EPA6 e	2.99E-02	EPA6 e	NA	EPA6 e	5.82E+05
alpha-BHC	319-84-6	5.00E-02	EPA6 e	1.25E-04	EPA6 e	4.54E+02	EPA6 e	NA
beta-BHC	319-85-7	5.41E-02	EPA6 e	1.35E-04	EPA6 e	4.82E+02	EPA6 e	NA
Chlordane	57-74-9	6.88E+00	EPA6 e	1.72E-02	EPA6 e	6.07E-01	EPA6 e	NA
DDE	72-55-9	1.43E+01	EPA6 e	3.58E-02	EPA6 e	NA	EPA6 e	5.53E+05
Delta-BHC	319-86-8	1.10E-01	EPA6 calc c	2.74E-04	EPA6 calc c	NA	NA	NA
Dieldrin	60-57-1	1.48E+00	EPA6 e	3.68E-03	EPA6 e	NA	EPA6 e	2.86E+04
Endothall	145-73-3	1.07E-06	EPA6 calc c	2.68E-09	EPA6 calc c	1.28E-01	EPA6 calc c	NA
Endrin	72-20-8	6.19E-01	EPA6 e	1.55E-03	EPA6 e	NA	EPA6 e	8.55E+03
gamma-BHC(Lindane)	58-89-9	4.17E-02	EPA6 calc c	1.04E-04	EPA6 calc c	3.96E+02	EPA6 calc c	NA
Heptachlor	76-44-8	8.22E-01	EPA6 e	2.05E-03	EPA6 e	5.52E+03	EPA6 e	NA
Isodrin	465-73-6	NA	NA	NA	NA	NA	NA	NA
Methoxychlor	72-43-5	2.67E-01	EPA6 e	6.66E-04	EPA6 e	NA	EPA6 e	3.16E+03
Silvex (2,4,5-TP)	93-72-1	9.38E-02	EPA6 calc c	2.34E-04	EPA6 calc c	NA	NA	NA
Toxaphene	8001-35-2	2.51E+00	EPA6 calc c	6.27E-03	EPA6 calc c	NA	NA	NA

NA = Value not available
** = Varies with pH; values given for a pH of 7.0.
a = Illustrated Handbook of Physical - Chemical Properties and Environmental Fate f
b = RAIS database: http://risk.lsd.ornl.gov/rap_hp.shtml
c = Equations derived from EPA Region 6 Guidance Appendix A-3
d = Handbook of Environmental degradation Rates. 1991.
e = EPA Region 6 Hazardous Waste Combustion Guidance
f = "A Critical Review of Henry's Law Constants for Chemicals of Environmental Im
g = Handbook of Environmental Fate and Exposure Data for Organic Chemicals
h = http://www.epa.gov/region09/waste/sfund/prg/.
i = http://esc.syres.com/efdb/Chemfate.htm
j = Hansch and Leo 1985 in Syracuse 1996

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors						
Constituent of Potential Concern	CAS Registry Number	Ba_{egg} (day/kg FW)	Source	$Ba_{chicken}$ (day/kg FW)	Source	BCF_{fish} (L/kg FW tissue)	Source	BAF_{fish} (L/kg FW)

k = Schwarzenbach et al. 1993
l = Tomlin 1994 in Syracuse 1996
m = http://www.speclab.com/compound/c959988.htm

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors				
Constituent of Potential Concern	CAS Registry Number	Source	$BSAF_{fish}$ (unitless)	Source		
Aromatic Halogenated Hydrocarbons						
2,3,4,6-Tetrachlorophenol	58-90-2	NA	NA	NA		
4-Chloro-3-methylphenol	59-50-7	NA	NA	NA		
Aromatic Nonhalogenated Hydrocarbons						
2-Nitrotoluene	88-72-2	NA	NA	NA		
4-Nitrobiphenyl	92-93-3	NA	NA	NA		
Benzaldehyde	100-52-7	EPA6 e	NA	EPA6 e		
Benzene	71-43-2	EPA6 e	NA	EPA6 e		
Benzyl alcohol	100-51-6	EPA6 e	NA	EPA6 e		
Ethyl Benzene	100-41-4	EPA6 e	NA	EPA6 e		
m-Xylene	108-38-3	EPA6 e	NA	EPA6 e		
o-Xylene	95-47-6	EPA6 e	NA	EPA6 e		
p-Xylene	106-42-3	EPA6 e	NA	EPA6 e		
Styrene	100-42-5	EPA6 e	NA	EPA6 e		
Toluene	108-88-3	EPA6 e	NA	EPA6 e		
Non-aromatic Nonhalogenated Hydrocarbons						
1,2-Epoxybutane	106-88-7	NA	NA	NA		
1,3-Butadiene	106-99-0	NA	NA	NA		
1,4-Dioxane	123-91-1	EPA6 e	NA	EPA6 e		
1-Methylpropyl alcohol	78-92-2	NA	NA	NA		
1-Nitropropane	108-03-2	NA	NA	NA		
2,2,4-Trimethlylpentane	540-84-1	NA	NA	NA		
2-Butanone	78-93-3	EPA6 e	NA	EPA6 e		
2-Butenaldehyde (2-Butenal)	4170-30-3	NA	NA	NA		
2-Ethoxyethanol	110-80-5	NA	NA	NA		
2-Heptanone	110-43-0	NA	NA	NA		
2-Hexanone	591-78-6	NA	NA	NA		
2-Methoxyethanol	109-86-4	NA	NA	NA		
2-Methyl-2-propanol	75-65-0	NA	NA	NA		

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors				
Constituent of Potential Concern	CAS Registry Number	Source	$BSAF_{fish}$ (unitless)	Source		
2-Methyl-2-propenenitrile	126-98-7	EPA6 e	NA	EPA6 e		
2-Methylaziridine	75-55-8	NA	NA	NA		
2-Methylpropyl alcohol	78-83-1	NA	NA	NA		
2-Pentanone	107-87-9	NA	NA	NA		
2-Propanone (Acetone)	67-64-1	EPA6 e	NA	EPA6 e		
2-Propene-1-ol	107-18-6	NA	NA	NA		
2-Propyl alcohol	67-63-0	NA	NA	NA		
3-Heptanone	106-35-4	NA	NA	NA		
3-Methyl-1-butanol	123-51-3	NA	NA	NA		
3-Methyl-2-butanone	563-80-4	NA	NA	NA		
3-Pentanone	96-22-0	NA	NA	NA		
4-Heptanone	123-19-3	NA	NA	NA		
4-Methyl-2-pentanone	108-10-1	EPA6 e	NA	EPA6 e		
4-Methyl-3-penten-2-one	141-79-7	NA	NA	NA		
5-Methyl-2-hexanone	110-12-3	NA	NA	NA		
Acetaldehyde	75-07-0	EPA6 e	NA	EPA6 e		
Acetamide	60-35-5	NA	NA	NA		
Acetic acid	64-19-7	NA	NA	NA		
Acetic acid ethyl ester	141-78-6	NA	NA	NA		
Acetic acid n-butyl ester	123-86-4	NA	NA	NA		
Acetonitrile	75-05-8	EPA6 e	NA	EPA6 e		
Acrolein	107-02-8	EPA6 e	NA	EPA6 e		
Acrylonitrile	107-13-1	EPA6 e	NA	EPA6 e		
Bis(isopropyl)ether	108-20-3	NA	NA	NA		
Butane	106-97-8	NA	NA	NA		
Carbon disulfide	75-15-0	EPA6 e	NA	EPA6 e		
Cyanogen	460-19-5	NA	NA	NA		
Cyclohexane	110-82-7	NA	NA	NA		
Cyclohexanone	108-94-1	NA	NA	NA		

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors				
Constituent of Potential Concern	CAS Registry Number	Source	$BSAF_{fish}$ (unitless)	Source		
Cyclohexene	110-83-8	NA	NA	NA		
Cyclopentane	287-92-3	NA	NA	NA		
Ethyl alcohol	64-17-5	NA	NA	NA		
Ethyl ether	60-29-7	NA	NA	NA		
Ethyl methacrylate	97-63-2	EPA6 e	NA	EPA6 e		
Formaldehyde	50-00-0	EPA6 e	NA	EPA6 e		
Formamide	75-12-7	NA	NA	NA		
Formic acid	64-18-6	EPA6 e	NA	EPA6 e		
Formic acid, methyl ester	107-31-3	NA	NA	NA		
Glycidylaldehyde	765-34-4	NA	NA	NA		
Methyl acetate	79-20-9	NA	NA	NA		
Methyl alcohol	67-56-1	EPA6 e	NA	EPA6 e		
Methyl isocyanate	624-83-9	NA	NA	NA		
Methyl methacrylate	80-62-6	NA	NA	NA		
Methyl tert-butyl ether	1634-04-4	NA	NA	NA		
Methylacetylene	74-99-7	NA	NA	NA		
Methylcyclohexane	108-87-2	NA	NA	NA		
N,N-Dimethylacetamide	127-19-5	NA	NA	NA		
n-Butyl alcohol	71-36-3	NA	NA	NA		
n-Heptane	142-82-5	NA	NA	NA		
n-Hexane	110-54-3	NA	NA	NA		
Nitromethane	75-52-5	NA	NA	NA		
n-Nonane	111-84-2	NA	NA	NA		
n-Octane	111-65-9	NA	NA	NA		
n-Pentane	109-66-0	NA	NA	NA		
n-Propionaldehyde	123-38-6	NA	NA	NA		
n-Propyl alcohol	71-23-8	NA	NA	NA		
n-Valeraldehyde	110-62-3	NA	NA	NA		
Oxirane	75-21-8	EPA6 e	NA	EPA6 e		

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors				
Constituent of Potential Concern	CAS Registry Number	Source	$BSAF_{fish}$ (unitless)	Source		
p-Cymene	99-87-6	NA	NA	NA		
Phosgene	75-44-5	NA	NA	NA		
Propargyl alcohol	107-19-7	NA	NA	NA		
Propionic acid	79-09-4	NA	NA	NA		
Propionitrile	107-12-0	NA	NA	NA		
Propylene glycol monomethyl ether	107-98-2	NA	NA	NA		
p-tert-Butyltoluene	98-51-1	NA	NA	NA		
Triethylamine	121-44-8	NA	NA	NA		
Trimethylamine	75-50-3	NA	NA	NA		
Vinyl acetate	108-05-4	EPA6 e	NA	EPA6 e		
Non-aromatic Halogenated Hydrocarbons						
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	NA	NA	NA		
1,1,1,2-Tetrachloroethane	630-20-6	EPA6 e	NA	EPA6 e		
1,1,1-Trichloroethane	71-55-6	EPA6 e	NA	EPA6 e		
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	NA	NA	NA		
1,1,2,2-Tetrachloroethane	79-34-5	EPA6 e	NA	EPA6 e		
1,1,2,2-Tetrachloroethene	127-18-4	EPA6 e	NA	EPA6 e		
1,1,2-Trichloro-1,2,2-trifluoroethane	76-13-1	NA	NA	NA		
1,1,2-Trichloroethane	79-00-5	EPA6 e	NA	EPA6 e		
1,1,2-Trichloroethylene	79-01-6	EPA6 e	NA	EPA6 e		
1,1-Dichloroethane	75-34-3	EPA6 e	NA	EPA6 e		
1,1-Dichloroethene	75-35-4	EPA6 e	NA	EPA6 e		
1,2,3-Trichloropropane	96-18-4	EPA6 e	NA	EPA6 e		
1,2-dibromo-3-chloropropane	96-12-8	EPA6 e	NA	EPA6 e		
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	NA	NA	NA		
1,2-Dichloroethane	107-06-2	EPA6 e	NA	EPA6 e		
1,2-Dichloroethylene	540-59-0	NA	NA	NA		
1,2-Dichloropropane	78-87-5	EPA6 e	NA	EPA6 e		
1,3-Dichloropropene	542-75-6	EPA6 e	NA	EPA6 e		

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors				
Constituent of Potential Concern	CAS Registry Number	Source	$BSAF_{fish}$ (unitless)	Source		
1,4-Dichloro-2-butene	764-41-0	NA	NA	NA		
1-Chloroethene	75-01-4	EPA6 e	NA	EPA6 e		
2,2-Dichloropropionic acid	75-99-0	NA	NA	NA		
2-Chloropropane	75-29-6	NA	NA	NA		
3-Chloropropene (allyl chloride)	107-05-1	NA	NA	NA		
Bromochloromethane	74-97-5	NA	NA	NA		
Bromodichloromethane	75-27-4	EPA6 e	NA	EPA6 e		
Bromoethene	593-60-2	NA	NA	NA		
Bromoform	75-25-2	EPA6 e	NA	EPA6 e		
Bromomethane	74-83-9	EPA6 e	NA	EPA6 e		
Carbon tetrachloride	56-23-5	EPA6 e	NA	EPA6 e		
Chlordibromomethane	124-48-1	EPA6 e	NA	EPA6 e		
Chlorodifluoromethane	75-45-6	EPA6 e	NA	EPA6 e		
Chloroethane	75-00-3	EPA6 e	NA	EPA6 e		
Chloroform	67-66-3	EPA6 e	NA	EPA6 e		
Chloromethane	74-87-3	EPA6 e	NA	EPA6 e		
Chloropentafluoroethane	76-15-3	NA	NA	NA		
cis-1,2-Dichloroethene	156-59-2	EPA6 e	NA	EPA6 e		
cis-1,3-Dichloropropene	10061-01-5	NA	NA	NA		
Cyanogen bromide	506-68-3	NA	NA	NA		
Cyanogen chloride	506-77-4	NA	NA	NA		
Dichlorodifluoromethane	75-71-8	EPA6 e	NA	EPA6 e		
Dichlorofluoromethane	75-43-4	NA	NA	NA		
Dichloromethane	75-09-2	EPA6 e	NA	EPA6 e		
Difluorodibromomethane	75-61-6	NA	NA	NA		
Hexafluoroacetone	684-16-2	NA	NA	NA		
Iodomethane	74-88-4	NA	NA	NA		
Methylene bromide	74-95-3	EPA6 e	NA	EPA6 e		
Pentachloroethane	76-01-7	NA	NA	NA		

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors				
Constituent of Potential Concern	CAS Registry Number	Source	$BSAF_{fish}$ (unitless)	Source		
trans-1,2-Dichloroethene	156-60-5	EPA6 e	NA	EPA6 e		
trans-1,3-Dichloropropene	10061-02-6	NA	NA	NA		
Trichloroacetic acid	76-03-9	NA	NA	NA		
Trichlorofluoroethane	27154-33-2	NA	NA	NA		
Trichlorofluoromethane	75-69-4	EPA6 e	NA	EPA6 e		
Trifluorobromomethane	75-63-8	NA	NA	NA		
Dioxin and Furan Compounds						
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	35822-46-9	EPA6 e	5.00E-03	EPA6 e		
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	EPA6 e	5.00E-03	EPA6 e		
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	EPA6 e	5.00E-03	EPA6 e		
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	39227-28-6	EPA6 e	4.00E-02	EPA6 e		
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	EPA6 e	4.00E-02	EPA6 e		
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	57653-85-7	EPA6 e	4.00E-02	EPA6 e		
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	EPA6 e	4.00E-02	EPA6 e		
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	19408-74-3	EPA6 e	4.00E-02	EPA6 e		
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	EPA6 e	4.00E-02	EPA6 e		
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	40321-76-4	EPA6 e	9.00E-02	EPA6 e		
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	EPA6 e	9.00E-02	EPA6 e		
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	EPA6 e	4.00E-02	EPA6 e		
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	EPA6 e	9.00E-02	EPA6 e		
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1746-01-6	EPA6 e	9.00E-02	EPA6 e		
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	EPA6 e	9.00E-02	EPA6 e		
Dibenzofuran	132-64-9	NA	NA	NA		
Octachlorodibenzo(p)dioxin	3268-87-9	EPA6 e	1.00E-04	EPA6 e		
Octachlorodibenzofuran	39001-02-0	EPA6 e	1.00E-04	EPA6 e		
PCBs						
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	NA	NA	NA		
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	NA	NA	NA		
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	NA	NA	NA		

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors				
Constituent of Potential Concern	CAS Registry Number	Source	$BSAF_{fish}$ (unitless)	Source		
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	NA	NA	NA		
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	NA	NA	NA		
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	NA	NA	NA		
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	NA	NA	NA		
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	NA	NA	NA		
2',3,4,4',5-Pentachlorobiphenyl	no cas #	NA	NA	NA		
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	NA	NA	NA		
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	NA	NA	NA		
3,3',4,4',5-Pentachlorobiphenyl	no cas #	NA	NA	NA		
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	NA	NA	NA		
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	NA	NA			
Polychlorinated biphenyls (PCBs)**	1336-36-3	RAIS b	NA	NA		
Phthalates						
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	EPA6 e	NA	EPA6 e		
Butylbenzyl phthalate	85-68-7	EPA6 e	NA	EPA6 e		
Dibutyl phthalate	84-74-2	EPA6 e	NA	EPA6 e		
Diethyl phthalate	84-66-2	EPA6 e	NA	EPA6 e		
Dimethyl phthalate	131-11-3	EPA6 e	NA	EPA6 e		
n-Dioctyl phthalate	117-84-0	EPA6 e	NA	EPA6 e		
Light Polycyclic Aromatic Hydrocarbons						
2-Chloronaphthalene	91-58-7	EPA6 e	NA	EPA6 e		
2-Methyl naphthalene	91-57-6	NA	NA	NA		
5-Nitroacenaphthene	602-87-9	NA	NA	NA		
Acenaphthene	83-32-9	EPA6 e	NA	EPA6 e		
Acenaphthylene	208-96-8	NA	NA	NA		
Anthracene	120-12-7	EPA6 e	NA	EPA6 e		
Fluorene	86-73-7	NA	NA	NA		
Indene	95-13-6	NA	NA	NA		
Napthalene	91-20-3	EPA6 e	NA	EPA6 e		

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors				
Constituent of Potential Concern	CAS Registry Number	Source	$BSAF_{fish}$ (unitless)	Source		
Phenanthrene	85-01-8	EPA6 e	NA	EPA6 e		
Pyrene	129-00-0	EPA6 e	NA	EPA6 e		
Heavy Polycyclic Aromatic Hydrocarbons						
3-Methylcholanthrene	56-49-5	NA	NA	NA		
5-Methylchrysene	3697-24-3	NA	NA	NA		
Benzo(a)anthracene	56-55-3	EPA6 e	NA	EPA6 e		
Benzo(a)pyrene	50-32-8	EPA6 e	NA	EPA6 e		
Benzo(b)fluoranthene	205-99-2	EPA6 e	NA	EPA6 e		
Benzo(e)pyrene	192-97-2	NA	NA	NA		
Benzo(g,h,i)perylene	191-24-2	NA	NA	NA		
Benzo(j)fluoranthene	205-82-3	NA	NA	NA		
Benzo(k)fluoranthene	207-08-9	EPA6 e	NA	EPA6 e		
Benzo[a,i]pyrene	191-30-0	NA	NA	NA		
Chrysene	218-01-9	EPA6 e	NA	EPA6 e		
Dibenz(a,h)anthracene	53-70-3	EPA6 e	NA	EPA6 e		
Dibenz[a,h]acridine	226-36-8	NA	NA	NA		
Dibenz[a,j]acridine	224-42-0	NA	NA	NA		
Dibenzo(a,e)fluoranthene	5385-75-1	NA	NA	NA		
Dibenzo(a,h)fluoranthene	no cas #	NA	NA	NA		
Dibenzo[a,e]pyrene	192-65-4	NA	NA	NA		
Dibenzo[a,h]pyrene	189-64-0	NA	NA	NA		
Dibenzo[a,i]pyrene	189-55-9	NA	NA	NA		
Fluoranthene	206-44-0	EPA6 e	NA	EPA6 e		
Hexachloronaphthalene	1335-87-1	NA	NA	NA		
Indeno(1,2,3-cd)pyrene	193-39-5	EPA6 e	NA	EPA6 e		
Octachloronaphthalene	2234-13-1	NA	NA	NA		
Pentachloronaphthalene	1321-64-8	NA	NA	NA		
Tetrachloronaphthalene	1335-88-2	NA	NA	NA		
Trichloronaphthalene	1321-65-9	NA	NA	NA		

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors				
Constituent of Potential Concern	CAS Registry Number	Source	$BSAF_{fish}$ (unitless)	Source		
Light Substituted Benzene Compounds						
1,2,3-Trichlorobenzene	87-61-6	EPA6 e	NA	EPA6 e		
1,2,4-Trichlorobenzene	120-82-1	EPA6 e	NA	EPA6 e		
1,2,4-Trimethlyl benzene	25551-13-7	NA	NA	NA		
1,2,4-Trimethyl benzene	95-63-6	NA	NA	NA		
1,2-Dichlorobenzene	95-50-1	EPA6 e	NA	EPA6 e		
1,3,5-Trimethyl benzene	108-67-8	EPA6 e	NA	EPA6 e		
1,3-Dichlorobenzene	541-73-1	EPA6 e	NA	EPA6 e		
1,3-Dinitrobenzene	99-65-0	EPA6 e	NA	EPA6 e		
1,4-Dichlorobenzene	106-46-7	EPA6 e	NA	EPA6 e		
1,4-Dinitrobenzene	100-25-4	NA	NA	NA		
2,4,5-Trichorophenol	95-95-4	EPA6 e	NA	EPA6 e		
2,4,6-Trichlorophenol	88-06-2	EPA6 e	NA	EPA6 e		
2,4-Dichlorophenol	120-83-2	EPA6 e	NA	EPA6 e		
2,4-Dimethylphenol	105-67-9	EPA6 e	NA	EPA6 e		
2,4-Dinitrophenol	51-28-5	EPA6 e	NA	EPA6 e		
2,4-Dinitrotoluene	121-14-2	EPA6 e	NA	EPA6 e		
2,6-Dinitrotoluene	606-20-2	EPA6 e	NA	EPA6 e		
2-Chlorotoluene	95-49-8	NA	NA	NA		
2-Chorophenol	95-57-8	EPA6 e	NA	EPA6 e		
2-Nitrophenol	88-75-5	EPA6 e	NA	EPA6 e		
4,6-Dinitro-o-cresol	534-52-1	NA	NA	NA		
4-Chlorotoluene	106-43-4	NA	NA	NA		
4-Nitrophenol	100-02-7	EPA6 e	NA	EPA6 e		
alpha-Methylstyrene	98-83-9	NA	NA	NA		
Aniline	62-53-3	EPA6 e	NA	EPA6 e		
Benzotrichloride	98-07-7	NA	NA	NA		
Benzyl Chloride	100-44-7	EPA6 e	NA	EPA6 e		
Bromobenzene	108-86-1	NA	NA	NA		

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors				
Constituent of Potential Concern	CAS Registry Number	Source	$BSAF_{fish}$ (unitless)	Source		
Chlorobenzene	108-90-7	EPA6 e	NA	EPA6 e		
Cumene	98-82-8	EPA6 e	NA	EPA6 e		
m-Cresol	108-39-4	EPA6 e	NA	EPA6 e		
n-Butyl benzene	104-51-8	NA	NA	NA		
Nitrobenzene	98-95-3	EPA6 e	NA	EPA6 e		
n-Propyl benzene	103-65-1	NA	NA	NA		
o-Cresol	95-48-7	EPA6 e	NA	EPA6 e		
o-Dinitrobenzene	528-29-0	NA	NA	NA		
o-Nitroaniline	88-74-4	EPA6 e	NA	EPA6 e		
o-Toluidine	95-53-4	EPA6 e	NA	EPA6 e		
p-Chloroaniline	106-47-8	EPA6 e	NA	EPA6 e		
p-Cresol	106-44-5	EPA6 e	NA	EPA6 e		
Phenol	108-95-2	EPA6 e	NA	EPA6 e		
p-Nitrochlorobenzene	100-00-5	NA	NA	NA		
p-Toluidine	106-49-0	NA	NA	NA		
sec-Butyl benzene	135-98-8	NA	NA	NA		
tert-Butyl benzene	98-06-6	NA	NA	NA		
Toluene-2,6-diamine	823-40-5	NA	NA	NA		
Other Light Semivolatile Compounds						
1,1 Dimethyl hydrazine	57-14-7	NA	NA	NA		
1,1-Biphenyl	92-52-4	NA	NA	NA		
1,2 Diphenylhydrazine	122-66-7	EPA6 e	NA	EPA6 e		
1,2-Dimethylhydrazine	540-73-8	NA	NA	NA		
1,3-Propane sultone	1120-71-4	NA	NA	NA		
2,4-Toluene diisocyanate	584-84-9	NA	NA	NA		
2-Chloroacetophenone	532-27-4	NA	NA	NA		
2-Propenoic acid	79-10-7	NA	NA	NA		
4,4-Methylenedianiline	101-77-9	NA	NA	NA		
Acetophenone	98-86-2	EPA6 e	NA	EPA6 e		

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors				
Constituent of Potential Concern	CAS Registry Number	Source	$BSAF_{fish}$ (unitless)	Source		
Benzoic acid	65-85-0	NA	NA	NA		
bis(2-Chloroethoxy)methane	111-91-1	NA	NA	NA		
bis(2-Chloroethyl) ether	111-44-4	EPA6 e	NA	EPA6 e		
Chlorocyclopentadiene	41851-50-7	NA	NA	NA		
Cyclohexanol	108-93-0	NA	NA	NA		
Dichloroisopropyl ether	108-60-1	NA	NA	NA		
Dichloromethyl ether	542-88-1	RAIS b	NA	NA		
Dichloropentadiene	no cas #	NA	NA	NA		
Dimethyl sulfate	77-78-1	NA	NA	NA		
Dimethylaniline	121-69-7	NA	NA	NA		
di-n-propylnitrosamine	621-64-7	EPA6 e	NA	EPA6 e		
Diphenyl ether	101-84-8	NA	NA	NA		
Epichlorohydrin	106-89-8	EPA6 e	NA	EPA6 e		
Ethyl carbomate (urethane)	51-79-6	NA	NA	NA		
Ethyl methane sulfanate	62-50-0	EPA6 e	NA	EPA6 e		
Ethylene dibromide	106-93-4	EPA6 e	NA	EPA6 e		
Ethylene glycol	107-21-1	NA	NA	NA		
Ethylene glycol monobutyl ether	111-76-2	NA	NA	NA		
Ethylene glycol monoethyl ether acetate	111-15-9	NA	NA	NA		
Ethylene thiourea	96-45-7	NA	NA	NA		
Furfural	98-01-1	NA	NA	NA		
Maleic hydrazide	123-33-1	NA	NA	NA		
Malononitrile	109-77-3	NA	NA	NA		
Methyl styrene (mixed isomers)	25013-15-4	NA	NA	NA		
Methylhydrazine	60-34-4	NA	NA	NA		
N,N-Diphenylamine	122-39-4	NA	NA	NA		
Nitric acid, propyl ester	627-13-4	NA	NA	NA		
N-Nitrosodi-n-butylamine	924-16-3	EPA6 e	NA	EPA6 e		
N-Nitrosomorpholine	59-89-2	NA	NA	NA		

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors				
Constituent of Potential Concern	CAS Registry Number	Source	$BSAF_{fish}$ (unitless)	Source		
N-Nitroso-N,N-dimethylamine	62-75-9	NA	NA	NA		
o-Anisidine	90-04-0	NA	NA	NA		
Oxalic acid	144-62-7	NA	NA	NA		
Phthalic Anhydride	85-44-9	EPA6 e	NA	EPA6 e		
p-Phthalic acid	100-21-0	NA	NA	NA		
Pyridine	110-86-1	EPA6 e	NA	EPA6 e		
Quinoline	91-22-5	NA	NA	NA		
Quinone	106-51-4	NA	NA	NA		
Safrole	94-59-7	EPA6 e	NA	EPA6 e		
Tetrahydrofuran	109-99-9	EPA6 e	NA	EPA6 e		
Other Heavy Semivolatile Compounds						
1,2,4,5-Tetrachlorobenzene	95-94-3	EPA6 e	NA	EPA6 e		
1,3,5-Trinitrobenzene	99-35-4	EPA6 e	NA	EPA6 e		
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	NA	NA	NA		
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	NA	NA	NA		
2-sec-Butyl-4,6-dinitrophenol	88-85-7	NA	NA	NA		
3,3-Dichlorobenzidine	91-94-1	EPA6 e	NA	EPA6 e		
3,3'-Dimethoxybenzidine	119-90-4	EPA6 e	NA	EPA6 e		
4-Bromophenylphenyl ether	101-55-3	NA	NA	NA		
Ammonium perfluorooctanoate	3825-26-1	NA	NA	NA		
Azobenzene	103-33-3	NA	NA	NA		
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	NA	NA	NA		
Captan	133-06-2	NA	NA	NA		
Chlorobenzilate	510-15-6	EPA6 e	NA	EPA6 e		
Dibutylphosphate	107-66-4	NA	NA	NA		
Dimethyl aminoazobenzene	60-11-7	NA	NA	NA		
Hexachlorobenzene	118-74-1	EPA6 e	NA	EPA6 e		
Hexachlorobutadiene	87-68-3	EPA6 e	NA	EPA6 e		
Hexachlorocyclopentadiene	77-47-4	EPA6 e	NA	EPA6 e		

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors				
Constituent of Potential Concern	CAS Registry Number	Source	$BSAF_{fish}$ (unitless)	Source		
Hexachloroethane	67-72-1	EPA6 e	NA	EPA6 e		
Hexachlorophene	70-30-4	EPA6 e	NA	EPA6 e		
Hexamethylene-1,5-diisocyanate	822-06-0	NA	NA	NA		
Mirex	2385-85-5	NA	NA	NA		
Nitrofen	1836-75-5	NA	NA	NA		
Pentachlorobenzene	608-93-5	EPA6 e	NA	EPA6 e		
Pentachloronitrobenzene	82-68-8	EPA6 e	NA	EPA6 e		
Pentachlorophenol*	87-86-5	EPA6 e	NA	EPA6 e		
Pentachlorophenol*	87-86-5	EPA6 e	NA	EPA6 e		
Pentachlorophenol*	87-86-5	EPA6 e	NA	EPA6 e		
Pentachlorophenol*	87-86-5	EPA6 e	NA	EPA6 e		
Pentachlorophenol*	87-86-5	EPA6 e	NA	EPA6 e		
Pentachlorophenol*	87-86-5	EPA6 e	NA	EPA6 e		
Pentachlorophenol*	87-86-5	EPA6 e	NA	EPA6 e		
Pentachlorophenol*	87-86-5	EPA6 e	NA	EPA6 e		
Pentachlorophenol*	87-86-5	EPA6 e	NA	EPA6 e		
Pentachlorophenol*	87-86-5	EPA6 e	NA	EPA6 e		
Pentachlorophenol*	87-86-5	EPA6 e	NA	EPA6 e		
Pentachlorophenol*	87-86-5	EPA6 e	NA	EPA6 e		
Picric acid	88-89-1	NA	NA	NA		
Pronamide	23950-58-5	EPA6 e	NA	EPA6 e		
Strychnine	57-24-9	EPA6 e	NA	EPA6 e		
Terphenyls	26140-60-3	NA	NA	NA		
Tributyl phosphate	126-73-8	NA	NA	NA		
Trifluralin	1582-09-8	NA	NA	NA		
Triphenylamine	603-34-9	NA	NA	NA		
Herbicides and Organopesticides						
2,4,5-T	93-76-5	NA	NA	NA		

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors				
Constituent of Potential Concern	CAS Registry Number	Source	$BSAF_{fish}$ (unitless)	Source		
2,4-D and esters	94-75-7	NA	NA	NA		
4,4-DDD	72-54-8	NA	NA	NA		
4,4-DDT	50-29-3	NA	NA	NA		
Aldrin	309-00-2	EPA6 e	NA	EPA6 e		
alpha-BHC	319-84-6	EPA6 e	NA	EPA6 e		
beta-BHC	319-85-7	EPA6 e	NA	EPA6 e		
Chlordane	57-74-9	EPA6 e	NA	EPA6 e		
DDE	72-55-9	EPA6 e	NA	EPA6 e		
Delta-BHC	319-86-8	NA	NA	NA		
Dieldrin	60-57-1	EPA6 e	NA	EPA6 e		
Endothall	145-73-3	NA	NA	NA		
Endrin	72-20-8	EPA6 e	NA	EPA6 e		
gamma-BHC(Lindane)	58-89-9	NA	NA	NA		
Heptachlor	76-44-8	EPA6 e	NA	EPA6 e		
Isodrin	465-73-6	NA	NA	NA		
Methoxychlor	72-43-5	EPA6 e	NA	EPA6 e		
Silvex (2,4,5-TP)	93-72-1	NA	NA	NA		
Toxaphene	8001-35-2	NA	NA	NA		

NA = Value not available
** = Varies with pH; values given for a pH of 7.0.
a = Illustrated Handbook of Physical - Chemical Properties and Environmental Fate f
b = RAIS database: http://risk.lsd.ornl.gov/rap_hp.shtml
c = Equations derived from EPA Region 6 Guidance Appendix A-3
d = Handbook of Environmental degradation Rates. 1991.
e = EPA Region 6 Hazardous Waste Combustion Guidance
f = "A Critical Review of Henry's Law Constants for Chemicals of Environmental In
g = Handbook of Environmental Fate and Exposure Data for Organic Chemicals
h = http://www.epa.gov/region09/waste/sfund/prg/.
i = http://esc.syrres.com/efdb/Chemfate.htm
j = Hansch and Leo 1985 in Syracuse 1996

Table B1-1. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Organic COPCs

		Bioconcentration / Bioaccumulation Factors				
Constituent of Potential Concern	CAS Registry Number	Source	$BSAF_{fish}$ (unitless)	Source		

k = Schwarzenbach et al. 1993
l = Tomlin 1994 in Syracuse 1996
m = http://www.speclab.com/compound/c959988.htm

Table B1-2. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Inorganic COPCs

		Mol. Wt.		Diffusivity				Partitioning Coefficients				
Constituent of Potential Concern	CAS Registry Number	MW (g/mol)	pH	D_a (cm^2/s)	Source	D_w (cm^2/s)	Source	Kd_s (cm^3/g)	Source	Kd_{sw} (L/kg)	Source	Kd_{bs} (cm^3/g)
Metals												
Aluminum	7429-90-5	26.98	NA	2.11E-01	EPA6 a	2.45E-05	EPA6 a	1.50E+03	EPA6 a	1.50E+03	EPA6 a	1.50E+03
Antimony	7440-36-0	121.75	6.8	7.73E-02	EPA6 a	8.96E-06	EPA6 a	4.50E+01	EPA6 a	4.50E+01	EPA6 a	4.50E+01
Arsenic*	7440-38-2	74.92	4.9	1.07E-01	EPA6 a	1.24E-05	EPA6 a	2.50E+01	EPA6 a	2.50E+01	EPA6 a	2.50E+01
Arsenic*	7440-38-2	74.92	6.8	1.07E-01	EPA6 a	1.24E-05	EPA6 a	2.90E+01	EPA6 a	2.90E+01	EPA6 a	2.90E+01
Arsenic*	7440-38-2	74.92	8	1.07E-01	EPA6 a	1.24E-05	EPA6 a	3.10E+01	EPA6 a	3.10E+01	EPA6 a	3.10E+01
Barium*	7440-39-3	137.33	4.9	7.14E-02	EPA6 a	8.27E-06	EPA6 a	1.10E+01	EPA6 a	1.10E+01	EPA6 a	1.10E+01
Barium*	7440-39-3	137.33	6.8	7.14E-02	EPA6 a	8.27E-06	EPA6 a	4.10E+01	EPA6 a	4.10E+01	EPA6 a	4.10E+01
Barium*	7440-39-3	137.33	8	7.14E-02	EPA6 a	8.27E-06	EPA6 a	5.20E+01	EPA6 a	5.20E+01	EPA6 a	5.20E+01
Beryllium*	7440-41-7	9.01	4.9	4.39E-01	EPA6 a	5.08E-05	EPA6 a	2.30E+01	EPA6 a	2.30E+01	EPA6 a	2.30E+01
Beryllium*	7440-41-7	9.01	6.8	4.39E-01	EPA6 a	5.08E-05	EPA6 a	7.90E+02	EPA6 a	7.90E+02	EPA6 a	7.90E+02
Beryllium*	7440-41-7	9.01	8	4.39E-01	EPA6 a	5.08E-05	EPA6 a	1.00E+05	EPA6 a	1.00E+05	EPA6 a	1.00E+05
Bismuth	7440-69-9	208.98	NA	5.40E-02	EPA6 calc d	6.25E-06	EPA6 calc d	2.00E+02	Baes b	2.00E+02	Baes b	2.00E+02
Boron	7440-42-8	10.81	NA	3.89E-01	EPA6 calc d	4.50E-05	EPA6 calc d	3.00E+00	Baes b	3.00E+00	Baes b	3.00E+00
Cadmium*	7440-43-9	112.41	4.9	8.16E-02	EPA6 a	9.45E-06	EPA6 a	1.50E+01	EPA6 a	1.50E+01	EPA6 a	1.50E+01
Cadmium*	7440-43-9	112.41	6.8	8.16E-02	EPA6 a	9.45E-06	EPA6 a	7.50E+01	EPA6 a	7.50E+01	EPA6 a	7.50E+01
Cadmium*	7440-43-9	112.41	8	8.16E-02	EPA6 a	9.45E-06	EPA6 a	4.30E+03	EPA6 a	4.30E+03	EPA6 a	4.30E+03
Calcium	7440-70-2	40.08	NA	1.62E-01	EPA6 calc d	1.88E-05	EPA6 calc d	4.00E+00	Baes b	4.00E+00	Baes b	4.00E+00
Chromium (and VI)*	18540-29-9	52.00	4.9	1.36E-01	EPA6 a	1.58E-05	EPA6 a	3.10E+01	EPA6 a	3.10E+01	EPA6 a	3.10E+01
Chromium (and VI)*	18540-29-9	52.00	6.8	1.36E-01	EPA6 a	1.58E-05	EPA6 a	1.90E+01	EPA6 a	1.90E+01	EPA6 a	1.90E+01
Chromium (and VI)*	18540-29-9	52.00	8	1.36E-01	EPA6 a	1.58E-05	EPA6 a	1.40E+01	EPA6 a	1.40E+01	EPA6 a	1.40E+01
Chromium (total)*	7440-47-3	52.00	4.9	1.01E-01	EPA6 a	4.63E-05	EPA6 a	1.20E+03	EPA6 a	1.20E+03	EPA6 a	1.20E+03
Chromium (total)*	7440-47-3	52.00	6.8	1.01E-01	EPA6 a	4.63E-05	EPA6 a	1.80E+06	EPA6 a	1.80E+06	EPA6 a	1.80E+06
Chromium (total)*	7440-47-3	52.00	8	1.01E-01	EPA6 a	4.63E-05	EPA6 a	4.30E+06	EPA6 a	4.30E+06	EPA6 a	4.30E+06
Cobalt	7440-48-4	58.93	NA	1.25E-01	EPA6 calc d	1.45E-05	EPA6 calc d	4.50E+01	Baes b	4.50E+01	Baes b	4.50E+01
Copper	7440-50-8	63.55	NA	1.19E-01	EPA6 calc d	1.38E-05	EPA6 calc d	3.50E+01	Baes b	3.50E+01	Baes b	3.50E+01
Iron	7439-89-6	55.85	NA	1.30E-01	EPA6 calc d	1.51E-05	EPA6 calc d	2.50E+01	Baes b	2.50E+01	Baes b	2.50E+01
Lead	7439-92-1	207.20	NA	5.43E-02	EPA6 a	6.28E-06	EPA6 a	9.00E+02	Baes b	9.00E+02	Baes b	9.00E+02

Table B1-2. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Inorganic COPCs

		Mol. Wt.		Diffusivity				Partitioning Coefficients				
Constituent of Potential Concern	CAS Registry Number	MW (g/mol)	pH	D_a (cm^2/s)	Source	D_w (cm^2/s)	Source	Kd_s (cm^3/g)	Source	Kd_{sw} (L/kg)	Source	Kd_{bs} (cm^3/g)
Lithium	7439-93-2	6.94	NA	5.22E-01	EPA6 calc d	6.05E-05	EPA6 calc d	3.00E+02	Baes b	3.00E+02	Baes b	3.00E+02
Magnesium	7439-95-4	24.31	NA	2.26E-01	EPA6 calc d	2.62E-05	EPA6 calc d	4.50E+00	Baes b	4.50E+00	Baes b	4.50E+00
Manganese	7439-96-5	54.94	NA	1.31E-01	EPA6 calc d	1.52E-05	EPA6 calc d	6.50E+01	Baes b	6.50E+01	Baes b	6.50E+01
Mercury	7439-97-6	200.59	NA	1.09E-02	EPA6 a	3.01E-05	EPA6 a	1.00E+03	EPA6 a	1.00E+03	EPA6 a	3.00E+03
Molybdenum	7439-98-7	95.94	NA	9.07E-02	EPA6 calc d	1.05E-05	EPA6 calc d	2.00E+01	Baes b	2.00E+01	Baes b	2.00E+01
Nickel*	7440-02-0	58.69	4.9	1.26E-01	EPA6 a	1.46E-05	EPA6 a	1.60E+01	EPA6 a	1.60E+01	EPA6 a	1.60E+01
Nickel*	7440-02-0	58.69	6.8	1.26E-01	EPA6 a	1.46E-05	EPA6 a	6.50E+01	EPA6 a	6.50E+01	EPA6 a	6.50E+01
Nickel*	7440-02-0	58.69	8	1.26E-01	EPA6 a	1.46E-05	EPA6 a	1.90E+03	EPA6 a	1.90E+03	EPA6 a	1.90E+03
Potassium	7440-09-7	39.10	NA	1.65E-01	EPA6 calc d	1.91E-05	EPA6 calc d	5.50E+00	Baes b	5.50E+00	Baes b	5.50E+00
Rhodium	7440-16-6	102.91	NA	8.65E-02	EPA6 calc d	1.00E-05	EPA6 calc d	6.00E+01	Baes b	6.00E+01	Baes b	6.00E+01
Selenium*	7782-49-2	78.96	4.9	1.03E-01	EPA6 a	1.20E-05	EPA6 a	1.80E+01	EPA6 a	1.80E+01	EPA6 a	1.80E+01
Selenium*	7782-49-2	78.96	6.8	1.03E-01	EPA6 a	1.20E-05	EPA6 a	5.00E+00	EPA6 a	5.00E+00	EPA6 a	5.00E+00
Selenium*	7782-49-2	78.96	8	1.03E-01	EPA6 a	1.20E-05	EPA6 a	2.20E+00	EPA6 a	2.20E+00	EPA6 a	2.20E+00
Silicon	7440-21-3	28.09	NA	2.06E-01	EPA6 calc d	2.38E-05	EPA6 calc d	3.00E+01	Baes b	3.00E+01	Baes b	3.00E+01
Silver*	7440-22-4	107.87	4.9	8.38E-02	EPA6 a	9.71E-06	EPA6 a	1.00E-01	EPA6 a	1.00E-01	EPA6 a	1.00E-01
Silver*	7440-22-4	107.87	6.8	8.38E-02	EPA6 a	9.71E-06	EPA6 a	8.30E+00	EPA6 a	8.30E+00	EPA6 a	8.30E+00
Silver*	7440-22-4	107.87	8	8.38E-02	EPA6 a	9.71E-06	EPA6 a	1.10E+02	EPA6 a	1.10E+02	EPA6 a	1.10E+02
Sodium	7440-23-5	22.99	NA	2.35E-01	EPA6 calc d	2.72E-05	EPA6 calc d	1.00E+02	Baes b	1.00E+02	Baes b	1.00E+02
Tantalum	7440-25-7	180.95	NA	5.94E-02	EPA6 calc d	6.88E-06	EPA6 calc d	6.50E+02	Baes b	6.50E+02	Baes b	6.50E+02
Thallium*	7440-28-0	204.38	4.9	5.48E-02	EPA6 a	6.34E-06	EPA6 a	4.40E+01	EPA6 a	4.40E+01	EPA6 a	4.40E+01
Thallium*	7440-28-0	204.38	6.8	5.48E-02	EPA6 a	6.34E-06	EPA6 a	7.10E+01	EPA6 a	7.10E+01	EPA6 a	7.10E+01
Thallium*	7440-28-0	204.38	8	5.48E-02	EPA6 a	6.34E-06	EPA6 a	9.60E+01	EPA6 a	9.60E+01	EPA6 a	9.60E+01
Tin	7440-31-5	118.71	NA	7.87E-02	EPA6 calc d	9.11E-06	EPA6 calc d	2.50E+02	Baes b	2.50E+02	Baes b	2.50E+02
Tungsten	7440-33-7	183.84	NA	5.88E-02	EPA6 calc d	6.80E-06	EPA6 calc d	1.50E+02	Baes b	1.50E+02	Baes b	1.50E+02
Uranium	7440-61-1	238.03	NA	4.95E-02	EPA6 calc d	5.73E-06	EPA6 calc d	4.50E+02	Baes b	4.50E+02	Baes b	4.50E+02
Vanadium	7440-62-2	50.94	NA	1.38E-01	EPA6 calc d	1.60E-05	EPA6 calc d	1.00E+03	Baes b	1.00E+03	Baes b	1.00E+03
Yttrium	7440-65-5	88.91	NA	9.54E-02	EPA6 calc d	1.10E-05	EPA6 calc d	5.00E+02	Baes b	5.00E+02	Baes b	5.00E+02
Zinc	7440-66-6	65.38	6.8	1.17E-01	EPA6 a	1.36E-05	EPA6 a	6.20E+01	EPA6 a	6.20E+01	EPA6 a	6.20E+01

Table B1-2. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Inorganic COPCs

		Mol. Wt.		Diffusivity				Partitioning Coefficients				
Constituent of Potential Concern	CAS Registry Number	MW (g/mol)	pH	D_a (cm^2/s)	Source	D_w (cm^2/s)	Source	Kd_s (cm^3/g)	Source	Kd_{sw} (L/kg)	Source	Kd_{bs} (cm^3/g)
Zirconium	7440-67-7	91.22	NA	9.38E-02	EPA6 calc d	1.09E-05	EPA6 calc d	3.00E+03	Baes b	3.00E+03	Baes b	3.00E+03
Non-Metal and Anions												
Ammonia/Ammonium	7664-41-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Bromide	24959-67-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Chloride	16887-00-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Cyanide	57-12-5	26.02	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Fluoride	16984-48-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Hydroxide	14280-30-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Iodine	7553-56-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Nitrate	14797-55-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Nitrite	14797-65-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Phosphate	14265-44-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Sulfate	14808-79-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Criteria Pollutants												
Carbon dioxide	124-38-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Nitrogen dioxide	10102-44-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Ozone	10028-15-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Particulate matter	no cas #	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Sulfur dioxide	7446-09-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

* These metals have varying dissociation constants depending on pH.
a = EPA Region 6 Hazardous Waste Combustion Facility Guidance
b = Baes et al
c = Ng et al
d = Calculation from EPA Region 6 Hazardous Waste Combustion Facility Guidance
e = Br_{ag} is a weighted average of $Br_{ag(fruit)}$ and $Br_{ag(veg)}$

Table B1-2. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Inorganic COPCs

		Plant Uptake Factors							
Constituent of Potential Concern	CAS Registry Number	Source	$Br_{root\ veg}$ ((ug/g DW plant) /(ug/g soil))	Source	$Br_{ag(fruit)}$ ((ug/g DW plant) /(ug/g soil))	Source	$Br_{ag(veg)}$ ((ug/g DW plant) /(ug/g soil))[e]	Source	Br_{ag} ((ug/g DW plant) /(ug/g soil))
Metals									
Aluminum	7429-90-5	EPA6 a	6.50E-04	Baes b	6.50E-04	Baes b	1.49E-03	EPA calc d	1.08E-03
Antimony	7440-36-0	EPA6 a	3.00E-02	Baes b	3.00E-02	Baes b	7.25E-02	EPA calc d	5.16E-02
Arsenic*	7440-38-2	EPA6 a	8.00E-03	EPA6 a	NA	NA	NA	NA	6.33E-03
Arsenic*	7440-38-2	EPA6 a	8.00E-03	EPA6 a	NA	NA	NA	NA	6.33E-03
Arsenic*	7440-38-2	EPA6 a	8.00E-03	EPA6 a	NA	NA	NA	NA	6.33E-03
Barium*	7440-39-3	EPA6 a	1.50E-02	EPA6 a	1.50E-02	Baes b	4.88E-02	EPA calc d	3.22E-02
Barium*	7440-39-3	EPA6 a	1.50E-02	EPA6 a	1.50E-02	Baes b	4.88E-02	EPA calc d	3.22E-02
Barium*	7440-39-3	EPA6 a	1.50E-02	EPA6 a	1.50E-02	Baes b	4.88E-02	EPA calc d	3.22E-02
Beryllium*	7440-41-7	EPA6 a	1.50E-03	EPA6 a	1.50E-03	EPA6 a	3.63E-03	EPA calc d	2.58E-03
Beryllium*	7440-41-7	EPA6 a	1.50E-03	EPA6 a	1.50E-03	EPA6 a	3.63E-03	EPA calc d	2.58E-03
Beryllium*	7440-41-7	EPA6 a	1.50E-03	EPA6 a	1.50E-03	EPA6 a	3.63E-03	EPA calc d	2.58E-03
Bismuth	7440-69-9	Baes b	5.00E-03	Baes b	5.00E-03	Baes b	1.25E-02	EPA calc d	8.81E-03
Boron	7440-42-8	Baes b	2.00E+00	Baes b	2.00E+00	Baes b	2.50E+00	EPA calc d	2.25E+00
Cadmium*	7440-43-9	EPA6 a	6.40E-02	EPA6 a	NA	NA	NA	NA	1.25E-01
Cadmium*	7440-43-9	EPA6 a	6.40E-02	EPA6 a	NA	NA	NA	NA	1.25E-01
Cadmium*	7440-43-9	EPA6 a	6.40E-02	EPA6 a	NA	NA	NA	NA	1.25E-01
Calcium	7440-70-2	Baes b	3.50E-01	Baes b	3.50E-01	Baes b	1.14E+00	EPA calc d	7.50E-01
Chromium (and VI)*	18540-29-9	EPA6 a	4.50E-03	Baes b	4.50E-03	Baes b	5.25E-03	EPA calc d	4.88E-03
Chromium (and VI)*	18540-29-9	EPA6 a	4.50E-03	Baes b	4.50E-03	Baes b	5.25E-03	EPA calc d	4.88E-03
Chromium (and VI)*	18540-29-9	EPA6 a	4.50E-03	Baes b	4.50E-03	Baes b	5.25E-03	EPA calc d	4.88E-03
Chromium (total)*	7440-47-3	EPA6 a	4.50E-03	EPA6 a	NA	NA	NA	NA	NA
Chromium (total)*	7440-47-3	EPA6 a	4.50E-03	EPA6 a	NA	NA	NA	NA	NA
Chromium (total)*	7440-47-3	EPA6 a	4.50E-03	EPA6 a	NA	NA	NA	NA	NA
Cobalt	7440-48-4	Baes b	7.00E-02	Baes b	7.00E-02	Baes b	5.75E-02	EPA calc d	6.36E-02
Copper	7440-50-8	Baes b	2.50E-01	Baes b	2.50E-01	Baes b	2.88E-01	EPA calc d	2.69E-01
Iron	7439-89-6	Baes b	1.00E-03	Baes b	1.00E-03	Baes b	1.75E-03	EPA calc d	1.38E-03
Lead	7439-92-1	Baes b	9.00E-03	Baes b	9.00E-03	Baes b	1.80E-02	EPA calc d	1.36E-02

Table B1-2. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Inorganic COPCs

		Plant Uptake Factors							
Constituent of Potential Concern	CAS Registry Number	Source	$Br_{root\ veg}$ ((ug/g DW plant) /(ug/g soil))	Source	$Br_{ag(fruit)}$ ((ug/g DW plant) /(ug/g soil))	Source	$Br_{ag(veg)}$ ((ug/g DW plant) /(ug/g soil))[e]	Source	Br_{ag} ((ug/g DW plant) /(ug/g soil))
Lithium	7439-93-2	Baes b	4.00E-03	Baes b	4.00E-03	Baes b	9.25E-03	EPA calc d	6.67E-03
Magnesium	7439-95-4	Baes b	5.50E-01	Baes b	5.50E-01	Baes b	6.63E-01	EPA calc d	6.07E-01
Manganese	7439-96-5	Baes b	5.00E-02	Baes b	5.00E-02	Baes b	1.00E-01	EPA calc d	7.54E-02
Mercury	7439-97-6	EPA6 a	2.00E-01	Baes b	2.00E-01	Baes b	3.75E-01	EPA calc d	2.89E-01
Molybdenum	7439-98-7	Baes b	6.00E-02	Baes b	6.00E-02	Baes b	1.08E-01	EPA calc d	8.42E-02
Nickel*	7440-02-0	EPA6 a	8.00E-03	EPA6 a	NA	NA	NA	NA	9.31E-03
Nickel*	7440-02-0	EPA6 a	8.00E-03	EPA6 a	NA	NA	NA	NA	9.31E-03
Nickel*	7440-02-0	EPA6 a	8.00E-03	EPA6 a	NA	NA	NA	NA	9.31E-03
Potassium	7440-09-7	Baes b	5.50E-01	Baes b	5.50E-01	Baes b	6.63E-01	EPA calc d	6.07E-01
Rhodium	7440-16-6	Baes b	4.00E-02	Baes b	4.00E-02	Baes b	6.75E-02	EPA calc d	5.40E-02
Selenium*	7782-49-2	EPA6 a	2.20E-02	EPA6 a	NA	NA	NA	NA	1.95E-02
Selenium*	7782-49-2	EPA6 a	2.20E-02	EPA6 a	NA	NA	NA	NA	1.95E-02
Selenium*	7782-49-2	EPA6 a	2.20E-02	EPA6 a	NA	NA	NA	NA	1.95E-02
Silicon	7440-21-3	Baes b	7.00E-02	Baes b	7.00E-02	Baes b	1.40E-01	EPA calc d	1.06E-01
Silver*	7440-22-4	EPA6 a	1.00E-01	EPA6 a	1.00E-01	Baes b	1.75E-01	EPA calc d	1.38E-01
Silver*	7440-22-4	EPA6 a	1.00E-01	EPA6 a	1.00E-01	Baes b	1.75E-01	EPA calc d	1.38E-01
Silver*	7440-22-4	EPA6 a	1.00E-01	EPA6 a	1.00E-01	Baes b	1.75E-01	EPA calc d	1.38E-01
Sodium	7440-23-5	Baes b	5.50E-02	Baes b	5.50E-02	Baes b	6.00E-02	EPA calc d	5.75E-02
Tantalum	7440-25-7	Baes b	2.50E-03	Baes b	2.50E-03	Baes b	4.38E-03	EPA calc d	3.45E-03
Thallium*	7440-28-0	EPA6 a	4.00E-04	EPA6 a	4.00E-04	Baes b	1.30E-03	EPA calc d	8.58E-04
Thallium*	7440-28-0	EPA6 a	4.00E-04	EPA6 a	4.00E-04	Baes b	1.30E-03	EPA calc d	8.58E-04
Thallium*	7440-28-0	EPA6 a	4.00E-04	EPA6 a	4.00E-04	Baes b	1.30E-03	EPA calc d	8.58E-04
Tin	7440-31-5	Baes b	6.00E-03	Baes b	6.00E-03	Baes b	1.20E-02	EPA calc d	9.05E-03
Tungsten	7440-33-7	Baes b	1.00E-02	Baes b	1.00E-02	Baes b	1.88E-02	EPA calc d	1.25E-02
Uranium	7440-61-1	Baes b	4.00E-03	Baes b	4.00E-03	Baes b	5.13E-03	EPA calc d	4.57E-03
Vanadium	7440-62-2	Baes b	3.00E-03	Baes b	3.00E-03	Baes b	3.63E-03	EPA calc d	3.32E-03
Yttrium	7440-65-5	Baes b	6.00E-03	Baes b	6.00E-03	Baes b	8.25E-03	EPA calc d	7.14E-03
Zinc	7440-66-6	EPA6 a	4.40E-02	Baes b	4.60E-02	EPA6 a	9.70E-02	EPA6 a	7.19E-02

Table B1-2. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Inorganic COPCs

		Plant Uptake Factors							
Constituent of Potential Concern	CAS Registry Number	Source	$Br_{root\ veg}$ ((ug/g DW plant) /(ug/g soil))	Source	$Br_{ag(fruit)}$ ((ug/g DW plant) /(ug/g soil))	Source	$Br_{ag(veg)}$ ((ug/g DW plant) /(ug/g soil))[e]	Source	Br_{ag} ((ug/g DW plant) /(ug/g soil))
Zirconium	7440-67-7	Baes b	5.00E-04	Baes b	5.00E-04	Baes b	8.75E-04	EPA calc d	1.02E-01
Non-Metal and Anions									
Ammonia/Ammonium	7664-41-7	NA	NA	NA	NA	NA	NA	NA	NA
Bromide	24959-67-9	NA	NA	NA	NA	NA	NA	NA	NA
Chloride	16887-00-6	NA	NA	NA	NA	NA	NA	NA	NA
Cyanide	57-12-5	NA	NA	NA	NA	NA	NA	NA	NA
Fluoride	16984-48-8	NA	NA	NA	NA	NA	NA	NA	NA
Hydroxide	14280-30-9	NA	NA	NA	NA	NA	NA	NA	NA
Iodine	7553-56-2	NA	NA	NA	NA	NA	NA	NA	NA
Nitrate	14797-55-8	NA	NA	NA	NA	NA	NA	NA	NA
Nitrite	14797-65-0	NA	NA	NA	NA	NA	NA	NA	NA
Phosphate	14265-44-2	NA	NA	NA	NA	NA	NA	NA	NA
Sulfate	14808-79-8	NA	NA	NA	NA	NA	NA	NA	NA
Criteria Pollutants									
Carbon dioxide	124-38-9	NA	NA	NA	NA	NA	NA	NA	NA
Nitrogen dioxide	10102-44-0	NA	NA	NA	NA	NA	NA	NA	NA
Ozone	10028-15-6	NA	NA	NA	NA	NA	NA	NA	NA
Particulate matter	no cas #	NA	NA	NA	NA	NA	NA	NA	NA
Sulfur dioxide	7446-09-5	NA	NA	NA	NA	NA	NA	NA	NA

* These metals have varying dissociation cc
a = EPA Region 6 Hazardous Waste Combı
b = Baes et al
c = Ng et al
d = Calculation from EPA Region 6 Hazard
e = Br_{ag} is a weighted average of $Br_{ag(fruit)}$ a

Table B1-2. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Inorganic COPCs

		Plant Uptake Factors					Bioaccumulation/Biotransfer Factors				
Constituent of Potential Concern	CAS Registry Number	Source	Br_{forage} ((ug/g DW plant) /(ug/g soil))	Source	Br_{grain} ((ug/g DW plant) /(ug/g soil))	Source	Ba_{milk} (day/kg FW)	Source	Ba_{beef} (day/kg FW)	Source	Ba_{pork} (day/kg FW)
Metals											
Aluminum	7429-90-5	EPA6 calc d	4.00E-03	Baes b	6.50E-04	Baes b	2.00E-04	Baes b	1.50E-03	Baes b	NA
Antimony	7440-36-0	EPA6 calc d	2.00E-01	Baes b	3.00E-02	Baes b	1.00E-04	Baes b	1.00E-03	Baes b	NA
Arsenic*	7440-38-2	EPA6 a	3.60E-02	EPA6 a	4.00E-03	EPA6 a	6.00E-03	Baes b	2.00E-03	Baes b	NA
Arsenic*	7440-38-2	EPA6 a	3.60E-02	EPA6 a	4.00E-03	EPA6 a	6.00E-03	Baes b	2.00E-03	Baes b	NA
Arsenic*	7440-38-2	EPA6 a	3.60E-02	EPA6 a	4.00E-03	EPA6 a	6.00E-03	Baes b	2.00E-03	Baes b	NA
Barium*	7440-39-3	EPA6 calc d	1.50E-01	EPA6 a	1.50E-02	EPA6 a	3.50E-04	EPA6 a	1.50E-04	EPA6 a	NA
Barium*	7440-39-3	EPA6 calc d	1.50E-01	EPA6 a	1.50E-02	EPA6 a	3.50E-04	EPA6 a	1.50E-04	EPA6 a	NA
Barium*	7440-39-3	EPA6 calc d	1.50E-01	EPA6 a	1.50E-02	EPA6 a	3.50E-04	EPA6 a	1.50E-04	EPA6 a	NA
Beryllium*	7440-41-7	EPA6 calc d	1.00E-02	EPA6 a	1.50E-03	EPA6 a	9.00E-07	EPA6 a	1.00E-03	EPA6 a	NA
Beryllium*	7440-41-7	EPA6 calc d	1.00E-02	EPA6 a	1.50E-03	EPA6 a	9.00E-07	EPA6 a	1.00E-03	EPA6 a	NA
Beryllium*	7440-41-7	EPA6 calc d	1.00E-02	EPA6 a	1.50E-03	EPA6 a	9.00E-07	EPA6 a	1.00E-03	EPA6 a	NA
Bismuth	7440-69-9	EPA6 calc d	3.50E-02	Baes b	5.00E-03	Baes b	5.00E-04	Baes b	4.00E-04	Baes b	NA
Boron	7440-42-8	EPA6 calc d	4.00E+00	Baes b	2.00E+00	Baes b	1.50E-03	Baes b	8.00E-04	Baes b	NA
Cadmium*	7440-43-9	EPA6 a	3.64E-01	EPA6 a	6.20E-02	EPA6 a	6.50E-06	EPA6 a	1.20E-04	EPA6 a	1.91E-04
Cadmium*	7440-43-9	EPA6 a	3.64E-01	EPA6 a	6.20E-02	EPA6 a	6.50E-06	EPA6 a	1.20E-04	EPA6 a	1.91E-04
Cadmium*	7440-43-9	EPA6 a	3.64E-01	EPA6 a	6.20E-02	EPA6 a	6.50E-06	EPA6 a	1.20E-04	EPA6 a	1.91E-04
Calcium	7440-70-2	EPA6 calc d	3.50E+00	Baes b	3.50E-01	Baes b	1.00E-02	Baes b	7.00E-04	Baes b	NA
Chromium (and VI)*	18540-29-9	EPA6 calc d	7.50E-03	Baes b	4.50E-03	Baes b	1.50E-03	Baes b	5.50E-03	Baes b	NA
Chromium (and VI)*	18540-29-9	EPA6 calc d	7.50E-03	Baes b	4.50E-03	Baes b	1.50E-03	Baes b	5.50E-03	Baes b	NA
Chromium (and VI)*	18540-29-9	EPA6 calc d	7.50E-03	Baes b	4.50E-03	Baes b	1.50E-03	Baes b	5.50E-03	Baes b	NA
Chromium (total)*	7440-47-3	NA	7.50E-03	EPA6 a	4.50E-03	EPA6 a	1.50E-03	EPA6 a	5.50E-03	EPA6 a	NA
Chromium (total)*	7440-47-3	NA	7.50E-03	EPA6 a	4.50E-03	EPA6 a	1.50E-03	EPA6 a	5.50E-03	EPA6 a	NA
Chromium (total)*	7440-47-3	NA	7.50E-03	EPA6 a	4.50E-03	EPA6 a	1.50E-03	EPA6 a	5.50E-03	EPA6 a	NA
Cobalt	7440-48-4	EPA6 calc d	2.00E-02	Baes b	7.00E-02	Baes b	2.00E-03	Baes b	2.00E-02	Baes b	1.70E-01
Copper	7440-50-8	EPA6 calc d	4.00E-01	Baes b	2.50E-01	Baes b	1.50E-03	Baes b	1.00E-02	Baes b	2.20E-02
Iron	7439-89-6	EPA6 calc d	4.00E-03	Baes b	1.00E-03	Baes b	2.50E-04	Baes b	2.00E-02	Baes b	2.60E-02
Lead	7439-92-1	EPA6 calc d	4.50E-02	Baes b	9.00E-03	Baes b	2.50E-04	Baes b	3.00E-04	Baes b	NA

Table B1-2. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Inorganic COPCs

		Plant Uptake Factors					Bioaccumulation/Biotransfer Factors				
Constituent of Potential Concern	CAS Registry Number	Source	Br_{forage} ((ug/g DW plant) /(ug/g soil))	Source	Br_{grain} ((ug/g DW plant) /(ug/g soil))	Source	Ba_{milk} (day/kg FW)	Source	Ba_{beef} (day/kg FW)	Source	Ba_{pork} (day/kg FW)
Lithium	7439-93-2	EPA6 calc d	2.50E-02	Baes b	4.00E-03	Baes b	2.00E-02	Baes b	1.00E-02	Baes b	NA
Magnesium	7439-95-4	EPA6 calc d	1.00E+00	Baes b	5.50E-01	Baes b	4.00E-03	Baes b	5.00E-03	Baes b	NA
Manganese	7439-96-5	EPA6 calc d	2.50E-01	Baes b	5.00E-02	Baes b	3.50E-04	Baes b	4.00E-04	Baes b	3.60E-03
Mercury	7439-97-6	EPA6 calc d	9.00E-01	Baes b	2.00E-01	Baes b	4.50E-04	Baes b	2.50E-01	Baes b	NA
Molybdenum	7439-98-7	EPA6 calc d	2.50E-01	Baes b	6.00E-02	Baes b	1.50E-03	Baes b	6.00E-03	Baes b	NA
Nickel*	7440-02-0	EPA6 a	3.20E-02	EPA6 a	6.00E-03	EPA6 a	1.00E-03	EPA6 a	6.00E-03	EPA6 a	NA
Nickel*	7440-02-0	EPA6 a	3.20E-02	EPA6 a	6.00E-03	EPA6 a	1.00E-03	EPA6 a	6.00E-03	EPA6 a	NA
Nickel*	7440-02-0	EPA6 a	3.20E-02	EPA6 a	6.00E-03	EPA6 a	1.00E-03	EPA6 a	6.00E-03	EPA6 a	NA
Potassium	7440-09-7	EPA6 calc d	1.00E+00	Baes b	5.50E-01	Baes b	7.00E-03	Baes b	2.00E-02	Baes b	NA
Rhodium	7440-16-6	EPA6 calc d	1.50E-01	Baes b	4.00E-02	Baes b	1.00E-02	Baes b	2.00E-03	Baes b	NA
Selenium*	7782-49-2	EPA6 a	1.60E-02	EPA6 a	2.00E-03	EPA6 a	5.86E-03	EPA6 a	2.27E-03	EPA6 a	3.70E-03
Selenium*	7782-49-2	EPA6 a	1.60E-02	EPA6 a	2.00E-03	EPA6 a	5.86E-03	EPA6 a	2.27E-03	EPA6 a	3.70E-03
Selenium*	7782-49-2	EPA6 a	1.60E-02	EPA6 a	2.00E-03	EPA6 a	5.86E-03	EPA6 a	2.27E-03	EPA6 a	3.70E-03
Silicon	7440-21-3	EPA6 calc d	3.50E-01	Baes b	7.00E-02	Baes b	2.00E-05	Baes b	4.00E-05	Baes b	NA
Silver*	7440-22-4	EPA6 calc d	4.00E-01	EPA6 a	1.00E-01	EPA6 a	2.00E-02	EPA6 a	3.00E-03	EPA6 a	NA
Silver*	7440-22-4	EPA6 calc d	4.00E-01	EPA6 a	1.00E-01	EPA6 a	2.00E-02	EPA6 a	3.00E-03	EPA6 a	NA
Silver*	7440-22-4	EPA6 calc d	4.00E-01	EPA6 a	1.00E-01	EPA6 a	2.00E-02	EPA6 a	3.00E-03	EPA6 a	NA
Sodium	7440-23-5	EPA6 calc d	7.50E-02	Baes b	5.50E-02	Baes b	3.50E-02	Baes b	5.50E-02	Baes b	NA
Tantalum	7440-25-7	EPA6 calc d	1.00E-02	Baes b	2.50E-03	Baes b	3.00E-04	Baes b	6.00E-04	Baes b	NA
Thallium*	7440-28-0	EPA6 calc d	4.00E-03	EPA6 a	4.00E-04	EPA6 a	2.00E-03	EPA6 a	4.00E-02	EPA6 a	NA
Thallium*	7440-28-0	EPA6 calc d	4.00E-03	EPA6 a	4.00E-04	EPA6 a	2.00E-03	EPA6 a	4.00E-02	EPA6 a	NA
Thallium*	7440-28-0	EPA6 calc d	4.00E-03	EPA6 a	4.00E-04	EPA6 a	2.00E-03	EPA6 a	4.00E-02	EPA6 a	NA
Tin	7440-31-5	EPA6 calc d	3.00E-02	Baes b	6.00E-03	Baes b	1.00E-03	Baes b	8.00E-02	Baes b	NA
Tungsten	7440-33-7	EPA6 calc d	4.50E-02	Baes b	1.00E-02	Baes b	3.00E-04	Baes b	4.50E-02	Baes b	NA
Uranium	7440-61-1	EPA6 calc d	8.50E-03	Baes b	4.00E-03	Baes b	6.00E-04	Baes b	2.00E-04	Baes b	4.00E-02
Vanadium	7440-62-2	EPA6 calc d	5.50E-03	Baes b	3.00E-03	Baes b	2.00E-05	Baes b	2.50E-03	Baes b	NA
Yttrium	7440-65-5	EPA6 calc d	1.50E-02	Baes b	6.00E-03	Baes b	2.00E-05	Baes b	3.00E-04	Baes b	NA
Zinc	7440-66-6	EPA6 calc d	2.50E-01	EPA6 a	5.40E-02	EPA6 a	3.25E-05	EPA6 a	9.00E-05	EPA6 a	1.28E-04

Table B1-2. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Inorganic COPCs

		Plant Uptake Factors					Bioaccumulation/Biotransfer Factors				
Constituent of Potential Concern	CAS Registry Number	Source	Br_{forage} ((ug/g DW plant) /(ug/g soil))	Source	Br_{grain} ((ug/g DW plant) /(ug/g soil))	Source	Ba_{milk} (day/kg FW)	Source	Ba_{beef} (day/kg FW)	Source	Ba_{pork} (day/kg FW)
Zirconium	7440-67-7	EPA6 calc d	2.00E-03	Baes b	5.00E-04	Baes b	3.00E-05	Baes b	5.50E-03	Baes b	NA
Non-Metal and Anions											
Ammonia/Ammonium	7664-41-7	NA	NA	NA	NA	NA	7.90E-09	NA	NA	NA	NA
Bromide	24959-67-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Chloride	16887-00-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Cyanide	57-12-5	NA	NA	NA	NA	NA	9.90E-08	NA	NA	NA	NA
Fluoride	16984-48-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Hydroxide	14280-30-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Iodine	7553-56-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Nitrate	14797-55-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Nitrite	14797-65-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Phosphate	14265-44-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Sulfate	14808-79-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Criteria Pollutants											
Carbon dioxide	124-38-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Nitrogen dioxide	10102-44-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Ozone	10028-15-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Particulate matter	no cas #	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Sulfur dioxide	7446-09-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

* These metals have varying dissociation cc
a = EPA Region 6 Hazardous Waste Combı
b = Baes et al
c = Ng et al
d = Calculation from EPA Region 6 Hazard
e = Br_{ag} is a weighted average of $Br_{ag(fruit)}$ a

Table B1-2. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Inorganic COPCs

		Bioaccumulation/Biotransfer Factors										
Constituent of Potential Concern	CAS Registry Number	Source	Ba_{egg} (day/kg FW)	Source	$Ba_{chicken}$ (day/kg FW)	Source	BCF_{fish} (L/kg FW tissue)	Source	BAF_{fish} (L/kg FW)	Source	$BSAF_{fish}$ (unitless)	Source
Metals												
Aluminum	7429-90-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Antimony	7440-36-0	NA	NA	NA	NA	NA	4.00E+01	EPA6 a	NA	NA	NA	NA
Arsenic*	7440-38-2	NA	NA	NA	NA	NA	2.00E+01	EPA6 a	NA	NA	NA	NA
Arsenic*	7440-38-2	NA	NA	NA	NA	NA	2.00E+01	EPA6 a	NA	NA	NA	NA
Arsenic*	7440-38-2	NA	NA	NA	NA	NA	2.00E+01	EPA6 a	NA	NA	NA	NA
Barium*	7440-39-3	NA	9.00E-01	Ng c	2.00E-02	Ng c	NA	NA	NA	NA	NA	NA
Barium*	7440-39-3	NA	9.00E-01	Ng c	2.00E-02	Ng c	NA	NA	NA	NA	NA	NA
Barium*	7440-39-3	NA	9.00E-01	Ng c	2.00E-02	Ng c	NA	NA	NA	NA	NA	NA
Beryllium*	7440-41-7	NA	NA	NA	NA	NA	4.20E+01	EPA6 a	NA	NA	NA	NA
Beryllium*	7440-41-7	NA	NA	NA	NA	NA	4.20E+01	EPA6 a	NA	NA	NA	NA
Beryllium*	7440-41-7	NA	NA	NA	NA	NA	4.20E+01	EPA6 a	NA	NA	NA	NA
Bismuth	7440-69-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Boron	7440-42-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Cadmium*	7440-43-9	EPA6 a	2.50E-03	EPA6 a	1.06E-01	EPA6 a	2.50E+02	EPA6 a	NA	NA	NA	NA
Cadmium*	7440-43-9	EPA6 a	2.50E-03	EPA6 a	1.06E-01	EPA6 a	2.50E+02	EPA6 a	NA	NA	NA	NA
Cadmium*	7440-43-9	EPA6 a	2.50E-03	EPA6 a	1.06E-01	EPA6 a	2.50E+02	EPA6 a	NA	NA	NA	NA
Calcium	7440-70-2	NA	4.40E-02	Ng c	4.40E-02	Ng c	NA	NA	NA	NA	NA	NA
Chromium (and VI)*	18540-29-9	NA	NA	NA	NA	NA	3.00E+00	EPA6 a	NA	NA	NA	NA
Chromium (and VI)*	18540-29-9	NA	NA	NA	NA	NA	3.00E+00	EPA6 a	NA	NA	NA	NA
Chromium (and VI)*	18540-29-9	NA	NA	NA	NA	NA	3.00E+00	EPA6 a	NA	NA	NA	NA
Chromium (total)*	7440-47-3	NA	NA	NA	NA	NA	2.83E+02	EPA6 a	NA	NA	NA	NA
Chromium (total)*	7440-47-3	NA	NA	NA	NA	NA	2.83E+02	EPA6 a	NA	NA	NA	NA
Chromium (total)*	7440-47-3	NA	NA	NA	NA	NA	2.83E+02	EPA6 a	NA	NA	NA	NA
Cobalt	7440-48-4	Ng c	2.00E-02	Ng c	5.30E-01	Ng c	NA	NA	3.00E+02	NA	NA	NA
Copper	7440-50-8	Ng c	4.90E-01	Ng c	5.10E-01	Ng c	NA	NA	2.00E+02	NA	NA	NA
Iron	7439-89-6	Ng c	1.30E+00	Ng c	1.50E+00	Ng c	NA	NA	2.00E+02	NA	NA	NA
Lead	7439-92-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

Table B1-2. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Inorganic COPCs

		Bioaccumulation/Biotransfer Factors										
Constituent of Potential Concern	CAS Registry Number	Source	Ba_{egg} (day/kg FW)	Source	$Ba_{chicken}$ (day/kg FW)	Source	BCF_{fish} (L/kg FW tissue)	Source	BAF_{fish} (L/kg FW)	Source	$BSAF_{fish}$ (unitless)	Source
Lithium	7439-93-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Magnesium	7439-95-4	NA	1.60E+00	Ng c	NA	NA	NA	NA	NA	NA	NA	NA
Manganese	7439-96-5	Ng c	6.50E-02	Ng c	5.10E-02	Ng c	NA	NA	NA	NA	NA	NA
Mercury	7439-97-6	NA	NA	NA	2.70E-02	Ng c	NA	NA	NA	NA	NA	NA
Molybdenum	7439-98-7	NA	5.00E-01	Ng c	5.00E-02	Ng c	NA	NA	NA	NA	NA	NA
Nickel*	7440-02-0	NA	NA	NA	NA	NA	3.07E+02	EPA6 a	NA	NA	NA	NA
Nickel*	7440-02-0	NA	NA	NA	NA	NA	3.07E+02	EPA6 a	NA	NA	NA	NA
Nickel*	7440-02-0	NA	NA	NA	NA	NA	3.07E+02	EPA6 a	NA	NA	NA	NA
Potassium	7440-09-7	NA	1.10E+00	Ng c	NA	NA	NA	NA	NA	NA	NA	NA
Rhodium	7440-16-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Selenium*	7782-49-2	EPA6 a	1.13E+00	EPA6 a	1.13E+00	EPA6 a	1.29E+02	EPA6 a	NA	NA	NA	NA
Selenium*	7782-49-2	EPA6 a	1.13E+00	EPA6 a	1.13E+00	EPA6 a	1.29E+02	EPA6 a	NA	NA	NA	NA
Selenium*	7782-49-2	EPA6 a	1.13E+00	EPA6 a	1.13E+00	EPA6 a	1.29E+02	EPA6 a	NA	NA	NA	NA
Silicon	7440-21-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Silver*	7440-22-4	NA	NA	NA	NA	NA	2.04E+02	EPA6 a	NA	NA	NA	NA
Silver*	7440-22-4	NA	NA	NA	NA	NA	2.04E+02	EPA6 a	NA	NA	NA	NA
Silver*	7440-22-4	NA	NA	NA	NA	NA	2.04E+02	EPA6 a	NA	NA	NA	NA
Sodium	7440-23-5	NA	6.10E+00	Ng c	NA	Ng c	NA	NA	NA	NA	NA	NA
Tantalum	7440-25-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Thallium*	7440-28-0	NA	NA	NA	NA	NA	1.40E+03	EPA6 a	NA	NA	NA	NA
Thallium*	7440-28-0	NA	NA	NA	NA	NA	1.40E+03	EPA6 a	NA	NA	NA	NA
Thallium*	7440-28-0	NA	NA	NA	NA	NA	1.40E+03	EPA6 a	NA	NA	NA	NA
Tin	7440-31-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Tungsten	7440-33-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Uranium	7440-61-1	Ng c	9.90E-01	Ng c	1.20E+00	Ng c	NA	NA	NA	NA	NA	NA
Vanadium	7440-62-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Yttrium	7440-65-5	NA	2.00E-03	Ng c	1.00E-02	Ng c	NA	NA	NA	NA	NA	NA
Zinc	7440-66-6	EPA6 a	8.75E-03	EPA6 a	8.75E-03	EPA6 a	NA	NA	NA	NA	NA	NA

Table B1-2. Physical/Chemical Parameters for Exposure Point Concentration Modelling for Inorganic COPCs

		Bioaccumulation/Biotransfer Factors										
Constituent of Potential Concern	CAS Registry Number	Source	Ba_{egg} (day/kg FW)	Source	$Ba_{chicken}$ (day/kg FW)	Source	BCF_{fish} (L/kg FW tissue)	Source	BAF_{fish} (L/kg FW)	Source	$BSAF_{fish}$ (unitless)	Source
Zirconium	7440-67-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Non-Metal and Anions												
Ammonia/Ammonium	7664-41-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Bromide	24959-67-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Chloride	16887-00-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Cyanide	57-12-5	NA	NA	NA	NA	NA	NA	NA	6.33E+02	NA	NA	NA
Fluoride	16984-48-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Hydroxide	14280-30-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Iodine	7553-56-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Nitrate	14797-55-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Nitrite	14797-65-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Phosphate	14265-44-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Sulfate	14808-79-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Criteria Pollutants												
Carbon dioxide	124-38-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Nitrogen dioxide	10102-44-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Ozone	10028-15-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Particulate matter	no cas #	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Sulfur dioxide	7446-09-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

* These metals have varying dissociation cc
a = EPA Region 6 Hazardous Waste Combı
b = Baes et al
c = Ng et al
d = Calculation from EPA Region 6 Hazard
e = Br_{ag} is a weighted average of $Br_{ag(fruit)}$ a

Table B1-3. Physical/Chemical Parameters for Exposure Point Concentration Modelling for ROPCs

			Diffusivity				Partitioning Coefficients					
Constituent of Potential Concern	CAS Registry Number	MW (g/mol)	D_a (cm^2/s)	Source	D_w (cm^2/s)	Source	Kd_s (cm^3/g)	Source	Kd_{sw} (L/Kg)	Source	Kd_{bs} (cm^3/g)	Source
Actinum-227	14952-40-0	227	5.11E-02	EPA6 calc a	5.91E-06	EPA6 calc a	1.50E+03	Baes b	1.50E+03	Baes b	1.50E+03	Baes b
Americium-241	14596-10-2	241	4.91E-02	EPA6 calc a	5.68E-06	EPA6 calc a	7.00E+02	Baes b	7.00E+02	Baes b	7.00E+02	Baes b
Americium-243	14993-75-0	243	4.88E-02	EPA6 calc a	5.65E-06	EPA6 calc a	7.00E+02	Baes b	7.00E+02	Baes b	7.00E+02	Baes b
Antimony-125	14234-35-6	125	7.60E-02	EPA6 calc a	8.80E-06	EPA6 calc a	4.50E+01	Baes b	4.50E+01	Baes b	4.50E+01	Baes b
Barium-137	13981-97-0	137	7.15E-02	EPA6 calc a	8.28E-06	EPA6 calc a	6.00E+01	Baes b	6.00E+01	Baes b	6.00E+01	Baes b
Cadmium-113	no cas #	113	8.13E-02	EPA6 calc a	9.41E-06	EPA6 calc a	6.50E+00	Baes b	6.50E+00	Baes b	6.50E+00	Baes b
Carbon-14	14762-75-5	14	3.27E-01	EPA6 calc a	3.79E-05	EPA6 calc a	NA	NA	NA	NA	NA	NA
Cesium-134	13967-70-9	134	7.26E-02	EPA6 calc a	8.40E-06	EPA6 calc a	1.00E+03	Baes b	1.00E+03	Baes b	1.00E+03	Baes b
Cesium-137	10045-97-3	137	7.15E-02	EPA6 calc a	8.28E-06	EPA6 calc a	1.00E+03	Baes b	1.00E+03	Baes b	1.00E+03	Baes b
Cobalt-60	10198-40-0	60	1.24E-01	EPA6 calc a	1.44E-05	EPA6 calc a	4.50E+01	Baes b	4.50E+01	Baes b	4.50E+01	Baes b
Curium-242	15510-73-3	242	4.89E-02	EPA6 calc a	5.67E-06	EPA6 calc a	2.00E+03	Baes b	2.00E+03	Baes b	2.00E+03	Baes b
Curium-243	15757-87-6	243	4.88E-02	EPA6 calc a	5.65E-06	EPA6 calc a	2.00E+03	Baes b	2.00E+03	Baes b	2.00E+03	Baes b
Curium-244	13981-15-2	244	4.87E-02	EPA6 calc a	5.63E-06	EPA6 calc a	2.00E+03	Baes b	2.00E+03	Baes b	2.00E+03	Baes b
Europium-152	14683-23-9	152	6.67E-02	EPA6 calc a	7.72E-06	EPA6 calc a	6.50E+02	Baes b	6.50E+02	Baes b	6.50E+02	Baes b
Europium-154	15585-10-1	154	6.61E-02	EPA6 calc a	7.66E-06	EPA6 calc a	6.50E+02	Baes b	6.50E+02	Baes b	6.50E+02	Baes b
Europium-155	14391-16-3	155	6.58E-02	EPA6 calc a	7.62E-06	EPA6 calc a	6.50E+02	Baes b	6.50E+02	Baes b	6.50E+02	Baes b
Iodine-129	15046-84-1	129	7.44E-02	EPA6 calc a	8.62E-06	EPA6 calc a	6.00E+01	Baes b	6.00E+01	Baes b	6.00E+01	Baes b
Neptunium-237	13994-20-2	237	4.96E-02	EPA6 calc a	5.74E-06	EPA6 calc a	3.00E+01	Baes b	3.00E+01	Baes b	3.00E+01	Baes b
Nickel-59	14336-70-0	59	1.25E-01	EPA6 calc a	1.45E-05	EPA6 calc a	1.50E+02	Baes b	1.50E+02	Baes b	1.50E+02	Baes b
Nickel-63	13981-37-8	63	1.20E-01	EPA6 calc a	1.39E-05	EPA6 calc a	1.50E+02	Baes b	1.50E+02	Baes b	1.50E+02	Baes b
Niobium-93	7440-03-1	93	9.26E-02	EPA6 calc a	1.07E-05	EPA6 calc a	3.50E+02	Baes b	3.50E+02	Baes b	3.50E+02	Baes b
Plutonium-238	13981-16-3	238	4.95E-02	EPA6 calc a	5.73E-06	EPA6 calc a	4.50E+03	Baes b	4.50E+03	Baes b	4.50E+03	Baes b
Plutonium-239	15117-48-3	239	4.93E-02	EPA6 calc a	5.71E-06	EPA6 calc a	4.50E+03	Baes b	4.50E+03	Baes b	4.50E+03	Baes b
Plutonium-240	14119-33-6	240	4.92E-02	EPA6 calc a	5.70E-06	EPA6 calc a	4.50E+03	Baes b	4.50E+03	Baes b	4.50E+03	Baes b
Plutonium-241	14119-32-5	241	4.91E-02	EPA6 calc a	5.68E-06	EPA6 calc a	4.50E+03	Baes b	4.50E+03	Baes b	4.50E+03	Baes b
Plutonium-242	13982-10-0	242	4.89E-02	EPA6 calc a	5.67E-06	EPA6 calc a	4.50E+03	Baes b	4.50E+03	Baes b	4.50E+03	Baes b
Protactinium-231	14331-85-2	231	5.05E-02	EPA6 calc a	5.84E-06	EPA6 calc a	2.50E+03	Baes b	2.50E+03	Baes b	2.50E+03	Baes b
Radium-226	13982-63-3	226	5.12E-02	EPA6 calc a	5.93E-06	EPA6 calc a	4.50E+02	Baes b	4.50E+02	Baes b	4.50E+02	Baes b
Radium-228	15262-20-1	228	5.09E-02	EPA6 calc a	5.89E-06	EPA6 calc a	4.50E+02	Baes b	4.50E+02	Baes b	4.50E+02	Baes b
Ruthenium-106	13967-48-1	106	8.48E-02	EPA6 calc a	9.82E-06	EPA6 calc a	3.50E+02	Baes b	3.50E+02	Baes b	3.50E+02	Baes b

Table B1-3. Physical/Chemical Parameters for Exposure Point Concentration Modelling for ROPCs

			Diffusivity				Partitioning Coefficients					
Constituent of Potential Concern	CAS Registry Number	MW (g/mol)	D_a (cm^2/s)	Source	D_w (cm^2/s)	Source	Kd_s (cm^3/g)	Source	Kd_{sw} (L/Kg)	Source	Kd_{bs} (cm^3/g)	Source
Samarium-151	15715-94-3	151	6.70E-02	EPA6 calc a	7.76E-06	EPA6 calc a	6.50E+02	Baes b	6.50E+02	Baes b	6.50E+02	Baes b
Selenium-79	no cas #	79	1.03E-01	EPA6 calc a	1.19E-05	EPA6 calc a	3.00E+02	Baes b	3.00E+02	Baes b	3.00E+02	Baes b
Strontium-90	10098-97-2	90	9.46E-02	EPA6 calc a	1.10E-05	EPA6 calc a	3.50E+01	Baes b	3.50E+01	Baes b	3.50E+01	Baes b
Technetium-99	14133-76-7	99	8.88E-02	EPA6 calc a	1.03E-05	EPA6 calc a	1.50E+00	Baes b	1.50E+00	Baes b	1.50E+00	Baes b
Thorium-229	15594-54-4	229	5.08E-02	EPA6 calc a	5.88E-06	EPA6 calc a	1.50E+05	Baes b	1.50E+05	Baes b	1.50E+05	Baes b
Thorium-232	7440-29-1	232	5.03E-02	EPA6 calc a	5.83E-06	EPA6 calc a	1.50E+05	Baes b	1.50E+05	Baes b	1.50E+05	Baes b
Tin-126	15832-50-5	126	7.56E-02	EPA6 calc a	8.75E-06	EPA6 calc a	2.50E+02	Baes b	2.50E+02	Baes b	2.50E+02	Baes b
Tritium	10028-17-8	3	9.13E-01	EPA6 calc a	1.06E-04	EPA6 calc a	NA	NA	NA	NA	NA	NA
Uranium-232	14158-29-3	232	5.03E-02	EPA6 calc a	5.83E-06	EPA6 calc a	4.50E+02	Baes b	4.50E+02	Baes b	4.50E+02	Baes b
Uranium-233	13968-55-3	233	5.02E-02	EPA6 calc a	5.81E-06	EPA6 calc a	4.50E+02	Baes b	4.50E+02	Baes b	4.50E+02	Baes b
Uranium-234	13966-29-5	234	5.00E-02	EPA6 calc a	5.79E-06	EPA6 calc a	4.50E+02	Baes b	4.50E+02	Baes b	4.50E+02	Baes b
Uranium-235	15117-96-1	235	4.99E-02	EPA6 calc a	5.78E-06	EPA6 calc a	4.50E+02	Baes b	4.50E+02	Baes b	4.50E+02	Baes b
Uranium-236	13982-70-2	236	4.98E-02	EPA6 calc a	5.76E-06	EPA6 calc a	4.50E+02	Baes b	4.50E+02	Baes b	4.50E+02	Baes b
Uranium-238	7440-61-1	238	4.95E-02	EPA6 calc a	5.73E-06	EPA6 calc a	4.50E+02	Baes b	4.50E+02	Baes b	4.50E+02	Baes b
Yttrium-90	10098-91-6	90	9.46E-02	EPA6 calc a	1.10E-05	EPA6 calc a	5.00E+02	Baes b	5.00E+02	Baes b	5.00E+02	Baes b
Zirconium-93	15751-77-6	93	9.26E-02	EPA6 calc a	1.07E-05	EPA6 calc a	3.00E+03	Baes b	3.00E+03	Baes b	3.00E+03	Baes b

a = Calculation from EPA Region 6 Hazardous Waste Combustion Facility Guidance (b)
b = Baes et al
c = RAIS database
d = Br_{ag} is a weighted average of $Br_{ag(fruit)}$ and $Br_{ag(veg)}$

Table B1-3. Physical/Chemical Parameters for Exposure Point Concentration Modelling for ROPCs

		Half - life	Plant Uptake Factors							
Constituent of Potential Concern	CAS Registry Number	$t_{1/2}$ (days)	Source	$Br_{root\ veg}$ ((ug/g DW plant) /(ug/g soil))	Source	$Br_{ag(fruit)}$ ((ug/g DW plant) /(ug/g soil))	Source	$Br_{ag(veg)}$ ((ug/g DW plant) /(ug/g soil))	Source	Br_{ag} ((ug/g DW plant) /(ug/g soil))[d]
Actinum-227	14952-40-0	7.96E+03	RAIS c	3.50E-04	Baes b	3.50E-04	Baes b	1.14E-03	EPA6 calc d	7.50E-04
Americium-241	14596-10-2	1.58E+05	RAIS c	2.50E-04	Baes b	2.50E-04	Baes b	1.56E-03	EPA6 calc d	9.17E-04
Americium-243	14993-75-0	2.69E+06	RAIS c	2.50E-04	Baes b	2.50E-04	Baes b	1.56E-03	EPA6 calc d	9.17E-04
Antimony-125	14234-35-6	1.01E+03	RAIS c	3.00E-02	Baes b	3.00E-02	Baes b	7.25E-02	EPA6 calc d	5.16E-02
Barium-137	13981-97-0	1.77E-03	RAIS c	1.50E-02	Baes b	1.50E-02	Baes b	4.88E-02	EPA6 calc d	3.22E-02
Cadmium-113	no cas #	NA	NA	1.50E-01	Baes b	1.50E-01	Baes b	2.50E-01	EPA6 calc d	2.01E-01
Carbon-14	14762-75-5	2.09E+06	RAIS c	NA	NA	NA	NA	NA	NA	NA
Cesium-134	13967-70-9	7.52E+02	RAIS c	3.00E-02	Baes b	3.00E-02	Baes b	4.25E-02	EPA6 calc d	3.64E-02
Cesium-137	10045-97-3	1.10E+04	RAIS c	3.00E-02	Baes b	3.00E-02	Baes b	4.25E-02	EPA6 calc d	3.64E-02
Cobalt-60	10198-40-0	1.92E+03	RAIS c	7.00E-03	Baes b	7.00E-03	Baes b	1.03E-02	EPA6 calc d	8.65E-03
Curium-242	15510-73-3	1.63E+02	RAIS c	1.50E-05	Baes b	1.50E-05	Baes b	2.24E-04	EPA6 calc d	1.21E-04
Curium-243	15757-87-6	1.04E+04	RAIS c	1.50E-05	Baes b	1.50E-05	Baes b	2.24E-04	EPA6 calc d	1.21E-04
Curium-244	13981-15-2	6.61E+03	RAIS c	1.50E-05	Baes b	1.50E-05	Baes b	2.24E-04	EPA6 calc d	1.21E-04
Europium-152	14683-23-9	4.96E+03	RAIS c	4.00E-03	Baes b	4.00E-03	Baes b	5.50E-03	EPA6 calc d	4.76E-03
Europium-154	15585-10-1	3.21E+03	RAIS c	4.00E-03	Baes b	4.00E-03	Baes b	5.50E-03	EPA6 calc d	4.76E-03
Europium-155	14391-16-3	1.81E+03	RAIS c	4.00E-03	Baes b	4.00E-03	Baes b	5.50E-03	EPA6 calc d	4.76E-03
Iodine-129	15046-84-1	5.73E+09	RAIS c	5.00E-02	Baes b	5.00E-02	Baes b	7.50E-02	EPA6 calc d	6.27E-02
Neptunium-237	13994-20-2	7.81E+08	RAIS c	1.00E-02	Baes b	1.00E-02	Baes b	3.25E-02	EPA6 calc d	2.14E-02
Nickel-59	14336-70-0	2.74E+07	RAIS c	3.20E-02	Baes b	3.20E-02	Baes b	2.55E-02	EPA6 calc d	2.87E-02
Nickel-63	13981-37-8	3.65E+04	RAIS c	6.00E-02	Baes b	6.00E-02	Baes b	6.00E-02	EPA6 calc d	6.00E-02
Niobium-93	7440-03-1	5.33E+03	RAIS c	5.00E-03	Baes b	5.00E-03	Baes b	8.75E-03	EPA6 calc d	6.91E-03
Plutonium-238	13981-16-3	3.20E+04	RAIS c	4.50E-05	Baes b	4.50E-05	Baes b	1.46E-04	EPA6 calc d	9.65E-05
Plutonium-239	15117-48-3	8.80E+06	RAIS c	4.50E-05	Baes b	4.50E-05	Baes b	1.46E-04	EPA6 calc d	9.65E-05
Plutonium-240	14119-33-6	2.40E+06	RAIS c	4.50E-05	Baes b	4.50E-05	Baes b	1.46E-04	EPA6 calc d	9.65E-05
Plutonium-241	14119-32-5	5.26E+03	RAIS c	4.50E-05	Baes b	4.50E-05	Baes b	1.46E-04	EPA6 calc d	9.65E-05
Plutonium-242	13982-10-0	1.37E+08	RAIS c	4.50E-05	Baes b	4.50E-05	Baes b	1.46E-04	EPA6 calc d	9.65E-05
Protactinium-231	14331-85-2	1.36E+07	RAIS c	2.50E-04	Baes b	2.50E-04	Baes b	8.13E-04	EPA6 calc d	5.36E-04
Radium-226	13982-63-3	5.84E+05	RAIS c	1.50E-03	Baes b	1.50E-03	Baes b	4.88E-03	EPA6 calc d	3.22E-03
Radium-228	15262-20-1	2.10E+03	RAIS c	1.50E-03	Baes b	1.50E-03	Baes b	4.88E-03	EPA6 calc d	3.22E-03
Ruthenium-106	13967-48-1	3.68E+02	RAIS c	2.00E-02	Baes b	2.00E-02	Baes b	3.38E-02	EPA6 calc d	2.70E-02

Table B1-3. Physical/Chemical Parameters for Exposure Point Concentration Modelling for ROPCs

		Half - life	Plant Uptake Factors							
Constituent of Potential Concern	CAS Registry Number	$t_{1/2}$ (days)	Source	$Br_{root\ veg}$ ((ug/g DW plant) /(ug/g soil))	Source	$Br_{ag(fruit)}$ ((ug/g DW plant) /(ug/g soil))	Source	$Br_{ag(veg)}$ ((ug/g DW plant) /(ug/g soil))	Source	Br_{ag} ((ug/g DW plant) /(ug/g soil))[d]
Samarium-151	15715-94-3	3.29E+04	RAIS c	4.00E-03	Baes b	4.00E-03	Baes b	5.50E-03	EPA6 calc d	4.76E-03
Selenium-79	no cas #	NA	NA	2.50E-02	Baes b	2.50E-02	Baes b	2.50E-02	EPA6 calc d	2.50E-02
Strontium-90	10098-97-2	1.04E+04	RAIS c	2.50E-01	Baes b	2.50E-01	Baes b	8.13E-01	EPA6 calc d	5.36E-01
Technetium-99	14133-76-7	7.77E+07	RAIS c	1.50E+00	Baes b	1.50E+00	Baes b	3.50E+00	EPA6 calc d	2.52E+00
Thorium-229	15594-54-4	2.68E+06	RAIS c	8.50E-05	Baes b	8.50E-05	Baes b	2.76E-04	EPA6 calc d	1.82E-04
Thorium-232	7440-29-1	5.15E+12	RAIS c	8.50E-05	Baes b	8.50E-05	Baes b	2.76E-04	EPA6 calc d	1.82E-04
Tin-126	15832-50-5	3.65E+07	RAIS c	6.00E-03	Baes b	6.00E-03	Baes b	1.20E-02	EPA6 calc d	9.05E-03
Tritium	10028-17-8	4.49E+03	RAIS c	NA	NA	NA	NA	NA	NA	NA
Uranium-232	14158-29-3	2.63E+04	RAIS c	4.00E-03	Baes b	4.00E-03	Baes b	5.13E-03	EPA6 calc d	4.57E-03
Uranium-233	13968-55-3	5.80E+07	RAIS c	4.00E-03	Baes b	4.00E-03	Baes b	5.13E-03	EPA6 calc d	4.57E-03
Uranium-234	13966-29-5	8.94E+07	RAIS c	4.00E-03	Baes b	4.00E-03	Baes b	5.13E-03	EPA6 calc d	4.57E-03
Uranium-235	15117-96-1	2.57E+11	RAIS c	4.00E-03	Baes b	4.00E-03	Baes b	5.13E-03	EPA6 calc d	4.57E-03
Uranium-236	13982-70-2	8.54E+09	RAIS c	4.00E-03	Baes b	4.00E-03	Baes b	5.13E-03	EPA6 calc d	4.57E-03
Uranium-238	7440-61-1	1.63E+12	RAIS c	4.00E-03	Baes b	4.00E-03	Baes b	5.13E-03	EPA6 calc d	4.57E-03
Yttrium-90	10098-91-6	2.67E+00	RAIS c	6.00E-03	Baes b	6.00E-03	Baes b	8.25E-03	EPA6 calc d	7.14E-03
Zirconium-93	15751-77-6	5.58E+08	RAIS c	5.00E-04	Baes b	5.00E-04	Baes b	8.75E-04	EPA6 calc d	6.91E-04

a = Calculation from EPA Region 6 H:
b = Baes et al
c = RAIS database
d = Br_{ag} is a weighted average of $Br_{ag(}$

Table B1-3. Physical/Chemical Parameters for Exposure Point Concentration Modelling for ROPCs

		Plant Uptake Factors					Bioconcentration/Bioaccumulation Factors				
Constituent of Potential Concern	CAS Registry Number	Source	Br_{forage} ((ug/g DW plant) /(ug/g soil))	Source	Br_{grain} ((ug/g DW plant) /(ug/g soil))	Source	Ba_{milk} (day/kg FW)	Source	Ba_{beef} (day/kg FW)	Source	Ba_{pork} (day/kg FW)
Actinum-227	14952-40-0	EPA6 calc d	3.50E-03	Baes b	3.50E-04	Baes b	2.00E-05	Baes b	2.50E-05	Baes b	NA
Americium-241	14596-10-2	EPA6 calc d	5.50E-03	Baes b	2.50E-04	Baes b	4.00E-07	Baes b	3.50E-06	Baes b	NA
Americium-243	14993-75-0	EPA6 calc d	5.50E-03	Baes b	2.50E-04	Baes b	4.00E-07	Baes b	3.50E-06	Baes b	NA
Antimony-125	14234-35-6	EPA6 calc d	2.00E-01	Baes b	3.00E-02	Baes b	1.00E-04	Baes b	1.00E-03	Baes b	NA
Barium-137	13981-97-0	EPA6 calc d	1.50E-01	Baes b	1.50E-02	Baes b	3.50E-04	Baes b	1.50E-04	Baes b	NA
Cadmium-113	no cas #	EPA6 calc d	5.50E-01	Baes b	1.50E-01	Baes b	1.00E-03	Baes b	5.50E-04	Baes b	NA
Carbon-14	14762-75-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Cesium-134	13967-70-9	EPA6 calc d	8.00E-02	Baes b	3.00E-02	Baes b	7.00E-03	Baes b	2.00E-02	Baes b	NA
Cesium-137	10045-97-3	EPA6 calc d	8.00E-02	Baes b	3.00E-02	Baes b	7.00E-03	Baes b	2.00E-02	Baes b	NA
Cobalt-60	10198-40-0	EPA6 calc d	2.00E-02	Baes b	7.00E-03	Baes b	2.00E-03	Baes b	2.00E-02	Baes b	NA
Curium-242	15510-73-3	EPA6 calc d	8.50E-04	Baes b	1.50E-05	Baes b	2.00E-03	Baes b	3.50E-06	Baes b	NA
Curium-243	15757-87-6	EPA6 calc d	8.50E-04	Baes b	1.50E-05	Baes b	2.00E-03	Baes b	3.50E-06	Baes b	NA
Curium-244	13981-15-2	EPA6 calc d	8.50E-04	Baes b	1.50E-05	Baes b	2.00E-03	Baes b	3.50E-06	Baes b	NA
Europium-152	14683-23-9	EPA6 calc d	1.00E-02	Baes b	4.00E-03	Baes b	2.00E-05	Baes b	5.00E-03	Baes b	NA
Europium-154	15585-10-1	EPA6 calc d	1.00E-02	Baes b	4.00E-03	Baes b	2.00E-05	Baes b	5.00E-03	Baes b	NA
Europium-155	14391-16-3	EPA6 calc d	1.00E-02	Baes b	4.00E-03	Baes b	2.00E-05	Baes b	5.00E-03	Baes b	NA
Iodine-129	15046-84-1	EPA6 calc d	1.50E-01	Baes b	5.00E-02	Baes b	1.00E-02	Baes b	7.00E-03	Baes b	NA
Neptunium-237	13994-20-2	EPA6 calc d	1.00E-01	Baes b	1.00E-02	Baes b	5.00E-06	Baes b	5.50E-05	Baes b	NA
Nickel-59	14336-70-0	EPA6 calc d	6.00E-03	Baes b	3.20E-02	Baes b	1.00E-02	Baes b	6.00E-03	Baes b	NA
Nickel-63	13981-37-8	EPA6 calc d	6.00E-02	Baes b	6.00E-02	Baes b	1.00E-02	Baes b	6.00E-03	Baes b	NA
Niobium-93	7440-03-1	EPA6 calc d	2.00E-02	Baes b	5.00E-03	Baes b	2.00E-02	Baes b	2.50E-01	Baes b	NA
Plutonium-238	13981-16-3	EPA6 calc d	4.50E-04	Baes b	4.50E-05	Baes b	1.00E-07	Baes b	5.00E-07	Baes b	NA
Plutonium-239	15117-48-3	EPA6 calc d	4.50E-04	Baes b	4.50E-05	Baes b	1.00E-07	Baes b	5.00E-07	Baes b	NA
Plutonium-240	14119-33-6	EPA6 calc d	4.50E-04	Baes b	4.50E-05	Baes b	1.00E-07	Baes b	5.00E-07	Baes b	NA
Plutonium-241	14119-32-5	EPA6 calc d	4.50E-04	Baes b	4.50E-05	Baes b	1.00E-07	Baes b	5.00E-07	Baes b	NA
Plutonium-242	13982-10-0	EPA6 calc d	4.50E-04	Baes b	4.50E-05	Baes b	1.00E-07	Baes b	5.00E-07	Baes b	NA
Protactinium-231	14331-85-2	EPA6 calc d	2.50E-03	Baes b	2.50E-04	Baes b	5.00E-06	Baes b	1.00E-05	Baes b	NA
Radium-226	13982-63-3	EPA6 calc d	1.50E-02	Baes b	1.50E-03	Baes b	4.50E-04	Baes b	2.50E-04	Baes b	NA
Radium-228	15262-20-1	EPA6 calc d	1.50E-02	Baes b	1.50E-03	Baes b	4.50E-04	Baes b	2.50E-04	Baes b	NA
Ruthenium-106	13967-48-1	EPA6 calc d	7.50E-02	Baes b	2.00E-02	Baes b	6.00E-07	Baes b	2.00E-03	Baes b	NA

Table B1-3. Physical/Chemical Parameters for Exposure Point Concentration Modelling for ROPCs

		Plant Uptake Factors					Bioconcentration/Bioaccumulation Factors				
Constituent of Potential Concern	CAS Registry Number	Source	Br_{forage} ((ug/g DW plant) /(ug/g soil))	Source	Br_{grain} ((ug/g DW plant) /(ug/g soil))	Source	Ba_{milk} (day/kg FW)	Source	Ba_{beef} (day/kg FW)	Source	Ba_{pork} (day/kg FW)
Samarium-151	15715-94-3	EPA6 calc d	1.00E-02	Baes b	4.00E-03	Baes b	2.00E-05	Baes b	5.00E-03	Baes b	NA
Selenium-79	no cas #	EPA6 calc d	2.50E-02	Baes b	2.50E-02	Baes b	4.00E-03	Baes b	1.50E-02	Baes b	NA
Strontium-90	10098-97-2	EPA6 calc d	2.50E+00	Baes b	2.50E-01	Baes b	1.50E-03	Baes b	3.00E-04	Baes b	NA
Technetium-99	14133-76-7	EPA6 calc d	9.50E+00	Baes b	1.50E+00	Baes b	1.00E-02	Baes b	8.50E-03	Baes b	NA
Thorium-229	15594-54-4	EPA6 calc d	8.50E-04	Baes b	8.50E-05	Baes b	5.00E-06	Baes b	6.00E-06	Baes b	NA
Thorium-232	7440-29-1	EPA6 calc d	8.50E-04	Baes b	8.50E-05	Baes b	5.00E-06	Baes b	6.00E-06	Baes b	NA
Tin-126	15832-50-5	EPA6 calc d	3.00E-02	Baes b	6.00E-03	Baes b	1.00E-03	Baes b	8.00E-02	Baes b	NA
Tritium	10028-17-8	NA	NA	NA	NA	NA	1.00E-03	RAIS c	NA	NA	NA
Uranium-232	14158-29-3	EPA6 calc d	8.50E-03	Baes b	4.00E-03	Baes b	6.00E-04	Baes b	2.00E-04	Baes b	NA
Uranium-233	13968-55-3	EPA6 calc d	8.50E-03	Baes b	4.00E-03	Baes b	6.00E-04	Baes b	2.00E-04	Baes b	NA
Uranium-234	13966-29-5	EPA6 calc d	8.50E-03	Baes b	4.00E-03	Baes b	6.00E-04	Baes b	2.00E-04	Baes b	NA
Uranium-235	15117-96-1	EPA6 calc d	8.50E-03	Baes b	4.00E-03	Baes b	6.00E-04	Baes b	2.00E-04	Baes b	NA
Uranium-236	13982-70-2	EPA6 calc d	8.50E-03	Baes b	4.00E-03	Baes b	6.00E-04	Baes b	2.00E-04	Baes b	NA
Uranium-238	7440-61-1	EPA6 calc d	8.50E-03	Baes b	4.00E-03	Baes b	6.00E-04	Baes b	2.00E-04	Baes b	NA
Yttrium-90	10098-91-6	EPA6 calc d	1.50E-02	Baes b	6.00E-03	Baes b	2.00E-05	Baes b	3.00E-04	Baes b	NA
Zirconium-93	15751-77-6	EPA6 calc d	2.00E-03	Baes b	5.00E-04	Baes b	3.00E-05	Baes b	5.50E-03	Baes b	NA

a = Calculation from EPA Region 6 H:
b = Baes et al
c = RAIS database
d = Br_{ag} is a weighted average of $Br_{ag(}$

Table B1-3. Physical/Chemical Parameters for Exposure Point Concentration Modelling for ROPCs

		Bioconcentration/Bioaccumulation Factors					
Constituent of Potential Concern	CAS Registry Number	Ba_{egg} (day/kg FW)	$Ba_{chicken}$ (day/kg FW)	BCF_{fish} (L/kg FW tissue)	BAF_{fish} (L/kg FW)	Source	$BSAF_{fish}$ (unitless)
Actinum-227	14952-40-0	NA	NA	NA	2.50E+01	RAIS c	NA
Americium-241	14596-10-2	NA	NA	NA	3.00E+01	RAIS c	NA
Americium-243	14993-75-0	NA	NA	NA	3.00E+01	RAIS c	NA
Antimony-125	14234-35-6	NA	NA	NA	1.00E-02	RAIS c	NA
Barium-137	13981-97-0	NA	NA	NA	4.00E+00	RAIS c	NA
Cadmium-113	no cas #	NA	NA	NA	2.00E+02	RAIS c	NA
Carbon-14	14762-75-5	NA	NA	NA	NA	NA	NA
Cesium-134	13967-70-9	NA	NA	NA	2.00E+03	RAIS c	NA
Cesium-137	10045-97-3	NA	NA	NA	2.00E+03	RAIS c	NA
Cobalt-60	10198-40-0	NA	NA	NA	3.00E+02	RAIS c	NA
Curium-242	15510-73-3	NA	NA	NA	3.00E+01	RAIS c	NA
Curium-243	15757-87-6	NA	NA	NA	3.00E+01	RAIS c	NA
Curium-244	13981-15-2	NA	NA	NA	3.00E+01	RAIS c	NA
Europium-152	14683-23-9	NA	NA	NA	5.00E+01	RAIS c	NA
Europium-154	15585-10-1	NA	NA	NA	5.00E+01	RAIS c	NA
Europium-155	14391-16-3	NA	NA	NA	5.00E+01	RAIS c	NA
Iodine-129	15046-84-1	NA	NA	NA	4.00E+01	RAIS c	NA
Neptunium-237	13994-20-2	NA	NA	NA	3.00E+01	RAIS c	NA
Nickel-59	14336-70-0	NA	NA	NA	1.00E+02	RAIS c	NA
Nickel-63	13981-37-8	NA	NA	NA	1.00E+02	RAIS c	NA
Niobium-93	7440-03-1	NA	NA	NA	3.00E+02	RAIS c	NA
Plutonium-238	13981-16-3	NA	NA	NA	3.00E+01	RAIS c	NA
Plutonium-239	15117-48-3	NA	NA	NA	3.00E+01	RAIS c	NA
Plutonium-240	14119-33-6	NA	NA	NA	3.00E+01	RAIS c	NA
Plutonium-241	14119-32-5	NA	NA	NA	3.00E+01	RAIS c	NA
Plutonium-242	13982-10-0	NA	NA	NA	3.00E+01	RAIS c	NA
Protactinium-231	14331-85-2	NA	NA	NA	1.00E+01	RAIS c	NA
Radium-226	13982-63-3	NA	NA	NA	5.00E+01	RAIS c	NA
Radium-228	15262-20-1	NA	NA	NA	5.00E+01	RAIS c	NA
Ruthenium-106	13967-48-1	NA	NA	NA	1.00E+01	RAIS c	NA

Table B1-3. Physical/Chemical Parameters for Exposure Point Concentration Modelling for ROPCs

		Bioconcentration/Bioaccumulation Factors					
Constituent of Potential Concern	CAS Registry Number	Ba_{egg} (day/kg FW)	$Ba_{chicken}$ (day/kg FW)	BCF_{fish} (L/kg FW tissue)	BAF_{fish} (L/kg FW)	Source	$BSAF_{fish}$ (unitless)
Samarium-151	15715-94-3	NA	NA	NA	NA	NA	NA
Selenium-79	no cas #	NA	NA	NA	NA	NA	NA
Strontium-90	10098-97-2	NA	NA	NA	6.00E+01	RAIS c	NA
Technetium-99	14133-76-7	NA	NA	NA	2.00E+01	RAIS c	NA
Thorium-229	15594-54-4	NA	NA	NA	1.00E+02	RAIS c	NA
Thorium-232	7440-29-1	NA	NA	NA	1.00E+02	RAIS c	NA
Tin-126	15832-50-5	NA	NA	NA	3.00E+03	RAIS c	NA
Tritium	10028-17-8	NA	NA	NA	1.00E+00	RAIS c	NA
Uranium-232	14158-29-3	NA	NA	NA	1.00E+01	RAIS c	NA
Uranium-233	13968-55-3	NA	NA	NA	1.00E+01	RAIS c	NA
Uranium-234	13966-29-5	NA	NA	NA	1.00E+01	RAIS c	NA
Uranium-235	15117-96-1	NA	NA	NA	1.00E+01	RAIS c	NA
Uranium-236	13982-70-2	NA	NA	NA	1.00E+01	RAIS c	NA
Uranium-238	7440-61-1	NA	NA	NA	1.00E+01	RAIS c	NA
Yttrium-90	10098-91-6	NA	NA	NA	3.00E+01	RAIS c	NA
Zirconium-93	15751-77-6	NA	NA	NA	3.00E+02	RAIS c	NA

a = Calculation from EPA Region 6 H:
b = Baes et al
c = RAIS database
d = Br_{ag} is a weighted average of $Br_{ag(}$

NOTES

COPC	=	Constituent of Potential Concern to include Organic, Inorganic, and Radiological Constituents
MW	=	Molecular weight of COPC (g/mole)
H	=	Henry's Law Constant (atm*m^3/mol)
D_a	=	Diffusivity of COPC in air (cm^2/s)
D_w	=	Diffusivity of COPC in water (cm^2/s)
K_{ow}	=	Octanol/water partitioning coefficient (unitless)
K_{oc}	=	Soil organic carbon-water partitioning coefficient (ml water/g soil)
Kd_s	=	Soil-water partitioning coefficient (ml water/g soil)
Kd_{sw}	=	Suspended sediment/surface water partitioning coefficient (L water/kg suspended sediment)
Kd_{bs}	=	Bed sediment/sediment pore water partitioning coefficient (L COPC/kg water body)
ksg	=	COPC soil loss due to biotic and abiotic degradation (year^{-1})
$t_{1/2}$	=	Half time of COPC in soil (days)
F_v	=	Fraction of COPC air concetration in vapor phase (unitless)
RCF	=	Root concentration factor (ug COPC/g DW plant)/(ug COPC/ml soil water) - (unitless)
$Br_{rootveg}$	=	Plant-soil bioconcentration factor for belowground produce (ug COPC/g DW plant)/(ug COPC/ml soil) - (unitless)
Br_{forage}	=	Plant-soil bioconcentration factor in forage and silage (ug COPC/g DW plant)/(ug COPC/ml soil) - (unitless)
$Br_{ag(fruit)}$	=	Plant-soil bioconcentration factor in abovegroud produce for fruit (ug COPC/g DW plant)/(ug COPC/ml soil) - (unitless)
$Br_{ag(veg)}$	=	Plant-soil bioconcentration factor in aboveground produce for vegetables (ug COPC/g DW plant)/(ug COPC/ml soil) - (unitless)
Br_{ag}	=	Plant-soil bioconcentration factor in abovegraound produce as a weighted average of $Br_{ag(fruit)}$ and $Br_{ag(veg)}$ (ug COPC/g DW plant)/(ug COPC/ml soil) - (unitless)
Br_{grain}	=	Plant-soil bioconcentration factor in grain (ug COPC/g DW plant)/(ug COPC/ml soil) - (unitless)
Bv_{ag}	=	COPC air-to-plant biotransfer factor for aboveground produce ((ug COPC/g DW plant)/(ug COPC/ml air) - (unitless)
Bv_{forage}	=	COPC air-to-plant biotransfer factor for forage and silage ((ug COPC/g DW plant)/(ug COPC/ml air) - (unitless)
Ba_{milk}	=	Biotransfer factor for milk (mg COPC/kg FW tissue)/(mg COPC/day) OR (day/kg FW tissue)
Ba_{beef}	=	Biotransfer factor for beef (mg COPC/kg FW tissue)/(mg COPC/day) OR (day/kg FW tissue)
Ba_{pork}	=	Biotransfer factor for pork (mg COPC/kg FW tissue)/(mg COPC/day) OR (day/kg FW tissue)
Ba_{egg}	=	Biotransfer factor for eggs (mg COPC/kg FW tissue)/(mg COPC/day) OR (day/kg FW tissue)
$Ba_{chicken}$	=	Biotransfer factor for chicken (mg COPC/kg FW tissue)/(mg COPC/day) OR (day/kg FW tissue)
BCF_{fish}	=	Bioconcentration factor in fish (L/kg FW) OR (unitless)
BAF_{fish}	=	Bioaccumulatin factor in fish (mg COPC/kg FW tissue)/(mg COPC/L total water column)
$BSAF_{fish}$	=	Biota-sediment accumulation factor in fish (mg COPC/kg lipid tissue)/(mg COPC/kg sediment) - (unitless)

References for Appendix B1

Baes CF, Sharp RD, Sjoreen AL, and Shor RW. 1984. *Reviw and Analysis of Parameters and Assessing Transport of Environmentally Released Radionuclides Through Agriculture*, ORNL-5786, 1984. Oak Ridge National Laboratory, Oak Ridge, Tennessee.

EPA Region 9 Preliminary Remediation Goals. http://www.epa.gov/region09/waste/sfund/prg/

EPA. 1998. *Peer Review Draft of Human Health Risk Assessment Protocol for Hazardous Waste Combustion Facilities*, EPA530-D-98-001A, July 1998. US Environmental Protection Agency, Washington D.C., USA.

Howard PH. 1993. *Handbook of Environmental Fate and Exposure Data for Organic Chemicals*, Voume 1: Large Production and Priority Pollutants, Volume II: Solvents, Volume III: Pesticides, Volume IV: Solvents 2, Volume V: Solvents 3. Lewis Publishers, Chelsea, Michigan, USA.

Howard PH, Boethling RS, Jarvis WF, Meylan WM, and Michalenko EM. 1991. *Handbook of Environmental Degradation Rates*. Lewis Publishers. Chelsea, Michigan, USA.

Mackay D, and. Shiu WY. 1981. *A Critical Review of Henry's Law Constants for Chemicals of Environmental Interest*, Journal of Physical Chemical Reference Data, volume 10, Number 4.

Mackay D, Shiu WY, and Ma KC. 1992. *Illustrated Handbook of Physical-Chemical Properties and Environmental Fate for Organic Chemicals. Volume I - Monoaromatic Hydrocarbons, Chlorobenzenes, and PCBs. Volume II - Polynuclear Aromatic Hydrocarbons, Polychlorinated Dioxins and Dibenzofurans. Volume III - Volatile Organic Chemicals. Volume IV - Herbicides and Pesticides*. Lewis Publishers. Boca Raton, Florida, USA.

Ng YC, Colsher CS, Thompson SE. 1982. *Transfer Coefficients for Assessing the Dose from Radionuclides in Meat and Eggs*, Lawrence Livermore National Laboratory, Livermore, California, USA

Risk Assessment Information System Database. http://risk.lsd.ornl.gov/rap_hp.shtml

Appendix B2

Supporting Equations for Media Constituents of Potential Concern Concentrations

Appendix B2

Supporting Equations for Media Constituents of Potential Concern Concentrations

Soil Concentration

The following equations can be used to calculate the constituent of potential concern (COPC) soil loss constants that are dependent on significant amounts of water, such as erosion, runoff, and leaching. These processes are not likely to require quantitative evaluation in the risk assessment due to the low precipitation and high evapotranspiration in the region.

The soil loss constant due to soil erosion (kse) can be calculated based on the following equation:

$$\text{kse} = \frac{0.1 \times X_e \times \text{SD} \times \text{ER}}{\text{BD} \times Z_s} \times \frac{\text{Kd}_s \times \text{BD}}{\theta_{sw} + (\text{Kd}_s \times \text{BD})}$$

where

kse = COPC soil loss constant due to soil erosion (1/yr)
0.1 = Units conversion factor (1,000 g per kg/10,000 cm^2 per m^2)
X_e = Unit soil loss as calculated from Universal Soil Loss Equation (kg/m^2-yr)
SD = Sediment delivery ratio (unitless)
ER = Soil enrichment ratio (unitless)
Kd_s = Soil-water partition coefficient (ml water/g soil)
BD = Soil bulk density (g soil/cm^3 soil)
Z_s = Soil mixing zone depth (cm)
θ_{sw} = Soil volumetric water content (ml water/cm^3 soil)

The unit soil loss (X_e) and sediment delivery ratio (SD) are used to account for losses of COPCs from soil, and are used to represent loads of COPCs that may migrate to surface water bodies. Therefore, further discussion of these parameters can be found in the surface water section (section 6). EPA 1998a recommends that the parameter for soil loss due to erosion be set to zero, based on the assumption that soil loss due to runoff from a site would be equal to soil gained due to run on to a site from adjacent areas. An additional assumption is that the soil run on contamination is equal to the soil runoff contamination.

The COPC loss constant due to surface runoff (ksr) can be calculated based on the following equation:

$$\text{ksr} = \frac{\text{RO} \times \text{CF}}{\theta_{sw} \times Z_s} \times \left(\frac{1}{1 + \left(\text{Kd}_s \times \text{BD} / \theta_{sw} \right)} \right)$$

where

ksr = COPC loss constant due to surface runoff (1/yr)
RO = Average annual surface runoff from pervious areas (cm/yr)
CF = Conversion factor of 1.0 (ml/cm^3)

θ_{sw} = Soil volumetric water content (ml water/cm^3 soil)
Z_s = Soil mixing zone depth (cm)
Kd_s = Soil-water partition coefficient (ml water/g soil)
BD = Soil bulk density (g soil/cm^3 soil)

The COPC loss constant due to leaching (ksl) can be calculated based on the following equation:

$$ksl = \left[\frac{P + I - RO - E_v}{\theta_{sw} \times Z_s \times \left(1 + \left(BD \times Kd_s / \theta_{sw}\right)\right)}\right](CF)$$

where

ksl = COPC loss constant due to leaching (1/yr)
P = Average annual precipitation (cm/yr)
I = Average annual irrigation (cm/yr)
RO = Average annual surface runoff from pervious areas (cm/yr)
E_v = Average annual evapotranspiration (cm/yr)
θ_{sw} = Soil volumetric water content (ml water/cm^3 soil)
Z_s = Soil mixing zone depth (cm)
BD = Soil bulk density (g soil/cm^3 soil)
Kd_s = Soil-water partition coefficient (ml water/g soil)
CF = Conversion factor of 1.0 (ml/cm^3)

Surface Water and Fish Concentrations

The following equations are not likely to be necessary in the calculation of surface water and fish concentrations, due to a lack of available water in the Hanford site. They are included in this appendix for completeness.

The runoff load from impervious surfaces (L_{ri}) can be calculated as follows:

$$L_{ri} = Q \times \left[F_v \times Dywwv + \left(1.0 - F_v\right) \times Dytwp \times A_I\right]$$

where

L_{ri} = Runoff load from impervious surfaces (g/yr)
Q = COPC emission rate (g/s)
F_v = Fraction of COPC in air concentration in vapor phase (unitless)
Dywwv = Unitized yearly (waterbody and watershed) average wet deposition from vapor phase (s/m^2/yr)
Dytwp = Unitized yearly (waterbody and watershed) average total (wet and dry) deposition from vapor phase (s/m^2/yr)
A_I = Impervious watershed area receiving COPC deposition (m^2)

The runoff load from pervious surfaces (L_{rp}) can be calculated with the following equation:

$$L_{rp} = RO \times (A_L - A_I) \frac{Cs \times BD}{\theta_{sw} + Kd_s \times BD} \times (0.01)(CF)$$

where

L_{rp}	=	Runoff load from previous surfaces (g/yr)
RO	=	Average annual surface runoff from pervious areas (cm/yr)
A_L	=	Total watershed area receiving COPC deposition (m^2)
A_I	=	Impervious watershed area receiving COPC deposition (m^2)
Cs	=	Average soil concentration over exposure duration (in watershed soils) (mass COPC/kg soil)
BD	=	Soil bulk density (g soil/cm^3 soil)
θ_{sw}	=	Soil volumetric water content (ml water/cm^3 soil)
Kd_s	=	Soil-water partition coefficient (ml water/g soil)
0.01	=	Units conversion factor (kg-cm^2/mg-m^2)
CF	=	Conversion factor of 1.0 (ml/cm^3)

The COPC load to surface water due to soil erosion will be calculated with the following equation:

$$L_E = X_e \times (A_L - A_I) \times SD \times ER \frac{Cs \times Kd_s \times BD}{\theta_{sw} + Kd_s \times BD} \times (0.001)$$

where

L_E	=	Soil erosion load (g/yr)
X_e	=	Unit soil loss (kg/m^2-yr)
A_L	=	Total watershed area evaluated receiving COPC deposition (m^2)
A_I	=	Impervious watershed area receiving COPC deposition (m^2)
SD	=	Sediment delivery ratio for watershed (unitless)
ER	=	Soil enrichment ratio (unitless)
Cs	=	Average soil concentration in watershed soils over exposure duration (mass COPC/kg soil)
Kd_s	=	Soil-water partition coefficient (ml water/g soil)
BD	=	Soil bulk density (g soil/cm^3 soil)
θ_{sw}	=	Soil volumetric water content (ml water/cm^3 soil)
0.001	=	Units conversion factor (g/mg)

The soil loss due to erosion (X_e) will be calculated with the universal soil loss equation (USLE):

$$X_e = RF \times K \times LS \times C \times PF \times \frac{907.18}{4047}$$

where

X_e	=	Unit soil loss (kg/m^2-yr)
RF	=	USLE rainfall (or erosivity) factor (1/yr)
K	=	USLE erodibility factor (ton/acre)
LS	=	USLE length-slope factor (unitless)
C	=	USLE cover management factor (unitless)
PF	=	USLE supporting practice factor (unitless)
907.18	=	Conversion factor (kg/ton)
4047	=	Conversion factor (m^2/acre)

Water Concentration

The fraction of total water body COPC concentration in the water column (f_{wc}) will be calculated with the following equations:

$$f_{wc} = \frac{(1 + Kd_{sw} \times TSS \times CF) \times d_{wc}/d_z}{\left((1 + Kd_{sw} \times TSS \times CF) \times d_{wc}/d_z\right) + (\theta_{bs} + Kd_{bs} \times C_{BS}) \times d_{bs}/d_z}$$

where

f_{wc}	=	Fraction of total water body COPC concentration in the water column (unitless)
Kd_{sw}	=	Suspended sediments/surface water partition coefficient (L water/kg suspended sediment)
TSS	=	Total suspended solids concentration (mg/L)
CF	=	Conversion factor of 1×10^{-6} (kg/mg)
d_{wc}	=	Depth of water column (m)
d_z	=	Total water body depth (m)
2_{bs}	=	Bed sediment porosity (L water/L sediment)
Kd_{bs}	=	Bed sediment/sediment pore water partition coefficient (L water/kg bottom sediment)
C_{BS}	=	Bed sediment concentration (g/cm^3)
d_{bs}	=	Depth of upper benthic sediment layer (m)

If measured data are not available, the total suspended solids (TSS) concentration will be calculated with the following equation:

$$TSS = \frac{X_e \times (A_L - A_I) \times SD \times CF}{Vf_x + D_{ss} \times A_W}$$

where

TSS	=	Total suspended solids concentration (mg/L)
X_e	=	Unit soil loss (kg/m^2-yr)
A_L	=	Total watershed area receiving COPC deposition (m^2)
A_I	=	Total impervious watershed area receiving COPC deposition (m^2)
SD	=	Sediment delivery ratio for watershed (unitless)
CF	=	Conversion factor of 1000 (mg - m^3/kg - L)
Vf_x	=	Average volumetric flow rate through water body (m^3/year)
Dss	=	Suspended solids deposition rate (m/yr)
A_W	=	Water body surface area (m^2)

The sediment delivery ratio (SD) will be calculated as follows:

$$SD = a \times (A_L)^{-b}$$

where

SD	=	Sediment delivery ratio [watershed (unitless)]
a	=	Empirical intercept coefficient (unitless)
A_L	=	Total watershed area evaluated which receives COPC deposition (m^2)
b	=	Empirical slope coefficient (unitless)

The bed sediment porosity (2_{bs}) will be calculated with the following equation:

$$\theta_{bs} = 0.6^{L_{water}/L_{sediment}}$$

The above equation assumes the bed sediment density is 2.65 kg/L and the bed sediment concentration is 1 kg/L.

The overall total waterbody COPC dissipation rate constant (k_{wt}) will be calculated with the following equation:

$$k_{wt} = f_{wc} \times k_v + f_{bs} \times k_b$$

where

k_{wt}	=	Total water body COPC dissipation rate constant (1/yr)
f_{wc}	=	Fraction of total water body COPC concentration in the water column (unitless)
k_v	=	Water column volatilization rate constant (1/yr)
f_{bs}	=	Fraction of total water body COPC concentration in benthic sediment (unitless)
k_b	=	Benthic burial rate constant (1/yr)

The water column volatilization rate constant (k_v) will be calculated with the following equation:

$$k_v = \frac{K_v}{d_z \times (1 + Kd_{sw} \times TSS \times CF)}$$

where

k_v	=	Water column volatilization rate constant (1/yr)
K_v	=	Overall COPC transfer rate coefficient (m/yr)
d_z	=	Total water body depth (m)
Kd_{sw}	=	Suspended sediments/surface water partition coefficient (L water/kg suspended sediments)
TSS	=	Total suspended sediments concentration (mg/L)
CF	=	Conversion factor of 1×10^{-6} (kg/mg)

The overall COPC transfer rate coefficient (K_V) will be calculated as follows:

$$K_v = \left(K_L^{-1} + \left(K_G \frac{H}{R \cdot T_{wk}} \right)^{-1} \right)^{-1} \times \theta^{T_{wk} - 293}$$

where

K_v	=	Overall COPC transfer rate coefficient (m/yr)
K_L	=	Liquid phase transfer coefficient (m/yr)
K_G	=	Gas phase transfer coefficient (m/yr)
H	=	Henry's Law constant (atm-m^3/mol)
R	=	Universal gas constant of 8.205×10^{-5} (atm-m^3/mol-°K)
T_{wk}	=	Water body temperature (°K)
2	=	Temperature correction factor of 1.026 (unitless)

For flowing water bodies the liquid phase transfer coefficient (K_L) will be calculated using the following equation:

$$K_L = \left(\sqrt{\frac{CF_1 \times D_w \times u}{d_z}}\right)(CF_2)$$

where

K_L	=	Liquid phase transfer coefficient (m/yr)
CF_1	=	Conversion factor of 1×10^{-4} (m^2/cm^2)
D_w	=	Diffusivity of COPC in water (cm^2/s)
u	=	Current velocity (m/s)
d_z	=	Total water body depth (m)
CF_2	=	Conversion factor of 3.1536×10^7 (s/yr)

The gas phase transfer coefficient (K_G) will be assumed to be equal to 36500 m/yr for flowing water bodies.

For quiescent lakes or ponds the liquid phase transfer coefficient (K_L) will be calculated using the following equation:

$$K_L = \left(C_d^{\,0.5} \times W\right) \times \left(\frac{\rho_a}{\rho_w}\right)^{0.5} \times \left(\frac{k^{0.33}}{\lambda_z}\right) \times \left(\frac{\mu_w}{\rho_w \times D_w}\right)^{-0.67} \times CF$$

where

K_L	=	Liquid phase transfer coefficient (m/yr)
C_d	=	Drag coefficient (unitless)
W	=	Average annual wind speed (m/s)
Δ_a	=	Density of air (g/cm^3)
Δ_w	=	Density of water (g/cm^3)
k	=	von Karman's constant (unitless)
8_z	=	Dimensionless viscous sublayer thickness (unitless)
μ_w	=	Viscosity of water corresponding to water temperature (g/cm - s)
D_w	=	Diffusivity of COPC in water (cm^2/s)
CF	=	Conversion factor of 3.1536×10^7 (s/yr)

The gas phase transfer coefficient (K_G) for a quiescent stream will be calculated with the following equation:

$$K_G = \left(C_d^{\,0.5} \times W\right) \times \left(\frac{k^{0.33}}{\lambda_z}\right) \times \left(\frac{\mu_a}{\rho_a \times D_a}\right)^{-0.67} \times CF$$

where

K_G	=	Gas phase transfer coefficient (m/yr)
C_d	=	Drag coefficient (unitless)
W	=	Average annual wind speed (m/s)
k	=	von Karman's constant (unitless)
8_z	=	Dimensionless viscous sublayer thickness (unitless)
$:_a$	=	Viscosity of air corresponding to air temperature (g/cm-s)
Δ_a	=	Density of air (g/cm^3)
D_a	=	Diffusivity of COPC in air (cm^2/s)
CF	=	Conversion factor of 3.1536×10^7 (s/yr)

The fraction of total water body COPC concentration in benthic sediment (f_{bs}) will be calculated as follows:

$$f_{bs} = 1 - f_{wc}$$

where

f_{bs}	=	Fraction of total water body COPC concentration in benthic sediment (unitless)
f_{wc}	=	Fraction of total water body COPC concentration in the water column (unitless).

References for Appendix B2

EPA. 1998a. *Human Health Risk Assessment Protocol for Hazardous Waste Combustion Facilities,* Peer Review Draft, EPA/530/D-98/001B. US Environmental Protection Agency, Washington, D.C., USA

Appendix C

Chemical-Specific Parameters for Ecological Risk Assessment

Appendix C-1

Lists of Common Vascular Plants, Mammals, Common Birds, Amphibians and Reptiles, Plant Species of Concern, and State Candidate Species

Table C1-1. Common Vascular Plants Found on the Hanford Site

(Taxonomy follows Hitchcock and Cronquist 1973)

A. Shrub-Steppe Species	Scientific Name
Shrub	
Big sagebrush	*Artemisia tridentata*
Bitterbrush	*Purshia tridentata*
Gray rabbitbrush	*Chrysothamnus nauseosus*
Green rabbitbrush	*Chrysothamnus viscidiflorus*
Snow buckwheat	*Eriogonum niveum*
Spiny hopsage	*Grayia (Atriplex) spinosa*
Threetip sagebrush	*Artemisia tripartita*
Perennial Grasses	
Bluebunch wheatgrass	*Agropyron spicatum*
Bottlebrush squirreltail	*Sitanion hystrix*
Crested wheatgrass	*Agropyron desertorum (cristatum)*[a]
Indian ricegrass	*Oryzopsis hymenoides*
Needle-and-thread grass	*Stipa comata*
Prairie Junegrass	*Koeleria cristata*
Sand dropseed	*Sporobolus cryptandrus*
Sandberg's bluegrass	*Poa sandbergii (secunda)*
Thickspike wheatgrass	*Agropyron dasytachyum*
Perennial Forbs	
Bastard toad flax	*Comandra umbellata*
Buckwheat milkvetch	*Astragalus caricinus*
Carey's balsamroot	*Balsamorhiza careyana*
Cusick's sunflower	*Helianthus cusickii*
Cutleaf ladysfoot mustard	*Thelypodium laciniatum*
Douglas' clusterlily	*Brodiaea douglasii*
Dune scurfpea	*Psoralea lanceolata*
Franklin's sandwort	*Arenaria franklinii*
Gray's desertparsley	*Lomatium grayi*
Hoary aster	*Machaeranthera canescens*
Hoary false yarrow	*Chaenactis douglasii*
Longleaf phlox	*Phlox longifolia*
Munro's globemallow	*Sphaeralcea munroana*
Pale evening primrose	*Oenothera pallida*
Sand beardtongue	*Penstemon acuminatus*
Stalked-pod milkvetch	*Astragalus sclerocarpus*
Threadleaf fleabane	*Erigeron filifolius*

Table C1-1. Common Vascular Plants Found on the Hanford Site

(Taxonomy follows Hitchcock and Cronquist 1973)

A. Shrub-Steppe Species	Scientific Name
Turpentine spring parsley	*Cymopteris terebinthinus*
Winged dock	*Rumex venosus*
Yarrow	*Achillea millefolium*
Yellow bell	*Fritillaria pudica*
Annual Forbs	
Annual Jacob's ladder	*Polemonium micranthum*
Blue mustard	*Chorispora tenella*[a]
Bur ragweed	*Ambrosia acanthicarpa*
Clasping pepperweed	*Lepidium perfoliatum*
Indian wheat	*Plantago patagonica*
Jagged chickweed	*Holosteum umbellatum*[a]
Jim Hill's tumblemustard	*Sisymbrium altissimum*[a]
Matted cryptantha	*Cryptantha circumscissa*
Pink microsteris	*Microsteris gracilis*
Prickly lettuce	*Lactuca serriola*[a]
Rough wallflower	*Erysimum asperum*
Russian thistle (tumbleweed)	*Salsola kali*[a]
Slender hawksbeard	*Crepis atrabarba*
Spring whitlowgrass	*Draba verna*[a]
Storksbill	*Erodium cicutarium*[a]
Tall willowherb	*Epilobium paniculatum*
Tarweed fiddleneck	*Amsinckia lycopsoides*
Threadleaf scorpion weed	*Phacelia linearis*
Western tansymustard	*Descurainia pinnata*
White cupseed	*Plectritis macrocera*
Whitestem stickleaf	*Mentzelia albicaulis*
Winged cryptantha	*Cryptantha pterocarya*
Yellow salsify	*Tragopogon dubius*[a]
Annual Grasses	
Cheatgrass	*Bromus tectorum*[a]
Slender sixweeks	*Festuca octoflora*
Small sixweeks	*Festuca microstachys*

Table C1-1. Common Vascular Plants Found on the Hanford Site

(Taxonomy follows Hitchcock and Cronquist 1973)

B. Riparian Species	Scientific Name
Trees and Shrubs	
Black cottonwood	*Populus trichocarpa*
Black locust	*Robinia pseudo-acacia*
Coyote willow	*Salix exigua*
Dogbane	*Apocynum cannabinum*
Peach, apricot, cherry	*Prunus* spp.
Peachleaf willow	*Salix amygdaloides*
Willow	*Salix* spp.
White Mulberry	*Morus alba*[a]
Perennial Grasses and Forbs	
Bentgrass	*Agrostis* spp. [b]
Blanket flower	*Gaillardia aristata*
Bulrushes	*Scirpus* spp.[b]
Cattail	*Typha latifolia*[b]
Columbia River gumweed	*Grindelia columbiana*
Hairy golden aster	*Heterotheca villosa*
Heartweed	*Polygonum persicaria*
Horsetails	*Equisetum* spp.
Horseweed tickseed	*Coreopsis atkinsoniana*
Lovegrass	*Eragrostis* spp. [b]
Lupine	*Lupinus* spp.
Meadow foxtail	*Alopecurus aequalis* [b]
Pacific sage	*Artemisia campestris*
Prairie sagebrush	*Artemisia ludoviciana*
Reed canary grass	*Phalaris arundinacea*[b]
Rushes	*Juncus* spp.
Russian knapweed	*Centaurea repens*[a]
Sedge	*Carex* spp.[b]
Water speedwell	*Veronica anagallis-aquatica*
Western goldenrod	*Solidago occidentalis*
Wild onion	*Allium* spp.
Wiregrass spikerush	*Eleocharis* spp.[b]
Aquatic Vascular	
Canadian waterweed	*Elodea canadensis*
Columbia yellowcress	*Rorippa columbiae*
Duckweed	*Lemna minor*

Table C1-1. Common Vascular Plants Found on the Hanford Site

(Taxonomy follows Hitchcock and Cronquist 1973)

B. Riparian Species	Scientific Name
Aquatic Vascular	
Pondweed	*Potamogeton* spp.
Spiked water milfoil	*Myriophyllum spicatum*
Watercress	*Rorippa nasturtium-aquaticum*

(a) Introduced
(b) Perennial grasses and graminoids

from Neitzel et al. 1998

Table C1-2. List of Mammals Occurring on the Hanford Site

Scientific Name	Common Name
Antrozous pallidus	pallid bat
Brachylagus idahoensis	pygmy rabbit
Canis latrans	coyote
Castor canadensis	beaver
Cervus elaphus	elk
Erethizon dorsatum	porcupine
Eutamias minimus	least chipmunk
Lagurus curtatus	sagebrush vole
Lasionycteris noctivagans	silver-haired bat
Lasiurus cinereus	hoary bat
Lepus californicus	black-tailed jackrabbit
Lepus townsendi	white-tailed jackrabbit
Lutra canadensis	river otter
Lynx rufus	bobcat
Marmota flaviventris	yellow-bellied marmot
Mephitis mephitis	striped skunk
Microtus montanus	montane meadow mouse
Mus musculus	house mouse
Mustela erminea	short-tailed weasel
Mustela frenata	long-tailed weasel
Mustela vison	mink
Myotis californicus	California brown bat
Myotis lucifugus	little brown bat
Myotis yumanensis	Yuma brown bat
Neotoma cinerea	bushy-tailed woodrat
Odocoileus hemionus	mule deer
Odocoileus virginianus	white-tailed deer
Ondatra zibethicus	muskrat
Onychomys leucogaster	northern grasshopper mouse
Perognathus parvus	Great Basin pocket mouse
Peromyscus maniculatus	deer mouse
Plecotus townsendii townsendii	Pacific western big-eared bat
Procyon lotor	raccoon
Rattus norvegicus	Norway rat
Reithrodontomys megalotis	western harvest mouse
Sorex merriami	Merriam shrew
Sorex vagrans	vagrant shrew
Spermophilus townsendii	Townsend ground squirrel
Sylvilagus nuttallii	Nuttall cottontail rabbit
Taxidea taxus	badger
Thomomys talpoides	northern pocket gopher

Source: Cushing 1992 in DOE 1996a

Table C1-3. Common Birds Occurring on the Hanford Site

Scientific Name	Common Name
Ageless phonics	red-winged blackbird
Alas aceta	northern pintail
Alas americana	American wigeon
Anas clypeata	northern shoveler
Anas platyrhynchos	mallard
Ardea herodias	great blue heron
Aythya americana	edhead
Branta canadensis	Canada goose
Bucephala albeola	bufflehead
Calidris mauri	western sandpiper
Calidris minutilla	least sandpiper
Carpodacus mexicanus	house finch
Charadrius vociferus	killdeer
Chordeiles minor	common nighthawk
*Columba livia*rock	dove
Corvus corax	common raven
Dendroica coronata	yellow-rumped warbler
Eremophila alpestris	horned lark
Fulica americana	American coot
Hirundo pyrrhonota	cliff swallow
Hirundo rustica	barn swallow
Junco hyemalis	dark-eyed junco
Larus californicus	California gull
Larus delawarensis	ring-billed gull
Limnodromus scolopaceus	long-billed dowitcher
Mergus merganser	common merganser
Numenius americanus	long-billed curlew
Passer domesticus	house sparrow
Pica pica	black-billed magpie
Podilymbus podiceps	pied-billed grebe
Sturnella neglecta	western meadowlark
Sturnus vulgaris	european starling
Turdus migratorius	American robin
Tyrannus verticalis	western kingbird
Zenaida macroura	mourning dove
Zonotrichia leucophrys	white-crowned sparrow

Source: Cushing 1992 in DOE 1996a

Table C1-4. Amphibians and Reptiles Occurring on the Hanford Site

Scientific Name	Common Name
Amphibians	
Bufo woodhouseii	Woodhouse toad
Hyla regilla	Pacific treefrog
Spea intermontana	Great Basin spadefoot
Rana catesbeiana	Bullfrog
Reptiles	
Chrysemys picta	painted turtle
Coluber constrictor	western yellow-bellied racer
Crotalus viridis	western rattlesnake
Hyspiglena torquata	desert night snake
Masticophis taeniatus	striped whipsnake
Phrynosoma douglassii	short-horned lizard
Pituophis melanoleucus	gopher snake
Sceloporus graciosus	sagebrush lizard
Uta stansburiana	side-blotched lizard

Source: Cushing 1992 in DOE 1996a

Table C1-5. Plant Species of Concern on the Hanford Site

COMMON NAME	SCIENTIFIC NAME	STATE LISTING[a]
Annual paintbrush	*Castilleja exilis*	R1
Awned half chaff sedge	*Lipocarpha (= Hemicarpha) aristulata*	R1
Basalt milk-vetch	*Astragalus conjunctus var. rikardii*	R1
Bristly combseed	*Pectocarya setosa*	W
Brittle prickly-pear	*Opuntia fragilis*	R1
Canadian St. John's wort	*Hypericum majus*	S
Chaffweed	*Centunculus minimus*	R1
Columbia River mugwort	*Artemesia lindleyana*	W
Crouching milkvetch	*Astragalus succumbens*	W
Desert dodder	*Cuscuta denticulata*	S
Desert evening-primrose	*Oenothera cespitosa*	S
False pimpernel	*Lindernia dubia anagallidea*	R2
Fuzzytongue penstemon	*Penstemon eriantherus whitedii*	R1
Geyer's milkvetch	*Astragalus geyeri*	S
Grand redstem	*Ammannia robusta*	R1
Gray cryptantha	*Cryptantha leucophaea*	S
Great Basin gilia	*Gilia leptomeria*	R1
Hedge hog cactus	*Pediocactus simpsonii var. robustio nigrispinus*	R1
Kittitas larkspur	*Delphinium multiplex*	W
Miner's candle	*Cryptantha scoparia*	R1
Palouse thistle	*Cirsium brevifolium*	W
Piper's daisy	*Erigeron piperianus*	S
Robinson's onion	*Allium robinsonii*	W
Rosy balsamroot	*Balsamorhiza rosea*	W
Rosy pussypaws	*Calyptridium roseum*	S
Scilla onion	*Allium scilloides*	W
Shining flatsedge	*Cyperus bipartitus (rivularis)*	S
Small-flowered evening-primrose	*Camissonia (= Oenothera) minor*	R1
Small-flowered nama	*Nama densum var. parviflorum*	R1
Smooth cliffbrake	*Pellaea glabella simplex*	W
Snake River cryptantha	*Cryptantha spiculifera (= C. interrupta)*	S
Southern mudwort	*Limosella acaulis*	W
Stalked-pod milkvetch	*Astragalus sclerocarpus*	W
Suksdorf's monkey flower	*Mimulus suksdorfii*	S
Toothcup	*Rotala ramosior*	R1
Winged combseed	*Pectocarya linearis*	R1

The following species have been reported as occurring on the Hanford site, but the known collections are questionable in terms of location or identification, and have not been recently collected on the Hanford site.

Coyote tobacco	*Nicotiana attenuata*	S
Dense sedge	*Carex densa*	S
Few-flowered collinsia	*Collinsia sparsiflora var. bruciae*	S
Medic milkvetch	*Astragalus speirocarpus*	W
Palouse milkvetch	*Astragalus arrectus*	S
Thompson's sandwort	*Arenaria franklinii thompsonii*	R2

(a) S = Sensitive (i.e., taxa vulnerable or declining) and could become endangered or threatened without active management or removal of threats.
R1 = Taxa for which there are insufficient data to support listing as threatened, endangered, or sensitive (formerly monitor group 1).
R2 = Taxa with unresolved taxanomic questions (formerly monitor group 2).
W = Taxa that are more abundant and/or less threatened than previously assumed (formerly monitor group 3).

from Neitzel et al. 1998

Table C1-6. Wildlife Species of Concern on the Hanford Site

Common Name	Scientific Name
Molluscs	
Columbia pebble snail	*Fluminicola (= Lithoglyphus) columbiana*
Shortfaced lanx	*Fisherola (= Lanx) nuttalli*
Insects	
Columbia River tiger beetle[b]	*Cicindela columbica*
Juniper hairstreak	*Mitoura siva*
Silver-bordered bog fritillary	*Boloria selene atrocastalis*
Birds	
Burrowing owl	*Athene cunicularia*
Common loon	*Gavia immer*
Flammulated owl[a]	*Otus flammeolus*
Golden eagle	*Aquila chrysaetos*
Lewis' woodpecker[a]	*Melanerpes lewis*
Loggerhead shrike	*Lanius ludovicianus*
Northern goshawk[a]	*Accipter gentilis*
Sage sparrow	*Amphispiza belli*
Sage thrasher	*Oreoscoptes montanus*
Western sage grouse[a]	*Centrocercus urophasianus phaios*
Merlin	*Falco columbarius*
Reptiles	
Striped whipsnake	*Masticophis taeniatus*
Mammals	
Merriam's shrew	*Sorex merriami*
Pacific (Townsend's) western big-eared bat[b]	*Corynorhinus townsendii* [c]
Washington ground squirrel	*Spermophilus washingtoni*

(a) Reported, but seldom observed on the Hanford site
(b) Probable, but not observed, on the Hanford site
(c) Also known as *Plecotus townsendii*

from Neitzel et al. 1998

Table C1-7. Federally or Washington State Listed Threatened and Endangered Species Occurring or Potentially Occurring on the Hanford Site

Common Name	Scientific Name	Federal	State
Plants			
Columbia milk-vetch	*Astragalus columbianus*		T
Columbia yellowcress	*Rorippa columbiae*		E
Dwarf evening primrose	*Oenothera pygmaea*		T
Hoover's desert parsley	*Lomatium tuberosum*		T
Loeflingia	*Loeflingia squarrosa* var. *squarrosa*		T
Northern wormwood[a]	*Artemisia campestris borealis* var. *wormskioldii*		E
Umtanum desert buckwheat	*Eriogonum codium*		E
White Bluffs bladderpod	*Lesquerella tuplashensis*		E
White eatonella	*Eatonella nivea*		T
Birds			
Aleutian Canada goose[b]	*Branta canadensis leucopareia*	T	E
American white pelican	*Pelecanus erythrorhychos*		E
Bald eagle	*Haliaeetus leucocephalus*	T	T
Ferruginous hawk	*Buteo regalis*		T
Peregrine falcon[b]	*Falco peregrinus*	E	E
Sandhill crane[b]	*Grus canadensis*		E
Mammals			
Pygmy rabbit[a]	*Brachylagus idahoensis*		E
Fish			
Steelhead			
(Upper Columbia River ESU)	*Oncorhynchus mykiss*	E	
(Middle Columbia River ESU)[c]		T	
Chinook Salmon			
(Upper Columbia River spring-run ESU)[d]	*Oncorhynchus tshawytscha*	E	

T = threatened
E = endangered
[a]Likely not currently occurring on the Hanford site
[b]Incidental occurrence
[c]64 FR 14517, March 25, 1999
[d]64 FR 14308, March 24, 1999

from Neitzel et al. 1998

References for Appendix C-1

64 FR 14308. *Endangered and Threatened Species: Threatened Status for Three Chinook Salmon Evolutionarily Significant Units in Washington and Oregon, and Endangered Status of One Chinook Salmon ESU in Washington; Final Rule* and *Partial 6-Month Extension on Final Listing Determinations for Four Evolutionarily Significant Units of West Coast Chinook Salmon; Proposed Rule*, Volume 64, Number 56, p 14308-14328, 24 March 1999. Federal Register

64 FR 14517. *Endangered and Threatened Species: Threatened Status for Two ESUs of Chum Salmon in Washington and Oregon, for Two ESUs of Steelhead in Washington and Oregon, and for Ozette Lake Sockeye Salmon in Washington; Rules*, Volume 64, Number 57, p 14507-14517, 25 March 1999. Federal Register.

DOE. 1996. *Draft Hanford Remedial Action Environmental Impact Statement and Comprehensive Land Use Plan*, DOE/EIS-0222D, August 1996. US Department of Energy, Washington, D.C., USA.

Neitzel DA and others. 1998. *Hanford Site National Environmental Policy Act (NEPA) Characterization*, PNL-6415, Revision 10, 1998. Pacific Northwest National Laboratory, Richland, Washington, USA.

Appendix C-2

Soil-to-Plant Transfer Factors, Terrestrial Bioaccumulation Factors, and Aquatic Bioaccumulation Factors for Ecological Constituents of Potential Concern

Table C2-1. Terrestrial Soil-to-Plant Transfer Factors (SPv) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPv	SAIC Compiled SPv	Notes	Comments	Recommended SPv[d]
Organic Compounds										
Aromatic Halogenated Hydrocarbons										
4-Chloro-3-methylphenol	59-50-7	3.10		1.25E-01	e		1.25E-01	e		1.25E-01
2,3,4,6-Tetrachlorophenol	58-90-2	4.42		2.16E-02	e		2.16E-02	e		2.16E-02
Aromatic Nonhalogenated Hydrocarbons										
2-Nitrotoluene	88-72-2	2.30		3.63E-01	e		3.63E-01	e		3.63E-01
4-Nitrobiphenyl	92-93-3	3.77		5.13E-02	e		5.13E-02	e		5.13E-02
Benzaldehyde	100-52-7	1.48		1.08E+00	e		1.08E+00	e		1.08E+00
Benzene	71-43-2	-4.99		5.94E+03	e		5.94E+03	e		5.94E+03
Benzyl alcohol	100-51-6	1.10		1.79E+00	e		1.79E+00	e		1.79E+00
Ethyl benzene	100-41-4	3.12		1.21E-01	e		1.21E-01	e		1.21E-01
m-Xylene	108-38-3	3.20		1.09E-01	e		1.09E-01	e		1.09E-01
o-Xylene	95-47-6	3.13		1.20E-01	e		1.20E-01	e		1.20E-01
p-Xylene	106-42-3	3.17		1.14E-01	e		1.14E-01	e		1.14E-01
Styrene	100-42-5	2.93		1.57E-01	e		1.57E-01	e		1.57E-01
Toluene	108-88-3	2.67		2.22E-01	e		2.22E-01	e		2.22E-01
Non-aromatic Nonhalogenated Hydrocarbons										
1,2-Epoxybutane	106-88-7	1.44		1.14E+00	e		1.14E+00	e		1.14E+00
1,3-Butadiene	106-99-0	1.90		6.18E-01	e		6.18E-01	e		6.18E-01
1,4-Dioxane	123-91-1	-0.27	5.53E+01	1.11E+01	f		1.11E+01	e		1.11E+01
1-Methylpropyl alcohol	78-92-2	0.61		3.44E+00	e		3.44E+00	e		3.44E+00
1-Nitropropane	108-03-2	0.87		2.43E+00	e		2.43E+00	e		2.43E+00
2,2,4-Trimethylpentane	540-84-1	5.02		9.72E-03	e		9.72E-03	e		9.72E-03
2-Butanone	78-93-3	0.28		5.33E+00	e		5.33E+00	e		5.33E+00
2-Butenaldehyde (2-Butenal)	4170-30-3	No data		No data			1.00E+00		Default	1.00E+00
2-Ethoxyethanol	110-80-5	-0.10		8.85E+00	e		8.85E+00	e		8.85E+00
2-Heptanone	110-43-0	1.98		5.55E-01	e		5.55E-01	e		5.55E-01
2-Hexanone	591-78-6	1.38		1.23E+00	e		1.23E+00	e		1.23E+00
2-Methoxyethanol	109-86-4	0.25		5.54E+00	e		5.54E+00	e		5.54E+00

Table C2-1. Terrestrial Soil-to-Plant Transfer Factors (SPv) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPv	SAIC Compiled SPv	Notes	Comments	Recommended SPv[d]
2-Methyl-2-propanol	75-65-0	0.35		4.86E+00	e		4.86E+00	e		4.86E+00
2-Methyl-2-propenenitrile	126-98-7	0.54		3.77E+00	e		3.77E+00	e		3.77E+00
2-Methylaziridine	75-55-8	-0.60		1.73E+01	e		1.73E+01	e		1.73E+01
2-Methylpropyl alcohol	78-83-1	0.76		2.82E+00	e		2.82E+00	e		2.82E+00
2-Pentanone	107-87-9	0.91		2.31E+00	e		2.31E+00	e		2.31E+00
2-Propanone (Acetone)	67-64-1	-0.22	5.20E+01	1.04E+01	f		1.04E+01	e		1.04E+01
2-Propene-1-ol	107-18-6	0.17		6.18E+00	e		6.18E+00	e		6.18E+00
2-Propyl alcohol	67-63-0	0.05		7.25E+00	e		7.25E+00	e		7.25E+00
3-Heptanone	106-35-4	No data		No data			1.00E+00		Default	1.00E+00
3-Methyl-1-butanol	123-51-3	No data		No data			1.00E+00		Default	1.00E+00
3-Methyl-2-butanone	563-80-4	No data		No data			1.00E+00		Default	1.00E+00
3-Pentanone	96-22-0	0.99		2.07E+00	e		2.07E+00	e		2.07E+00
4-Heptanone	123-19-3	No data		No data			1.00E+00		Default	1.00E+00
4-Methyl-2-pentanone	108-10-1	1.19		1.59E+00	e		1.59E+00	e		1.59E+00
4-Methyl-3-penten-2-one	141-79-7	No data		No data			1.00E+00		Default	1.00E+00
5-Methyl-2-hexanone	110-12-3	No data		No data			1.00E+00		Default	1.00E+00
Acetaldehyde	75-07-0	-0.22		1.04E+01	e		1.04E+01	e		1.04E+01
Acetamide	60-35-5	-1.26		4.14E+01	e		4.14E+01	e		4.14E+01
Acetic acid	64-19-7	-0.17		9.71E+00	e		9.71E+00	e		9.71E+00
Acetic acid ethyl ester	141-78-6	0.73		2.93E+00	e		2.93E+00	e		2.93E+00
Acetic acid n-butyl ester	123-86-4	1.73		7.75E-01	e		7.75E-01	e		7.75E-01
Acetonitrile	75-05-8	-0.34		1.22E+01	e		1.22E+01	e		1.22E+01
Acrolein	107-02-8	-0.01		7.84E+00	e		7.84E+00	e		7.84E+00
Acrylonitrile	107-13-1	0.25	2.78E+01	5.55E+00	f		5.55E+00	e		5.55E+00
Bis(isopropyl)ether	108-20-3	No data		No data			1.00E+00		Default	1.00E+00
Butane	106-97-8	2.89		1.65E-01	e		1.65E-01	e		1.65E-01
Carbon disulfide	75-15-0	2.00		5.41E-01	e		5.41E-01	e		5.41E-01
Cyanogen	460-19-5	0.81		2.64E+00	e		2.64E+00	e		2.64E+00
Cyclohexane	110-82-7	3.44		7.96E-02	e		7.96E-02	e		7.96E-02

Table C2-1. Terrestrial Soil-to-Plant Transfer Factors (SPv) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPv	SAIC Compiled SPv	Notes	Comments	Recommended SPv[d]
Cyclohexanone	108-94-1	0.81		2.64E+00	e		2.64E+00	e		2.64E+00
Cyclohexene	110-83-8	2.86		1.72E-01	e		1.72E-01	e		1.72E-01
Cyclopentane	287-92-3	3.00		1.43E-01	e		1.43E-01	e		1.43E-01
Ethyl alcohol	64-17-5	0.31		5.13E+00	e		5.13E+00	e		5.13E+00
Ethyl ether	60-29-7	0.89		2.37E+00	e		2.37E+00	e		2.37E+00
Ethyl methacrylate	97-63-2	1.59		9.33E-01	e		9.33E-01	e		9.33E-01
Formaldehyde	50-00-0	0.34	2.46E+01	4.91E+00	f		4.91E+00	e		4.91E+00
Formamide	75-12-7	-1.51		5.78E+01	e		5.78E+01	e		5.78E+01
Formic acid	64-18-6	-0.54		1.58E+01	e		1.58E+01	e		1.58E+01
Formic acid, methyl ester	107-31-3	-0.26		1.10E+01	e		1.10E+01	e		1.10E+01
Glycidylaldehyde	765-34-4	-0.73		2.05E+01	e		2.05E+01	e		2.05E+01
Methyl acetate	79-20-9	0.18		6.10E+00	e		6.10E+00	e		6.10E+00
Methyl alcohol	67-56-1	-0.71		1.99E+01	e		1.99E+01	e		1.99E+01
Methyl isocyanate	624-83-9	No data		No data			NA		Reacts with water	NA
Methyl methacrylate	80-62-6	0.79		2.71E+00	e		2.71E+00	e		2.71E+00
Methyl tert-butyl ether	1634-04-4	0.94		2.22E+00	e		2.22E+00	e		2.22E+00
Methylacetylene	74-99-7	0.94		2.22E+00	e		2.22E+00	e		2.22E+00
Methylcyclohexane	108-87-2	4.10		3.31E-02	e		3.31E-02	e		3.31E-02
N,N-Dimethylacetamide	127-19-5	No data		No data			1.00E+00		Default	1.00E+00
n-Butyl alcohol	71-36-3	0.88		2.40E+00	e		2.40E+00	e		2.40E+00
n-Heptane	142-82-5	4.66		1.57E-02	e		1.57E-02	e		1.57E-02
n-Hexane	110-54-3	4.11		3.26E-02	e		3.26E-02	e		3.26E-02
Nitromethane	75-52-5	-0.35		1.23E+01	e		1.23E+01	e		1.23E+01
n-Nonane	111-84-2	5.65		4.20E-03	e		4.20E-03	e		4.20E-03
n-Octane	111-65-9	4.00		3.78E-02	e		3.78E-02	e		3.78E-02
n-Pentane	109-66-0	3.21		1.08E-01	e		1.08E-01	e		1.08E-01
n-Propionaldehyde	123-38-6	0.59		3.53E+00	e		3.53E+00	e		3.53E+00
n-Propyl alcohol	71-23-8	0.25		5.55E+00	e		5.55E+00	e		5.55E+00
n-Valeraldehyde	110-62-3	No data		No data	e		1.00E+00		Default	1.00E+00

Table C2-1. Terrestrial Soil-to-Plant Transfer Factors (SPv) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPv	SAIC Compiled SPv	Notes	Comments	Recommended SPv[d]
Oxirane	75-21-8	-0.30		1.15E+01	e		1.15E+01	e		1.15E+01
p-Cymene	99-87-6	4.10		3.31E-02	e		3.31E-02	e		3.31E-02
Phosgene	75-44-5	No data		No data	e		1.00E+00		Default	1.00E+00
Propargyl alcohol	107-19-7	0.26		5.48E+00	e		5.48E+00	e		5.48E+00
Propionic acid	79-09-4	0.33		4.99E+00	e		4.99E+00	e		4.99E+00
Propionitrile	107-12-0	0.16		6.26E+00	e		6.26E+00	e		6.26E+00
Propylene gylcol monomethyl ether	107-98-2	-0.18		9.84E+00	e		9.84E+00	e		9.84E+00
p-tert-Butyltoluene	98-51-1	No data		No data			1.00E+00		Default	1.00E+00
Triethylamine	121-44-8	0.16		6.26E+00	e		6.26E+00	e		6.26E+00
Trimethylamine	75-50-3	0.16		6.26E+00	e		6.26E+00	e		6.26E+00
Vinyl acetate	108-05-4	0.70		3.06E+00	e		3.06E+00	e		3.06E+00
Non-aromatic Halogenated Hydrocarbons										
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	No data		No data			1.00E+00		Default	1.00E+00
1,1,1,2-Tetrachloroethane	630-20-6	2.63		2.34E-01	e		2.34E-01	e		2.34E-01
1,1,1-Trichloroethane	71-55-6	2.42		3.09E-01	e		3.09E-01	e		3.09E-01
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	3.73		5.41E-02	e		5.41E-02	e		5.41E-02
1,1,2,2-Tetrachloroethane	79-34-5	4.64		1.60E-02	e		1.60E-02	e		1.60E-02
1,1,2,2-Tetrachloroethene	127-18-4	2.55		2.62E-01	e		2.62E-01	e		2.62E-01
1,1,2-Trichloroethane	79-00-5	2.10		4.75E-01	e		4.75E-01	e		4.75E-01
1,1,2-Trichloroethylene	79-01-6	2.43		3.04E-01	e		3.04E-01	e		3.04E-01
1,1-Dichloroethane	75-34-3	1.79		7.13E-01	e		7.13E-01	e		7.13E-01
1,1-Dichloroethene	75-35-4	2.12		4.61E-01	e		4.61E-01	e		4.61E-01
1,2,2-Trichloro-1,1,2-trifluoroethane	76-13-1	3.16		1.15E-01	e		1.15E-01	e		1.15E-01
1,2,3-Trichloropropane	96-18-4	2.25		3.88E-01	e		3.88E-01	e		3.88E-01
1,2-Dibromo-3-chloropropane	96-12-8	2.34		3.44E-01	e		3.44E-01	e		3.44E-01
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	2.82		1.82E-01	e		1.82E-01	e		1.82E-01
1,2-Dichloroethane	107-06-2	1.46		1.11E+00	e		1.11E+00	e		1.11E+00
1,2-Dichloroethylene	540-59-0	0.48		4.09E+00	e		4.09E+00	e		4.09E+00
1,2-Dichloropropane	78-87-5	2.25		3.88E-01	e		3.88E-01	e		3.88E-01

Table C2-1. Terrestrial Soil-to-Plant Transfer Factors (SPv) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPv	SAIC Compiled SPv	Notes	Comments	Recommended SPv[d]
1,3-Dichloropropene	542-75-6	1.75		7.56E-01	e		7.56E-01	e		7.56E-01
1,4-Dichloro-2-butene	764-41-0	0.87		2.42E+00	e		2.42E+00	e		2.42E+00
1-Chloroethene	75-01-4	1.15		1.68E+00	e		1.68E+00	e		1.68E+00
2,2-Dichloropropionic acid	75-99-0	0.78		2.75E+00	e		2.75E+00	e		2.75E+00
2-Chloropropane	75-29-6	1.90		6.18E-01	e		6.18E-01	e		6.18E-01
3-Chloropropene (allyl chloride)	107-05-1	0.95		2.17E+00	e		2.17E+00	e		2.17E+00
Bromochloromethane	74-97-5	1.41		1.19E+00	e		1.19E+00	e		1.19E+00
Bromodichloromethane	75-27-4	2.03		5.23E-01	e		5.23E-01	e		5.23E-01
Bromoethene	593-60-2	1.07		1.87E+00	e		1.87E+00	e		1.87E+00
Bromoform	75-25-2	2.35		3.39E-01	e		3.39E-01	e		3.39E-01
Bromomethane	74-83-9	1.11		1.76E+00	e		1.76E+00	e		1.76E+00
Carbon tetrachloride	56-23-5	2.72	1.04E+00	2.08E-01	f		2.08E-01	e		2.08E-01
Chlorodibromomethane	124-48-1	2.18		4.28E-01	e		4.28E-01	e		4.28E-01
Chlorodifluoromethane	75-45-6	1.08		1.84E+00	e		1.84E+00	e		1.84E+00
Chloroethane	75-00-3	3.10		1.25E-01	e		1.25E-01	e		1.25E-01
Chloroform	67-66-3	1.95	2.90E+00	5.80E-01	f		5.78E-01	e		5.80E-01
Chloromethane	74-87-3	0.90		2.33E+00	e		2.33E+00	e		2.33E+00
Chloropentafluoroethane	76-15-3	No data		No data			1.00E+00		Default	1.00E+00
cis-1,2-Dichloroethene	156-59-2	1.98		5.54E-01	e		5.54E-01	e		5.54E-01
cis-1,3-Dichloropropene	10061-01-5	No data		No data			1.00E+00		Default	1.00E+00
Cyanogen bromide	506-68-3	No data		No data			1.00E+00		Default	1.00E+00
Cyanogen chloride	506-77-4	0.20		5.94E+00	e		5.94E+00	e		5.94E+00
Dichlorodifluoromethane	75-71-8	2.16		4.38E-01	e		4.38E-01	e		4.38E-01
Dichlorofluoromethane	75-43-4	No data		No data			1.00E+00		Default	1.00E+00
Dichloromethane	75-09-2	1.26		1.46E+00	e		1.46E+00	e		1.46E+00
Difluorodibromomethane	75-61-6	No data		No data			1.00E+00		Default	1.00E+00
Hexafluoroacetone	684-16-2	No data		No data			1.00E+00		Default	1.00E+00
Iodomethane	74-88-4	1.69		8.17E-01	e		8.17E-01	e		8.17E-01
Methylene bromide	74-95-3	1.62		8.97E-01	e		8.97E-01	e		8.97E-01

Table C2-1. Terrestrial Soil-to-Plant Transfer Factors (SPv) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPv	SAIC Compiled SPv	Notes	Comments	Recommended SPv[d]
Pentachloroethane	76-01-7	3.05		1.34E-01	e		1.34E-01	e		1.34E-01
trans-1,2-Dichloroethene	156-60-5	1.98		5.54E-01	e		5.54E-01	e		5.54E-01
trans-1,3-Dichloropropene	10061-02-6	2.06		4.99E-01	e		4.99E-01	e		4.99E-01
Trichloroacetic acid	76-03-9	1.33		1.32E+00	e		1.32E+00	e		1.32E+00
Trichlorofluoroethane	27154-33-2	No data		No data			1.00E+00		Default	1.00E+00
Trichlorofluoromethane	75-69-4	2.53		2.67E-01	e		2.67E-01	e		2.67E-01
Trifluorobromomethane	75-63-8	1.86		6.52E-01	e		6.52E-01	e		6.52E-01
Dioxin and Furan Compounds										
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	35822-46-9	8.20	2.90E-04	5.80E-05	g		5.80E-05	g		5.80E-05
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	7.92	6.20E-05	1.24E-05	g		1.24E-05	g		1.24E-05
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	7.92	2.20E-03	4.40E-04	g		4.40E-04	g		4.40E-04
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	39227-28-6	7.79	1.70E-03	3.40E-04	g		3.40E-04	g		3.40E-04
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	7.25	4.30E-04	8.60E-05	g		8.60E-05	g		8.60E-05
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	57653-85-7	7.25	6.70E-04	1.34E-04	g		1.34E-04	g		1.34E-04
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	7.25	1.10E-03	2.20E-04	g		2.20E-04	g		2.20E-04
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	19408-74-3	7.25	7.80E-04	1.56E-04	g		1.56E-04	g		1.56E-04
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	7.25	3.50E-03	7.00E-04	g		7.00E-04	g		7.00E-04
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	40321-76-4	6.64	5.20E-03	1.04E-03	g		1.04E-03	g		1.04E-03
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	6.79	1.10E-03	2.20E-04	g		2.20E-04	g		2.20E-04
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	7.25	3.80E-03	7.60E-04	g		7.60E-04	g		7.60E-04
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	6.92	9.00E-03	1.80E-03	g		1.80E-03	g		1.80E-03
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1746-01-6	6.64	5.62E-03	1.12E-03	e		1.12E-03	e		1.12E-03
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	6.53	4.50E-03	9.00E-04	g		9.00E-04	g		9.00E-04
Dibenzofuran	132-64-9	4.33		2.43E-02	e		2.43E-02	e		2.43E-02
Octachlorodibenzo(p)dioxin	3268-87-9	8.20	6.70E-05	1.34E-05	g		1.34E-05	g		1.34E-05
Octachlorodibenzofuran	39001-02-0	8.78	9.00E-05	1.80E-05	g		1.80E-05	g		1.80E-05
PCBs										
Polychlorinated biphenyls (PCBs)[h]	1336-36-3	7.31	1.00E-02	2.00E-03	f		4.63E-04	e		2.00E-03
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	7.08		6.26E-04	e		6.26E-04	e		6.26E-04

Table C2-1. Terrestrial Soil-to-Plant Transfer Factors (SPv) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPv	SAIC Compiled SPv	Notes	Comments	Recommended SPv[d]
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	7.12		5.94E-04	e		5.94E-04	e		5.94E-04
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	No data		No data	e		1.00E+00		Default	1.00E+00
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	No data		No data	e		1.00E+00		Default	1.00E+00
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	No data		No data	e		1.00E+00		Default	1.00E+00
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	No data		No data	e		1.00E+00		Default	1.00E+00
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	No data		No data	e		1.00E+00		Default	1.00E+00
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	No data		No data	e		1.00E+00		Default	1.00E+00
2',3,4,4',5-Pentachlorobiphenyl	no cas #	No data		No data	e		1.00E+00		Default	1.00E+00
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	7.12		5.94E-04	e		5.94E-04	e		5.94E-04
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	7.41		4.05E-04	e		4.05E-04	e		4.05E-04
3,3',4,4',5-Pentachlorobiphenyl	no cas #	No data		No data	e		1.00E+00		Default	1.00E+00
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	No data		No data	e		1.00E+00		Default	1.00E+00
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	No data		No data	e		1.00E+00		Default	1.00E+00
Phthalates										
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	5.20	3.80E-02	7.60E-03	f		7.60E-03	e		7.60E-03
Butylbenzyl phthalate	85-68-7	4.41		2.18E-02	e		2.18E-02	e		2.18E-02
Dibutyl phthalate	84-74-2	4.72		1.45E-02	e		1.45E-02	e		1.45E-02
Diethyl phthalate	84-66-2	4.44		2.11E-02	e		2.11E-02	e		2.11E-02
Dimethylphthalate	131-11-3	1.63		8.81E-01	e		8.81E-01	e		8.81E-01
n-Dioctyl phthalate	117-84-0	9.33	1.57E-04	3.14E-05	f		3.13E-05	e		3.14E-05
Light Polycyclic Aromatic Hydrocarbons (molecular weight <200 g/mole)										
2-Chloronapthalene	91-58-7	4.07		3.45E-02	e		3.45E-02	e		3.45E-02
2-Methyl naphthalene	91-57-6	3.86		4.55E-02	e		4.55E-02	e		4.55E-02
5-Nitroacenaphthene	602-87-9	No data		No data			1.00E+00		Default	1.00E+00
Acenaphthene	83-32-9	3.96		3.96E-02	e		3.96E-02	e		3.96E-02
Acenaphthylene	208-96-8	4.07		3.44E-02	e		3.44E-02	e		3.44E-02
Anthracene	120-12-7	4.47		2.02E-02	e		2.02E-02	e		2.02E-02
Fluorene	86-73-7	4.18		2.97E-02	e		2.97E-02	e		2.97E-02
Indene	95-13-6	No data		No data			1.00E+00		Default	1.00E+00

Table C2-1. Terrestrial Soil-to-Plant Transfer Factors (SPv) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPv	SAIC Compiled SPv	Notes	Comments	Recommended SPv[d]
Naphthalene	91-20-3	3.37		8.70E-02	e		8.70E-02	e		8.70E-02
Phenanthrene	85-01-8	4.55		1.82E-02	e		1.82E-02	e		1.82E-02
Pyrene	129-00-0	5.00		9.98E-03	e		9.98E-03	e		9.98E-03
Heavy Polycyclic Aromatic Hydrocarbons (molecular weight >200 g/mole)										
3-Methylcholanthrene	56-49-5	7.11		6.02E-04	e		6.02E-04	e		6.02E-04
5-Methylchrysene	3697-24-3	No data		No data			1.00E+00		Default	1.00E+00
Benzo(a)anthracene	56-55-3	5.68	2.02E-02	4.04E-03	f		4.04E-03	e		4.04E-03
Benzo(a)pyrene	50-32-8	6.13	1.11E-02	2.22E-03	f		2.22E-03	e		2.22E-03
Benzo(b)fluoranthene	205-99-2	6.20	1.01E-02	2.02E-03	f		2.02E-03	e		2.02E-03
Benzo(e)pyrene	192-97-2	7.40		4.09E-04	e		4.09E-04	e		4.09E-04
Benzo(g,h,i)perylene	191-24-2	7.10		6.10E-04	e		6.10E-04	e		6.10E-04
Benzo(j)fluoranthene	205-82-3	6.44		1.47E-03	e		1.47E-03	e		1.47E-03
Benzo(k)fluoranthene	207-08-9	6.20	1.01E-02	2.02E-03	f		2.01E-03	e		2.02E-03
Benzo[a,i]pyrene	191-30-0	No data		No data			1.00E+00		Default	1.00E+00
Chrysene	218-01-9	5.74	1.87E-02	3.74E-03	f		3.73E-03	e		3.74E-03
Dibenz(a,h)anthracene	53-70-3	6.55	6.40E-03	1.28E-03	f		1.27E-03	e		1.28E-03
Dibenz[a,h]acridine	226-36-8	No data		No data			1.00E+00		Default	1.00E+00
Dibenz[a,j]acridine	224-42-0	No data		No data			1.00E+00		Default	1.00E+00
Dibenzo(a,e)fluoranthene	5385-75-1	No data		No data			1.00E+00		Default	1.00E+00
Dibenzo(a,h)fluoranthene	no cas #	No data		No data			1.00E+00		Default	1.00E+00
Dibenzo[a,e]pyrene	192-65-4	No data		No data			1.00E+00		Default	1.00E+00
Dibenzo[a,h]pyrene	189-64-0	No data		No data			1.00E+00		Default	1.00E+00
Dibenzo[a,i]pyrene	189-55-9	7.29		4.74E-04	e		4.74E-04	e		4.74E-04
Fluoranthene	206-44-0	5.08		8.94E-03	e		8.94E-03	e		8.94E-03
Hexachloronaphthalene	1335-87-1	7.59		3.18E-04	e		3.18E-04	e		3.18E-04
Indeno(1,2,3-cd)pyrene	193-39-5	6.91	3.90E-03	7.80E-04	f		7.80E-04	e		7.80E-04
Octachloronaphthalene	2234-13-1	6.42		1.51E-03	e		1.51E-03	e		1.51E-03
Pentachloronaphthalene	1321-64-8	No data		No data			1.00E+00		Default	1.00E+00
Tetrachloronaphthalene	1335-88-2	No data		No data			1.00E+00		Default	1.00E+00

Table C2-1. Terrestrial Soil-to-Plant Transfer Factors (SPv) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPv	SAIC Compiled SPv	Notes	Comments	Recommended SPv[d]
Trichloronaphthalene	1321-65-9	No data		No data			1.00E+00		Default	1.00E+00
Light Substituted Benzene Compounds (MW <200 g/mole)										
1,2,3-Trichlorobenzene	87-61-6	4.05		3.55E-02	e		3.55E-02	e		3.55E-02
1,2,4-Trichlorobenzene	120-82-1	3.99		3.84E-02	e		3.84E-02	e		3.84E-02
1,2,4-Trimethyl benzene	95-63-6	3.65		6.02E-02	e		6.02E-02	e		6.02E-02
1,2-Dichlorobenzene	95-50-1	3.45		7.90E-02	e		7.90E-02	e		7.90E-02
1,3,5-Trimethyl benzene	108-67-8	3.42		8.17E-02	e		8.17E-02	e		8.17E-02
1,3-Dichlorobenzene	541-73-1	3.53		7.06E-02	e		7.06E-02	e		7.06E-02
1,3-Dinitrobenzene	99-65-0	1.49	5.32E+00	1.06E+00	f		1.06E+00	e		1.06E+00
1,4-Dichlorobenzene	106-46-7	3.41		8.26E-02	e		8.26E-02	e		8.26E-02
1,4-Dinitrobenzene	100-25-4	1.50		1.05E+00	e		1.05E+00	e		1.05E+00
2,4,5-Trichlorophenol	95-95-4	3.87		4.49E-02	e		4.49E-02	e		4.49E-02
2,4,6-Trichlorophenol	88-06-2	3.71		5.54E-02	e		5.54E-02	e		5.54E-02
2,4-Dichlorophenol	120-83-2	3.04		1.36E-01	e		1.36E-01	e		1.36E-01
2,4-Dimethylphenol	105-67-9	2.36		3.35E-01	e		3.35E-01	e		3.35E-01
2,4-Dinitrophenol	51-28-5	1.52		1.03E+00	e		1.03E+00	e		1.03E+00
2,4-Dinitrotoluene	121-14-2	2.00	2.72E+00	5.44E-01	f		5.44E-01	e		5.44E-01
2,6-Dinitrotoluene	606-20-2	1.89	3.15E+00	6.30E-01	f		6.29E-01	e		6.30E-01
2-Chlorophenol	95-57-8	2.16		4.36E-01	e		4.36E-01	e		4.36E-01
2-Chlorotoluene	95-49-8	3.54		7.00E-02	e		7.00E-02	e		7.00E-02
2-Nitrophenol	88-75-5	1.79		7.15E-01	e		7.15E-01	e		7.15E-01
4,6-Dinitro-o-cresol	534-52-1	2.85		1.74E-01	e		1.74E-01	e		1.74E-01
4-Chlorotoluene	106-43-4	3.33		9.21E-02	e		9.21E-02	e		9.21E-02
4-Nitrophenol	100-02-7	1.91		6.10E-01	e		6.10E-01	e		6.10E-01
alpha-Methylstyrene	98-83-9	3.46		7.71E-02	e		7.71E-02	e		7.71E-02
Aniline	62-53-3	0.98		2.10E+00	e		2.10E+00	e		2.10E+00
Benzotrichloride	98-07-7	2.92		1.59E-01	e		1.59E-01	e		1.59E-01
Benzyl chloride	100-44-7	0.36		4.79E+00	e		4.79E+00	e		4.79E+00
Bromobenzene	108-86-1	2.99		1.45E-01	e		1.45E-01	e		1.45E-01

Table C2-1. Terrestrial Soil-to-Plant Transfer Factors (SPv) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPv	SAIC Compiled SPv	Notes	Comments	Recommended SPv[d]
Chlorobenzene	108-90-7	2.79		1.89E-01	e		1.89E-01	e		1.89E-01
Cumene	98-82-8	3.61		6.32E-02	e		6.32E-02	e		6.32E-02
m-Cresol	108-39-4	1.96		5.71E-01	e		5.71E-01	e		5.71E-01
n-Butyl benzene	104-51-8	4.28		2.60E-02	e		2.60E-02	e		2.60E-02
Nitrobenzene	98-95-3	1.83	3.38E+00	6.76E-01	f		6.76E-01	e		6.76E-01
n-Propyl benzene	103-65-1	3.69		5.70E-02	e		5.70E-02	e		5.70E-02
o-Cresol	95-48-7	2.02		5.26E-01	e		5.26E-01	e		5.26E-01
o-Dinitrobenzene	528-29-0	1.60		9.21E-01	e		9.21E-01	e		9.21E-01
o-Nitroaniline	88-74-4	1.85		6.60E-01	e		6.60E-01	e		6.60E-01
o-Toluidine	95-53-4	1.34		1.30E+00	e		1.30E+00	e		1.30E+00
p-Chloroaniline	106-47-8	1.87		6.44E-01	e		6.44E-01	e		6.44E-01
p-Cresol	106-44-5	1.94		5.86E-01	e		5.86E-01	e		5.86E-01
Phenol	108-95-2	1.48		1.08E+00	e		1.08E+00	e		1.08E+00
p-Nitrochlorobenzene	100-00-5	2.39		3.22E-01	e		3.22E-01	e		3.22E-01
p-Toluidine	106-49-0	1.40		1.20E+00	e		1.20E+00	e		1.20E+00
sec-Butyl benzene	135-98-8	4.57		1.77E-02	e		1.77E-02	e		1.77E-02
tert-Butyl benzene	98-06-6	4.11		3.26E-02	e		3.26E-02	e		3.26E-02
Toluene-2,6-diamine	823-40-5	1.45		1.12E+00	e		1.12E+00	e		1.12E+00
Trimethyl benzene	25551-13-7	3.40		8.39E-02	e		8.39E-02	e		8.39E-02
Other Light Semivolatile Compounds (molecular weight <200 g/mole)										
1,1'-Biphenyl	92-52-4	3.90		4.31E-02	e		4.31E-02	e		4.31E-02
1,1-Dimethylhydrazine	57-14-7	0.60		3.51E+00	e		3.51E+00	e		3.51E+00
1,2-Dimethylhydrazine	540-73-8	-1.37		4.78E+01	e		4.78E+01	e		4.78E+01
1,2-Diphenylhydrazine	122-66-7	2.94		1.55E-01	e		1.55E-01	e		1.55E-01
1,3-Propane sultone	1120-71-4	-0.52		1.55E+01	e		1.55E+01	e		1.55E+01
2,4-Toluene diisocyante	584-84-9	No data		No data			NA		Decomposes rapidly in water	NA
2-Chloroacetophenone	532-27-4	2.59		2.45E-01	e		2.45E-01	e		2.45E-01
2-Propenoic acid	79-10-7	0.33		4.99E+00	e		4.99E+00	e		4.99E+00

Table C2-1. Terrestrial Soil-to-Plant Transfer Factors (SPv) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPv	SAIC Compiled SPv	Notes	Comments	Recommended SPv[d]
4,4-Methylenedianiline	101-77-9	3.38		8.62E-02	e		8.62E-02	e		8.62E-02
Acetophenone	98-86-2	1.64		8.73E-01	e		8.73E-01	e		8.73E-01
Benzoic acid	65-85-0	1.86		6.52E-01	e		6.52E-01	e		6.52E-01
bis(2-Chloroethoxy)methane	111-91-1	7.59		3.18E-04	e		3.18E-04	e		3.18E-04
bis(2-Chloroethyl) ether	111-44-4	1.30		1.37E+00	e		1.37E+00	e		1.37E+00
Chlorocyclopentadiene	41851-50-7	2.43		3.05E-01	e		3.05E-01	e		3.05E-01
Cyclohexanol	108-93-0	1.23		1.51E+00	e		1.51E+00	e		1.51E+00
Dichloroisopropyl ether	108-60-1	2.58		2.50E-01	e		2.50E-01	e		2.50E-01
Dichloromethyl ether	542-88-1	-0.38		1.28E+01	e		1.28E+01	e		1.28E+01
Dichloropentadiene	no cas #	No data		No data			1.00E+00		Default	1.00E+00
Dimethyl sulfate	77-78-1	0.32		5.07E+00	e		5.07E+00	e		5.07E+00
Dimethylaniline	121-69-7	-0.88		2.50E+01	e		2.50E+01	e		2.50E+01
Di-n-propylnitrosamine	621-64-7	1.38		1.23E+00	e		1.23E+00	e		1.23E+00
Diphenyl ether	101-84-8	4.21		2.86E-02	e		2.86E-02	e		2.86E-02
Epichlorohydrin	106-89-8	0.25		5.55E+00	e		5.55E+00	e		5.55E+00
Ethyl carbamate (urethane)	51-79-6	-0.15		9.46E+00	e		9.46E+00	e		9.46E+00
Ethyl methanesulfonate	62-50-0	0.05		7.25E+00	e		7.25E+00	e		7.25E+00
Ethylene dibromide	106-93-4	1.75		7.55E-01	e		7.55E-01	e		7.55E-01
Ethylene glycol	107-21-1	-0.91		2.62E+01	e		2.62E+01	e		2.62E+01
Ethylene glycol monobutyl ether	111-76-2	1.55		9.78E-01	e		9.78E-01	e		9.78E-01
Ethylene glycol monoethyl ether acetate	111-15-9	0.62		3.40E+00	e		3.40E+00	e		3.40E+00
Ethylene thiourea	96-45-7	-0.64		1.82E+01	e		1.82E+01	e		1.82E+01
Furfural	98-01-1	0.96		2.16E+00	e		2.16E+00	e		2.16E+00
Maleic hydrazide	123-33-1	-0.74		2.07E+01	e		2.07E+01	e		2.07E+01
Malononitrile	109-77-3	0.04		7.34E+00	e		7.34E+00	e		7.34E+00
Methyl styrene (mixed isomers)	25013-15-4	3.35		8.97E-02	e		8.97E-02	e		8.97E-02
Methylhydrazine	60-34-4	-1.06		3.17E+01	e		3.17E+01	e		3.17E+01
N,N-Diphenylamine	122-39-4	3.50		7.35E-02	e		7.35E-02	e		7.35E-02
Nitric acid, propyl ester	627-13-4	No data		No data			1.00E+00		Default	1.00E+00

Table C2-1. Terrestrial Soil-to-Plant Transfer Factors (SPv) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPv	SAIC Compiled SPv	Notes	Comments	Recommended SPv[d]
N-Nitrosodi-n-butylamine	924-16-3	2.41		3.13E-01	e		3.13E-01	e		3.13E-01
N-Nitrosomorpholine	59-89-2	0.98		2.10E+00	e		2.10E+00	e		2.10E+00
N-Nitroso-N,N-dimethylamine	62-75-9	-0.47		1.45E+01	e		1.45E+01	e		1.45E+01
o-Anisidine	90-04-0	1.18		1.61E+00	e		1.61E+00	e		1.61E+00
Oxalic acid	144-62-7	No data		No data			1.00E+00		Default	1.00E+00
Phthalic anhydride	85-44-9	-0.60		1.73E+01	e		1.73E+01	e		1.73E+01
p-Phthalic acid	100-21-0	0.82		2.59E+00	e		2.59E+00	e		2.59E+00
Pyridine	110-86-1	0.67		3.17E+00	e		3.17E+00	e		3.17E+00
Quinoline	91-22-5	2.03		5.20E-01	e		5.20E-01	e		5.20E-01
Quinone	106-51-4	0.20		5.94E+00	e		5.94E+00	e		5.94E+00
Safrole	94-59-7	2.66		2.25E-01	e		2.25E-01	e		2.25E-01
Tetrahydrofuran	109-99-9	0.45		4.27E+00	e		4.27E+00	e		4.27E+00
Other Heavy Semivolatile Compounds (molecular weight >200 g/mole)										
1,2,4,5-Tetrachlorobenzene	95-94-3	4.64		1.61E-02	e		1.61E-02	e		1.61E-02
1,3,5-Trinitrobenzene	99-35-4	1.18		1.61E+00	e		1.61E+00	e		1.61E+00
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	4.17		3.01E-02	e		3.01E-02	e		3.01E-02
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	-2.70		2.82E+02	e		2.82E+02	e		2.82E+02
2-sec-Butyl-4,6-dinitrophenol	88-85-7	3.14		1.19E-01	e		1.19E-01	e		1.19E-01
3,3'-Dimethoxybenzidine	119-90-4	1.81		6.96E-01	e		6.96E-01	e		6.96E-01
3,3-Dichlorobenzidine	91-94-1	3.58		6.65E-02	e		6.65E-02	e		6.65E-02
4-Bromophenylphenyl ether	101-55-3	5.00		9.98E-03	e		9.98E-03	e		9.98E-03
Ammonium perfluorooctanoate	3825-26-1	No data		No data			1.00E+00		Default	1.00E+00
Azobenzene	103-33-3	3.82		4.80E-02	e		4.80E-02	e		4.80E-02
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	No data		No data			1.00E+00		Default	1.00E+00
Captan	133-06-2	2.35		3.39E-01	e		3.39E-01	e		3.39E-01
Chlorobenzilate	510-15-6	4.38		2.28E-02	e		2.28E-02	e		2.28E-02
Dibutylphosphate	107-66-4	No data		No data			1.00E+00		Default	1.00E+00
Dimethyl aminoazobenzene	60-11-7	4.58		1.74E-02	e		1.74E-02	e		1.74E-02
Hexachlorobenzene	118-74-1	5.50	2.55E-02	5.10E-03	f		5.11E-03	e		5.10E-03

Table C2-1. Terrestrial Soil-to-Plant Transfer Factors (SPv) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPv	SAIC Compiled SPv	Notes	Comments	Recommended SPv[d]
Hexachlorobutadiene	87-68-3	4.73	7.14E-02	1.43E-02	f		1.43E-02	e		1.43E-02
Hexachlorocyclopentadiene	77-47-4	5.04	5.65E-02	1.13E-02	f		9.46E-03	e		1.13E-02
Hexachloroethane	67-72-1	3.98		3.85E-02	e		3.85E-02	e		3.85E-02
Hexachlorophene	70-30-4	7.54		3.39E-04	e		3.39E-04	e		3.39E-04
Hexamethylene-1,5-diisocyanate	822-06-0	1.27		1.42E+00	e		1.42E+00	e		1.42E+00
Mirex	2385-85-5	6.89		8.07E-04	e		8.07E-04	e		8.07E-04
Nitrofen	1836-75-5	5.53		4.93E-03	e		4.93E-03	e		4.93E-03
Pentachlorobenzene	608-93-5	5.09	4.40E-02	8.80E-03	f		8.89E-03	e		8.80E-03
Pentachloronitrobenzene	82-68-8	4.64	8.00E-02	1.61E-02	f		1.61E-02	e		1.61E-02
Pentachlorophenol	87-86-5	5.08	4.49E-02	8.98E-03	f		8.98E-03	e		8.98E-03
Picric acid	88-89-1	2.03		5.20E-01	e		5.20E-01	e		5.20E-01
Pronamide	23950-58-5	3.51		7.24E-02	e		7.24E-02	e		7.24E-02
Strychnine	57-24-9	1.93		5.94E-01	e		5.94E-01	e		5.94E-01
Terphenyls	26140-60-3	No data		No data			1.00E+00		Default	1.00E+00
Tributyl phosphate	126-73-8	4.00		3.78E-02	e		3.78E-02	e		3.78E-02
Trifluralin	1582-09-8	5.34		6.35E-03	e		6.35E-03	e		6.35E-03
Triphenylamine	603-34-9	No data		No data			1.00E+00		Default	1.00E+00
Herbicides and Organochlorinated Pesticides										
2,4,5-T	93-76-5	3.36		8.85E-02	e		8.85E-02	e		8.85E-02
2,4-D and esters	94-75-7	2.81		1.84E-01	e		1.84E-01	e		1.84E-01
4,4-DDD	72-54-8	6.20		2.02E-03	e		2.02E-03	e		2.02E-03
4,4-DDE	72-55-9	6.26	9.37E-03	1.87E-03	f		1.88E-03	e		1.87E-03
4,4-DDT	50-29-3	6.00		2.64E-03	e		2.64E-03	e		2.64E-03
Aldrin	309-00-2	6.18		2.08E-03	e		2.08E-03	e		2.08E-03
alpha-BHC	319-84-6	3.80		4.93E-02	e		4.93E-02	e		4.93E-02
beta-BHC	319-85-7	3.83		4.71E-02	e		4.71E-02	e		4.71E-02
Chlordane	57-74-9	5.94		2.87E-03	e		2.87E-03	e		2.87E-03
Delta-BHC	319-86-8	4.14		3.13E-02	e		3.13E-02	e		3.13E-02
Dieldrin	60-57-1	5.27		6.97E-03	e		6.97E-03	e		6.97E-03

Table C2-1. Terrestrial Soil-to-Plant Transfer Factors (SPv) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPv	SAIC Compiled SPv	Notes	Comments	Recommended SPv[d]
Endothall	145-73-3	-0.87		2.47E+01	e		2.47E+01	e		2.47E+01
Endrin	72-20-8	4.89		1.15E-02	e		1.15E-02	e		1.15E-02
gamma-BHC (Lindane)	58-89-9	3.72		5.48E-02	e		5.48E-02	e		5.48E-02
Heptachlor	76-44-8	5.02	4.89E-02	9.78E-03	f		9.75E-03	e		9.78E-03
Isodrin	465-73-6	3.55		6.87E-02	e		6.87E-02	e		6.87E-02
Methoxychlor	72-43-5	4.53		1.87E-02	e		1.87E-02	e		1.87E-02
Silvex (2,4,5-TP)	93-72-1	4.07		3.43E-02	e		3.43E-02	e		3.43E-02
Toxaphene	8001-35-2	5.50		5.13E-03	e		5.13E-03	e		5.13E-03
Inorganic Chemicals and Compounds										
Metals										
Aluminum	7429-90-5	NA	4.00E-03	8.00E-04	f		8.00E-04	i		8.00E-04
Antimony	7440-36-0	NA	2.00E-01	4.00E-02	f		4.00E-02	i		4.00E-02
Arsenic	7440-38-2	NA	3.60E-02	7.20E-03	f		8.00E-03	i		7.20E-03
Barium	7440-39-3	NA	1.50E-01	3.00E-02	f		3.00E-02	i		3.00E-02
Beryllium	7440-41-7	NA	1.00E-02	2.00E-03	f		2.00E-03	i		2.00E-03
Bismuth	7440-69-9	NA		7.00E-03	i		7.00E-03	i		7.00E-03
Boron	7440-42-8	NA		8.00E-01	i		8.00E-01	i		8.00E-01
Cadmium	7440-43-9	NA	3.64E-01	7.28E-02	f		1.10E-01	i		7.28E-02
Calcium	7440-70-2	NA		7.00E-01	i		7.00E-01	i		7.00E-01
Chromium (and VI)	18540-29-9	NA	7.50E-03	1.50E-03	f		1.50E-03	i		1.50E-03
Cobalt	7440-48-4	NA		4.00E-03	i		4.00E-03	i		4.00E-03
Copper	7440-50-8	NA	4.00E-01	8.00E-02	f		8.00E-02	i		8.00E-02
Iron	7439-89-6	NA		8.00E-04	i		8.00E-04	i		8.00E-04
Lead	7439-92-1	NA	4.50E-02	9.00E-03	f		9.00E-03	i		9.00E-03
Lithium	7439-93-2	NA		5.00E-03	i		5.00E-03	i		5.00E-03
Magnesium	7439-95-4	NA		2.00E-01	i		2.00E-01	i		2.00E-01
Manganese	7439-96-5	NA		5.00E-02	i		5.00E-02	i		5.00E-02
Mercury	7439-97-6	NA	3.75E-02	7.50E-03	f		1.80E-01	i		7.50E-03
Molybdenum	7439-98-7	NA		5.00E-02	i		5.00E-02	i		5.00E-02

Table C2-1. Terrestrial Soil-to-Plant Transfer Factors (SPv) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPv	SAIC Compiled SPv	Notes	Comments	Recommended SPv[d]
Nickel	7440-02-0	NA	3.20E-02	6.40E-03	f		1.20E-02	i		6.40E-03
Potassium	7440-09-7	NA		2.00E-01	i		2.00E-01	i		2.00E-01
Rhodium	7440-16-6	NA		3.00E-02	i		3.00E-02	i		3.00E-02
Selenium	7782-49-2	NA	1.60E-02	3.20E-03	f		5.00E-03	i		3.20E-03
Silicon	7440-21-3	NA		7.00E-02	i		7.00E-02	i		7.00E-02
Silver	7440-22-4	NA	4.00E-01	8.00E-02	f		8.00E-02	i		8.00E-02
Sodium	7440-23-5	NA		1.50E-02	i		1.50E-02	i		1.50E-02
Strontium	7440-24-6	NA		5.00E-01	i		5.00E-01	i		5.00E-01
Tantalum	7440-25-7	NA		2.00E-03	i		2.00E-03	i		2.00E-03
Thallium	7440-28-0	NA	4.00E-03	8.00E-04	f		8.00E-04	i		8.00E-04
Tin	7440-31-5	NA		6.00E-03	i		6.00E-03	i		6.00E-03
Tungsten	7440-33-7	NA		9.00E-03	i		9.00E-03	i		9.00E-03
Uranium	7440-61-1	NA		1.70E-03	i		1.70E-03	i		1.70E-03
Vanadium	7440-62-2	NA		1.10E-03	i		1.10E-03	i		1.10E-03
Yttrium	7440-65-5	NA		3.00E-03	i		3.00E-03	i		3.00E-03
Zinc	7440-66-6	NA	1.20E-12	2.40E-13	f		3.00E-01	i		2.40E-13
Zirconium	7440-67-7	NA		4.00E-04	i		4.00E-04	i		4.00E-04
Non-metals and Anions										
Ammonia/Ammonium	7664-41-7	NA		2.27E-02	j					2.27E-02
Bromide	24959-67-9	NA		3.00E-01	i		3.00E-01	i		3.00E-01
Chloride	16887-00-6	NA		1.40E+01	i		1.40E+01	i		1.40E+01
Cyanide	57-12-5	NA		2.27E-02	j		1.00E+00	k		2.27E-02
Fluoride	16984-48-8	NA		1.20E-02	i		1.20E-02	i		1.20E-02
Hydroxide	14280-30-9	NA		NA					Depends on pH	NA
Iodine	7553-56-2	NA		3.00E-02	i		3.00E-02	i		3.00E-02
Nitrate	14797-55-8	NA		2.27E-02	j					2.27E-02
Nitrite	14797-65-0	NA		2.27E-02	j					2.27E-02
Phosphate	14265-44-2	NA		7.00E-01	i		7.00E-01	i		7.00E-01
Phosphorus	7723-14-0	NA		7.00E-01	i		7.00E-01	i		7.00E-01

Table C2-1. Terrestrial Soil-to-Plant Transfer Factors (SPv) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPv	SAIC Compiled SPv	Notes	Comments	Recommended SPv[d]
Sulfate	14808-79-8	NA		3.00E-01	i		3.00E-01	i		3.00E-01
Total Sulfur	63705-05-5	NA		3.00E-01	i		3.00E-01	i		3.00E-01
Priority Pollutants										
Carbon Dioxide	124-38-9	NA		2.27E-02	j					2.27E-02
Nitrogen Dioxide	10102-44-0	NA		2.27E-02	j					2.27E-02
Ozone	10028-15-6	NA		2.27E-02	j					2.27E-02
Particulate Matter	No CAS #	NA		NA						NA
Sulfur Dioxode	7446-09-5	NA		2.27E-02	j					2.27E-02
Radionuclides										
Americium-241	1596-10-2	NA		1.10E-03	i		1.10E-03	i		1.10E-03
Antimony-125	14234-35-6	NA		4.00E-02	i		4.00E-02	i		4.00E-02
Barium-137	13981-97-0	NA		3.00E-02	i		3.00E-02	i		3.00E-02
Cadmium-113	None	NA		1.10E-01	i		1.10E-01	i		1.10E-01
Cesium-134	13967-70-9	NA		1.60E-02	i		1.60E-02	i		1.60E-02
Cesium-137	10045-97-3	NA		1.60E-02	i		1.60E-02	i		1.60E-02
Europium-154	15585-10-1	NA		2.00E-03	i		2.00E-03	i		2.00E-03
Europium-155	14391-16-3	NA		2.00E-03	i		2.00E-03	i		2.00E-03
Nickel-63	13981-37-8	NA		1.20E-02	i		1.20E-02	i		1.20E-02
Plutonium-239	15117-48-3	NA		9.00E-05	i		9.00E-05	i		9.00E-05
Plutonium-241	14119-32-5	NA		9.00E-05	i		9.00E-05	i		9.00E-05
Samarium-151	15715-94-3	NA		2.00E-03	i		2.00E-03	i		2.00E-03
Strontium-90	10098-97-2	NA		5.00E-01	i		5.00E-01	i		5.00E-01
Technetium-99	14133-79-7	NA		1.90E+00	i		1.90E+00	i		1.90E+00
Tritium	10028-17-8	NA		2.27E-02	j					2.27E-02
Yttrium-90	10098-91-6	NA		3.00E-03	i		3.00E-03	i		3.00E-03
Uranium-232	14158-29-3	NA		1.70E-03	i		1.70E-03	i		1.70E-03
Uranium-233	13968-55-3	NA		1.70E-03	i		1.70E-03	i		1.70E-03
Uranium-234	13966-29-5	NA		1.70E-03	i		1.70E-03	i		1.70E-03
Uranium-235	15117-96-1	NA		1.70E-03	i		1.70E-03	i		1.70E-03

Table C2-1. Terrestrial Soil-to-Plant Transfer Factors (SPv) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPv	SAIC Compiled SPv	Notes	Comments	Recommended SPv[d]
Uranium-236	13982-70-2	NA		1.70E-03	i		1.70E-03	i		1.70E-03
Uranium-238	7440-61-1	NA		1.70E-03	i		1.70E-03	i		1.70E-03
Actinium-227	14952-40-0	NA		7.00E-04	i		7.00E-04	i		7.00E-04
Americium-243	14993-75-0	NA		1.10E-03	i		1.10E-03	i		1.10E-03
Carbon-14	14762-75-5	NA		2.27E-02	j					2.27E-02
Cobalt-60	10198-40-0	NA		4.00E-03	i		4.00E-03	i		4.00E-03
Curium-242	15510-73-3	NA		1.70E-04	i		1.70E-04	i		1.70E-04
Curium-243	15757-87-6	NA		1.70E-04	i		1.70E-04	i		1.70E-04
Curium-244	13981-15-2	NA		1.70E-04	i		1.70E-04	i		1.70E-04
Europium-152	14683-23-9	NA		2.00E-03	i		2.00E-03	i		2.00E-03
Iodine-129	15046-84-1	NA		3.00E-02	i		3.00E-02	i		3.00E-02
Neptunium-237	13994-20-2	NA		2.00E-02	i		2.00E-02	i		2.00E-02
Nickel-59	14336-70-0	NA	3.20E-02	6.40E-03	f		1.20E-02	i		6.40E-03
Niobium-93	None	NA		4.00E-03	i		4.00E-03	i		4.00E-03
Plutonium-238	13981-16-3	NA		9.00E-05	i		9.00E-05	i		9.00E-05
Plutonium-240	14119-33-6	NA		9.00E-05	i		9.00E-05	i		9.00E-05
Plutonium-242	13982-10-0	NA		9.00E-05	i		9.00E-05	i		9.00E-05
Protactinium-231	14331-85-2	NA		5.00E-04	i		5.00E-04	i		5.00E-04
Radium-226	13982-63-3	NA		3.00E-03	i		3.00E-03	i		3.00E-03
Radium-228	15262-20-1	NA		3.00E-03	i		3.00E-03	i		3.00E-03
Ruthenium-106	13967-48-1	NA		1.50E-02	i		1.50E-02	i		1.50E-02
Selenium-79	None	NA		5.00E-03	i		5.00E-03	i		5.00E-03
Thorium-229	15594-54-4	NA		1.70E-04	i		1.70E-04	i		1.70E-04
Thorium-232	7440-29-1	NA		1.70E-04	i		1.70E-04	i		1.70E-04
Tin-126	15832-50-5	NA		6.00E-03	i		6.00E-03	i		6.00E-03
Zirconium-93	15751-77-6	NA		4.00E-04	i		4.00E-04	i		4.00E-04

NA = Not applicable

[a] $\log_{10}$ of K_{ow} values in Table 4.1

[b] Published in Appendix C of EPA (1999), Table C-2

Table C2-1. Terrestrial Soil-to-Plant Transfer Factors (SPv) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPv	SAIC Compiled SPv	Notes	Comments	Recommended SPv[d]

[c] Calculated or chosen as described in Appendix C, Sect. C-1.2 of EPA (1999), then multiplied by 0.2 to adjust to wet weight of plants

[d] Selection criteria described in Sect. 8.2.4.3

[e] Wet weight values calculated by using log K_{ow}: log SPv = 1.588 - 0.578 x log K_{ow} (Travis and Arms 1988) for dry weight and multiplying by 0.2 to adjust to wet weight

[f] EPA (1999) values published in Table C-2 multiplied by 0.2 to adjust to wet weight of plants

[g] Calculated from value for TCDD in Appendix C of EPA (1999), Table C-2 by multiplying by BEF, then by 0.2 to adjust to wet weight of plants

[h] Value for Aroclor 1254 from Appendix C of EPA (1999), Table C-2, was used for PCB mixtures

[i] Baes and others. (1984) (Figure 2.1) values multiplied by 0.2 to adjust to wet weight of plants

[j] Average of values for other inorganics published in Appendix C of EPA (1999), Table C-2, then multiplied by 0.2 to adjust to wet weight of plants

[k] HAZWRAP (1994), Table 4.4

Table C2-2. Terrestrial Soil-to-Plant Transfer Factors (SPr) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPr	SAIC Compiled SPr	Notes	Comments	Recommended SPr[d]
Organic Compounds										
Aromatic Halogenated Hydrocarbons										
4-Chloro-3-methylphenol	59-50-7	3.10		1.25E-01	e		1.25E-01	e		1.25E-01
2,3,4,6-Tetrachlorophenol	58-90-2	4.42		2.16E-02	e		2.16E-02	e		2.16E-02
Aromatic Nonhalogenated Hydrocarbons										
2-Nitrotoluene	88-72-2	2.30		3.63E-01	e		3.63E-01	e		3.63E-01
4-Nitrobiphenyl	92-93-3	3.77		5.13E-02	e		5.13E-02	e		5.13E-02
Benzaldehyde	100-52-7	1.48		1.08E+00	e		1.08E+00	e		1.08E+00
Benzene	71-43-2	-4.99		5.94E+03	e		5.94E+03	e		5.94E+03
Benzyl alcohol	100-51-6	1.10		1.79E+00	e		1.79E+00	e		1.79E+00
Ethyl benzene	100-41-4	3.12		1.21E-01	e		1.21E-01	e		1.21E-01
m-Xylene	108-38-3	3.20		1.09E-01	e		1.09E-01	e		1.09E-01
o-Xylene	95-47-6	3.13		1.20E-01	e		1.20E-01	e		1.20E-01
p-Xylene	106-42-3	3.17		1.14E-01	e		1.14E-01	e		1.14E-01
Styrene	100-42-5	2.93		1.57E-01	e		1.57E-01	e		1.57E-01
Toluene	108-88-3	2.67		2.22E-01	e		2.22E-01	e		2.22E-01
Non-aromatic Nonhalogenated Hydrocarbons										
1,2-Epoxybutane	106-88-7	1.44		1.14E+00	e		1.14E+00	e		1.14E+00
1,3-Butadiene	106-99-0	1.90		6.18E-01	e		6.18E-01	e		6.18E-01
1,4-Dioxane	123-91-1	-0.27	5.53E+01	1.11E+01	f		1.11E+01	e		1.11E+01
1-Methylpropyl alcohol	78-92-2	0.61		3.44E+00	e		3.44E+00	e		3.44E+00
1-Nitropropane	108-03-2	0.87		2.43E+00	e		2.43E+00	e		2.43E+00
2,2,4-Trimethylpentane	540-84-1	5.02		9.72E-03	e		9.72E-03	e		9.72E-03
2-Butanone	78-93-3	0.28		5.33E+00	e		5.33E+00	e		5.33E+00
2-Butenaldehyde (2-Butenal)	4170-30-3	No data		No data			1.00E+00		Default	1.00E+00
2-Ethoxyethanol	110-80-5	-0.10		8.85E+00	e		8.85E+00	e		8.85E+00
2-Heptanone	110-43-0	1.98		5.55E-01	e		5.55E-01	e		5.55E-01

Table C2-2. Terrestrial Soil-to-Plant Transfer Factors (SPr) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPr	SAIC Compiled SPr	Notes	Comments	Recommended SPr[d]
2-Hexanone	591-78-6	1.38		1.23E+00	e		1.23E+00	e		1.23E+00
2-Methoxyethanol	109-86-4	0.25		5.54E+00	e		5.54E+00	e		5.54E+00
2-Methyl-2-propanol	75-65-0	0.35		4.86E+00	e		4.86E+00	e		4.86E+00
2-Methyl-2-propenenitrile	126-98-7	0.54		3.77E+00	e		3.77E+00	e		3.77E+00
2-Methylaziridine	75-55-8	-0.60		1.73E+01	e		1.73E+01	e		1.73E+01
2-Methylpropyl alcohol	78-83-1	0.76		2.82E+00	e		2.82E+00	e		2.82E+00
2-Pentanone	107-87-9	0.91		2.31E+00	e		2.31E+00	e		2.31E+00
2-Propanone (Acetone)	67-64-1	-0.22	5.20E+01	1.04E+01	f		1.04E+01	e		1.04E+01
2-Propene-1-ol	107-18-6	0.17		6.18E+00	e		6.18E+00	e		6.18E+00
2-Propyl alcohol	67-63-0	0.05		7.25E+00	e		7.25E+00	e		7.25E+00
3-Heptanone	106-35-4	No data		No data			1.00E+00		Default	1.00E+00
3-Methyl-1-butanol	123-51-3	No data		No data			1.00E+00		Default	1.00E+00
3-Methyl-2-butanone	563-80-4	No data		No data			1.00E+00		Default	1.00E+00
3-Pentanone	96-22-0	0.99		2.07E+00	e		2.07E+00	e		2.07E+00
4-Heptanone	123-19-3	No data		No data			1.00E+00		Default	1.00E+00
4-Methyl-2-pentanone	108-10-1	1.19		1.59E+00	e		1.59E+00	e		1.59E+00
4-Methyl-3-penten-2-one	141-79-7	No data		No data			1.00E+00		Default	1.00E+00
5-Methyl-2-hexanone	110-12-3	No data		No data			1.00E+00		Default	1.00E+00
Acetaldehyde	75-07-0	-0.22		1.04E+01	e		1.04E+01	e		1.04E+01
Acetamide	60-35-5	-1.26		4.14E+01	e		4.14E+01	e		4.14E+01
Acetic acid	64-19-7	-0.17		9.71E+00	e		9.71E+00	e		9.71E+00
Acetic acid ethyl ester	141-78-6	0.73		2.93E+00	e		2.93E+00	e		2.93E+00
Acetic acid n-butyl ester	123-86-4	1.73		7.75E-01	e		7.75E-01	e		7.75E-01
Acetonitrile	75-05-8	-0.34		1.22E+01	e		1.22E+01	e		1.22E+01
Acrolein	107-02-8	-0.01		7.84E+00	e		7.84E+00	e		7.84E+00
Acrylonitrile	107-13-1	0.25	2.78E+01	5.55E+00	f		5.55E+00	e		5.55E+00
Bis(isopropyl)ether	108-20-3	No data		No data			1.00E+00		Default	1.00E+00

Table C2-2. Terrestrial Soil-to-Plant Transfer Factors (SPr) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPr	SAIC Compiled SPr	Notes	Comments	Recommended SPr[d]
Butane	106-97-8	2.89		1.65E-01	e		1.65E-01	e		1.65E-01
Carbon disulfide	75-15-0	2.00		5.41E-01	e		5.41E-01	e		5.41E-01
Cyanogen	460-19-5	0.81		2.64E+00	e		2.64E+00	e		2.64E+00
Cyclohexane	110-82-7	3.44		7.96E-02	e		7.96E-02	e		7.96E-02
Cyclohexanone	108-94-1	0.81		2.64E+00	e		2.64E+00	e		2.64E+00
Cyclohexene	110-83-8	2.86		1.72E-01	e		1.72E-01	e		1.72E-01
Cyclopentane	287-92-3	3.00		1.43E-01	e		1.43E-01	e		1.43E-01
Ethyl alcohol	64-17-5	0.31		5.13E+00	e		5.13E+00	e		5.13E+00
Ethyl ether	60-29-7	0.89		2.37E+00	e		2.37E+00	e		2.37E+00
Ethyl methacrylate	97-63-2	1.59		9.33E-01	e		9.33E-01	e		9.33E-01
Formaldehyde	50-00-0	0.34	2.46E+01	4.91E+00	f		4.91E+00	e		4.91E+00
Formamide	75-12-7	-1.51		5.78E+01	e		5.78E+01	e		5.78E+01
Formic acid	64-18-6	-0.54		1.58E+01	e		1.58E+01	e		1.58E+01
Formic acid, methyl ester	107-31-3	-0.26		1.10E+01	e		1.10E+01	e		1.10E+01
Glycidylaldehyde	765-34-4	-0.73		2.05E+01	e		2.05E+01	e		2.05E+01
Methyl acetate	79-20-9	0.18		6.10E+00	e		6.10E+00	e		6.10E+00
Methyl alcohol	67-56-1	-0.71		1.99E+01	e		1.99E+01	e		1.99E+01
Methyl isocyanate	624-83-9	No data		No data			NA		Reacts with water	NA
Methyl methacrylate	80-62-6	0.79		2.71E+00	e		2.71E+00	e		2.71E+00
Methyl tert-butyl ether	1634-04-4	0.94		2.22E+00	e		2.22E+00	e		2.22E+00
Methylacetylene	74-99-7	0.94		2.22E+00	e		2.22E+00	e		2.22E+00
Methylcyclohexane	108-87-2	4.10		3.31E-02	e		3.31E-02	e		3.31E-02
N,N-Dimethylacetamide	127-19-5	No data		No data			1.00E+00		Default	1.00E+00
n-Butyl alcohol	71-36-3	0.88		2.40E+00	e		2.40E+00	e		2.40E+00
n-Heptane	142-82-5	4.66		1.57E-02	e		1.57E-02	e		1.57E-02
n-Hexane	110-54-3	4.11		3.26E-02	e		3.26E-02	e		3.26E-02
Nitromethane	75-52-5	-0.35		1.23E+01	e		1.23E+01	e		1.23E+01

Table C2-2. Terrestrial Soil-to-Plant Transfer Factors (SPr) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPr	SAIC Compiled SPr	Notes	Comments	Recommended SPr[d]
n-Nonane	111-84-2	5.65		4.20E-03	e		4.20E-03	e		4.20E-03
n-Octane	111-65-9	4.00		3.78E-02	e		3.78E-02	e		3.78E-02
n-Pentane	109-66-0	3.21		1.08E-01	e		1.08E-01	e		1.08E-01
n-Propionaldehyde	123-38-6	0.59		3.53E+00	e		3.53E+00	e		3.53E+00
n-Propyl alcohol	71-23-8	0.25		5.55E+00	e		5.55E+00	e		5.55E+00
n-Valeraldehyde	110-62-3	No data		No data	e		1.00E+00		Default	1.00E+00
Oxirane	75-21-8	-0.30		1.15E+01	e		1.15E+01	e		1.15E+01
p-Cymene	99-87-6	4.10		3.31E-02	e		3.31E-02	e		3.31E-02
Phosgene	75-44-5	No data		No data	e		1.00E+00		Default	1.00E+00
Propargyl alcohol	107-19-7	0.26		5.48E+00	e		5.48E+00	e		5.48E+00
Propionic acid	79-09-4	0.33		4.99E+00	e		4.99E+00	e		4.99E+00
Propionitrile	107-12-0	0.16		6.26E+00	e		6.26E+00	e		6.26E+00
Propylene gylcol monomethyl ether	107-98-2	-0.18		9.84E+00	e		9.84E+00	e		9.84E+00
p-tert-Butyltoluene	98-51-1	No data		No data			1.00E+00		Default	1.00E+00
Triethylamine	121-44-8	0.16		6.26E+00	e		6.26E+00	e		6.26E+00
Trimethylamine	75-50-3	0.16		6.26E+00	e		6.26E+00	e		6.26E+00
Vinyl acetate	108-05-4	0.70		3.06E+00	e		3.06E+00	e		3.06E+00
Non-aromatic Halogenated Hydrocarbons										
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	No data		No data			1.00E+00		Default	1.00E+00
1,1,1,2-Tetrachloroethane	630-20-6	2.63		2.34E-01	e		2.34E-01	e		2.34E-01
1,1,1-Trichloroethane	71-55-6	2.42		3.09E-01	e		3.09E-01	e		3.09E-01
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	3.73		5.41E-02	e		5.41E-02	e		5.41E-02
1,1,2,2-Tetrachloroethane	79-34-5	4.64		1.60E-02	e		1.60E-02	e		1.60E-02
1,1,2,2-Tetrachloroethene	127-18-4	2.55		2.62E-01	e		2.62E-01	e		2.62E-01
1,1,2-Trichloroethane	79-00-5	2.10		4.75E-01	e		4.75E-01	e		4.75E-01
1,1,2-Trichloroethylene	79-01-6	2.43		3.04E-01	e		3.04E-01	e		3.04E-01
1,1-Dichloroethane	75-34-3	1.79		7.13E-01	e		7.13E-01	e		7.13E-01

Table C2-2. Terrestrial Soil-to-Plant Transfer Factors (SPr) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPr	SAIC Compiled SPr	Notes	Comments	Recommended SPr[d]
1,1-Dichloroethene	75-35-4	2.12		4.61E-01	e		4.61E-01	e		4.61E-01
1,2,2-Trichloro-1,1,2-trifluoroethane	76-13-1	3.16		1.15E-01	e		1.15E-01	e		1.15E-01
1,2,3-Trichloropropane	96-18-4	2.25		3.88E-01	e		3.88E-01	e		3.88E-01
1,2-Dibromo-3-chloropropane	96-12-8	2.34		3.44E-01	e		3.44E-01	e		3.44E-01
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	2.82		1.82E-01	e		1.82E-01	e		1.82E-01
1,2-Dichloroethane	107-06-2	1.46		1.11E+00	e		1.11E+00	e		1.11E+00
1,2-Dichloroethylene	540-59-0	0.48		4.09E+00	e		4.09E+00	e		4.09E+00
1,2-Dichloropropane	78-87-5	2.25		3.88E-01	e		3.88E-01	e		3.88E-01
1,3-Dichloropropene	542-75-6	1.75		7.56E-01	e		7.56E-01	e		7.56E-01
1,4-Dichloro-2-butene	764-41-0	0.87		2.42E+00	e		2.42E+00	e		2.42E+00
1-Chloroethene	75-01-4	1.15		1.68E+00	e		1.68E+00	e		1.68E+00
2,2-Dichloropropionic acid	75-99-0	0.78		2.75E+00	e		2.75E+00	e		2.75E+00
2-Chloropropane	75-29-6	1.90		6.18E-01	e		6.18E-01	e		6.18E-01
3-Chloropropene (allyl chloride)	107-05-1	0.95		2.17E+00	e		2.17E+00	e		2.17E+00
Bromochloromethane	74-97-5	1.41		1.19E+00	e		1.19E+00	e		1.19E+00
Bromodichloromethane	75-27-4	2.03		5.23E-01	e		5.23E-01	e		5.23E-01
Bromoethene	593-60-2	1.07		1.87E+00	e		1.87E+00	e		1.87E+00
Bromoform	75-25-2	2.35		3.39E-01	e		3.39E-01	e		3.39E-01
Bromomethane	74-83-9	1.11		1.76E+00	e		1.76E+00	e		1.76E+00
Carbon tetrachloride	56-23-5	2.72	1.04E+00	2.08E-01	f		2.08E-01	e		2.08E-01
Chlorodibromomethane	124-48-1	2.18		4.28E-01	e		4.28E-01	e		4.28E-01
Chlorodifluoromethane	75-45-6	1.08		1.84E+00	e		1.84E+00	e		1.84E+00
Chloroethane	75-00-3	3.10		1.25E-01	e		1.25E-01	e		1.25E-01
Chloroform	67-66-3	1.95	2.90E+00	5.80E-01	f		5.78E-01	e		5.80E-01
Chloromethane	74-87-3	0.90		2.33E+00	e		2.33E+00	e		2.33E+00
Chloropentafluoroethane	76-15-3	No data		No data			1.00E+00		Default	1.00E+00
cis-1,2-Dichloroethene	156-59-2	1.98		5.54E-01	e		5.54E-01	e		5.54E-01

Table C2-2. Terrestrial Soil-to-Plant Transfer Factors (SPr) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPr	SAIC Compiled SPr	Notes	Comments	Recommended SPr[d]
cis-1,3-Dichloropropene	10061-01-5	No data		No data			1.00E+00		Default	1.00E+00
Cyanogen bromide	506-68-3	No data		No data			1.00E+00		Default	1.00E+00
Cyanogen chloride	506-77-4	0.20		5.94E+00	e		5.94E+00	e		5.94E+00
Dichlorodifluoromethane	75-71-8	2.16		4.38E-01	e		4.38E-01	e		4.38E-01
Dichlorofluoromethane	75-43-4	No data		No data			1.00E+00		Default	1.00E+00
Dichloromethane	75-09-2	1.26		1.46E+00	e		1.46E+00	e		1.46E+00
Difluorodibromomethane	75-61-6	No data		No data			1.00E+00		Default	1.00E+00
Hexafluoroacetone	684-16-2	No data		No data			1.00E+00		Default	1.00E+00
Iodomethane	74-88-4	1.69		8.17E-01	e		8.17E-01	e		8.17E-01
Methylene bromide	74-95-3	1.62		8.97E-01	e		8.97E-01	e		8.97E-01
Pentachloroethane	76-01-7	3.05		1.34E-01	e		1.34E-01	e		1.34E-01
trans-1,2-Dichloroethene	156-60-5	1.98		5.54E-01	e		5.54E-01	e		5.54E-01
trans-1,3-Dichloropropene	10061-02-6	2.06		4.99E-01	e		4.99E-01	e		4.99E-01
Trichloroacetic acid	76-03-9	1.33		1.32E+00	e		1.32E+00	e		1.32E+00
Trichlorofluoroethane	27154-33-2	No data		No data			1.00E+00		Default	1.00E+00
Trichlorofluoromethane	75-69-4	2.53		2.67E-01	e		2.67E-01	e		2.67E-01
Trifluorobromomethane	75-63-8	1.86		6.52E-01	e		6.52E-01	e		6.52E-01
Dioxin and Furan Compounds										
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	35822-46-9	8.20	2.90E-04	5.80E-05	g		5.80E-05	g		5.80E-05
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	7.92	6.20E-05	1.24E-05	g		1.24E-05	g		1.24E-05
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	7.92	2.20E-03	4.40E-04	g		4.40E-04	g		4.40E-04
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	39227-28-6	7.79	1.70E-03	3.40E-04	g		3.40E-04	g		3.40E-04
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	7.25	4.30E-04	8.60E-05	g		8.60E-05	g		8.60E-05
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	57653-85-7	7.25	6.70E-04	1.34E-04	g		1.34E-04	g		1.34E-04
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	7.25	1.10E-03	2.20E-04	g		2.20E-04	g		2.20E-04
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	19408-74-3	7.25	7.80E-04	1.56E-04	g		1.56E-04	g		1.56E-04
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	7.25	3.50E-03	7.00E-04	g		7.00E-04	g		7.00E-04

Table C2-2. Terrestrial Soil-to-Plant Transfer Factors (SPr) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPr	SAIC Compiled SPr	Notes	Comments	Recommended SPr[d]
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	40321-76-4	6.64	5.20E-03	1.04E-03	g		1.04E-03	g		1.04E-03
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	6.79	1.10E-03	2.20E-04	g		2.20E-04	g		2.20E-04
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	7.25	3.80E-03	7.60E-04	g		7.60E-04	g		7.60E-04
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	6.92	9.00E-03	1.80E-03	g		1.80E-03	g		1.80E-03
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1746-01-6	6.64	5.62E-03	1.12E-03	e		1.12E-03	e		1.12E-03
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	6.53	4.50E-03	9.00E-04	g		9.00E-04	g		9.00E-04
Dibenzofuran	132-64-9	4.33		2.43E-02	e		2.43E-02	e		2.43E-02
Octachlorodibenzo(p)dioxin	3268-87-9	8.20	6.70E-05	1.34E-05	g		1.34E-05	g		1.34E-05
Octachlorodibenzofuran	39001-02-0	8.78	9.00E-05	1.80E-05	g		1.80E-05	g		1.80E-05
PCBs										
Polychlorinated biphenyls (PCBs)[h]	1336-36-3	7.31	1.00E-02	2.00E-03	f		4.63E-04	e		2.00E-03
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	7.08		6.26E-04	e		6.26E-04	e		6.26E-04
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	7.12		5.94E-04	e		5.94E-04	e		5.94E-04
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	No data		No data	e		1.00E+00		Default	1.00E+00
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	No data		No data	e		1.00E+00		Default	1.00E+00
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	No data		No data	e		1.00E+00		Default	1.00E+00
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	No data		No data	e		1.00E+00		Default	1.00E+00
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	No data		No data	e		1.00E+00		Default	1.00E+00
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	No data		No data	e		1.00E+00		Default	1.00E+00
2',3,4,4',5-Pentachlorobiphenyl	no cas #	No data		No data	e		1.00E+00		Default	1.00E+00
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	7.12		5.94E-04	e		5.94E-04	e		5.94E-04
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	7.41		4.05E-04	e		4.05E-04	e		4.05E-04
3,3',4,4',5-Pentachlorobiphenyl	no cas #	No data		No data	e		1.00E+00		Default	1.00E+00
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	No data		No data	e		1.00E+00		Default	1.00E+00
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	No data		No data	e		1.00E+00		Default	1.00E+00
Phthalates										
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	5.20	3.80E-02	7.60E-03	f		7.60E-03	e		7.60E-03

Table C2-2. Terrestrial Soil-to-Plant Transfer Factors (SPr) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPr	SAIC Compiled SPr	Notes	Comments	Recommended SPr[d]
Butylbenzyl phthalate	85-68-7	4.41		2.18E-02	e		2.18E-02	e		2.18E-02
Dibutyl phthalate	84-74-2	4.72		1.45E-02	e		1.45E-02	e		1.45E-02
Diethyl phthalate	84-66-2	4.44		2.11E-02	e		2.11E-02	e		2.11E-02
Dimethylphthalate	131-11-3	1.63		8.81E-01	e		8.81E-01	e		8.81E-01
n-Dioctyl phthalate	117-84-0	9.33	1.57E-04	3.14E-05	f		3.13E-05	e		3.14E-05
Light Polycyclic Aromatic Hydrocarbons (molecular weight <200 g/mole)										
2-Chloronapthalene	91-58-7	4.07		3.45E-02	e		3.45E-02	e		3.45E-02
2-Methyl naphthalene	91-57-6	3.86		4.55E-02	e		4.55E-02	e		4.55E-02
5-Nitroacenaphthene	602-87-9	No data		No data			1.00E+00		Default	1.00E+00
Acenaphthene	83-32-9	3.96		3.96E-02	e		3.96E-02	e		3.96E-02
Acenaphthylene	208-96-8	4.07		3.44E-02	e		3.44E-02	e		3.44E-02
Anthracene	120-12-7	4.47		2.02E-02	e		2.02E-02	e		2.02E-02
Fluorene	86-73-7	4.18		2.97E-02	e		2.97E-02	e		2.97E-02
Indene	95-13-6	No data		No data			1.00E+00		Default	1.00E+00
Naphthalene	91-20-3	3.37		8.70E-02	e		8.70E-02	e		8.70E-02
Phenanthrene	85-01-8	4.55		1.82E-02	e		1.82E-02	e		1.82E-02
Pyrene	129-00-0	5.00		9.98E-03	e		9.98E-03	e		9.98E-03
Heavy Polycyclic Aromatic Hydrocarbons (molecular weight >200 g/mole)										
3-Methylcholanthrene	56-49-5	7.11		6.02E-04	e		6.02E-04	e		6.02E-04
5-Methylchrysene	3697-24-3	No data		No data			1.00E+00		Default	1.00E+00
Benzo(a)anthracene	56-55-3	5.68	2.02E-02	4.04E-03	f		4.04E-03	e		4.04E-03
Benzo(a)pyrene	50-32-8	6.13	1.11E-02	2.22E-03	f		2.22E-03	e		2.22E-03
Benzo(b)fluoranthene	205-99-2	6.20	1.01E-02	2.02E-03	f		2.02E-03	e		2.02E-03
Benzo(e)pyrene	192-97-2	7.40		4.09E-04	e		4.09E-04	e		4.09E-04
Benzo(g,h,i)perylene	191-24-2	7.10		6.10E-04	e		6.10E-04	e		6.10E-04
Benzo(j)fluoranthene	205-82-3	6.44		1.47E-03	e		1.47E-03	e		1.47E-03
Benzo(k)fluoranthene	207-08-9	6.20	1.01E-02	2.02E-03	f		2.30E-03	e		2.02E-03

Table C2-2. Terrestrial Soil-to-Plant Transfer Factors (SPr) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPr	SAIC Compiled SPr	Notes	Comments	Recommended SPr[d]
Benzo[a,i]pyrene	191-30-0	No data		No data			1.00E+00		Default	1.00E+00
Chrysene	218-01-9	5.74	1.87E-02	3.74E-03	f		3.73E-03	e		3.74E-03
Dibenz(a,h)anthracene	53-70-3	6.55	6.40E-03	1.28E-03	f		1.27E-03	e		1.28E-03
Dibenz[a,h]acridine	226-36-8	No data		No data			1.00E+00		Default	1.00E+00
Dibenz[a,j]acridine	224-42-0	No data		No data			1.00E+00		Default	1.00E+00
Dibenzo(a,e)fluoranthene	5385-75-1	No data		No data			1.00E+00		Default	1.00E+00
Dibenzo(a,h)fluoranthene	no cas #	No data		No data			1.00E+00		Default	1.00E+00
Dibenzo[a,e]pyrene	192-65-4	No data		No data			1.00E+00		Default	1.00E+00
Dibenzo[a,h]pyrene	189-64-0	No data		No data			1.00E+00		Default	1.00E+00
Dibenzo[a,i]pyrene	189-55-9	7.29		4.74E-04	e		4.74E-04	e		4.74E-04
Fluoranthene	206-44-0	5.08		8.94E-03	e		8.94E-03	e		8.94E-03
Hexachloronaphthalene	1335-87-1	7.59		3.18E-04	e		3.18E-04	e		3.18E-04
Indeno(1,2,3-cd)pyrene	193-39-5	6.91	3.90E-03	7.80E-04	f		7.80E-04	e		7.80E-04
Octachloronaphthalene	2234-13-1	6.42		1.51E-03	e		1.51E-03	e		1.51E-03
Pentachloronaphthalene	1321-64-8	No data		No data			1.00E+00		Default	1.00E+00
Tetrachloronaphthalene	1335-88-2	No data		No data			1.00E+00		Default	1.00E+00
Trichloronaphthalene	1321-65-9	No data		No data			1.00E+00		Default	1.00E+00
Light Substituted Benzene Compounds (MW <200 g/mole)										
1,2,3-Trichlorobenzene	87-61-6	4.05		3.55E-02	e		3.55E-02	e		3.55E-02
1,2,4-Trichlorobenzene	120-82-1	3.99		3.84E-02	e		3.84E-02	e		3.84E-02
1,2,4-Trimethyl benzene	95-63-6	3.65		6.02E-02	e		6.02E-02	e		6.02E-02
1,2-Dichlorobenzene	95-50-1	3.45		7.90E-02	e		7.90E-02	e		7.90E-02
1,3,5-Trimethyl benzene	108-67-8	3.42		8.17E-02	e		8.17E-02	e		8.17E-02
1,3-Dichlorobenzene	541-73-1	3.53		7.06E-02	e		7.06E-02	e		7.06E-02
1,3-Dinitrobenzene	99-65-0	1.49	5.32E+00	1.06E+00	f		1.06E+00	e		1.06E+00
1,4-Dichlorobenzene	106-46-7	3.41		8.26E-02	e		8.26E-02	e		8.26E-02
1,4-Dinitrobenzene	100-25-4	1.50		1.05E+00	e		1.05E+00	e		1.05E+00

Table C2-2. Terrestrial Soil-to-Plant Transfer Factors (SPr) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPr	SAIC Compiled SPr	Notes	Comments	Recommended SPr[d]
2,4,5-Trichlorophenol	95-95-4	3.87		4.49E-02	e		4.49E-02	e		4.49E-02
2,4,6-Trichlorophenol	88-06-2	3.71		5.54E-02	e		5.54E-02	e		5.54E-02
2,4-Dichlorophenol	120-83-2	3.04		1.36E-01	e		1.36E-01	e		1.36E-01
2,4-Dimethylphenol	105-67-9	2.36		3.35E-01	e		3.35E-01	e		3.35E-01
2,4-Dinitrophenol	51-28-5	1.52		1.03E+00	e		1.03E+00	e		1.03E+00
2,4-Dinitrotoluene	121-14-2	2.00	2.72E+00	5.44E-01	f		5.44E-01	e		5.44E-01
2,6-Dinitrotoluene	606-20-2	1.89	3.15E+00	6.30E-01	f		6.29E-01	e		6.30E-01
2-Chlorophenol	95-57-8	2.16		4.36E-01	e		4.36E-01	e		4.36E-01
2-Chlorotoluene	95-49-8	3.54		7.00E-02	e		7.00E-02	e		7.00E-02
2-Nitrophenol	88-75-5	1.79		7.15E-01	e		7.15E-01	e		7.15E-01
4,6-Dinitro-o-cresol	534-52-1	2.85		1.74E-01	e		1.74E-01	e		1.74E-01
4-Chlorotoluene	106-43-4	3.33		9.21E-02	e		9.21E-02	e		9.21E-02
4-Nitrophenol	100-02-7	1.91		6.10E-01	e		6.10E-01	e		6.10E-01
alpha-Methylstyrene	98-83-9	3.46		7.71E-02	e		7.71E-02	e		7.71E-02
Aniline	62-53-3	0.98		2.10E+00	e		2.10E+00	e		2.10E+00
Benzotrichloride	98-07-7	2.92		1.59E-01	e		1.59E-01	e		1.59E-01
Benzyl chloride	100-44-7	0.36		4.79E+00	e		4.79E+00	e		4.79E+00
Bromobenzene	108-86-1	2.99		1.45E-01	e		1.45E-01	e		1.45E-01
Chlorobenzene	108-90-7	2.79		1.89E-01	e		1.89E-01	e		1.89E-01
Cumene	98-82-8	3.61		6.32E-02	e		6.32E-02	e		6.32E-02
m-Cresol	108-39-4	1.96		5.71E-01	e		5.71E-01	e		5.71E-01
n-Butyl benzene	104-51-8	4.28		2.60E-02	e		2.60E-02	e		2.60E-02
Nitrobenzene	98-95-3	1.83	3.38E+00	6.76E-01	f		6.76E-01	e		6.76E-01
n-Propyl benzene	103-65-1	3.69		5.70E-02	e		5.70E-02	e		5.70E-02
o-Cresol	95-48-7	2.02		5.26E-01	e		5.26E-01	e		5.26E-01
o-Dinitrobenzene	528-29-0	1.60		9.21E-01	e		9.21E-01	e		9.21E-01
o-Nitroaniline	88-74-4	1.85		6.60E-01	e		6.60E-01	e		6.60E-01

Table C2-2. Terrestrial Soil-to-Plant Transfer Factors (SPr) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPr	SAIC Compiled SPr	Notes	Comments	Recommended SPr[d]
o-Toluidine	95-53-4	1.34		1.30E+00	e		1.30E+00	e		1.30E+00
p-Chloroaniline	106-47-8	1.87		6.44E-01	e		6.44E-01	e		6.44E-01
p-Cresol	106-44-5	1.94		5.86E-01	e		5.86E-01	e		5.86E-01
Phenol	108-95-2	1.48		1.08E+00	e		1.08E+00	e		1.08E+00
p-Nitrochlorobenzene	100-00-5	2.39		3.22E-01	e		3.22E-01	e		3.22E-01
p-Toluidine	106-49-0	1.40		1.20E+00	e		1.20E+00	e		1.20E+00
sec-Butyl benzene	135-98-8	4.57		1.77E-02	e		1.77E-02	e		1.77E-02
tert-Butyl benzene	98-06-6	4.11		3.26E-02	e		3.26E-02	e		3.26E-02
Toluene-2,6-diamine	823-40-5	1.45		1.12E+00	e		1.12E+00	e		1.12E+00
Trimethyl benzene	25551-13-7	3.40		8.39E-02	e		8.39E-02	e		8.39E-02
Other Light Semivolatile Compounds (molecular weight <200 g/mole)										
1,1'-Biphenyl	92-52-4	3.90		4.31E-02	e		4.31E-02	e		4.31E-02
1,1-Dimethylhydrazine	57-14-7	0.60		3.51E+00	e		3.51E+00	e		3.51E+00
1,2-Dimethylhydrazine	540-73-8	-1.37		4.78E+01	e		4.78E+01	e		4.78E+01
1,2-Diphenylhydrazine	122-66-7	2.94		1.55E-01	e		1.55E-01	e		1.55E-01
1,3-Propane sultone	1120-71-4	-0.52		1.55E+01	e		1.55E+01	e		1.55E+01
2,4-Toluene diisocyante	584-84-9	No data		No data			NA		Decomposes rapidly in water	NA
2-Chloroacetophenone	532-27-4	2.59		2.45E-01	e		2.45E-01	e		2.45E-01
2-Propenoic acid	79-10-7	0.33		4.99E+00	e		4.99E+00	e		4.99E+00
4,4-Methylenedianiline	101-77-9	3.38		8.62E-02	e		8.62E-02	e		8.62E-02
Acetophenone	98-86-2	1.64		8.73E-01	e		8.73E-01	e		8.73E-01
Benzoic acid	65-85-0	1.86		6.52E-01	e		6.52E-01	e		6.52E-01
bis(2-Chloroethoxy)methane	111-91-1	7.59		3.18E-04	e		3.18E-04	e		3.18E-04
bis(2-Chloroethyl) ether	111-44-4	1.30		1.37E+00	e		1.37E+00	e		1.37E+00
Chlorocyclopentadiene	41851-50-7	2.43		3.05E-01	e		3.05E-01	e		3.05E-01
Cyclohexanol	108-93-0	1.23		1.51E+00	e		1.51E+00	e		1.51E+00
Dichloroisopropyl ether	108-60-1	2.58		2.50E-01	e		2.50E-01	e		2.50E-01

Table C2-2. Terrestrial Soil-to-Plant Transfer Factors (SPr) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPr	SAIC Compiled SPr	Notes	Comments	Recommended SPr[d]
Dichloromethyl ether	542-88-1	-0.38		1.28E+01	e		1.28E+01	e		1.28E+01
Dichloropentadiene	no cas #	No data		No data			1.00E+00		Default	1.00E+00
Dimethyl sulfate	77-78-1	0.32		5.07E+00	e		5.07E+00	e		5.07E+00
Dimethylaniline	121-69-7	-0.88		2.50E+01	e		2.50E+01	e		2.50E+01
Di-n-propylnitrosamine	621-64-7	1.38		1.23E+00	e		1.23E+00	e		1.23E+00
Diphenyl ether	101-84-8	4.21		2.86E-02	e		2.86E-02	e		2.86E-02
Epichlorohydrin	106-89-8	0.25		5.55E+00	e		5.55E+00	e		5.55E+00
Ethyl carbamate (urethane)	51-79-6	-0.15		9.46E+00	e		9.46E+00	e		9.46E+00
Ethyl methanesulfonate	62-50-0	0.05		7.25E+00	e		7.25E+00	e		7.25E+00
Ethylene dibromide	106-93-4	1.75		7.55E-01	e		7.55E-01	e		7.55E-01
Ethylene glycol	107-21-1	-0.91		2.62E+01	e		2.62E+01	e		2.62E+01
Ethylene glycol monobutyl ether	111-76-2	1.55		9.78E-01	e		9.78E-01	e		9.78E-01
Ethylene glycol monoethyl ether acetate	111-15-9	0.62		3.40E+00	e		3.40E+00	e		3.40E+00
Ethylene thiourea	96-45-7	-0.64		1.82E+01	e		1.82E+01	e		1.82E+01
Furfural	98-01-1	0.96		2.16E+00	e		2.16E+00	e		2.16E+00
Maleic hydrazide	123-33-1	-0.74		2.07E+01	e		2.07E+01	e		2.07E+01
Malononitrile	109-77-3	0.04		7.34E+00	e		7.34E+00	e		7.34E+00
Methyl styrene (mixed isomers)	25013-15-4	3.35		8.97E-02	e		8.97E-02	e		8.97E-02
Methylhydrazine	60-34-4	-1.06		3.17E+01	e		3.17E+01	e		3.17E+01
N,N-Diphenylamine	122-39-4	3.50		7.35E-02	e		7.35E-02	e		7.35E-02
Nitric acid, propyl ester	627-13-4	No data		No data			1.00E+00		Default	1.00E+00
N-Nitrosodi-n-butylamine	924-16-3	2.41		3.13E-01	e		3.13E-01	e		3.13E-01
N-Nitrosomorpholine	59-89-2	0.98		2.10E+00	e		2.10E+00	e		2.10E+00
N-Nitroso-N,N-dimethylamine	62-75-9	-0.47		1.45E+01	e		1.45E+01	e		1.45E+01
o-Anisidine	90-04-0	1.18		1.61E+00	e		1.61E+00	e		1.61E+00
Oxalic acid	144-62-7	No data		No data			1.00E+00		Default	1.00E+00
Phthalic anhydride	85-44-9	-0.60		1.73E+01	e		1.73E+01	e		1.73E+01

Table C2-2. Terrestrial Soil-to-Plant Transfer Factors (SPr) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPr	SAIC Compiled SPr	Notes	Comments	Recommended SPr[d]
p-Phthalic acid	100-21-0	0.82		2.59E+00	e		2.59E+00	e		2.59E+00
Pyridine	110-86-1	0.67		3.17E+00	e		3.17E+00	e		3.17E+00
Quinoline	91-22-5	2.03		5.20E-01	e		5.20E-01	e		5.20E-01
Quinone	106-51-4	0.20		5.94E+00	e		5.94E+00	e		5.94E+00
Safrole	94-59-7	2.66		2.25E-01	e		2.25E-01	e		2.25E-01
Tetrahydrofuran	109-99-9	0.45		4.27E+00	e		4.27E+00	e		4.27E+00
Other Heavy Semivolatile Compounds (molecular weight >200 g/mole)										
1,2,4,5-Tetrachlorobenzene	95-94-3	4.64		1.61E-02	e		1.61E-02	e		1.61E-02
1,3,5-Trinitrobenzene	99-35-4	1.18		1.61E+00	e		1.61E+00	e		1.61E+00
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	4.17		3.01E-02	e		3.01E-02	e		3.01E-02
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	-2.70		2.82E+02	e		2.82E+02	e		2.82E+02
2-sec-Butyl-4,6-dinitrophenol	88-85-7	3.14		1.19E-01	e		1.19E-01	e		1.19E-01
3,3'-Dimethoxybenzidine	119-90-4	1.81		6.96E-01	e		6.96E-01	e		6.96E-01
3,3-Dichlorobenzidine	91-94-1	3.58		6.65E-02	e		6.65E-02	e		6.65E-02
4-Bromophenylphenyl ether	101-55-3	5.00		9.98E-03	e		9.98E-03	e		9.98E-03
Ammonium perfluorooctanoate	3825-26-1	No data		No data			1.00E+00		Default	1.00E+00
Azobenzene	103-33-3	3.82		4.80E-02	e		4.80E-02	e		4.80E-02
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	No data		No data			1.00E+00		Default	1.00E+00
Captan	133-06-2	2.35		3.39E-01	e		3.39E-01	e		3.39E-01
Chlorobenzilate	510-15-6	4.38		2.28E-02	e		2.28E-02	e		2.28E-02
Dibutylphosphate	107-66-4	No data		No data			1.00E+00		Default	1.00E+00
Dimethyl aminoazobenzene	60-11-7	4.58		1.74E-02	e		1.74E-02	e		1.74E-02
Hexachlorobenzene	118-74-1	5.50	2.55E-02	5.10E-03	f		5.11E-03	e		5.10E-03
Hexachlorobutadiene	87-68-3	4.73	7.14E-02	1.43E-02	f		1.43E-02	e		1.43E-02
Hexachlorocyclopentadiene	77-47-4	5.04	5.65E-02	1.13E-02	f		9.46E-03	e		1.13E-02
Hexachloroethane	67-72-1	3.98		3.85E-02	e		3.85E-02	e		3.85E-02
Hexachlorophene	70-30-4	7.54		3.39E-04	e		3.39E-04	e		3.39E-04

Table C2-2. Terrestrial Soil-to-Plant Transfer Factors (SPr) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPr	SAIC Compiled SPr	Notes	Comments	Recommended SPr[d]
Hexamethylene-1,5-diisocyanate	822-06-0	1.27		1.42E+00	e		1.42E+00	e		1.42E+00
Mirex	2385-85-5	6.89		8.07E-04	e		8.07E-04	e		8.07E-04
Nitrofen	1836-75-5	5.53		4.93E-03	e		4.93E-03	e		4.93E-03
Pentachlorobenzene	608-93-5	5.09	4.40E-02	8.80E-03	f		8.89E-03	e		8.80E-03
Pentachloronitrobenzene	82-68-8	4.64	8.00E-02	1.61E-02	f		1.61E-02	e		1.61E-02
Pentachlorophenol	87-86-5	5.08	4.49E-02	8.98E-03	f		8.98E-03	e		8.98E-03
Picric acid	88-89-1	2.03		5.20E-01	e		5.20E-01	e		5.20E-01
Pronamide	23950-58-5	3.51		7.24E-02	e		7.24E-02	e		7.24E-02
Strychnine	57-24-9	1.93		5.94E-01	e		5.94E-01	e		5.94E-01
Terphenyls	26140-60-3	No data		No data			1.00E+00		Default	1.00E+00
Tributyl phosphate	126-73-8	4.00		3.78E-02	e		3.78E-02	e		3.78E-02
Trifluralin	1582-09-8	5.34		6.35E-03	e		6.35E-03	e		6.35E-03
Triphenylamine	603-34-9	No data		No data			1.00E+00		Default	1.00E+00
Herbicides and Organochlorinated Pesticides										
2,4,5-T	93-76-5	3.36		8.85E-02	e		8.85E-02	e		8.85E-02
2,4-D and esters	94-75-7	2.81		1.84E-01	e		1.84E-01	e		1.84E-01
4,4-DDD	72-54-8	6.20		2.02E-03	e		2.02E-03	e		2.02E-03
4,4-DDE	72-55-9	6.26	9.37E-03	1.87E-03	f		1.88E-03	e		1.87E-03
4,4-DDT	50-29-3	6.00		2.64E-03	e		2.64E-03	e		2.64E-03
Aldrin	309-00-2	6.18		2.08E-03	e		2.08E-03	e		2.08E-03
alpha-BHC	319-84-6	3.80		4.93E-02	e		4.93E-02	e		4.93E-02
beta-BHC	319-85-7	3.83		4.71E-02	e		4.71E-02	e		4.71E-02
Chlordane	57-74-9	5.94		2.87E-03	e		2.87E-03	e		2.87E-03
Delta-BHC	319-86-8	4.14		3.13E-02	e		3.13E-02	e		3.13E-02
Dieldrin	60-57-1	5.27		6.97E-03	e		6.97E-03	e		6.97E-03
Endothall	145-73-3	-0.87		2.47E+01	e		2.47E+01	e		2.47E+01
Endrin	72-20-8	4.89		1.15E-02	e		1.15E-02	e		1.15E-02

Table C2-2. Terrestrial Soil-to-Plant Transfer Factors (SPr) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPr	SAIC Compiled SPr	Notes	Comments	Recommended SPr[d]
gamma-BHC (Lindane)	58-89-9	3.72		5.48E-02	e		5.48E-02	e		5.48E-02
Heptachlor	76-44-8	5.02	4.89E-02	9.78E-03	f		9.75E-03	e		9.78E-03
Isodrin	465-73-6	3.55		6.87E-02	e		6.87E-02	e		6.87E-02
Methoxychlor	72-43-5	4.53		1.87E-02	e		1.87E-02	e		1.87E-02
Silvex (2,4,5-TP)	93-72-1	4.07		3.43E-02	e		3.43E-02	e		3.43E-02
Toxaphene	8001-35-2	5.50		5.13E-03	e		5.13E-03	e		5.13E-03
Inorganic Chemicals and Compounds										
Metals										
Aluminum	7429-90-5	NA	4.00E-03	8.00E-04	f		1.30E-04	i		1.30E-04
Antimony	7440-36-0	NA	2.00E-01	4.00E-02	f		6.00E-03	i		6.00E-03
Arsenic	7440-38-2	NA	3.60E-02	7.20E-03	f		1.20E-03	i		1.20E-03
Barium	7440-39-3	NA	1.50E-01	3.00E-02	f		3.00E-03	i		3.00E-03
Beryllium	7440-41-7	NA	1.00E-02	2.00E-03	f		3.00E-04	i		3.00E-04
Bismuth	7440-69-9	NA		7.00E-03	i		1.00E-03	i		1.00E-03
Boron	7440-42-8	NA		8.00E-01	i		4.00E-01	i		4.00E-01
Cadmium	7440-43-9	NA	3.64E-01	7.28E-02	f		3.00E-02	i		3.00E-02
Calcium	7440-70-2	NA		7.00E-01	i		7.00E-02	i		7.00E-02
Chromium (and VI)	18540-29-9	NA	7.50E-03	1.50E-03	f		9.00E-04	i		9.00E-04
Cobalt	7440-48-4	NA		4.00E-03	i		1.40E-03	i		1.40E-03
Copper	7440-50-8	NA	4.00E-01	8.00E-02	f		5.00E-02	i		5.00E-02
Iron	7439-89-6	NA		8.00E-04	i		2.00E-04	i		2.00E-04
Lead	7439-92-1	NA	4.50E-02	9.00E-03	f		1.80E-03	i		1.80E-03
Lithium	7439-93-2	NA		5.00E-03	i		8.00E-04	i		8.00E-04
Magnesium	7439-95-4	NA		2.00E-01	i		1.10E-01	i		1.10E-01
Manganese	7439-96-5	NA		5.00E-02	i		1.00E-02	i		1.00E-02
Mercury	7439-97-6	NA	3.75E-02	7.50E-03	f		4.00E-02	i		4.00E-02
Molybdenum	7439-98-7	NA		5.00E-02	i		1.20E-02	i		1.20E-02

Table C2-2. Terrestrial Soil-to-Plant Transfer Factors (SPr) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPr	SAIC Compiled SPr	Notes	Comments	Recommended SPr[d]
Nickel	7440-02-0	NA	3.20E-02	6.40E-03	f		1.20E-02	i		1.20E-02
Potassium	7440-09-7	NA		2.00E-01	i		1.10E-01	i		1.10E-01
Rhodium	7440-16-6	NA		3.00E-02	i		8.00E-03	i		8.00E-03
Selenium	7782-49-2	NA	1.60E-02	3.20E-03	f		5.00E-03	i		5.00E-03
Silicon	7440-21-3	NA		7.00E-02	i		1.40E-02	i		1.40E-02
Silver	7440-22-4	NA	4.00E-01	8.00E-02	f		2.00E-02	i		2.00E-02
Sodium	7440-23-5	NA		1.50E-02	i		1.10E-02	i		1.10E-02
Strontium	7440-24-6	NA		5.00E-01	i		5.00E-02	i		5.00E-02
Tantalum	7440-25-7	NA		2.00E-03	i		5.00E-04	i		5.00E-04
Thallium	7440-28-0	NA	4.00E-03	8.00E-04	f		8.00E-05	i		8.00E-05
Tin	7440-31-5	NA		6.00E-03	i		1.20E-03	i		1.20E-03
Tungsten	7440-33-7	NA		9.00E-03	i		2.00E-03	i		2.00E-03
Uranium	7440-61-1	NA		1.70E-03	i		8.00E-04	i		8.00E-04
Vanadium	7440-62-2	NA		1.10E-03	i		6.00E-04	i		6.00E-04
Yttrium	7440-65-5	NA		3.00E-03	i		1.20E-03	i		1.20E-03
Zinc	7440-66-6	NA	1.20E-12	2.40E-13	f		1.80E-01	i		1.80E-01
Zirconium	7440-67-7	NA		4.00E-04	i		1.00E-04	i		1.00E-04
Non-metals and Anions										
Ammonia/Ammonium	7664-41-7	NA		2.27E-02	j					2.27E-02
Bromide	24959-67-9	NA		3.00E-01	i		3.00E-01	i		3.00E-01
Chloride	16887-00-6	NA		1.40E+01	i		1.40E+01	i		1.40E+01
Cyanide	57-12-5	NA		2.27E-02	j		0.00E+00	k		2.27E-02
Fluoride	16984-48-8	NA		1.20E-02	i		1.20E-03	i		1.20E-02
Hydroxide	14280-30-9	NA		NA					Depends on pH	NA
Iodine	7553-56-2	NA		3.00E-02	i		1.00E-02	i		3.00E-02
Nitrate	14797-55-8	NA		2.27E-02	j					2.27E-02
Nitrite	14797-65-0	NA		2.27E-02	j					2.27E-02

Table C2-2. Terrestrial Soil-to-Plant Transfer Factors (SPr) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPr	SAIC Compiled SPr	Notes	Comments	Recommended SPr[d]
Phosphate	14265-44-2	NA		7.00E-01	i		7.00E-01	i		7.00E-01
Phosphorus	7723-14-0	NA		7.00E-01	i		7.00E-01	i		7.00E-01
Sulfate	14808-79-8	NA		3.00E-01	i		3.00E-01	i		3.00E-01
Total Sulfur	63705-05-5	NA		3.00E-01	i		3.00E-01	i		3.00E-01
Priority Pollutants										
Carbon Dioxide	124-38-9	NA		2.27E-02	j					2.27E-02
Nitrogen Dioxide	10102-44-0	NA		2.27E-02	j					2.27E-02
Ozone	10028-15-6	NA		2.27E-02	j					2.27E-02
Particulate Matter	No CAS #	NA		NA						NA
Sulfur Dioxode	7446-09-5	NA		2.27E-02	j					2.27E-02
Radionuclides										
Americium-241	1596-10-2	NA		1.10E-03	i		5.00E-05	i		5.00E-05
Antimony-125	14234-35-6	NA		4.00E-02	i		6.00E-03	i		6.00E-03
Barium-137	13981-97-0	NA		3.00E-02	i		3.00E-03	i		3.00E-03
Cadmium-113	None	NA		1.10E-01	i		3.00E-02	i		3.00E-02
Cesium-134	13967-70-9	NA		1.60E-02	i		6.00E-03	i		6.00E-03
Cesium-137	10045-97-3	NA		1.60E-02	i		6.00E-03	i		6.00E-03
Europium-154	15585-10-1	NA		2.00E-03	i		8.00E-04	i		8.00E-04
Europium-155	14391-16-3	NA		2.00E-03	i		8.00E-04	i		8.00E-04
Nickel-63	13981-37-8	NA		1.20E-02	i		1.20E-02	i		1.20E-02
Plutonium-239	15117-48-3	NA		9.00E-05	i		9.00E-06	i		9.00E-06
Plutonium-241	14119-32-5	NA		9.00E-05	i		9.00E-06	i		9.00E-06
Samarium-151	15715-94-3	NA		2.00E-03	i		8.00E-04	i		8.00E-04
Strontium-90	10098-97-2	NA		5.00E-01	i		5.00E-02	i		5.00E-02
Technetium-99	14133-79-7	NA		1.90E+00	i		3.00E-01	i		3.00E-01
Tritium	10028-17-8	NA		2.27E-02	j					2.27E-02
Yttrium-90	10098-91-6	NA		3.00E-03	i		1.20E-03	i		1.20E-03

Table C2-2. Terrestrial Soil-to-Plant Transfer Factors (SPr) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPr	SAIC Compiled SPr	Notes	Comments	Recommended SPr[d]
Uranium-232	14158-29-3	NA		1.70E-03	i		8.00E-04	i		8.00E-04
Uranium-233	13968-55-3	NA		1.70E-03	i		8.00E-04	i		8.00E-04
Uranium-234	13966-29-5	NA		1.70E-03	i		8.00E-04	i		8.00E-04
Uranium-235	15117-96-1	NA		1.70E-03	i		8.00E-04	i		8.00E-04
Uranium-236	13982-70-2	NA		1.70E-03	i		8.00E-04	i		8.00E-04
Uranium-238	7440-61-1	NA		1.70E-03	i		8.00E-04	i		8.00E-04
Actinium-227	14952-40-0	NA		7.00E-04	i		7.00E-05	i		7.00E-05
Americium-243	14993-75-0	NA		1.10E-03	i		5.00E-05	i		5.00E-05
Carbon-14	14762-75-5	NA		2.27E-02	j					2.27E-02
Cobalt-60	10198-40-0	NA		4.00E-03	i		1.40E-03	i		1.40E-03
Curium-242	15510-73-3	NA		1.70E-04	i		3.00E-06	i		3.00E-06
Curium-243	15757-87-6	NA		1.70E-04	i		3.00E-06	i		3.00E-06
Curium-244	13981-15-2	NA		1.70E-04	i		3.00E-06	i		3.00E-06
Europium-152	14683-23-9	NA		2.00E-03	i		8.00E-04	i		8.00E-04
Iodine-129	15046-84-1	NA		3.00E-02	i		1.00E-02	i		1.00E-02
Neptunium-237	13994-20-2	NA		2.00E-02	i		2.00E-03	i		2.00E-03
Nickel-59	14336-70-0	NA	3.20E-02	6.40E-03	f		1.20E-02	i		1.20E-02
Niobium-93	None	NA		4.00E-03	i		1.00E-03	i		1.00E-03
Plutonium-238	13981-16-3	NA		9.00E-05	i		9.00E-06	i		9.00E-06
Plutonium-240	14119-33-6	NA		9.00E-05	i		9.00E-06	i		9.00E-06
Plutonium-242	13982-10-0	NA		9.00E-05	i		9.00E-06	i		9.00E-06
Protactinium-231	14331-85-2	NA		5.00E-04	i		5.00E-05	i		5.00E-05
Radium-226	13982-63-3	NA		3.00E-03	i		3.00E-04	i		3.00E-04
Radium-228	15262-20-1	NA		3.00E-03	i		3.00E-04	i		3.00E-04
Ruthenium-106	13967-48-1	NA		1.50E-02	i		4.00E-03	i		4.00E-03
Selenium-79	None	NA		5.00E-03	i		5.00E-03	i		5.00E-03
Thorium-229	15594-54-4	NA		1.70E-04	i		1.70E-05	i		1.70E-05

Table C2-2. Terrestrial Soil-to-Plant Transfer Factors (SPr) for Ecological Receptors (mg/kg tissue wet weight/mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SPv[b] (per dry weight of plants)	Calculated from EPA (1999) SPv[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SPr	SAIC Compiled SPr	Notes	Comments	Recommended SPr[d]
Thorium-232	7440-29-1	NA		1.70E-04	i		1.70E-05	i		1.70E-05
Tin-126	15832-50-5	NA		6.00E-03	i		1.20E-03	i		1.20E-03
Zirconium-93	15751-77-6	NA		4.00E-04	i		1.00E-04	i		1.00E-04

NA = Not applicable

[a] $\log_{10}$ of K_{ow} values in Table 4.1

[b] Published in Appendix C of EPA (1999), Table C-2

[c] Calculated or chosen as described in Appendix C, Sect. C-1.2 of EPA (1999), then multiplied by 0.2 to adjust to wet weight of plants

[d] Selection criteria described in Sect. 8.2.4.3

[e] Wet weight values calculated by using log K_{ow}: log SPv = 1.588 - 0.578 x log K_{ow} (Travis and Arms 1988) for dry weight and multiplying by 0.2 to adjust to wet weight

[f] EPA (1999) values published in Table C-2 multiplied by 0.2 to adjust to wet weight of plants

[g] Calculated from value for TCDD in Appendix C of EPA (1999), Table C-2 by multiplying by BEF, then multiplying by 0.2 to adjust to wet weight of plants

[h] Value for Aroclor 1254 from Appendix C of EPA (1999), Table C-2, was used for PCB mixtures

[i] Baes and others. (1984) (Figure 2.2) values multiplied by 0.2 to adjust to wet weight of plants

[j] Average of values for other inorganics published in Appendix C of EPA (1999), Table C-2, then multiplied by 0.2 to adjust to wet weight of plants

[k] HAZWRAP (1994), Table 4.4

Table C2-3. Terrestrial Air-to-Plant Transfer Factors (APv) for Ecological Receptors (mg/kg tissue wet weight/mg/m^3 air)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	H[b] (atm x m^3/mol)	EPA (1999) APv[c] (kg air/kg dry wt)	APv Calculated by EPA (1999) Methods[d] (kg air/kg dry wt)	APv Adjusted to Wet Weight of Plants and Volume of Air (m^3 air/kg wet wt)[e]	Ecology Guidance APv	Default APv	Comments	Recommended APv[f]
Organic Compounds										
Aromatic Halogenated Hydrocarbons										
4-Chloro-3-methylphenol	59-50-7	3.10	2.50E-06		4.47E+01	7.51E+00				7.51E+00
2,3,4,6-Tetrachlorophenol	58-90-2	4.42	3.51E-06		8.11E+02	1.36E+02				1.36E+02
Aromatic Nonhalogenated Hydrocarbons										
2-Nitrotoluene	88-72-2	2.30	5.75E-05		2.73E-01	4.60E-02				4.60E-02
4-Nitrobiphenyl	92-93-3	3.77	3.54E-06		1.63E+02	2.75E+01				2.75E+01
Benzaldehyde	100-52-7	1.48	4.18E-05		5.00E-02	8.41E-03				8.41E-03
Benzene	71-43-2	-4.99	1.78E+03		1.52E-16	2.55E-17				2.55E-17
Benzyl alcohol	100-51-6	1.10	3.78E-07		2.20E+00	3.69E-01				3.69E-01
Ethyl benzene	100-41-4	3.12	7.73E-03		1.53E-02	2.58E-03				2.58E-03
m-Xylene	108-38-3	3.20	9.26E-06		1.55E+01	2.60E+00				2.60E+00
o-Xylene	95-47-6	3.13	6.73E-06		1.79E+01	3.01E+00				3.01E+00
p-Xylene	106-42-3	3.17	6.89E-06		1.93E+01	3.24E+00				3.24E+00
Styrene	100-42-5	2.93	3.33E-03		2.21E-02	3.71E-03				3.71E-03
Toluene	108-88-3	2.67	6.13E-03		6.32E-03	1.06E-03				1.06E-03
Non-aromatic Nonhalogenated Hydrocarbons										
1,2-Epoxybutane	106-88-7	1.44	4.61E-04		4.16E-03	6.98E-04				6.98E-04
1,3-Butadiene	106-99-0	1.90	7.46E+00		7.91E-07	1.33E-07				1.33E-07
1,4-Dioxane	123-91-1	-0.27	4.89E-06	5.93E-03		9.96E-04				9.96E-04
1-Methylpropyl alcohol	78-92-2	0.61	1.02E-05		2.44E-02	4.10E-03				4.10E-03
1-Nitropropane	108-03-2	0.87	No data		No data	No data		1.00E+00		1.00E+00
2,2,4-Trimethylpentane	540-84-1	5.02	3.02E+00		4.11E-03	6.90E-04				6.90E-04
2-Butanone	78-93-3	0.28	3.61E-05		3.08E-03	5.18E-04				5.18E-04
2-Butenaldehyde (2-Butenal)	4170-30-3	No data	No data		No data	No data		1.00E+00		1.00E+00
2-Ethoxyethanol	110-80-5	-0.10	5.13E-02		8.52E-07	1.43E-07				1.43E-07
2-Heptanone	110-43-0	1.98	1.45E-04		4.97E-02	8.34E-03				8.34E-03
2-Hexanone	591-78-6	1.38	9.60E-05		1.72E-02	2.89E-03				2.89E-03
2-Methoxyethanol	109-86-4	0.25	8.24E-07		1.26E-01	2.11E-02				2.11E-02

Table C2-3. Terrestrial Air-to-Plant Transfer Factors (APv) for Ecological Receptors (mg/kg tissue wet weight/mg/m^3 air)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	H[b] (atm x m^3/mol)	EPA (1999) APv[c] (kg air/kg dry wt)	APv Calculated by EPA (1999) Methods[d] (kg air/kg dry wt)	APv Adjusted to Wet Weight of Plants and Volume of Air (m^3 air/kg wet wt)[e]	Ecology Guidance APv	Default APv	Comments	Recommended APv[f]
2-Methyl-2-propanol	75-65-0	0.35	1.20E-05		1.10E-02	1.84E-03				1.84E-03
2-Methyl-2-propenenitrile	126-98-7	0.54	2.39E-04		8.80E-04	1.48E-04				1.48E-04
2-Methylaziridine	75-55-8	-0.60	9.74E-06		1.31E-03	2.20E-04				2.20E-04
2-Methylpropyl alcohol	78-83-1	0.76	1.20E-05		3.00E-02	5.04E-03				5.04E-03
2-Pentanone	107-87-9	0.91	6.46E-05		8.06E-03	1.36E-03				1.36E-03
2-Propanone (Acetone)	67-64-1	-0.22	2.88E-05	1.13E-03		1.89E-04				1.89E-04
2-Propene-1-ol	107-18-6	0.17	5.01E-06		1.69E-02	2.84E-03				2.84E-03
2-Propyl alcohol	67-63-0	0.05	8.12E-06		7.78E-03	1.31E-03				1.31E-03
3-Heptanone	106-35-4	No data	No data		No data	No data		1.00E+00		1.00E+00
3-Methyl-1-butanol	123-51-3	No data	No data		No data	No data		1.00E+00		1.00E+00
3-Methyl-2-butanone	563-80-4	No data	No data		No data	No data		1.00E+00		1.00E+00
3-Pentanone	96-22-0	0.99	8.05E-05		7.87E-03	1.32E-03				1.32E-03
4-Heptanone	123-19-3	No data	No data		No data	No data		1.00E+00		1.00E+00
4-Methyl-2-pentanone	108-10-1	1.19	1.25E-04		8.28E-03	1.39E-03				1.39E-03
4-Methyl-3-penten-2-one	141-79-7	No data	No data		No data	No data		1.00E+00		1.00E+00
5-Methyl-2-hexanone	110-12-3	No data	No data		No data	No data		1.00E+00		1.00E+00
Acetaldehyde	75-07-0	-0.22	7.89E-05		4.13E-04	6.93E-05				6.93E-05
Acetamide	60-35-5	-1.26	3.50E-09		7.28E-01	1.22E-01				1.22E-01
Acetic acid	64-19-7	-0.17	3.02E-07		1.22E-01	2.05E-02				2.05E-02
Acetic acid ethyl ester	141-78-6	0.73	1.35E-04		2.48E-03	4.17E-04				4.17E-04
Acetic acid n-butyl ester	123-86-4	1.73	3.31E-04		1.17E-02	1.97E-03				1.97E-03
Acetonitrile	75-05-8	-0.34	3.79E+01		6.40E-10	1.08E-10				1.08E-10
Acrolein	107-02-8	-0.01	9.34E-05		5.86E-04	9.84E-05				9.84E-05
Acrylonitrile	107-13-1	0.25	9.90E-05	1.04E-03		1.75E-04				1.75E-04
Bis(isopropyl)ether	108-20-3	No data	No data		No data	No data		1.00E+00		1.00E+00
Butane	106-97-8	2.89	9.59E+01		6.97E-07	1.17E-07				1.17E-07
Carbon disulfide	75-15-0	2.00	1.27E-02		5.94E-04	9.98E-05				9.98E-05
Cyanogen	460-19-5	0.81	4.96E-03		8.17E-05	1.37E-05				1.37E-05
Cyclohexane	110-82-7	3.44	1.95E-01		1.32E-03	2.22E-04				2.22E-04

Table C2-3. Terrestrial Air-to-Plant Transfer Factors (APv) for Ecological Receptors (mg/kg tissue wet weight/mg/m^3 air)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	H[b] (atm x m^3/mol)	EPA (1999) APv[c] (kg air/kg dry wt)	APv Calculated by EPA (1999) Methods[d] (kg air/kg dry wt)	APv Adjusted to Wet Weight of Plants and Volume of Air (m^3 air/kg wet wt)[e]	Ecology Guidance APv	Default APv	Comments	Recommended APv[f]
Cyclohexanone	108-94-1	0.81	1.20E-05		3.38E-02	5.69E-03				5.69E-03
Cyclohexene	110-83-8	2.86	3.98E-02		1.56E-03	2.62E-04				2.62E-04
Cyclopentane	287-92-3	3.00	1.86E-01		4.70E-04	7.90E-05				7.90E-05
Ethyl alcohol	64-17-5	0.31	6.29E-06		1.90E-02	3.19E-03				3.19E-03
Ethyl ether	60-29-7	0.89	1.29E-03		3.85E-04	6.47E-05				6.47E-05
Ethyl methacrylate	97-63-2	1.59	1.38E-04		2.00E-02	3.36E-03				3.36E-03
Formaldehyde	50-00-0	0.34	2.78E-04	4.65E-04		7.82E-05				7.82E-05
Formamide	75-12-7	-1.51	1.39E-09		9.91E-01	1.67E-01				1.67E-01
Formic acid	64-18-6	-0.54	2.49E-06		6.01E-03	1.01E-03				1.01E-03
Formic acid, methyl ester	107-31-3	-0.26	2.23E-04		1.31E-04	2.20E-05				2.20E-05
Glycidylaldehyde	765-34-4	-0.73	5.11E-07		1.83E-02	3.07E-03				3.07E-03
Methyl acetate	79-20-9	0.18	8.71E-05		9.98E-04	1.68E-04				1.68E-04
Methyl alcohol	67-56-1	-0.71	1.44E-04		6.80E-05	1.14E-05				1.14E-05
Methyl isocyanate	624-83-9	No data	No data		No data	No data			Reacts with water	1.00E+00
Methyl methacrylate	80-62-6	0.79	3.18E-04		1.22E-03	2.05E-04				2.05E-04
Methyl tert-butyl ether	1634-04-4	0.94	5.89E-04		9.52E-04	1.60E-04				1.60E-04
Methylacetylene	74-99-7	0.94	No data		No data	No data		1.00E+00		1.00E+00
Methylcyclohexane	108-87-2	4.10	4.36E-01		2.98E-03	5.00E-04				5.00E-04
N,N-Dimethylacetamide	127-19-5	No data	No data		No data	No data		1.00E+00		1.00E+00
n-Butyl alcohol	71-36-3	0.88	8.51E-06		5.68E-02	9.54E-03				9.54E-03
n-Heptane	142-82-5	4.66	2.28E+00		2.25E-03	3.79E-04				3.79E-04
n-Hexane	110-54-3	4.11	1.82E+00		7.32E-04	1.23E-04				1.23E-04
Nitromethane	75-52-5	-0.35	No data		No data	No data		1.00E+00		1.00E+00
n-Nonane	111-84-2	5.65	4.95E+00		1.17E-02	1.97E-03				1.97E-03
n-Octane	111-65-9	4.00	3.24E+00		3.14E-04	5.28E-05				5.28E-05
n-Pentane	109-66-0	3.21	1.26E+00		1.16E-04	1.96E-05				1.96E-05
n-Propionaldehyde	123-38-6	0.59	7.41E-05		3.20E-03	5.38E-04				5.38E-04
n-Propyl alcohol	71-23-8	0.25	6.85E-06		1.51E-02	2.53E-03				2.53E-03

Table C2-3. Terrestrial Air-to-Plant Transfer Factors (APv) for Ecological Receptors (mg/kg tissue wet weight/mg/m^3 air)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	H[b] (atm x m^3/mol)	EPA (1999) APv[c] (kg air/kg dry wt)	APv Calculated by EPA (1999) Methods[d] (kg air/kg dry wt)	APv Adjusted to Wet Weight of Plants and Volume of Air (m^3 air/kg wet wt)[e]	Ecology Guidance APv	Default APv	Comments	Recommended APv[f]
n-Valeraldehyde	110-62-3	No data	No data		No data	No data		1.00E+00		1.00E+00
Oxirane	75-21-8	-0.30	1.67E-04		1.60E-04	2.69E-05				2.69E-05
p-Cymene	99-87-6	4.10	7.92E-03		1.64E-01	2.76E-02				2.76E-02
Phosgene	75-44-5	No data	No data		No data	No data		1.00E+00		1.00E+00
Propargyl alcohol	107-19-7	0.26	8.56E-06		1.24E-02	2.08E-03				2.08E-03
Propionic acid	79-09-4	0.33	4.46E-07		2.82E-01	4.73E-02				4.73E-02
Propionitrile	107-12-0	0.16	3.71E-05		2.23E-03	3.74E-04				3.74E-04
Propylene gylcol monomethyl ether	107-98-2	-0.18	1.04E-06		3.46E-02	5.81E-03				5.81E-03
p-tert-Butyltoluene	98-51-1	No data	No data		No data	No data		1.00E+00		1.00E+00
Triethylamine	121-44-8	0.16	1.04E-04		7.93E-04	1.33E-04				1.33E-04
Trimethylamine	75-50-3	0.16	6.61E-05		1.25E-03	2.10E-04				2.10E-04
Vinyl acetate	108-05-4	0.70	5.50E-04		5.64E-04	9.48E-05				9.48E-05
Non-aromatic Halogenated Hydrocarbons										
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	No data	1.00E-02		No data	No data		1.00E+00		1.00E+00
1,1,1,2-Tetrachloroethane	630-20-6	2.63	2.44E-03		1.45E-02	2.44E-03				2.44E-03
1,1,1-Trichloroethane	71-55-6	2.42	1.86E-02		1.14E-03	1.92E-04				1.92E-04
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	3.73	9.77E-02		5.37E-03	9.02E-04				9.02E-04
1,1,2,2-Tetrachloroethane	79-34-5	4.64	3.72E-04		1.32E+01	2.23E+00				2.23E+00
1,1,2,2-Tetrachloroethene	127-18-4	2.55	1.73E-02		1.66E-03	2.79E-04				2.79E-04
1,1,2-Trichloroethane	79-00-5	2.10	1.00E-03		9.56E-03	1.61E-03				1.61E-03
1,1,2-Trichloroethylene	79-01-6	2.43	1.06E-02		2.06E-03	3.46E-04				3.46E-04
1,1-Dichloroethane	75-34-3	1.79	5.75E-03		7.88E-04	1.32E-04				1.32E-04
1,1-Dichloroethene	75-35-4	2.12	2.55E-02		3.97E-04	6.68E-05				6.68E-05
1,2,2-Trichloro-1,1,2-trifluoroethane	76-13-1	3.16	5.28E-01		2.46E-04	4.13E-05				4.13E-05
1,2,3-Trichloropropane	96-18-4	2.25	3.80E-04		3.67E-02	6.16E-03				6.16E-03
1,2-Dibromo-3-chloropropane	96-12-8	2.34	1.97E-04		8.82E-02	1.48E-02				1.48E-02
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	2.82	8.20E-04		6.87E-02	1.15E-02				1.15E-02
1,2-Dichloroethane	107-06-2	1.46	1.27E-03		1.59E-03	2.67E-04				2.67E-04
1,2-Dichloroethylene	540-59-0	0.48	1.50E-01		1.21E-06	2.03E-07				2.03E-07

Table C2-3. Terrestrial Air-to-Plant Transfer Factors (APv) for Ecological Receptors (mg/kg tissue wet weight/mg/m^3 air)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	H[b] (atm x m^3/mol)	EPA (1999) APv[c] (kg air/kg dry wt)	APv Calculated by EPA (1999) Methods[d] (kg air/kg dry wt)	APv Adjusted to Wet Weight of Plants and Volume of Air (m^3 air/kg wet wt)[e]	Ecology Guidance APv	Default APv	Comments	Recommended APv[f]
1,2-Dichloropropane	78-87-5	2.25	2.81E-03		4.96E-03	8.33E-04				8.33E-04
1,3-Dichloropropene	542-75-6	1.75	2.94E-03		1.38E-03	2.32E-04				2.32E-04
1,4-Dichloro-2-butene	764-41-0	0.87	1.06E-02		4.47E-05	7.52E-06				7.52E-06
1-Chloroethene	75-01-4	1.15	3.15E-01		2.95E-06	4.96E-07				4.96E-07
2,2-Dichloropropionic acid	75-99-0	0.78	6.43E-08		5.86E+00	9.84E-01				9.84E-01
2-Chloropropane	75-29-6	1.90	1.64E-02		3.61E-04	6.06E-05				6.06E-05
3-Chloropropene (allyl chloride)	107-05-1	0.95	1.09E-02		5.33E-05	8.96E-06				8.96E-06
Bromochloromethane	74-97-5	1.41	1.06E+03		1.67E-09	2.81E-10				2.81E-10
Bromodichloromethane	75-27-4	2.03	3.17E-03		2.53E-03	4.25E-04				4.25E-04
Bromoethene	593-60-2	1.07	2.76E-01		2.76E-06	4.65E-07				4.65E-07
Bromoform	75-25-2	2.35	6.16E-04		2.89E-02	4.85E-03				4.85E-03
Bromomethane	74-83-9	1.11	1.41E-02		6.09E-05	1.02E-05				1.02E-05
Carbon tetrachloride	56-23-5	2.72	2.87E-02	1.52E-03		2.56E-04				2.56E-04
Chlorodibromomethane	124-48-1	2.18	1.21E-03		9.59E-03	1.61E-03				1.61E-03
Chlorodifluoromethane	75-45-6	1.08	1.68E-01		4.69E-06	7.88E-07				7.88E-07
Chloroethane	75-00-3	3.10	1.80E+00		6.22E-05	1.05E-05				1.05E-05
Chloroform	67-66-3	1.95	4.03E-03	1.65E-03		2.78E-04				2.78E-04
Chloromethane	74-87-3	0.90	4.52E-02		1.13E-05	1.90E-06				1.90E-06
Chloropentafluoroethane	76-15-3	No data	No data		No data	No data		1.00E+00		1.00E+00
cis-1,2-Dichloroethene	156-59-2	1.98	4.51E-03		1.60E-03	2.69E-04				2.69E-04
cis-1,3-Dichloropropene	10061-01-5	No data	2.40E-01		No data	No data		1.00E+00		1.00E+00
Cyanogen bromide	506-68-3	No data	No data		No data	No data		1.00E+00		1.00E+00
Cyanogen chloride	506-77-4	0.20	No data		No data	No data		1.00E+00		1.00E+00
Dichlorodifluoromethane	75-71-8	2.16	2.58E+00		4.31E-06	7.24E-07				7.24E-07
Dichlorofluoromethane	75-43-4	No data	No data		No data	No data		1.00E+00		1.00E+00
Dichloromethane	75-09-2	1.26	2.38E-03		5.10E-04	8.57E-05				8.57E-05
Difluorodibromomethane	75-61-6	No data	No data		No data	No data		1.00E+00		1.00E+00
Hexafluoroacetone	684-16-2	No data	No data		No data	No data		1.00E+00		1.00E+00
Iodomethane	74-88-4	1.69	5.49E-03		6.42E-04	1.08E-04				1.08E-04

Table C2-3. Terrestrial Air-to-Plant Transfer Factors (APv) for Ecological Receptors (mg/kg tissue wet weight/mg/m^3 air)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	H[b] (atm x m^3/mol)	EPA (1999) APv[c] (kg air/kg dry wt)	APv Calculated by EPA (1999) Methods[d] (kg air/kg dry wt)	APv Adjusted to Wet Weight of Plants and Volume of Air (m^3 air/kg wet wt)[e]	Ecology Guidance APv	Default APv	Comments	Recommended APv[f]
Methylene bromide	74-95-3	1.62	2.64E-02		1.12E-04	1.89E-05				1.89E-05
Pentachloroethane	76-01-7	3.05	2.45E-03		4.03E-02	6.78E-03				6.78E-03
trans-1,2-Dichloroethene	156-60-5	1.98	7.44E-03		9.70E-04	1.63E-04				1.63E-04
trans-1,3-Dichloropropene	10061-02-6	2.06	No data		No data	No data		1.00E+00		1.00E+00
Trichloroacetic acid	76-03-9	1.33	1.80E-07		8.11E+00	1.36E+00				1.36E+00
Trichlorofluoroethane	27154-33-2	No data	No data		No data	No data		1.00E+00		1.00E+00
Trichlorofluoromethane	75-69-4	2.53	1.37E-01		2.03E-04	3.40E-05				3.40E-05
Trifluorobromomethane	75-63-8	1.86	No data		No data	No data		1.00E+00		1.00E+00
Dioxin and Furan Compounds										
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	35822-46-9	8.20	7.50E-06	9.10E+05		1.53E+05				1.53E+05
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	7.92	5.30E-05	8.30E+05		1.39E+05				1.39E+05
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	7.92	5.30E-05	8.30E+05		1.39E+05				1.39E+05
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	39227-28-6	7.79	1.20E-05	5.20E+05		8.74E+04				8.74E+04
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	7.25	1.40E-05	1.62E+05		2.72E+04				2.72E+04
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	57653-85-7	7.25	1.20E-05	5.20E+05		8.74E+04				8.74E+04
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	7.25	6.10E-06	1.62E+05		2.72E+04				2.72E+04
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	19408-74-3	7.25	1.20E-05	5.20E+05		8.74E+04				8.74E+04
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	7.25	1.00E-05	1.62E+05		2.72E+04				2.72E+04
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	40321-76-4	6.64	2.60E-06	2.39E+05		4.02E+04				4.02E+04
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	6.79	6.20E-06	9.75E+04		1.64E+04				1.64E+04
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	7.25	1.00E-05	1.62E+05		2.72E+04				2.72E+04
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	6.92	6.20E-06	9.75E+04		1.64E+04				1.64E+04
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1746-01-6	6.64	1.60E-05	6.55E+04		1.10E+04				1.10E+04
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	6.53	8.60E-06	4.57E+04		7.68E+03				7.68E+03
Dibenzofuran	132-64-9	4.33	1.30E-05		1.76E+02	2.95E+01				2.95E+01
Octachlorodibenzo(p)dioxin	3268-87-9	8.20	6.76E-06	2.36E+06		3.97E+05				3.97E+05
Octachlorodibenzofuran	39001-02-0	8.78	1.90E-06	2.28E+06		3.83E+05				3.83E+05
PCBs										
Polychlorinated biphenyls (PCBs)[f]	1336-36-3	7.31		3.09E+02		5.19E+01				5.19E+01

Table C2-3. Terrestrial Air-to-Plant Transfer Factors (APv) for Ecological Receptors (mg/kg tissue wet weight/mg/m^3 air)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	H[b] (atm x m^3/mol)	EPA (1999) APv[c] (kg air/kg dry wt)	APv Calculated by EPA (1999) Methods[d] (kg air/kg dry wt)	APv Adjusted to Wet Weight of Plants and Volume of Air (m^3 air/kg wet wt)[e]	Ecology Guidance APv	Default APv	Comments	Recommended APv[f]
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	7.08	1.91E-04		1.02E+04	1.71E+03				1.71E+03
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	7.12	3.01E-04		7.10E+03	1.19E+03				1.19E+03
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	No data	No data		No data	No data		1.00E+00		1.00E+00
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	No data	No data		No data	No data		1.00E+00		1.00E+00
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	No data	No data		No data	No data		1.00E+00		1.00E+00
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	No data	No data		No data	No data		1.00E+00		1.00E+00
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	No data	No data		No data	No data		1.00E+00		1.00E+00
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	No data	No data		No data	No data		1.00E+00		1.00E+00
2',3,4,4',5-Pentachlorobiphenyl	no cas #	No data	No data		No data	No data		1.00E+00		1.00E+00
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	7.12	No data		No data	No data		1.00E+00		1.00E+00
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	7.41	5.92E-05		7.32E+04	1.23E+04				1.23E+04
3,3',4,4',5-Pentachlorobiphenyl	no cas #	No data	No data		No data	No data		1.00E+00		1.00E+00
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	No data	No data		No data	No data		1.00E+00		1.00E+00
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	No data	No data		No data	No data		1.00E+00		1.00E+00
Phthalates										
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	5.20	1.10E-08	2.33E+03		3.92E+02				3.92E+02
Butylbenzyl phthalate	85-68-7	4.41	1.91E-06		1.47E+03	2.47E+02				2.47E+02
Dibutyl phthalate	84-74-2	4.72	1.43E-06		4.16E+03	6.99E+02				6.99E+02
Diethyl phthalate	84-66-2	4.44	5.48E-07		5.41E+03	9.09E+02				9.09E+02
Dimethylphthalate	131-11-3	1.63	1.01E-07		3.04E+01	5.11E+00				5.11E+00
n-Dioctyl phthalate	117-84-0	9.33	7.68E-07	6.30E+08		1.06E+08				1.06E+08
Light Polycyclic Aromatic Hydrocarbons (molecular weight <200 g/mole)										
2-Chloronapthalene	91-58-7	4.07	1.43E-04		8.41E+00	1.41E+00				1.41E+00
2-Methyl naphthalene	91-57-6	3.86	4.01E-04		1.80E+00	3.02E-01				3.02E-01
5-Nitroacenaphthene	602-87-9	No data	No data		No data	No data		1.00E+00		1.00E+00
Acenaphthene	83-32-9	3.96	2.00E-04		4.66E+00	7.84E-01				7.84E-01
Acenaphthylene	208-96-8	4.07	1.18E-04		1.02E+01	1.72E+00				1.72E+00
Anthracene	120-12-7	4.47	1.11E-04		2.90E+01	4.87E+00				4.87E+00
Fluorene	86-73-7	4.18	8.42E-05		1.88E+01	3.16E+00				3.16E+00

Table C2-3. Terrestrial Air-to-Plant Transfer Factors (APv) for Ecological Receptors (mg/kg tissue wet weight/mg/m^3 air)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	H[b] (atm x m^3/mol)	EPA (1999) APv[c] (kg air/kg dry wt)	APv Calculated by EPA (1999) Methods[d] (kg air/kg dry wt)	APv Adjusted to Wet Weight of Plants and Volume of Air (m^3 air/kg wet wt)[e]	Ecology Guidance APv	Default APv	Comments	Recommended APv[f]
Indene	95-13-6	No data	No data		No data	No data		1.00E+00		1.00E+00
Naphthalene	91-20-3	3.37	4.82E-04		4.53E-01	7.62E-02				7.62E-02
Phenanthrene	85-01-8	4.55	1.88E-01		2.08E-02	3.50E-03				3.50E-03
Pyrene	129-00-0	5.00	1.14E-08		1.04E+06	1.74E+05				1.74E+05
Heavy Polycyclic Aromatic Hydrocarbons (molecular weight >200 g/mole)										
3-Methylcholanthrene	56-49-5	7.11	1.24E-07		1.68E+07	2.82E+06				2.82E+06
5-Methylchrysene	3697-24-3	No data	No data		No data	No data		1.00E+00		1.00E+00
Benzo(a)anthracene	56-55-3	5.68	3.62E-06	1.72E+04		2.90E+03				2.90E+03
Benzo(a)pyrene	50-32-8	6.13	8.36E-07	2.25E+05		3.78E+04				3.78E+04
Benzo(b)fluoranthene	205-99-2	6.20	6.18E-06	3.65E+05		6.13E+04				6.13E+04
Benzo(e)pyrene	192-97-2	7.40	4.62E-07		9.19E+06	1.55E+06				1.55E+06
Benzo(g,h,i)perylene	191-24-2	7.10	7.02E-06		2.90E+05	4.87E+04				4.87E+04
Benzo(j)fluoranthene	205-82-3	6.44	No data		No data	No data		1.00E+00		1.00E+00
Benzo(k)fluoranthene	207-08-9	6.20	4.15E-07	5.40E+05		9.08E+04		1.00E+00		9.08E+04
Benzo[a,i]pyrene	191-30-0	No data	No data		No data	No data				1.00E+00
Chrysene	218-01-9	5.74	1.21E-06	5.97E+04		1.00E+04				1.00E+04
Dibenz(a,h)anthracene	53-70-3	6.55	1.12E-08	4.68E+07		7.87E+06				7.87E+06
Dibenz[a,h]acridine	226-36-8	No data	No data		No data	No data		1.00E+00		1.00E+00
Dibenz[a,j]acridine	224-42-0	No data	No data		No data	No data		1.00E+00		1.00E+00
Dibenzo(a,e)fluoranthene	5385-75-1	No data	No data		No data	No data		1.00E+00		1.00E+00
Dibenzo(a,h)fluoranthene	no cas #	No data	No data		No data	No data		1.00E+00		1.00E+00
Dibenzo[a,e]pyrene	192-65-4	No data	No data		No data	No data		1.00E+00		1.00E+00
Dibenzo[a,h]pyrene	189-64-0	No data	No data		No data	No data		1.00E+00		1.00E+00
Dibenzo[a,i]pyrene	189-55-9	7.29	1.41E-08		2.30E+08	3.87E+07				3.87E+07
Fluoranthene	206-44-0	5.08	9.33E-06		1.55E+03	2.61E+02				2.61E+02
Hexachloronaphthalene	1335-87-1	7.59	1.41E-04		4.80E+04	8.07E+03				8.07E+03
Indeno(1,2,3-cd)pyrene	193-39-5	6.91	4.86E-09	2.67E+08		4.49E+07				4.49E+07
Octachloronaphthalene	2234-13-1	6.42	1.02E-04		3.77E+03	6.33E+02				6.33E+02
Pentachloronaphthalene	1321-64-8	No data	No data		No data	No data		1.00E+00		1.00E+00

Table C2-3. Terrestrial Air-to-Plant Transfer Factors (APv) for Ecological Receptors (mg/kg tissue wet weight/mg/m^3 air)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	H[b] (atm x m^3/mol)	EPA (1999) APv[c] (kg air/kg dry wt)	APv Calculated by EPA (1999) Methods[d] (kg air/kg dry wt)	APv Adjusted to Wet Weight of Plants and Volume of Air (m^3 air/kg wet wt)[e]	Ecology Guidance APv	Default APv	Comments	Recommended APv[f]
Tetrachloronaphthalene	1335-88-2	No data	No data		No data	No data		1.00E+00		1.00E+00
Trichloronaphthalene	1321-65-9	No data	No data		No data	No data		1.00E+00		1.00E+00
Light Substituted Benzene Compounds (MW <200 g/mole)										
1,2,3-Trichlorobenzene	87-61-6	4.05	2.84E-03		4.00E-01	6.73E-02				6.73E-02
1,2,4-Trichlorobenzene	120-82-1	3.99	2.61E-03		3.78E-01	6.36E-02				6.36E-02
1,2,4-Trimethyl benzene	95-63-6	3.65	5.75E-03		7.49E-02	1.26E-02				1.26E-02
1,2-Dichlorobenzene	95-50-1	3.45	2.11E-03		1.24E-01	2.08E-02				2.08E-02
1,3,5-Trimethyl benzene	108-67-8	3.42	7.81E-03		3.14E-02	5.28E-03				5.28E-03
1,3-Dichlorobenzene	541-73-1	3.53	1.11E+02		2.89E-06	4.86E-07				4.86E-07
1,3-Dinitrobenzene	99-65-0	1.49	1.25E-07	1.43E-01		2.40E-02				2.40E-02
1,4-Dichlorobenzene	106-46-7	3.41	2.80E-03		8.58E-02	1.44E-02				1.44E-02
1,4-Dinitrobenzene	100-25-4	1.50	No data		No data	No data		1.00E+00		1.00E+00
2,4,5-Trichlorophenol	95-95-4	3.87	5.64E-06		1.31E+02	2.20E+01				2.20E+01
2,4,6-Trichlorophenol	88-06-2	3.71	4.06E-06		1.24E+02	2.08E+01				2.08E+01
2,4-Dichlorophenol	120-83-2	3.04	2.38E-07		4.03E+02	6.78E+01				6.78E+01
2,4-Dimethylphenol	105-67-9	2.36	4.27E-09		4.27E+03	7.17E+02				7.17E+02
2,4-Dinitrophenol	51-28-5	1.52	4.82E-09		4.80E+02	8.07E+01				8.07E+01
2,4-Dinitrotoluene	121-14-2	2.00	1.46E-07	5.10E+01		8.57E+00				8.57E+00
2,6-Dinitrotoluene	606-20-2	1.89	1.30E-07	4.41E+01		7.41E+00				7.41E+00
2-Chlorophenol	95-57-8	2.16	1.66E-05		6.75E-01	1.13E-01				1.13E-01
2-Chlorotoluene	95-49-8	3.54	6.41E-02		5.09E-03	8.55E-04				8.55E-04
2-Nitrophenol	88-75-5	1.79	1.46E-05		3.09E-01	5.19E-02				5.19E-02
4,6-Dinitro-o-cresol	534-52-1	2.85	1.40E-06		4.32E+01	7.25E+00				7.25E+00
4-Chlorotoluene	106-43-4	3.33	No data		No data	No data		1.00E+00		1.00E+00
4-Nitrophenol	100-02-7	1.91	7.32E-09		8.26E+02	1.39E+02				1.39E+02
alpha-Methylstyrene	98-83-9	3.46	5.91E-03		4.62E-02	7.76E-03				7.76E-03
Aniline	62-53-3	0.98	2.28E-06		2.71E-01	4.56E-02				4.56E-02
Benzotrichloride	98-07-7	2.92	9.81E-04		7.33E-02	1.23E-02				1.23E-02
Benzyl chloride	100-44-7	0.36	4.13E-04		3.29E-04	5.52E-05				5.52E-05

Table C2-3. Terrestrial Air-to-Plant Transfer Factors (APv) for Ecological Receptors (mg/kg tissue wet weight/mg/m^3 air)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	H[b] (atm x m^3/mol)	EPA (1999) APv[c] (kg air/kg dry wt)	APv Calculated by EPA (1999) Methods[d] (kg air/kg dry wt)	APv Adjusted to Wet Weight of Plants and Volume of Air (m^3 air/kg wet wt)[e]	Ecology Guidance APv	Default APv	Comments	Recommended APv[f]
Bromobenzene	108-86-1	2.99	2.09E-03		4.09E-02	6.87E-03				6.87E-03
Chlorobenzene	108-90-7	2.79	4.38E-03		1.19E-02	2.01E-03				2.01E-03
Cumene	98-82-8	3.61	1.29E-02		3.05E-02	5.13E-03				5.13E-03
m-Cresol	108-39-4	1.96	8.93E-07		7.63E+00	1.28E+00				1.28E+00
n-Butyl benzene	104-51-8	4.28	1.29E-02		1.57E-01	2.64E-02				2.64E-02
Nitrobenzene	98-95-3	1.83	2.06E-05	2.43E-01		4.08E-02				4.08E-02
n-Propyl benzene	103-65-1	3.69	1.10E-02		4.33E-02	7.28E-03				7.28E-03
o-Cresol	95-48-7	2.02	1.62E-06		4.90E+00	8.24E-01				8.24E-01
o-Dinitrobenzene	528-29-0	1.60	No data		No data	No data		1.00E+00		1.00E+00
o-Nitroaniline	88-74-4	1.85	1.17E-06		4.46E+00	7.50E-01				7.50E-01
o-Toluidine	95-53-4	1.34	2.43E-06		6.15E-01	1.03E-01				1.03E-01
p-Chloroaniline	106-47-8	1.87	1.17E-06		4.68E+00	7.86E-01				7.86E-01
p-Cresol	106-44-5	1.94	7.99E-07		8.13E+00	1.37E+00				1.37E+00
Phenol	108-95-2	1.48	5.95E-07		3.51E+00	5.91E-01				5.91E-01
p-Nitrochlorobenzene	100-00-5	2.39	No data		No data	No data		1.00E+00		1.00E+00
p-Toluidine	106-49-0	1.40	6.50E-06		2.66E-01	4.48E-02				4.48E-02
sec-Butyl benzene	135-98-8	4.57	1.39E-02		2.97E-01	4.99E-02				4.99E-02
tert-Butyl benzene	98-06-6	4.11	1.19E-02		1.12E-01	1.88E-02				1.88E-02
Toluene-2,6-diamine	823-40-5	1.45	1.24E-11		1.60E+05	2.68E+04				2.68E+04
Trimethyl benzene	25551-13-7	3.40	5.75E-03		4.06E-02	6.82E-03				6.82E-03
Other Light Semivolatile Compounds (molecular weight <200 g/mole)										
1,1'-Biphenyl	92-52-4	3.90	2.77E-04		2.87E+00	4.82E-01				4.82E-01
1,1-Dimethylhydrazine	57-14-7	0.60	1.64E-06		1.47E-01	2.47E-02				2.47E-02
1,2-Dimethylhydrazine	540-73-8	-1.37	6.95E-08		2.81E-02	4.72E-03				4.72E-03
1,2-Diphenylhydrazine	122-66-7	2.94	1.28E-07		5.90E+02	9.92E+01				9.92E+01
1,3-Propane sultone	1120-71-4	-0.52	2.36E-06		6.57E-03	1.10E-03				1.10E-03
2,4-Toluene diisocyante	584-84-9	No data	No data		No data	No data			Decomposes rapidly in water	1.00E+00
2-Chloroacetophenone	532-27-4	2.59	8.71E-07		3.72E+01	6.25E+00				6.25E+00

Table C2-3. Terrestrial Air-to-Plant Transfer Factors (APv) for Ecological Receptors (mg/kg tissue wet weight/mg/m^3 air)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	H[b] (atm x m^3/mol)	EPA (1999) APv[c] (kg air/kg dry wt)	APv Calculated by EPA (1999) Methods[d] (kg air/kg dry wt)	APv Adjusted to Wet Weight of Plants and Volume of Air (m^3 air/kg wet wt)[e]	Ecology Guidance APv	Default APv	Comments	Recommended APv[f]
2-Propenoic acid	79-10-7	0.33	4.46E-07		2.82E-01	4.73E-02				4.73E-02
4,4-Methylenedianiline	101-77-9	3.38	2.81E-08		7.90E+03	1.33E+03				1.33E+03
Acetophenone	98-86-2	1.64	1.03E-05		3.03E-01	5.09E-02				5.09E-02
Benzoic acid	65-85-0	1.86	1.54E-06		3.47E+00	5.84E-01				5.84E-01
bis(2-Chloroethoxy)methane	111-91-1	7.59	7.00E-09		9.67E+08	1.63E+08				1.63E+08
bis(2-Chloroethyl) ether	111-44-4	1.30	2.13E-05		6.37E-02	1.07E-02				1.07E-02
Chlorocyclopentadiene	41851-50-7	2.43	2.24E-02		9.66E-04	1.62E-04				1.62E-04
Cyclohexanol	108-93-0	1.23	5.75E-06		1.98E-01	3.33E-02				3.33E-02
Dichloroisopropyl ether	108-60-1	2.58	1.13E-04		2.77E-01	4.65E-02				4.65E-02
Dichloromethyl ether	542-88-1	-0.38	3.70E-03		5.95E-06	9.99E-07				9.99E-07
Dichloropentadiene	no cas #	No data	No data		No data	No data		1.00E+00		1.00E+00
Dimethyl sulfate	77-78-1	0.32	4.01E-06		3.04E-02	5.11E-03				5.11E-03
Dimethylaniline	121-69-7	-0.88	1.39E-05		4.66E-04	7.83E-05				7.83E-05
Di-n-propylnitrosamine	621-64-7	1.38	5.43E-08		3.04E+01	5.10E+00				5.10E+00
Diphenyl ether	101-84-8	4.21	2.49E-04		6.85E+00	1.15E+00				1.15E+00
Epichlorohydrin	106-89-8	0.25	3.08E-05		3.35E-03	5.63E-04				5.63E-04
Ethyl carbamate (urethane)	51-79-6	-0.15	No data		No data	No data		1.00E+00		1.00E+00
Ethyl methanesulfonate	62-50-0	0.05	8.87E-08		7.11E-01	1.19E-01				1.19E-01
Ethylene dibromide	106-93-4	1.75	4.47E-04		9.13E-03	1.53E-03				1.53E-03
Ethylene glycol	107-21-1	-0.91	1.80E-09		3.30E+00	5.54E-01				5.54E-01
Ethylene glycol monobutyl ether	111-76-2	1.55	5.26E-07		4.81E+00	8.08E-01				8.08E-01
Ethylene glycol monoethyl ether acetate	111-15-9	0.62	1.77E-06		1.44E-01	2.42E-02				2.42E-02
Ethylene thiourea	96-45-7	-0.64	1.52E-10		7.62E+01	1.28E+01				1.28E+01
Furfural	98-01-1	0.96	3.71E-06		1.58E-01	2.66E-02				2.66E-02
Maleic hydrazide	123-33-1	-0.74	6.59E-03		1.38E-06	2.32E-07				2.32E-07
Malononitrile	109-77-3	0.04	No data		No data	No data		1.00E+00		1.00E+00
Methyl styrene (mixed isomers)	25013-15-4	3.35	No data		No data	No data		1.00E+00		1.00E+00
Methylhydrazine	60-34-4	-1.06	No data		No data	No data		1.00E+00		1.00E+00
N,N-Diphenylamine	122-39-4	3.50	3.47E-07		8.61E+02	1.45E+02				1.45E+02

Table C2-3. Terrestrial Air-to-Plant Transfer Factors (APv) for Ecological Receptors (mg/kg tissue wet weight/mg/m^3 air)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	H[b] (atm x m^3/mol)	EPA (1999) APv[c] (kg air/kg dry wt)	APv Calculated by EPA (1999) Methods[d] (kg air/kg dry wt)	APv Adjusted to Wet Weight of Plants and Volume of Air (m^3 air/kg wet wt)[e]	Ecology Guidance APv	Default APv	Comments	Recommended APv[f]
Nitric acid, propyl ester	627-13-4	No data	No data		No data	No data		1.00E+00		1.00E+00
N-Nitrosodi-n-butylamine	924-16-3	2.41	5.47E-05		3.77E-01	6.33E-02				6.33E-02
N-Nitrosomorpholine	59-89-2	0.98	4.18E-08		1.48E+01	2.49E+00				2.49E+00
N-Nitroso-N,N-dimethylamine	62-75-9	-0.47	7.80E-07		2.26E-02	3.80E-03				3.80E-03
o-Anisidine	90-04-0	1.18	1.38E-06		7.31E-01	1.23E-01				1.23E-01
Oxalic acid	144-62-7	No data	No data		No data	No data		1.00E+00		1.00E+00
Phthalic anhydride	85-44-9	-0.60	6.28E-09		2.03E+00	3.42E-01				3.42E-01
p-Phthalic acid	100-21-0	0.82	1.04E-08		4.04E+01	6.79E+00				6.79E+00
Pyridine	110-86-1	0.67	6.86E-03		4.21E-05	7.08E-06				7.08E-06
Quinoline	91-22-5	2.03	2.70E-07		3.00E+01	5.05E+00				5.05E+00
Quinone	106-51-4	0.20	4.79E-04		1.91E-04	3.20E-05				3.20E-05
Safrole	94-59-7	2.66	1.19E-05		3.20E+00	5.37E-01				5.37E-01
Tetrahydrofuran	109-99-9	0.45	No data		No data	No data		1.00E+00		1.00E+00
Other Heavy Semivolatile Compounds (molecular weight >200 g/mole)										
1,2,4,5-Tetrachlorobenzene	95-94-3	4.64	1.18E-03		4.13E+00	6.95E-01				6.95E-01
1,3,5-Trinitrobenzene	99-35-4	1.18	8.66E-08		1.16E+01	1.95E+00				1.95E+00
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	4.17	No data		No data	No data		1.00E+00		1.00E+00
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	-2.70	No data		No data	No data		1.00E+00		1.00E+00
2-sec-Butyl-4,6-dinitrophenol	88-85-7	3.14	4.56E-07		2.71E+02	4.55E+01				4.55E+01
3,3'-Dimethoxybenzidine	119-90-4	1.81	3.36E-10		1.41E+04	2.37E+03				2.37E+03
3,3-Dichlorobenzidine	91-94-1	3.58	2.08E-08		1.72E+04	2.90E+03				2.90E+03
4-Bromophenylphenyl ether	101-55-3	5.00	1.17E-04		1.01E+02	1.70E+01				1.70E+01
Ammonium perfluorooctanoate	3825-26-1	No data	No data		No data	No data		1.00E+00		1.00E+00
Azobenzene	103-33-3	3.82	4.98E-06		1.31E+02	2.21E+01				2.21E+01
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfid	96-69-5	No data	No data		No data	No data		1.00E+00		1.00E+00
Captan	133-06-2	2.35	5.94E-06		2.99E+00	5.03E-01				5.03E-01
Chlorobenzilate	510-15-6	4.38	7.24E-08		3.57E+04	6.00E+03				6.00E+03
Dibutylphosphate	107-66-4	No data	No data		No data	No data		1.00E+00		1.00E+00
Dimethyl aminoazobenzene	60-11-7	4.58	4.00E-10		1.05E+07	1.77E+06				1.77E+06

Table C2-3. Terrestrial Air-to-Plant Transfer Factors (APv) for Ecological Receptors (mg/kg tissue wet weight/mg/m^3 air)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	H[b] (atm x m^3/mol)	EPA (1999) APv[c] (kg air/kg dry wt)	APv Calculated by EPA (1999) Methods[d] (kg air/kg dry wt)	APv Adjusted to Wet Weight of Plants and Volume of Air (m^3 air/kg wet wt)[e]	Ecology Guidance APv	Default APv	Comments	Recommended APv[f]
Hexachlorobenzene	118-74-1	5.50	5.35E-04	7.57E+01		1.27E+01				1.27E+01
Hexachlorobutadiene	87-68-3	4.73	2.39E-02	2.55E-01		4.29E-02				4.29E-02
Hexachlorocyclopentadiene	77-47-4	5.04	1.72E-02	5.47E-01		9.19E-02				9.19E-02
Hexachloroethane	67-72-1	3.98	3.60E-03		2.72E-01	4.58E-02				4.58E-02
Hexachlorophene	70-30-4	7.54	4.88E-10		1.23E+10	2.06E+09				2.06E+09
Hexamethylene-1,5-diisocyanate	822-06-0	1.27	2.66E-07		4.77E+00	8.02E-01				8.02E-01
Mirex	2385-85-5	6.89	5.27E-04		2.31E+03	3.88E+02				3.88E+02
Nitrofen	1836-75-5	5.53	1.69E-06		2.56E+04	4.31E+03				4.31E+03
Pentachlorobenzene	608-93-5	5.09	2.43E-02	6.04E-01		1.02E-01				1.02E-01
Pentachloronitrobenzene	82-68-8	4.64	2.86E-02	1.71E-01		2.87E-02				2.87E-02
Pentachlorophenol	87-86-5	5.08	1.41E-05	1.02E+03		1.71E+02				1.71E+02
Picric acid	88-89-1	2.03	3.79E-13		2.14E+07	3.60E+06				3.60E+06
Pronamide	23950-58-5	3.51	9.05E-06		3.38E+01	5.69E+00				5.69E+00
Strychnine	57-24-9	1.93	4.90E-13		1.30E+07	2.18E+06				2.18E+06
Terphenyls	26140-60-3	No data	No data		No data	No data		1.00E+00		1.00E+00
Tributyl phosphate	126-73-8	4.00	1.50E-07		6.78E+03	1.14E+03				1.14E+03
Trifluralin	1582-09-8	5.34	3.99E-05		6.82E+02	1.15E+02				1.15E+02
Triphenylamine	603-34-9	No data	No data		No data	No data		1.00E+00		1.00E+00
Herbicides and Organochlorinated Pesticides										
2,4,5-T	93-76-5	3.36	8.70E-09		2.43E+04	4.09E+03				4.09E+03
2,4-D and esters	94-75-7	2.81	1.37E-10		4.02E+05	6.76E+04				6.76E+04
4,4-DDD	72-54-8	6.20	8.43E-05		2.66E+03	4.47E+02				4.47E+02
4,4-DDE	72-55-9	6.26	1.24E-04	2.08E+03		3.50E+02				3.50E+02
4,4-DDT	50-29-3	6.00	1.30E-05		1.06E+04	1.78E+03				1.78E+03
Aldrin	309-00-2	6.18	1.35E-07		1.58E+06	2.65E+05				2.65E+05
alpha-BHC	319-84-6	3.80	6.78E-06		9.17E+01	1.54E+01				1.54E+01
beta-BHC	319-85-7	3.83	3.46E-07		1.95E+03	3.28E+02				3.28E+02
Chlordane	57-74-9	5.94	2.64E-05		4.46E+03	7.49E+02				7.49E+02
Delta-BHC	319-86-8	4.14	8.17E-07		1.75E+03	2.95E+02				2.95E+02

Table C2-3. Terrestrial Air-to-Plant Transfer Factors (APv) for Ecological Receptors (mg/kg tissue wet weight/mg/m^3 air)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	H[b] (atm x m^3/mol)	EPA (1999) APv[c] (kg air/kg dry wt)	APv Calculated by EPA (1999) Methods[d] (kg air/kg dry wt)	APv Adjusted to Wet Weight of Plants and Volume of Air (m^3 air/kg wet wt)[e]	Ecology Guidance APv	Default APv	Comments	Recommended APv[f]
Dieldrin	60-57-1	5.27	3.51E-09		6.52E+06	1.10E+06				1.10E+06
Endothall	145-73-3	-0.87	2.59E-10		2.56E+01	4.29E+00				4.29E+00
Endrin	72-20-8	4.89	1.19E-06		7.61E+03	1.28E+03				1.28E+03
gamma-BHC (Lindane)	58-89-9	3.72	1.56E-06		3.27E+02	5.50E+01				5.50E+01
Heptachlor	76-44-8	5.02	5.87E-06	2.09E+03		3.51E+02				3.51E+02
Isodrin	465-73-6	3.55	1.04E-05		3.24E+01	5.45E+00				5.45E+00
Methoxychlor	72-43-5	4.53	6.33E-06		5.84E+02	9.82E+01				9.82E+01
Silvex (2,4,5-TP)	93-72-1	4.07	6.23E-03		1.95E-01	3.28E-02				3.28E-02
Toxaphene	8001-35-2	5.50	4.51E-04		8.92E+01	1.50E+01				1.50E+01
Inorganic Chemicals and Compounds										
Metals										
Aluminum	7429-90-5	NA	NA	0.00E+00						0.00E+00
Antimony	7440-36-0	NA	NA	0.00E+00						0.00E+00
Arsenic	7440-38-2	NA	NA	0.00E+00						0.00E+00
Barium	7440-39-3	NA	NA	0.00E+00						0.00E+00
Beryllium	7440-41-7	NA	NA	0.00E+00						0.00E+00
Bismuth	7440-69-9	NA	NA	0.00E+00						0.00E+00
Boron	7440-42-8	NA	NA	0.00E+00						0.00E+00
Cadmium	7440-43-9	NA	NA	0.00E+00						0.00E+00
Calcium	7440-70-2	NA	NA	0.00E+00						0.00E+00
Chromium (and VI)	18540-29-9	NA	NA	0.00E+00						0.00E+00
Cobalt	7440-48-4	NA	NA	0.00E+00						0.00E+00
Copper	7440-50-8	NA	NA	0.00E+00						0.00E+00
Iron	7439-89-6	NA	NA	0.00E+00						0.00E+00
Lead	7439-92-1	NA	NA	0.00E+00						0.00E+00
Lithium	7439-93-2	NA	NA	0.00E+00						0.00E+00
Magnesium	7439-95-4	NA	NA	0.00E+00						0.00E+00
Manganese	7439-96-5	NA	NA	0.00E+00						0.00E+00
Mercury	7439-97-6	NA	NA	1.80E+03		3.03E+02				3.03E+02

Table C2-3. Terrestrial Air-to-Plant Transfer Factors (APv) for Ecological Receptors (mg/kg tissue wet weight/mg/m^3 air)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	H[b] (atm x m^3/mol)	EPA (1999) APv[c] (kg air/kg dry wt)	APv Calculated by EPA (1999) Methods[d] (kg air/kg dry wt)	APv Adjusted to Wet Weight of Plants and Volume of Air (m^3 air/kg wet wt)[e]	Ecology Guidance APv	Default APv	Comments	Recommended APv[f]
Molybdenum	7439-98-7	NA	NA	0.00E+00						0.00E+00
Nickel	7440-02-0	NA	NA	0.00E+00						0.00E+00
Potassium	7440-09-7	NA	NA	0.00E+00						0.00E+00
Rhodium	7440-16-6	NA	NA	0.00E+00						0.00E+00
Selenium	7782-49-2	NA	NA	0.00E+00						0.00E+00
Silicon	7440-21-3	NA	NA	0.00E+00						0.00E+00
Silver	7440-22-4	NA	NA	0.00E+00						0.00E+00
Sodium	7440-23-5	NA	NA	0.00E+00						0.00E+00
Strontium	7440-24-6	NA	NA	0.00E+00						0.00E+00
Tantalum	7440-25-7	NA	NA	0.00E+00						0.00E+00
Thallium	7440-28-0	NA	NA	0.00E+00						0.00E+00
Tin	7440-31-5	NA	NA	0.00E+00						0.00E+00
Tungsten	7440-33-7	NA	NA	0.00E+00						0.00E+00
Uranium	7440-61-1	NA	NA	0.00E+00						0.00E+00
Vanadium	7440-62-2	NA	NA	0.00E+00						0.00E+00
Yttrium	7440-65-5	NA	NA	0.00E+00						0.00E+00
Zinc	7440-66-6	NA	NA	0.00E+00						0.00E+00
Zirconium	7440-67-7	NA	NA	0.00E+00						0.00E+00
Non-metals and Anions										
Ammonia/Ammonium	7664-41-7	NA	NA	0.00E+00						0.00E+00
Bromide	24959-67-9	NA	NA	0.00E+00						0.00E+00
Chloride	16887-00-6	NA	NA	0.00E+00						0.00E+00
Cyanide	57-12-5	NA	NA	0.00E+00						0.00E+00
Fluoride	16984-48-8	NA	NA	0.00E+00						0.00E+00
Hydroxide	14280-30-9	NA	NA	0.00E+00						0.00E+00
Iodine	7553-56-2	NA	NA	0.00E+00						0.00E+00
Nitrate	14797-55-8	NA	NA	0.00E+00						0.00E+00
Nitrite	14797-65-0	NA	NA	0.00E+00						0.00E+00
Phosphate	14265-44-2	NA	NA	0.00E+00						0.00E+00

Table C2-3. Terrestrial Air-to-Plant Transfer Factors (APv) for Ecological Receptors (mg/kg tissue wet weight/mg/m^3 air)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	H[b] (atm x m^3/mol)	EPA (1999) APv[c] (kg air/kg dry wt)	APv Calculated by EPA (1999) Methods[d] (kg air/kg dry wt)	APv Adjusted to Wet Weight of Plants and Volume of Air (m^3 air/kg wet wt)[e]	Ecology Guidance APv	Default APv	Comments	Recommended APv[f]
Phosphorus	7723-14-0	NA	NA	0.00E+00						0.00E+00
Sulfate	14808-79-8	NA	NA	0.00E+00						0.00E+00
Total Sulfur	63705-05-5	NA	NA	0.00E+00						0.00E+00
Priority Pollutants										
Carbon Dioxide	124-38-9	NA	NA					1.00E+00		1.00E+00
Nitrogen Dioxide	10102-44-0	NA	NA					1.00E+00		1.00E+00
Ozone	10028-15-6	NA	NA					1.00E+00		1.00E+00
Particulate Matter	No CAS #	NA	NA	NA						NA
Sulfur Dioxode	7446-09-5	NA	NA					1.00E+00		1.00E+00
Radionuclides										
Americium-241	1596-10-2	NA	NA	0.00E+00						0.00E+00
Antimony-125	14234-35-6	NA	NA	0.00E+00						0.00E+00
Barium-137	13981-97-0	NA	NA	0.00E+00						0.00E+00
Cadmium-113	None	NA	NA	0.00E+00						0.00E+00
Cesium-134	13967-70-9	NA	NA	0.00E+00						0.00E+00
Cesium-137	10045-97-3	NA	NA	0.00E+00						0.00E+00
Europium-154	15585-10-1	NA	NA	0.00E+00						0.00E+00
Europium-155	14391-16-3	NA	NA	0.00E+00						0.00E+00
Nickel-63	13981-37-8	NA	NA	0.00E+00						0.00E+00
Plutonium-239	15117-48-3	NA	NA	0.00E+00						0.00E+00
Plutonium-241	14119-32-5	NA	NA	0.00E+00						0.00E+00
Samarium-151	15715-94-3	NA	NA	0.00E+00						0.00E+00
Strontium-90	10098-97-2	NA	NA	0.00E+00						0.00E+00
Technetium-99	14133-79-7	NA	NA	0.00E+00						0.00E+00
Tritium	10028-17-8	NA	NA	0.00E+00						0.00E+00
Yttrium-90	10098-91-6	NA	NA	0.00E+00						0.00E+00
Uranium-232	14158-29-3	NA	NA	0.00E+00						0.00E+00
Uranium-233	13968-55-3	NA	NA	0.00E+00						0.00E+00
Uranium-234	13966-29-5	NA	NA	0.00E+00						0.00E+00

Table C2-3. Terrestrial Air-to-Plant Transfer Factors (APv) for Ecological Receptors (mg/kg tissue wet weight/mg/m^3 air)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	H[b] (atm x m^3/mol)	EPA (1999) APv[c] (kg air/kg dry wt)	APv Calculated by EPA (1999) Methods[d] (kg air/kg dry wt)	APv Adjusted to Wet Weight of Plants and Volume of Air (m^3 air/kg wet wt)[e]	Ecology Guidance APv	Default APv	Comments	Recommended APv[f]
Uranium-235	15117-96-1	NA	NA	0.00E+00						0.00E+00
Uranium-236	13982-70-2	NA	NA	0.00E+00						0.00E+00
Uranium-238	7440-61-1	NA	NA	0.00E+00						0.00E+00
Actinium-227	14952-40-0	NA	NA	0.00E+00						0.00E+00
Americium-243	14993-75-0	NA	NA	0.00E+00						0.00E+00
Carbon-14	14762-75-5	NA	NA	0.00E+00						0.00E+00
Cobalt-60	10198-40-0	NA	NA	0.00E+00						0.00E+00
Curium-242	15510-73-3	NA	NA	0.00E+00						0.00E+00
Curium-243	15757-87-6	NA	NA	0.00E+00						0.00E+00
Curium-244	13981-15-2	NA	NA	0.00E+00						0.00E+00
Europium-152	14683-23-9	NA	NA	0.00E+00						0.00E+00
Iodine-129	15046-84-1	NA	NA	0.00E+00						0.00E+00
Neptunium-237	13994-20-2	NA	NA	0.00E+00						0.00E+00
Nickel-59	14336-70-0	NA	NA	0.00E+00						0.00E+00
Niobium-93	None	NA	NA	0.00E+00						0.00E+00
Plutonium-238	13981-16-3	NA	NA	0.00E+00						0.00E+00
Plutonium-240	14119-33-6	NA	NA	0.00E+00						0.00E+00
Plutonium-242	13982-10-0	NA	NA	0.00E+00						0.00E+00
Protactinium-231	14331-85-2	NA	NA	0.00E+00						0.00E+00
Radium-226	13982-63-3	NA	NA	0.00E+00						0.00E+00
Radium-228	15262-20-1	NA	NA	0.00E+00						0.00E+00
Ruthenium-106	13967-48-1	NA	NA	0.00E+00						0.00E+00
Selenium-79	None	NA	NA	0.00E+00						0.00E+00
Thorium-229	15594-54-4	NA	NA	0.00E+00						0.00E+00
Thorium-232	7440-29-1	NA	NA	0.00E+00						0.00E+00
Tin-126	15832-50-5	NA	NA	0.00E+00						0.00E+00
Zirconium-93	15751-77-6	NA	NA	0.00E+00						0.00E+00

NA = Not applicable

[a] $\log_{10}$ of K_{ow} values in Table 4.1

Table C2-3. Terrestrial Air-to-Plant Transfer Factors (APv) for Ecological Receptors (mg/kg tissue wet weight/mg/m^3 air)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	H[b] (atm x m^3/mol)	EPA (1999) APv[c] (kg air/kg dry wt)	APv Calculated by EPA (1999) Methods[d] (kg air/kg dry wt)	APv Adjusted to Wet Weight of Plants and Volume of Air (m^3 air/kg wet wt)[e]	Ecology Guidance APv	Default APv	Comments	Recommended APv[f]

[b] H values from Appendix B Table B-1-1

[c] Data published in Appendix C of EPA (1999), Table C-7

[d] Calculated as described in Appendix C, Sect. C-1.7 of EPA (1999): $APv = \rho_{air} \times B_{vol} / ((1-f_{water}) \times \rho_{forage} \times 100)$ and $\log(B_{vol}) = 1.605 \log K_{ow} - \log (H/RT) - 1.654$

where ρ_{air} = density of air, 1.19 g/L

B_{vol} = volumetric air-to-plant transfer

f_{water} = fraction of water in plants

ρ_{forage} = density of forage, 770 g/L

100 = uncertainty factor

R = gas-law constant, 8.025E-5 atm m^3/(deg mole)

T = temperature, 298 deg K

e Adjusted by dividing by the density of air (1.19 kg/m^3) and multiplying by 0.2 to adjust from dry weight to wet weight of plants

[f] Selection criteria described in Sect. 8.2.4.3

[g] Value for Aroclor 1254 from Appendix C of EPA (1999), Table C-7, was used for PCB mixtures

Table C2-4. Terrestrial Soil-to-Invertebrate Transfer Factors (BAF-S) for Ecological Receptors (mg/kg tissue wet weight / mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) BAF-S[b]	BAF-S Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance BAF-S	SAIC Compiled BAF-S	Notes	Comments	Recommended BAF-S[d]
Organic Compounds										
Aromatic Halogenated Hydrocarbons										
4-Chloro-3-methylphenol	59-50-7	3.10		2.47E+01	e		5.00E-02	f		2.47E+01
2,3,4,6-Tetrachlorophenol	58-90-2	4.42		2.98E+02	e					2.98E+02
Aromatic Nonhalogenated Hydrocarbons										
2-Nitrotoluene	88-72-2	2.30		5.47E+00	e					5.47E+00
4-Nitrobiphenyl	92-93-3	3.77		8.74E+01	e					8.74E+01
Benzaldehyde	100-52-7	1.48		1.16E+00	e					1.16E+00
Benzene	71-43-2	-4.99		5.84E-06	e		5.00E-02	f		5.84E-06
Benzyl alcohol	100-51-6	1.10		5.69E-01	e					5.69E-01
Ethyl benzene	100-41-4	3.12		2.58E+01	e		5.00E-02	f		2.58E+01
m-Xylene	108-38-3	3.20		2.99E+01	e		5.00E-02	f		2.99E+01
o-Xylene	95-47-6	3.13		2.62E+01	e		5.00E-02	f		2.62E+01
p-Xylene	106-42-3	3.17		2.82E+01	e		5.00E-02	f		2.82E+01
Styrene	100-42-5	2.93		1.79E+01	e					1.79E+01
Toluene	108-88-3	2.67		1.09E+01	e		5.00E-02	f		1.09E+01
Non-aromatic Nonhalogenated Hydrocarbons										
1,2-Epoxybutane	106-88-7	1.44		1.08E+00	e					1.08E+00
1,3-Butadiene	106-99-0	1.90		2.57E+00	e					2.57E+00
1,4-Dioxane	123-91-1	-0.27	4.31E-02		e					4.31E-02
1-Methylpropyl alcohol	78-92-2	0.61		2.26E-01	e					2.26E-01
1-Nitropropane	108-03-2	0.87		3.69E-01	e					3.69E-01
2,2,4-Trimethylpentane	540-84-1	5.02		9.23E+02	e					9.23E+02
2-Butanone	78-93-3	0.28		1.21E-01	e		5.00E-02	f		1.21E-01
2-Butenaldehyde (2-Butenal)	4170-30-3	No data		No data			1.00E+01		Default	1.00E+01
2-Ethoxyethanol	110-80-5	-0.10		5.92E-02	e					5.92E-02
2-Heptanone	110-43-0	1.98		2.99E+00	e					2.99E+00
2-Hexanone	591-78-6	1.38		9.64E-01	e					9.64E-01
2-Methoxyethanol	109-86-4	0.25		1.15E-01	e					1.15E-01
2-Methyl-2-propanol	75-65-0	0.35		1.38E-01	e					1.38E-01

Table C2-4. Terrestrial Soil-to-Invertebrate Transfer Factors (BAF-S) for Ecological Receptors (mg/kg tissue wet weight / mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BAF-S [b]	BAF-S Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BAF-S	SAIC Compiled BAF-S	Notes	Comments	Recommended BAF-S [d]
2-Methyl-2-propenenitrile	126-98-7	0.54		1.98E-01	e					1.98E-01
2-Methylaziridine	75-55-8	-0.60		2.29E-02	e					2.29E-02
2-Methylpropyl alcohol	78-83-1	0.76		3.00E-01	e					3.00E-01
2-Pentanone	107-87-9	0.91		3.97E-01	e					3.97E-01
2-Propanone (Acetone)	67-64-1	-0.22	4.70E-02		e		5.00E-02	f		4.70E-02
2-Propene-1-ol	107-18-6	0.17		9.85E-02	e					9.85E-02
2-Propyl alcohol	67-63-0	0.05		7.85E-02	e					7.85E-02
3-Heptanone	106-35-4	No data		No data			1.00E+01		Default	1.00E+01
3-Methyl-1-butanol	123-51-3	No data		No data			1.00E+01		Default	1.00E+01
3-Methyl-2-butanone	563-80-4	No data		No data			1.00E+01		Default	1.00E+01
3-Pentanone	96-22-0	0.99		4.62E-01	e					4.62E-01
4-Heptanone	123-19-3	No data		No data			1.00E+01		Default	1.00E+01
4-Methyl-2-pentanone	108-10-1	1.19		6.74E-01	e		5.00E-02	f		6.74E-01
4-Methyl-3-penten-2-one	141-79-7	No data		No data			1.00E+01		Default	1.00E+01
5-Methyl-2-hexanone	110-12-3	No data		No data			1.00E+01		Default	1.00E+01
Acetaldehyde	75-07-0	-0.22		4.72E-02	e					4.72E-02
Acetamide	60-35-5	-1.26		6.64E-03	e					6.64E-03
Acetic acid	64-19-7	-0.17		5.19E-02	e					5.19E-02
Acetic acid ethyl ester	141-78-6	0.73		2.83E-01	e					2.83E-01
Acetic acid n-butyl ester	123-86-4	1.73		1.87E+00	e					1.87E+00
Acetonitrile	75-05-8	-0.34		3.76E-02	e					3.76E-02
Acrolein	107-02-8	-0.01		7.03E-02	e					7.03E-02
Acrylonitrile	107-13-1	0.25	1.15E-01		e					1.15E-01
Bis(isopropyl)ether	108-20-3	No data		No data			1.00E+01		Default	1.00E+01
Butane	106-97-8	2.89		1.66E+01	e					1.66E+01
Carbon disulfide	75-15-0	2.00		3.10E+00	e					3.10E+00
Cyanogen	460-19-5	0.81		3.28E-01	e					3.28E-01
Cyclohexane	110-82-7	3.44		4.69E+01	e					4.69E+01
Cyclohexanone	108-94-1	0.81		3.29E-01	e					3.29E-01
Cyclohexene	110-83-8	2.86		1.57E+01	e					1.57E+01

Table C2-4. Terrestrial Soil-to-Invertebrate Transfer Factors (BAF-S) for Ecological Receptors (mg/kg tissue wet weight / mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) BAF-S[b]	BAF-S Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance BAF-S	SAIC Compiled BAF-S	Notes	Comments	Recommended BAF-S[d]
Cyclopentane	287-92-3	3.00		2.05E+01	e					2.05E+01
Ethyl alcohol	64-17-5	0.31		1.28E-01	e					1.28E-01
Ethyl ether	60-29-7	0.89		3.83E-01	e					3.83E-01
Ethyl methacrylate	97-63-2	1.59		1.43E+00	e					1.43E+00
Formaldehyde	50-00-0	0.34	1.36E-01		e					1.36E-01
Formamide	75-12-7	-1.51		4.14E-03	e					4.14E-03
Formic acid	64-18-6	-0.54		2.59E-02	e					2.59E-02
Formic acid, methyl ester	107-31-3	-0.26		4.34E-02	e					4.34E-02
Glycidylaldehyde	765-34-4	-0.73		1.80E-02	e					1.80E-02
Methyl acetate	79-20-9	0.18		1.00E-01	e					1.00E-01
Methyl alcohol	67-56-1	-0.71		1.87E-02	e					1.87E-02
Methyl isocyanate	624-83-9	No data		No data					Reacts with water	NA
Methyl methacrylate	80-62-6	0.79		3.17E-01	e					3.17E-01
Methyl tert-butyl ether	1634-04-4	0.94		4.21E-01	e					4.21E-01
Methylacetylene	74-99-7	0.94		4.21E-01	e					4.21E-01
Methylcyclohexane	108-87-2	4.10		1.63E+02	e					1.63E+02
N,N-Dimethylacetamide	127-19-5	No data		No data			1.00E+01		Default	1.00E+01
n-Butyl alcohol	71-36-3	0.88		3.76E-01	e					3.76E-01
n-Heptane	142-82-5	4.66		4.68E+02	e					4.68E+02
n-Hexane	110-54-3	4.11		1.66E+02	e					1.66E+02
Nitromethane	75-52-5	-0.35		3.69E-02	e					3.69E-02
n-Nonane	111-84-2	5.65		3.03E+03	e					3.03E+03
n-Octane	111-65-9	4.00		1.35E+02	e					1.35E+02
n-Pentane	109-66-0	3.21		3.04E+01	e					3.04E+01
n-Propionaldehyde	123-38-6	0.59		2.17E-01	e					2.17E-01
n-Propyl alcohol	71-23-8	0.25		1.14E-01	e					1.14E-01
n-Valeraldehyde	110-62-3	No data		No data			1.00E+01		Default	1.00E+01
Oxirane	75-21-8	-0.30		4.06E-02	e					4.06E-02
p-Cymene	99-87-6	4.10		1.63E+02	e					1.63E+02

Table C2-4. Terrestrial Soil-to-Invertebrate Transfer Factors (BAF-S) for Ecological Receptors (mg/kg tissue wet weight / mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BAF-S [b]	BAF-S Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BAF-S	SAIC Compiled BAF-S	Notes	Comments	Recommended BAF-S [d]
Phosgene	75-44-5	No data		No data			1.00E+01		Default	1.00E+01
Propargyl alcohol	107-19-7	0.26		1.17E-01	e					1.17E-01
Propionic acid	79-09-4	0.33		1.33E-01	e					1.33E-01
Propionitrile	107-12-0	0.16		9.66E-02	e					9.66E-02
Propylene gylcol monomethyl ether	107-98-2	-0.18		5.09E-02	e					5.09E-02
p-tert-Butyltoluene	98-51-1	No data		No data			1.00E+01		Default	1.00E+01
Triethylamine	121-44-8	0.16		9.66E-02	e					9.66E-02
Trimethylamine	75-50-3	0.16		9.66E-02	e					9.66E-02
Vinyl acetate	108-05-4	0.70		2.67E-01	e					2.67E-01
Non-aromatic Halogenated Hydrocarbons										
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	No data		No data			1.00E+01		Default	1.00E+01
1,1,1,2-Tetrachloroethane	630-20-6	2.63		1.02E+01	e					1.02E+01
1,1,1-Trichloroethane	71-55-6	2.42		6.88E+00	e					6.88E+00
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	3.73		8.11E+01	e					8.11E+01
1,1,2,2-Tetrachloroethane	79-34-5	4.64		4.54E+02	e		5.00E-02	f		4.54E+02
1,1,2,2-Tetrachloroethene	127-18-4	2.55		8.68E+00	e		5.00E-02	f		8.68E+00
1,1,2-Trichloroethane	79-00-5	2.10		3.73E+00	e					3.73E+00
1,1,2-Trichloroethylene	79-01-6	2.43		7.02E+00	e		5.00E-02	f		7.02E+00
1,1-Dichloroethane	75-34-3	1.79		2.10E+00	e					2.10E+00
1,1-Dichloroethene	75-35-4	2.12		3.90E+00	e					3.90E+00
1,2,2-Trichloro-1,1,2-trifluoroethane	76-13-1	3.16		2.77E+01	e					2.77E+01
1,2,3-Trichloropropane	96-18-4	2.25		4.98E+00	e					4.98E+00
1,2-Dibromo-3-chloropropane	96-12-8	2.34		5.90E+00	e					5.90E+00
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	2.82		1.46E+01	e					1.46E+01
1,2-Dichloroethane	107-06-2	1.46		1.13E+00	e		5.00E-02	f		1.13E+00
1,2-Dichloroethylene	540-59-0	0.48		1.77E-01	e					1.77E-01
1,2-Dichloropropane	78-87-5	2.25		4.98E+00	e					4.98E+00
1,3-Dichloropropene	542-75-6	1.75		1.93E+00	e					1.93E+00
1,4-Dichloro-2-butene	764-41-0	0.87		3.70E-01	e					3.70E-01
1-Chloroethene	75-01-4	1.15		6.20E-01	e					6.20E-01

Table C2-4. Terrestrial Soil-to-Invertebrate Transfer Factors (BAF-S) for Ecological Receptors (mg/kg tissue wet weight / mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BAF-S [b]	BAF-S Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BAF-S	SAIC Compiled BAF-S	Notes	Comments	Recommended BAF-S [d]
2,2-Dichloropropionic acid	75-99-0	0.78		3.10E-01	e					3.10E-01
2-Chloropropane	75-29-6	1.90		2.57E+00	e					2.57E+00
3-Chloropropene (allyl chloride)	107-05-1	0.95		4.32E-01	e					4.32E-01
Bromochloromethane	74-97-5	1.41		1.02E+00	e					1.02E+00
Bromodichloromethane	75-27-4	2.03		3.26E+00	e					3.26E+00
Bromoethene	593-60-2	1.07		5.34E-01	e					5.34E-01
Bromoform	75-25-2	2.35		6.01E+00	e					6.01E+00
Bromomethane	74-83-9	1.11		5.84E-01	e					5.84E-01
Carbon tetrachloride	56-23-5	2.72	1.20E+01		e					1.20E+01
Chlorodibromomethane	124-48-1	2.18		4.33E+00	e					4.33E+00
Chlorodifluoromethane	75-45-6	1.08		5.47E-01	e					5.47E-01
Chloroethane	75-00-3	3.10		2.47E+01	e					2.47E+01
Chloroform	67-66-3	1.95	2.82E+00		e		5.00E-02	f		2.82E+00
Chloromethane	74-87-3	0.90		3.92E-01	e					3.92E-01
Chloropentafluoroethane	76-15-3	No data		No data			1.00E+01		Default	1.00E+01
cis-1,2-Dichloroethene	156-59-2	1.98		3.00E+00	e		5.00E-02	f		3.00E+00
cis-1,3-Dichloropropene	10061-01-5	No data		No data			1.00E+01		Default	1.00E+01
Cyanogen bromide	506-68-3	No data		No data			1.00E+01		Default	1.00E+01
Cyanogen chloride	506-77-4	0.20		1.04E-01	e					1.04E-01
Dichlorodifluoromethane	75-71-8	2.16		4.19E+00	e					4.19E+00
Dichlorofluoromethane	75-43-4	No data		No data			1.00E+01		Default	1.00E+01
Dichloromethane	75-09-2	1.26		7.62E-01	e					7.62E-01
Difluorodibromomethane	75-61-6	No data		No data			1.00E+01		Default	1.00E+01
Hexafluoroacetone	684-16-2	No data		No data			1.00E+01		Default	1.00E+01
Iodomethane	74-88-4	1.69		1.73E+00	e					1.73E+00
Methylene bromide	74-95-3	1.62		1.52E+00	e					1.52E+00
Pentachloroethane	76-01-7	3.05		2.25E+01	e					2.25E+01
trans-1,2-Dichloroethene	156-60-5	1.98		3.00E+00	e		5.00E-02	f		3.00E+00
trans-1,3-Dichloropropene	10061-02-6	2.06		3.48E+00	e					3.48E+00
Trichloroacetic acid	76-03-9	1.33		8.78E-01	e					8.78E-01

Table C2-4. Terrestrial Soil-to-Invertebrate Transfer Factors (BAF-S) for Ecological Receptors (mg/kg tissue wet weight / mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) BAF-S[b]	BAF-S Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance BAF-S	SAIC Compiled BAF-S	Notes	Comments	Recommended BAF-S[d]
Trichlorofluoroethane	27154-33-2	No data		No data			1.00E+01		Default	1.00E+01
Trichlorofluoromethane	75-69-4	2.53		8.46E+00	e					8.46E+00
Trifluorobromomethane	75-63-8	1.86		2.38E+00	e					2.38E+00
Dioxin and Furan Compounds										
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	35822-46-9	8.20	8.10E-02		g		3.70E+05	e		8.10E-02
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	7.92	1.70E-02		g		2.19E+05	e		1.70E-02
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	7.92	6.20E-01		g		2.19E+05	e		6.20E-01
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	39227-28-6	7.79	4.90E-01		g		1.71E+05	e		4.90E-01
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	7.25	1.21E-01		g		6.20E+04	e		1.21E-01
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	57653-85-7	7.25	1.90E-01		g		6.20E+04	e		1.90E-01
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	7.25	3.00E-01		g		6.20E+04	e		3.00E-01
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	19408-74-3	7.25	2.20E-01		g		6.20E+04	e		2.20E-01
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	7.25	1.00E+00		g		6.20E+04	e		1.00E+00
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	40321-76-4	6.64	1.46E+00		g		1.96E+04	e		1.46E+00
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	6.79	3.20E-01		g		2.60E+04	e		3.20E-01
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	7.25	1.07E+00		g		6.20E+04	e		1.07E+00
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	6.92	2.54E+00		g		3.32E+04	e		2.54E+00
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1746-01-6	6.64	1.59E+00				1.96E+04	e		1.59E+00
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	6.53	1.27E+00		g		1.59E+04	e		1.27E+00
Dibenzofuran	132-64-9	4.33		2.51E+02	e		5.00E-02	e		2.51E+02
Octachlorodibenzo(p)dioxin	3268-87-9	8.20	1.90E-02		g		3.71E+05	e		1.90E-02
Octachlorodibenzofuran	39001-02-0	8.78	2.50E-02		g		1.11E+06	e		2.50E-02
PCBs										
Polychlorinated biphenyls (PCBs)[h]	1336-36-3	7.31	1.13E+00				5.80E+00	f		1.13E+00
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	7.08		4.49E+04	e					4.49E+04
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	7.12		4.84E+04	e					4.84E+04
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	No data		No data			1.00E+01		Default	1.00E+01
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	No data		No data			1.00E+01		Default	1.00E+01
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	No data		No data			1.00E+01		Default	1.00E+01
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	No data		No data			1.00E+01		Default	1.00E+01

Table C2-4. Terrestrial Soil-to-Invertebrate Transfer Factors (BAF-S) for Ecological Receptors (mg/kg tissue wet weight / mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) BAF-S[b]	BAF-S Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance BAF-S	SAIC Compiled BAF-S	Notes	Comments	Recommended BAF-S[d]
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	No data		No data			1.00E+01		Default	1.00E+01
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	No data		No data			1.00E+01		Default	1.00E+01
2',3,4,4',5-Pentachlorobiphenyl	no cas #	No data		No data			1.00E+01		Default	1.00E+01
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	7.12		4.84E+04	e					4.84E+04
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	7.41		8.34E+04	e					8.34E+04
3,3',4,4',5-Pentachlorobiphenyl	no cas #	No data		No data			1.00E+01		Default	1.00E+01
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	No data		No data			1.00E+01		Default	1.00E+01
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	No data		No data			1.00E+01		Default	1.00E+01
Phthalates										
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	5.20	1.31E+03		e		5.00E-02	f		1.31E+03
Butylbenzyl phthalate	85-68-7	4.41		2.94E+02	e		5.00E-02	f		2.94E+02
Dibutyl phthalate	84-74-2	4.72		5.25E+02	e		5.00E-02	f		5.25E+02
Diethyl phthalate	84-66-2	4.44		3.07E+02	e		5.00E-02	f		3.07E+02
Dimethylphthalate	131-11-3	1.63		1.56E+00	e					1.56E+00
n-Dioctyl phthalate	117-84-0	9.33	3.13E+06		e		5.00E-02	f		3.13E+06
Light Polycyclic Aromatic Hydrocarbons (molecular weight <200 g/mole)										
2-Chloronapthalene	91-58-7	4.07		1.53E+02	e					1.53E+02
2-Methyl naphthalene	91-57-6	3.86		1.04E+02	e		5.00E-02	f		1.04E+02
5-Nitroacenaphthene	602-87-9	No data		No data			1.00E+01		Default	1.00E+01
Acenaphthene	83-32-9	3.96		1.26E+02	e					1.26E+02
Acenaphthylene	208-96-8	4.07		1.54E+02	e					1.54E+02
Anthracene	120-12-7	4.47		3.27E+02	e		5.00E-02	f		3.27E+02
Fluorene	86-73-7	4.18		1.89E+02	e					1.89E+02
Indene	95-13-6	No data		No data			1.00E+01		Default	1.00E+01
Naphthalene	91-20-3	3.37		4.13E+01	e		5.00E-02	f		4.13E+01
Phenanthrene	85-01-8	4.55		3.81E+02	e		5.00E-02	f		3.81E+02
Pyrene	129-00-0	5.00		8.89E+02	e		5.00E-02	f		8.89E+02
Heavy Polycyclic Aromatic Hydrocarbons (molecular weight >200 g/mole)										
3-Methylcholanthrene	56-49-5	7.11		4.75E+04	e					4.75E+04
5-Methylchrysene	3697-24-3	No data		No data			1.00E+01		Default	1.00E+01

Table C2-4. Terrestrial Soil-to-Invertebrate Transfer Factors (BAF-S) for Ecological Receptors (mg/kg tissue wet weight / mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) BAF-S[b]	BAF-S Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance BAF-S	SAIC Compiled BAF-S	Notes	Comments	Recommended BAF-S[d]
Benzo(a)anthracene	56-55-3	5.68	3.00E-02				5.00E-02	f		3.00E-02
Benzo(a)pyrene	50-32-8	6.13	7.00E-02				5.00E-02	f		7.30E-02
Benzo(b)fluoranthene	205-99-2	6.20	7.00E-02				5.00E-02	f		7.00E-02
Benzo(e)pyrene	192-97-2	7.40		8.21E+04	e					8.21E+04
Benzo(g,h,i)perylene	191-24-2	7.10		4.67E+04	e		5.00E-02	f		4.67E+04
Benzo(j)fluoranthene	205-82-3	6.44		1.34E+04	e					1.34E+04
Benzo(k)fluoranthene	207-08-9	6.20	8.00E-02				5.00E-02	f		8.00E-02
Benzo[a,i]pyrene	191-30-0	No data		No data			1.00E+01		Default	1.00E+01
Chrysene	218-01-9	5.74	4.00E-02				5.00E-02	f		4.00E-02
Dibenz(a,h)anthracene	53-70-3	6.55	7.00E-02				5.00E-02	f		7.00E-02
Dibenz[a,h]acridine	226-36-8	No data		No data			1.00E+01		Default	1.00E+01
Dibenz[a,j]acridine	224-42-0	No data		No data			1.00E+01		Default	1.00E+01
Dibenzo(a,e)fluoranthene	5385-75-1	No data		No data			1.00E+01		Default	1.00E+01
Dibenzo(a,h)fluoranthene	no cas #	No data		No data			1.00E+01		Default	1.00E+01
Dibenzo[a,e]pyrene	192-65-4	No data		No data			1.00E+01		Default	1.00E+01
Dibenzo[a,h]pyrene	189-64-0	No data		No data			1.00E+01		Default	1.00E+01
Dibenzo[a,i]pyrene	189-55-9	7.29		6.68E+04	e					6.68E+04
Fluoranthene	206-44-0	5.08		1.04E+03	e		5.00E-02	f		1.04E+03
Hexachloronaphthalene	1335-87-1	7.59		1.18E+05	e					1.18E+05
Indeno(1,2,3-cd)pyrene	193-39-5	6.91	8.00E-02				5.00E-02	f		8.00E-02
Octachloronaphthalene	2234-13-1	6.42		1.29E+04	e					1.29E+04
Pentachloronaphthalene	1321-64-8	No data		No data			1.00E+01		Default	1.00E+01
Tetrachloronaphthalene	1335-88-2	No data		No data			1.00E+01		Default	1.00E+01
Trichloronaphthalene	1321-65-9	No data		No data			1.00E+01		Default	1.00E+01
Light Substituted Benzene Compounds (MW <200 g/mole)										
1,2,3-Trichlorobenzene	87-61-6	4.05		1.47E+02	e					1.47E+02
1,2,4-Trichlorobenzene	120-82-1	3.99		1.32E+02	e					1.32E+02
1,2,4-Trimethyl benzene	95-63-6	3.65		6.97E+01	e					6.97E+01
1,2-Dichlorobenzene	95-50-1	3.45		4.74E+01	e					4.74E+01
1,3,5-Trimethyl benzene	108-67-8	3.42		4.52E+01	e					4.52E+01

Table C2-4. Terrestrial Soil-to-Invertebrate Transfer Factors (BAF-S) for Ecological Receptors (mg/kg tissue wet weight / mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) BAF-S[b]	BAF-S Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance BAF-S	SAIC Compiled BAF-S	Notes	Comments	Recommended BAF-S[d]
1,3-Dichlorobenzene	541-73-1	3.53		5.56E+01	e					5.56E+01
1,3-Dinitrobenzene	99-65-0	1.49	1.19E+00		e					1.19E+00
1,4-Dichlorobenzene	106-46-7	3.41		4.45E+01	e					4.45E+01
1,4-Dinitrobenzene	100-25-4	1.50		1.21E+00	e					1.21E+00
2,4,5-Trichlorophenol	95-95-4	3.87		1.06E+02	e					1.06E+02
2,4,6-Trichlorophenol	88-06-2	3.71		7.83E+01	e					7.83E+01
2,4-Dichlorophenol	120-83-2	3.04		2.20E+01	e					2.20E+01
2,4-Dimethylphenol	105-67-9	2.36		6.12E+00	e					6.12E+00
2,4-Dinitrophenol	51-28-5	1.52		1.25E+00	e					1.25E+00
2,4-Dinitrotoluene	121-14-2	2.00	3.08E+00		e					3.08E+00
2,6-Dinitrotoluene	606-20-2	1.89	2.51E+00		e		5.00E-02	f		2.51E+00
2-Chlorophenol	95-57-8	2.16		4.21E+00	e					4.21E+00
2-Chlorotoluene	95-49-8	3.54		5.63E+01	e					5.63E+01
2-Nitrophenol	88-75-5	1.79		2.09E+00	e					2.09E+00
4,6-Dinitro-o-cresol	534-52-1	2.85		1.54E+01	e					1.54E+01
4-Chlorotoluene	106-43-4	3.33		3.81E+01	e					3.81E+01
4-Nitrophenol	100-02-7	1.91		2.62E+00	e		5.00E-02	f		2.62E+00
alpha-Methylstyrene	98-83-9	3.46		4.90E+01	e					4.90E+01
Aniline	62-53-3	0.98		4.54E-01	e					4.54E-01
Benzotrichloride	98-07-7	2.92		1.76E+01	e					1.76E+01
Benzyl chloride	100-44-7	0.36		1.41E-01	e					1.41E-01
Bromobenzene	108-86-1	2.99		2.01E+01	e					2.01E+01
Chlorobenzene	108-90-7	2.79		1.38E+01	e		5.00E-02	f		1.38E+01
Cumene	98-82-8	3.61		6.50E+01	e					6.50E+01
m-Cresol	108-39-4	1.96		2.87E+00	e					2.87E+00
n-Butyl benzene	104-51-8	4.28		2.29E+02	e					2.29E+02
Nitrobenzene	98-95-3	1.83	2.26E+00		e					2.26E+00
n-Propyl benzene	103-65-1	3.69		7.52E+01	e					7.52E+01
o-Cresol	95-48-7	2.02		3.23E+00	e					3.23E+00
o-Dinitrobenzene	528-29-0	1.60		1.46E+00	e					1.46E+00

Table C2-4. Terrestrial Soil-to-Invertebrate Transfer Factors (BAF-S) for Ecological Receptors (mg/kg tissue wet weight / mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) BAF-S[b]	BAF-S Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance BAF-S	SAIC Compiled BAF-S	Notes	Comments	Recommended BAF-S[d]
o-Nitroaniline	88-74-4	1.85		2.34E+00	e					2.34E+00
o-Toluidine	95-53-4	1.34		8.95E-01	e					8.95E-01
p-Chloroaniline	106-47-8	1.87		2.43E+00	e					2.43E+00
p-Cresol	106-44-5	1.94		2.77E+00	e					2.77E+00
Phenol	108-95-2	1.48		1.16E+00	e		5.00E-02	f		1.16E+00
p-Nitrochlorobenzene	100-00-5	2.39		6.48E+00	e					6.48E+00
p-Toluidine	106-49-0	1.40		1.00E+00	e					1.00E+00
sec-Butyl benzene	135-98-8	4.57		3.95E+02	e					3.95E+02
tert-Butyl benzene	98-06-6	4.11		1.66E+02	e					1.66E+02
Toluene-2,6-diamine	823-40-5	1.45		1.11E+00	e					1.11E+00
Trimethyl benzene	25551-13-7	3.40		4.35E+01	e					4.35E+01
Other Light Semivolatile Compounds (molecular weight <200 g/mole)										
1,1'-Biphenyl	92-52-4	3.90		1.12E+02	e					1.12E+02
1,1-Dimethylhydrazine	57-14-7	0.60		2.20E-01	e					2.20E-01
1,2-Dimethylhydrazine	540-73-8	-1.37		5.42E-03	e					5.42E-03
1,2-Diphenylhydrazine	122-66-7	2.94		1.83E+01	e					1.83E+01
1,3-Propane sultone	1120-71-4	-0.52		2.66E-02	e					2.66E-02
2,4-Toluene diisocyante	584-84-9	No data		No data					Decomposes rapidly in water	NA
2-Chloroacetophenone	532-27-4	2.59		9.53E+00	e					9.53E+00
2-Propenoic acid	79-10-7	0.33		1.33E-01	e					1.33E-01
4,4-Methylenedianiline	101-77-9	3.38		4.19E+01	e					4.19E+01
Acetophenone	98-86-2	1.64		1.58E+00	e					1.58E+00
Benzoic acid	65-85-0	1.86		2.38E+00	e					2.38E+00
bis(2-Chloroethoxy)methane	111-91-1	7.59		1.18E+05	e					1.18E+05
bis(2-Chloroethyl) ether	111-44-4	1.30		8.31E-01	e					8.31E-01
Chlorocyclopentadiene	41851-50-7	2.43		6.99E+00	e					6.99E+00
Cyclohexanol	108-93-0	1.23		7.27E-01	e					7.27E-01
Dichloroisopropyl ether	108-60-1	2.58		9.27E+00	e					9.27E+00

Table C2-4. Terrestrial Soil-to-Invertebrate Transfer Factors (BAF-S) for Ecological Receptors (mg/kg tissue wet weight / mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BAF-S [b]	BAF-S Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BAF-S	SAIC Compiled BAF-S	Notes	Comments	Recommended BAF-S [d]
Dichloromethyl ether	542-88-1	-0.38		3.49E-02	e					3.49E-02
Dichloropentadiene	no cas #	No data		No data			1.00E+01		Default	1.00E+01
Dimethyl sulfate	77-78-1	0.32		1.30E-01	e					1.30E-01
Dimethylaniline	121-69-7	-0.88		1.36E-02	e					1.36E-02
Di-n-propylnitrosamine	621-64-7	1.38		9.65E-01	e					9.65E-01
Diphenyl ether	101-84-8	4.21		2.00E+02	e					2.00E+02
Epichlorohydrin	106-89-8	0.25		1.15E-01	e					1.15E-01
Ethyl carbamate (urethane)	51-79-6	-0.15		5.38E-02	e					5.38E-02
Ethyl methanesulfonate	62-50-0	0.05		7.84E-02	e					7.84E-02
Ethylene dibromide	106-93-4	1.75		1.94E+00	e					1.94E+00
Ethylene glycol	107-21-1	-0.91		1.27E-02	e					1.27E-02
Ethylene glycol monobutyl ether	111-76-2	1.55		1.34E+00	e					1.34E+00
Ethylene glycol monoethyl ether acetate	111-15-9	0.62		2.30E-01	e					2.30E-01
Ethylene thiourea	96-45-7	-0.64		2.13E-02	e					2.13E-02
Furfural	98-01-1	0.96		4.37E-01	e					4.37E-01
Maleic hydrazide	123-33-1	-0.74		1.77E-02	e					1.77E-02
Malononitrile	109-77-3	0.04		7.70E-02	e					7.70E-02
Methyl styrene (mixed isomers)	25013-15-4	3.35		3.96E+01	e					3.96E+01
Methylhydrazine	60-34-4	-1.06		9.68E-03	e					9.68E-03
N,N-Diphenylamine	122-39-4	3.50		5.25E+01	e					5.25E+01
Nitric acid, propyl ester	627-13-4	No data		No data			1.00E+01		Default	1.00E+01
N-Nitrosodi-n-butylamine	924-16-3	2.41		6.73E+00	e					6.73E+00
N-Nitrosomorpholine	59-89-2	0.98		4.54E-01	e					4.54E-01
N-Nitroso-N,N-dimethylamine	62-75-9	-0.47		2.94E-02	e					2.94E-02
o-Anisidine	90-04-0	1.18		6.61E-01	e					6.61E-01
Oxalic acid	144-62-7	No data		No data			1.00E+01		Default	1.00E+01
Phthalic anhydride	85-44-9	-0.60		2.30E-02						2.30E-02
p-Phthalic acid	100-21-0	0.82		3.37E-01	e					3.37E-01
Pyridine	110-86-1	0.67		2.53E-01	e					2.53E-01
Quinoline	91-22-5	2.03		3.29E+00	e					3.29E+00

Table C2-4. Terrestrial Soil-to-Invertebrate Transfer Factors (BAF-S) for Ecological Receptors (mg/kg tissue wet weight / mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) BAF-S[b]	BAF-S Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance BAF-S	SAIC Compiled BAF-S	Notes	Comments	Recommended BAF-S[d]
Quinone	106-51-4	0.20		1.04E-01	e					1.04E-01
Safrole	94-59-7	2.66		1.08E+01	e					1.08E+01
Tetrahydrofuran	109-99-9	0.45		1.66E-01	e					1.66E-01
Other Heavy Semivolatile Compounds (molecular weight >200 g/mole)										
1,2,4,5-Tetrachlorobenzene	95-94-3	4.64		4.51E+02	e					4.51E+02
1,3,5-Trinitrobenzene	99-35-4	1.18		6.60E-01	e					6.60E-01
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	4.17		1.86E+02	e					1.86E+02
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	-2.70		4.39E-04	e					4.39E-04
2-sec-Butyl-4,6-dinitrophenol	88-85-7	3.14		2.66E+01	e					2.66E+01
3,3'-Dimethoxybenzidine	119-90-4	1.81		2.17E+00	e					2.17E+00
3,3-Dichlorobenzidine	91-94-1	3.58		6.05E+01	e					6.05E+01
4-Bromophenylphenyl ether	101-55-3	5.00		8.89E+02	e					8.89E+02
Ammonium perfluorooctanoate	3825-26-1	No data		No data			1.00E+01		Default	1.00E+01
Azobenzene	103-33-3	3.82		9.61E+01	e					9.61E+01
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	No data		No data			1.00E+01		Default	1.00E+01
Captan	133-06-2	2.35		6.01E+00	e					6.01E+00
Chlorobenzilate	510-15-6	4.38		2.76E+02	e					2.76E+02
Dibutylphosphate	107-66-4	No data		No data			1.00E+01		Default	1.00E+01
Dimethyl aminoazobenzene	60-11-7	4.58		4.03E+02	e					4.03E+02
Hexachlorobenzene	118-74-1	5.50	2.29E+03		e					2.29E+03
Hexachlorobutadiene	87-68-3	4.73	5.35E+02		e					5.35E+02
Hexachlorocyclopentadiene	77-47-4	5.04	7.45E+02		e					9.59E+02
Hexachloroethane	67-72-1	3.98		1.31E+02	e					1.31E+02
Hexachlorophene	70-30-4	7.54		1.07E+05	e					1.07E+05
Hexamethylene-1,5-diisocyanate	822-06-0	1.27		7.89E-01	e					7.89E-01
Mirex	2385-85-5	6.89		3.14E+04	e					3.14E+04
Nitrofen	1836-75-5	5.53		2.42E+03	e					2.42E+03
Pentachlorobenzene	608-93-5	5.09	1.05E+03		e					1.05E+03
Pentachloronitrobenzene	82-68-8	4.64	4.51E+02		e					4.51E+02
Pentachlorophenol	87-86-5	5.08	1.03E+03		e					1.03E+03

Table C2-4. Terrestrial Soil-to-Invertebrate Transfer Factors (BAF-S) for Ecological Receptors (mg/kg tissue wet weight / mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) BAF-S[b]	BAF-S Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance BAF-S	SAIC Compiled BAF-S	Notes	Comments	Recommended BAF-S[d]
Picric acid	88-89-1	2.03		3.29E+00	e					3.29E+00
Pronamide	23950-58-5	3.51		5.36E+01	e					5.36E+01
Strychnine	57-24-9	1.93		2.72E+00	e					2.72E+00
Terphenyls	26140-60-3	No data		No data			1.00E+01		Default	1.00E+01
Tributyl phosphate	126-73-8	4.00		1.35E+02	e					1.35E+02
Trifluralin	1582-09-8	5.34		1.69E+03	e					1.69E+03
Triphenylamine	603-34-9	No data		No data			1.00E+01		Default	1.00E+01
Herbicides and Organochlorinated Pesticides										
2,4,5-T	93-76-5	3.36		4.03E+01	e					4.03E+01
2,4-D and esters	94-75-7	2.81		1.43E+01	e					1.43E+01
4,4-DDD	72-54-8	6.20		8.55E+03	e		3.30E+00	f		8.55E+03
4,4-DDE	72-55-9	6.26	1.26E+00				1.70E+00	f		1.26E+00
4,4-DDT	50-29-3	6.00		1.26E+00	e		5.70E-01	f		1.26E+00
Aldrin	309-00-2	6.18		8.21E+03	e		5.60E-01	f		8.21E+03
alpha-BHC	319-84-6	3.80		9.24E+01	e		2.60E+00	f		9.24E+01
beta-BHC	319-85-7	3.83		9.85E+01	e		2.60E+00	f		9.85E+01
Chlordane	57-74-9	5.94		5.21E+03	e		1.60E+00	f		5.21E+03
Delta-BHC	319-86-8	4.14		1.76E+02	e		2.60E+00	f		1.76E+02
Dieldrin	60-57-1	5.27		1.48E+03	e		5.50E+00	f		1.48E+03
Endothall	145-73-3	-0.87		1.39E-02	e					1.39E-02
Endrin	72-20-8	4.89		7.25E+02	e		1.90E+00	f		7.25E+02
gamma-BHC (Lindane)	58-89-9	3.72		7.96E+01	e		2.60E+00	f		7.96E+01
Heptachlor	76-44-8	5.02	1.40E+00				1.00E+00	f		1.26E+00
Isodrin	465-73-6	3.55		5.77E+01	e		5.50E+00	f		5.77E+01
Methoxychlor	72-43-5	4.53		3.64E+02	e		5.70E-01	f		3.64E+02
Silvex (2,4,5-TP)	93-72-1	4.07		1.55E+02	e					1.55E+02
Toxaphene	8001-35-2	5.50		2.28E+03	e					2.28E+03
Inorganic Chemicals and Compounds										
Metals										
Aluminum	7429-90-5	NA	2.20E-01		i		7.50E-02			2.20E-01

Table C2-4. Terrestrial Soil-to-Invertebrate Transfer Factors (BAF-S) for Ecological Receptors (mg/kg tissue wet weight / mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) BAF-S[b]	BAF-S Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance BAF-S	SAIC Compiled BAF-S	Notes	Comments	Recommended BAF-S[d]
Antimony	7440-36-0	NA	2.20E-01		i		8.10E-02	j		8.10E-02
Arsenic	7440-38-2	NA	1.10E-01				5.23E-01	k		1.10E-01
Barium	7440-39-3	NA	2.20E-01		i		1.05E-01	j		1.05E-01
Beryllium	7440-41-7	NA	2.20E-01		i		4.50E-02	j		4.50E-02
Bismuth	7440-69-9	NA		2.20E-01	i					2.20E-01
Boron	7440-42-8	NA		2.20E-01	i					2.20E-01
Cadmium	7440-43-9	NA	9.60E-01				4.07E+01	k		9.60E-01
Calcium	7440-70-2	NA		2.20E-01	i		5.42E-01	j		5.42E-01
Chromium (and VI)	18540-29-9	NA	1.00E-02				3.16E+00	k		1.00E-02
Cobalt	7440-48-4	NA		2.20E-01	i		1.20E-01	j		1.20E-01
Copper	7440-50-8	NA	4.00E-02				1.53E+00	k		4.00E-02
Iron	7439-89-6	NA		2.20E-01	i		4.20E-02	j		4.20E-02
Lead	7439-92-1	NA	3.00E-02				See note l			See note l
Lithium	7439-93-2	NA		2.20E-01	i		2.28E-01	j		2.28E-01
Magnesium	7439-95-4	NA		2.20E-01	i		2.40E-01	j		2.40E-01
Manganese	7439-96-5	NA		2.20E-01	i		1.24E-01	k		1.24E-01
Mercury	7439-97-6	NA	4.00E-02				2.06E+01	k		4.00E-02
Molybdenum	7439-98-7	NA		2.20E-01	i					2.20E-01
Nickel	7440-02-0	NA	2.00E-02				4.73E+00	k		2.00E-02
Potassium	7440-09-7	NA		2.20E-01	i					2.20E-01
Rhodium	7440-16-6	NA		2.20E-01	i					2.20E-01
Selenium	7782-49-2	NA	2.20E-01		i		1.37E+01	k		1.37E+01
Silicon	7440-21-3	NA		2.20E-01	i					2.20E-01
Silver	7440-22-4	NA	2.20E-01		i		9.26E-01	j		9.26E-01
Sodium	7440-23-5	NA		2.20E-01	i		4.53E+00	j		4.53E+00
Strontium	7440-24-6	NA		2.20E-01	i					2.20E-01
Tantalum	7440-25-7	NA		2.20E-01	i					2.20E-01
Thallium	7440-28-0	NA	2.20E-01		i		1.70E-02	j		1.70E-02
Tin	7440-31-5	NA		2.20E-01	i					2.20E-01
Tungsten	7440-33-7	NA		2.20E-01	i					2.20E-01

Table C2-4. Terrestrial Soil-to-Invertebrate Transfer Factors (BAF-S) for Ecological Receptors (mg/kg tissue wet weight / mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BAF-S [b]	BAF-S Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BAF-S	SAIC Compiled BAF-S	Notes	Comments	Recommended BAF-S [d]
Uranium	7440-61-1	NA		2.20E-01	i		3.30E-02	j		3.30E-02
Vanadium	7440-62-2	NA		2.20E-01	i		6.60E-02	j		6.60E-02
Yttrium	7440-65-5	NA		2.20E-01	i					2.20E-01
Zinc	7440-66-6	NA	5.60E-01				4.95E+01	k		5.60E-01
Zirconium	7440-67-7	NA		2.20E-01	i					2.20E-01
Non-metals and Anions										
Ammonia/Ammonium	7664-41-7	NA					1.00E+00		Default	1.00E+00
Bromide	24959-67-9	NA		2.20E-01	i					2.20E-01
Chloride	16887-00-6	NA		2.20E-01	i					2.20E-01
Cyanide	57-12-5	NA	1.12E+00							1.12E+00
Fluoride	16984-48-8	NA		2.20E-01	i					2.20E-01
Hydroxide	14280-30-9	NA		NA					Depends on pH	NA
Iodine	7553-56-2	NA		2.20E-01	i					2.20E-01
Nitrate	14797-55-8	NA		2.20E-01	i					2.20E-01
Nitrite	14797-65-0	NA		2.20E-01	i					2.20E-01
Phosphate	14265-44-2	NA		2.20E-01	i					2.20E-01
Phosphorus	7723-14-0	NA		2.20E-01	i					2.20E-01
Sulfate	14808-79-8	NA		2.20E-01	i					2.20E-01
Total Sulfur	63705-05-5	NA		2.20E-01	i					2.20E-01
Priority Pollutants										
Carbon Dioxide	124-38-9	NA		NA						NA
Nitrogen Dioxide	10102-44-0	NA					1.00E+00		Default	1.00E+00
Ozone	10028-15-6	NA					1.00E+00		Default	1.00E+00
Particulate Matter	No CAS #	NA		NA						1.00E+00
Sulfur Dioxode	7446-09-5	NA					1.00E+00		Default	NA
Radionuclides										
Americium-241	1596-10-2	NA		2.20E-01	i		3.12E+00	j		3.12E+00
Antimony-125	14234-35-6	NA	2.20E-01		i		8.10E-02	j		8.10E-02
Barium-137	13981-97-0	NA	2.20E-01		i		1.05E-01	j		1.05E-01
Cadmium-113	None	NA	9.60E-01				3.16E+00	k		3.16E+00

Table C2-4. Terrestrial Soil-to-Invertebrate Transfer Factors (BAF-S) for Ecological Receptors (mg/kg tissue wet weight / mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BAF-S [b]	BAF-S Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BAF-S	SAIC Compiled BAF-S	Notes	Comments	Recommended BAF-S [d]
Cesium-134	13967-70-9	NA		2.20E-01	i					2.20E-01
Cesium-137	10045-97-3	NA		2.20E-01	i					2.20E-01
Europium-154	15585-10-1	NA		2.20E-01	i					2.20E-01
Europium-155	14391-16-3	NA		2.20E-01	i					2.20E-01
Nickel-63	13981-37-8	NA	2.00E-02				4.73E+00	k		2.00E-02
Plutonium-239	15117-48-3	NA		2.20E-01	i		3.92E+00	j		3.92E+00
Plutonium-241	14119-32-5	NA		2.20E-01	i		3.92E+00	j		3.92E+00
Samarium-151	15715-94-3	NA		2.20E-01	i					2.20E-01
Strontium-90	10098-97-2	NA		2.20E-01	i					2.20E-01
Technetium-99	14133-79-7	NA		2.20E-01	i					2.20E-01
Tritium	10028-17-8	NA		2.20E-01	i					2.20E-01
Yttrium-90	10098-91-6	NA		2.20E-01	i					2.20E-01
Uranium-232	14158-29-3	NA		2.20E-01	i		3.30E-02	j		3.30E-02
Uranium-233	13968-55-3	NA		2.20E-01	i		3.30E-02	j		3.30E-02
Uranium-234	13966-29-5	NA		2.20E-01	i		3.30E-02	j		3.30E-02
Uranium-235	15117-96-1	NA		2.20E-01	i		3.30E-02	j		3.30E-02
Uranium-236	13982-70-2	NA		2.20E-01	i		3.30E-02	j		3.30E-02
Uranium-238	7440-61-1	NA		2.20E-01	i		3.30E-02	j		3.30E-02
Actinium-227	14952-40-0	NA		2.20E-01	i					2.20E-01
Americium-243	14993-75-0	NA		2.20E-01	i		3.12E+00	j		3.12E+00
Carbon-14	14762-75-5	NA		2.20E-01	i					2.20E-01
Cobalt-60	10198-40-0	NA		2.20E-01	i		1.20E-01	j		1.20E-01
Curium-242	15510-73-3	NA		2.20E-01	i		1.18E+00	j		1.18E+00
Curium-243	15757-87-6	NA		2.20E-01	i		1.18E+00	j		1.18E+00
Curium-244	13981-15-2	NA		2.20E-01	i		1.18E+00	j		1.18E+00
Europium-152	14683-23-9	NA		2.20E-01	i					2.20E-01
Iodine-129	15046-84-1	NA		2.20E-01	i					2.20E-01
Neptunium-237	13994-20-2	NA		2.20E-01	i					2.20E-01
Nickel-59	14336-70-0	NA	2.00E-02				4.73E+00	k		2.00E-02
Niobium-93	None	NA		2.20E-01	i					2.20E-01

Table C2-4. Terrestrial Soil-to-Invertebrate Transfer Factors (BAF-S) for Ecological Receptors (mg/kg tissue wet weight / mg/kg soil)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BAF-S [b]	BAF-S Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BAF-S	SAIC Compiled BAF-S	Notes	Comments	Recommended BAF-S [d]
Plutonium-238	13981-16-3	NA		2.20E-01	i		9.25E+00	j		9.25E+00
Plutonium-240	14119-33-6	NA		2.20E-01	i		3.92E+00	j		3.92E+00
Plutonium-242	13982-10-0	NA		2.20E-01	i		3.92E+00	j		3.92E+00
Protactinium-231	14331-85-2	NA		2.20E-01	i					2.20E-01
Radium-226	13982-63-3	NA		2.20E-01	i					2.20E-01
Radium-228	15262-20-1	NA		2.20E-01	i					2.20E-01
Ruthenium-106	13967-48-1	NA		2.20E-01	i					2.20E-01
Selenium-79	None	NA	2.20E-01		i		1.37E+01	k		1.37E+01
Thorium-229	15594-54-4	NA		2.20E-01	i					2.20E-01
Thorium-232	7440-29-1	NA		2.20E-01	i					2.20E-01
Tin-126	15832-50-5	NA		2.20E-01	i					2.20E-01
Zirconium-93	15751-77-6	NA		2.20E-01	i					2.20E-01

NA = Not applicable

[a] $\log_{10}$ of K_{ow} values in Table 4.1

[b] Data published in Appendix C of EPA (1999), Table C-1

[c] Calculated or chosen as described in Appendix C, Sect. C-1.1 of EPA (1999)

[d] Selection criteria described in Sect. 8.2.4.3

[e] Calculated by using log K_{ow}: log BAF-S = 0.819 x log K_{ow} -1.146 (Southworth, Beauchamp, and Schmieder 1978)

[f] HAZWRAP (1994), Table 4.4

[g] Calculated from value for TCDD in Appendix C of EPA (1999), Table C-1, by multiplying by BEF

[h] Value for Aroclor 1254 from Appendix C of EPA (1999), Table C-1, was used for PCB mixtures

[i] Average of values for other inorganics as published in Appendix C of EPA (1999), Table C-1

[j] DOE (1996), Appendix G, Table G.39 (mean values)

[k] Sample and others. (1998), Table 11 (90th percentile values)

[l] Calculated by using existing concentrations of lead and calcium: BAF = 0.2 x C_{wormPb} / C_{soilPb}
where log C_{wormPb} = 1.16 + (0.916 x log C_{soilPb}) - (0.326 x log C_{soilCa}) (Corp and Morgan 1991)

Table C2-5. Terrestrial Animal-to-Animal Transfer Factors (Ba) for Ecological Receptors (mg/kg tissue/mg ingested/day)

Constituent of Potential Concern	CAS Registry Number	k	EPA (1999) Ba, Mammals[b]	Ba Calculated by EPA (1999) Methods[c]	Ecology Guidance Preferred Ba	SAIC Compiled Ba	Notes	Comments	Recommended Mammal Ba[d]
Organic Compounds									
Aromatic Halogenated Hydrocarbons									
4-Chloro-3-methylphenol	59-50-7	3.10		3.16E-05		3.16E-05	e		3.16E-05
2,3,4,6-Tetrachlorophenol	58-90-2	4.42		6.61E-04		6.61E-04	e		6.61E-04
Aromatic Nonhalogenated Hydrocarbons									
2-Nitrotoluene	88-72-2	2.30		5.01E-06		5.01E-06	e		5.01E-06
4-Nitrobiphenyl	92-93-3	3.77		1.48E-04		1.48E-04	e		1.48E-04
Benzaldehyde	100-52-7	1.48		7.54E-07		7.54E-07	e		7.54E-07
Benzene	71-43-2	-4.99		2.56E-13		2.56E-13	e		2.56E-13
Benzyl alcohol	100-51-6	1.10		3.16E-07		3.16E-07	e		3.16E-07
Ethyl benzene	100-41-4	3.12		3.34E-05		3.34E-05	e		3.34E-05
m-Xylene	108-38-3	3.20		3.99E-05		3.99E-05	e		3.99E-05
o-Xylene	95-47-6	3.13		3.39E-05		3.39E-05	e		3.39E-05
p-Xylene	106-42-3	3.17		3.72E-05		3.72E-05	e		3.72E-05
Styrene	100-42-5	2.93		2.13E-05		2.13E-05	e		2.13E-05
Toluene	108-88-3	2.67		1.17E-05		1.17E-05	e		1.17E-05
Non-aromatic Nonhalogenated Hydrocarbons									
1,2-Epoxybutane	106-88-7	1.44		6.94E-07		6.94E-07	e		6.94E-07
1,3-Butadiene	106-99-0	1.90		2.00E-06		2.00E-06	e		2.00E-06
1,4-Dioxane	123-91-1	-0.27	1.36E-08			1.36E-08	e		1.36E-08
1-Methylpropyl alcohol	78-92-2	0.61		1.02E-07		1.02E-07	e		1.02E-07
1-Nitropropane	108-03-2	0.87		1.86E-07		1.86E-07	e		1.86E-07
2,2,4-Trimethylpentane	540-84-1	5.02		2.63E-03		2.63E-03	e		2.63E-03
2-Butanone	78-93-3	0.28		4.80E-08		4.80E-08	e		4.80E-08
2-Butenaldehyde (2-Butenal)	4170-30-3	No data		No data		1.00E+01		Default	1.00E+01
2-Ethoxyethanol	110-80-5	-0.10		2.00E-08		2.00E-08	e		2.00E-08
2-Heptanone	110-43-0	1.98		2.40E-06		2.40E-06	e		2.40E-06
2-Hexanone	591-78-6	1.38		6.03E-07		6.03E-07	e		6.03E-07
2-Methoxyethanol	109-86-4	0.25		4.49E-08		4.49E-08	e		4.49E-08

Table C2-5. Terrestrial Animal-to-Animal Transfer Factors (Ba) for Ecological Receptors (mg/kg tissue/mg ingested/day)

Constituent of Potential Concern	CAS Registry Number	k	EPA (1999) Ba, Mammals[b]	Ba Calculated by EPA (1999) Methods[c]	Ecology Guidance Preferred Ba	SAIC Compiled Ba	Notes	Comments	Recommended Mammal Ba[d]
2-Methyl-2-propanol	75-65-0	0.35		5.62E-08		5.62E-08	e		5.62E-08
2-Methyl-2-propenenitrile	126-98-7	0.54		8.72E-08		8.72E-08	e		8.72E-08
2-Methylaziridine	75-55-8	-0.60		6.27E-09		6.27E-09	e		6.27E-09
2-Methylpropyl alcohol	78-83-1	0.76		1.45E-07		1.45E-07	e		1.45E-07
2-Pentanone	107-87-9	0.91		2.04E-07		2.04E-07	e		2.04E-07
2-Propanone (Acetone)	67-64-1	-0.22	1.51E-08			1.51E-08	e		1.51E-08
2-Propene-1-ol	107-18-6	0.17		3.72E-08		3.72E-08	e		3.72E-08
2-Propyl alcohol	67-63-0	0.05		2.82E-08		2.82E-08	e		2.82E-08
3-Heptanone	106-35-4	No data		No data		1.00E+01		Default	1.00E+01
3-Methyl-1-butanol	123-51-3	No data		No data		1.00E+01		Default	1.00E+01
3-Methyl-2-butanone	563-80-4	No data		No data		1.00E+01		Default	1.00E+01
3-Pentanone	96-22-0	0.99		2.45E-07		2.45E-07	e		2.45E-07
4-Heptanone	123-19-3	No data		No data		1.00E+01		Default	1.00E+01
4-Methyl-2-pentanone	108-10-1	1.19		3.89E-07		3.89E-07	e		3.89E-07
4-Methyl-3-penten-2-one	141-79-7	No data		No data		1.00E+01		Default	1.00E+01
5-Methyl-2-hexanone	110-12-3	No data		No data		1.00E+01		Default	1.00E+01
Acetaldehyde	75-07-0	-0.22		1.51E-08		1.51E-08	e		1.51E-08
Acetamide	60-35-5	-1.26		1.38E-09		1.38E-09	e		1.38E-09
Acetic acid	64-19-7	-0.17		1.70E-08		1.70E-08	e		1.70E-08
Acetic acid ethyl ester	141-78-6	0.73		1.35E-07		1.35E-07	e		1.35E-07
Acetic acid n-butyl ester	123-86-4	1.73		1.35E-06		1.35E-06	e		1.35E-06
Acetonitrile	75-05-8	-0.34		1.15E-08		1.15E-08	e		1.15E-08
Acrolein	107-02-8	-0.01		2.46E-08		2.46E-08	e		2.46E-08
Acrylonitrile	107-13-1	0.25	4.46E-08			4.47E-08	e		4.46E-08
Bis(isopropyl)ether	108-20-3	No data		No data		1.00E+01		Default	1.00E+01
Butane	106-97-8	2.89		1.95E-05		1.95E-05	e		1.95E-05
Carbon disulfide	75-15-0	2.00		2.51E-06		2.51E-06	e		2.51E-06
Cyanogen	460-19-5	0.81		1.61E-07		1.61E-07	e		1.61E-07
Cyclohexane	110-82-7	3.44		6.92E-05		6.92E-05	e		6.92E-05

Table C2-5. Terrestrial Animal-to-Animal Transfer Factors (Ba) for Ecological Receptors (mg/kg tissue/mg ingested/day)

Constituent of Potential Concern	CAS Registry Number	k	EPA (1999) Ba, Mammals[b]	Ba Calculated by EPA (1999) Methods[c]	Ecology Guidance Preferred Ba	SAIC Compiled Ba	Notes	Comments	Recommended Mammal Ba[d]
Cyclohexanone	108-94-1	0.81		1.62E-07		1.62E-07	e		1.62E-07
Cyclohexene	110-83-8	2.86		1.82E-05		1.82E-05	e		1.82E-05
Cyclopentane	287-92-3	3.00		2.51E-05		2.51E-05	e		2.51E-05
Ethyl alcohol	64-17-5	0.31		5.13E-08		5.13E-08	e		5.13E-08
Ethyl ether	60-29-7	0.89		1.95E-07		1.95E-07	e		1.95E-07
Ethyl methacrylate	97-63-2	1.59		9.77E-07		9.77E-07	e		9.77E-07
Formaldehyde	50-00-0	0.34	5.54E-08			5.53E-08	e		5.54E-08
Formamide	75-12-7	-1.51		7.76E-10		7.76E-10	e		7.76E-10
Formic acid	64-18-6	-0.54		7.28E-09		7.28E-09	e		7.28E-09
Formic acid, methyl ester	107-31-3	-0.26		1.37E-08		1.37E-08	e		1.37E-08
Glycidylaldehyde	765-34-4	-0.73		4.68E-09		4.68E-09	e		4.68E-09
Methyl acetate	79-20-9	0.18		3.80E-08		3.80E-08	e		3.80E-08
Methyl alcohol	67-56-1	-0.71		4.90E-09		4.90E-09	e		4.90E-09
Methyl isocyanate	624-83-9	No data		No data		NA		Reacts with water	NA
Methyl methacrylate	80-62-6	0.79		1.55E-07		1.55E-07	e		1.55E-07
Methyl tert-butyl ether	1634-04-4	0.94		2.19E-07		2.19E-07	e		2.19E-07
Methylacetylene	74-99-7	0.94		2.19E-07		2.19E-07	e		2.19E-07
Methylcyclohexane	108-87-2	4.10		3.16E-04		3.16E-04	e		3.16E-04
N,N-Dimethylacetamide	127-19-5	No data		No data		1.00E+01		Default	1.00E+01
n-Butyl alcohol	71-36-3	0.88		1.91E-07		1.91E-07	e		1.91E-07
n-Heptane	142-82-5	4.66		1.15E-03		1.15E-03	e		1.15E-03
n-Hexane	110-54-3	4.11		3.24E-04		3.24E-04	e		3.24E-04
Nitromethane	75-52-5	-0.35		1.12E-08		1.12E-08	e		1.12E-08
n-Nonane	111-84-2	5.65		1.12E-02		1.12E-02	e		1.12E-02
n-Octane	111-65-9	4.00		2.51E-04		2.51E-04	e		2.51E-04
n-Pentane	109-66-0	3.21		4.07E-05		4.07E-05	e		4.07E-05
n-Propionaldehyde	123-38-6	0.59		9.77E-08		9.77E-08	e		9.77E-08
n-Propyl alcohol	71-23-8	0.25		4.47E-08		4.47E-08	e		4.47E-08

Table C2-5. Terrestrial Animal-to-Animal Transfer Factors (Ba) for Ecological Receptors (mg/kg tissue/mg ingested/day)

Constituent of Potential Concern	CAS Registry Number	k	EPA (1999) Ba, Mammals[b]	Ba Calculated by EPA (1999) Methods[c]	Ecology Guidance Preferred Ba	SAIC Compiled Ba	Notes	Comments	Recommended Mammal Ba[d]
n-Valeraldehyde	110-62-3	No data		No data		1.00E+01		Default	1.00E+01
Oxirane	75-21-8	-0.30		1.26E-08		1.26E-08	e		1.26E-08
p-Cymene	99-87-6	4.10		3.16E-04		3.16E-04	e		3.16E-04
Phosgene	75-44-5	No data		No data		1.00E+01		Default	1.00E+01
Propargyl alcohol	107-19-7	0.26		4.57E-08		4.57E-08	e		4.57E-08
Propionic acid	79-09-4	0.33		5.37E-08		5.37E-08	e		5.37E-08
Propionitrile	107-12-0	0.16		3.63E-08		3.63E-08	e		3.63E-08
Propylene gylcol monomethyl ether	107-98-2	-0.18		1.66E-08		1.66E-08	e		1.66E-08
p-tert-Butyltoluene	98-51-1	No data		No data		1.00E+01		Default	1.00E+01
Triethylamine	121-44-8	0.16		3.63E-08		3.63E-08	e		3.63E-08
Trimethylamine	75-50-3	0.16		3.63E-08		3.63E-08	e		3.63E-08
Vinyl acetate	108-05-4	0.70		1.26E-07		1.26E-07	e		1.26E-07
Non-aromatic Halogenated Hydrocarbons									
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	No data		No data		1.00E+01		Default	1.00E+01
1,1,1,2-Tetrachloroethane	630-20-6	2.63		1.07E-05		1.07E-05	e		1.07E-05
1,1,1-Trichloroethane	71-55-6	2.42		6.63E-06		6.63E-06	e		1.00E+01
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	3.73		1.35E-04		1.35E-04	e		1.35E-04
1,1,2,2-Tetrachloroethane	79-34-5	4.64		1.11E-03		1.11E-03	e		1.11E-03
1,1,2,2-Tetrachloroethene	127-18-4	2.55		8.82E-06		8.82E-06	e		1.00E+01
1,1,2-Trichloroethane	79-00-5	2.10		3.14E-06		3.14E-06	e		3.14E-06
1,1,2-Trichloroethylene	79-01-6	2.43		6.81E-06		6.81E-06	e		1.00E+01
1,1-Dichloroethane	75-34-3	1.79		1.56E-06		1.56E-06	e		1.56E-06
1,1-Dichloroethene	75-35-4	2.12		3.32E-06		3.32E-06	e		3.32E-06
1,2,2-Trichloro-1,1,2-trifluoroethane	76-13-1	3.16		3.63E-05		3.63E-05	e		3.63E-05
1,2,3-Trichloropropane	96-18-4	2.25		4.47E-06		4.47E-06	e		4.47E-06
1,2-Dibromo-3-chloropropane	96-12-8	2.34		5.50E-06		5.50E-06	e		5.50E-06
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	2.82		1.66E-05		1.66E-05	e		1.66E-05
1,2-Dichloroethane	107-06-2	1.46		7.28E-07		7.28E-07	e		7.28E-07
1,2-Dichloroethylene	540-59-0	0.48		7.59E-08		7.59E-08	e		7.59E-08

Table C2-5. Terrestrial Animal-to-Animal Transfer Factors (Ba) for Ecological Receptors (mg/kg tissue/mg ingested/day)

Constituent of Potential Concern	CAS Registry Number	k	EPA (1999) Ba, Mammals[b]	Ba Calculated by EPA (1999) Methods[c]	Ecology Guidance Preferred Ba	SAIC Compiled Ba	Notes	Comments	Recommended Mammal Ba[d]
1,2-Dichloropropane	78-87-5	2.25		4.47E-06		4.47E-06	e		4.47E-06
1,3-Dichloropropene	542-75-6	1.75		1.41E-06		1.41E-06	e		1.41E-06
1,4-Dichloro-2-butene	764-41-0	0.87		1.87E-07		1.87E-07	e		1.87E-07
1-Chloroethene	75-01-4	1.15		3.52E-07		3.52E-07	e		3.52E-07
2,2-Dichloropropionic acid	75-99-0	0.78		1.51E-07		1.51E-07	e		1.51E-07
2-Chloropropane	75-29-6	1.90		2.00E-06		2.00E-06	e		2.00E-06
3-Chloropropene (allyl chloride)	107-05-1	0.95		2.26E-07		2.26E-07	e		2.26E-07
Bromochloromethane	74-97-5	1.41		6.46E-07		6.46E-07	e		6.46E-07
Bromodichloromethane	75-27-4	2.03		2.66E-06		2.66E-06	e		2.66E-06
Bromoethene	593-60-2	1.07		2.93E-07		2.93E-07	e		2.93E-07
Bromoform	75-25-2	2.35		5.63E-06		5.63E-06	e		5.63E-06
Bromomethane	74-83-9	1.11		3.27E-07		3.27E-07	e		3.27E-07
Carbon tetrachloride	56-23-5	2.72		1.31E-05		1.31E-05	e		1.31E-05
Chlorodibromomethane	124-48-1	2.18		3.77E-06		3.77E-06	e		3.77E-06
Chlorodifluoromethane	75-45-6	1.08		3.01E-07		3.01E-07	e		3.01E-07
Chloroethane	75-00-3	3.10		3.16E-05		3.16E-05	e		3.16E-05
Chloroform	67-66-3	1.95	2.24E-06			2.24E-06	e		2.24E-06
Chloromethane	74-87-3	0.90		2.01E-07		2.01E-07	e		2.01E-07
Chloropentafluoroethane	76-15-3	No data		No data		1.00E+01		Default	1.00E+01
cis-1,2-Dichloroethene	156-59-2	1.98		2.41E-06		2.41E-06	e		2.41E-06
cis-1,3-Dichloropropene	10061-01-5	No data		No data		1.00E+01		Default	1.00E+01
Cyanogen bromide	506-68-3	No data		No data		1.00E+01		Default	1.00E+01
Cyanogen chloride	506-77-4	0.20		3.98E-08		3.98E-08	e		3.98E-08
Dichlorodifluoromethane	75-71-8	2.16		3.62E-06		3.62E-06	e		3.62E-06
Dichlorofluoromethane	75-43-4	No data		No data		1.00E+01		Default	1.00E+01
Dichloromethane	75-09-2	1.26		4.52E-07		4.52E-07	e		4.52E-07
Difluorodibromomethane	75-61-6	No data		No data		1.00E+01		Default	1.00E+01
Hexafluoroacetone	684-16-2	No data		No data		1.00E+01		Default	1.00E+01
Iodomethane	74-88-4	1.69		1.23E-06		1.23E-06	e		1.23E-06

Table C2-5. Terrestrial Animal-to-Animal Transfer Factors (Ba) for Ecological Receptors (mg/kg tissue/mg ingested/day)

Constituent of Potential Concern	CAS Registry Number	k	EPA (1999) Ba, Mammals[b]	Ba Calculated by EPA (1999) Methods[c]	Ecology Guidance Preferred Ba	SAIC Compiled Ba	Notes	Comments	Recommended Mammal Ba[d]
Methylene bromide	74-95-3	1.62		1.05E-06		1.05E-06	e		1.05E-06
Pentachloroethane	76-01-7	3.05		2.82E-05		2.82E-05	e		2.82E-05
trans-1,2-Dichloroethene	156-60-5	1.98		2.41E-06		2.41E-06	e		2.41E-06
trans-1,3-Dichloropropene	10061-02-6	2.06		2.88E-06		2.88E-06	e		2.88E-06
Trichloroacetic acid	76-03-9	1.33		5.37E-07		5.37E-07	e		5.37E-07
Trichlorofluoroethane	27154-33-2	No data		No data		1.00E+01		Default	1.00E+01
Trichlorofluoromethane	75-69-4	2.53		8.54E-06		8.54E-06	e		8.54E-06
Trifluorobromomethane	75-63-8	1.86		1.82E-06		1.82E-06	e		1.82E-06
Dioxin and Furan Compounds									
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	35822-46-9	8.20	2.77E-03			3.97E+00	e		2.77E-03
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	7.92	5.99E-04			2.09E+00	e		5.99E-04
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	7.92	2.12E-02			2.09E+00	e		2.12E-02
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	39227-28-6	7.79	1.69E-02			1.55E+00	e		1.69E-02
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	7.25	4.12E-03			4.47E-01	e		4.12E-03
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	57653-85-7	7.25	6.52E-03			4.47E-01	e		6.52E-03
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	7.25	1.04E-02			4.47E-01	e		1.04E-02
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	19408-74-3	7.25	7.60E-03			4.47E-01	e		7.60E-03
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	7.25	3.42E-03			4.47E-01	e		3.42E-02
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	40321-76-4	6.64	4.98E-02			1.10E-01	e		4.98E-02
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	6.79	1.19E-02			1.55E-01	e		1.19E-02
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	7.25	3.64E-02			4.47E-01	e		3.64E-02
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	6.92	8.69E-02			2.09E-01	e		8.69E-02
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1746-01-6	6.64	5.43E-02			1.10E-01	e		5.43E-02
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	6.53	4.34E-02			8.52E-02	e		4.34E-02
Dibenzofuran	132-64-9	4.33		5.37E-04		5.37E-04	e		5.37E-04
Octachlorodibenzo(p)dioxin	3268-87-9	8.20	6.52E-04			3.98E+00	e		6.52E-04
Octachlorodibenzofuran	39001-02-0	8.78	8.69E-04			1.52E+01	e		8.70E-04
PCBs									
Polychlorinated biphenyls (PCBs)[f]	1336-36-3	7.31	4.04E-02			5.10E-01	e		4.04E-02

Table C2-5. Terrestrial Animal-to-Animal Transfer Factors (Ba) for Ecological Receptors (mg/kg tissue/mg ingested/day)

Constituent of Potential Concern	CAS Registry Number	k	EPA (1999) Ba, Mammals[b]	Ba Calculated by EPA (1999) Methods[c]	Ecology Guidance Preferred Ba	SAIC Compiled Ba	Notes	Comments	Recommended Mammal Ba[d]
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	7.08		3.02E-01		3.02E-01	e		3.02E-01
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	7.12		3.31E-01		3.31E-01	e		3.31E-01
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	No data		No data		1.00E+01		Default	1.00E+01
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	No data		No data		1.00E+01		Default	1.00E+01
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	No data		No data		1.00E+01		Default	1.00E+01
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	No data		No data		1.00E+01		Default	1.00E+01
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	No data		No data		1.00E+01		Default	1.00E+01
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	No data		No data		1.00E+01		Default	1.00E+01
2',3,4,4',5-Pentachlorobiphenyl	no cas #	No data		No data		1.00E+01		Default	1.00E+01
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	7.12		3.31E-01		3.31E-01	e		3.31E-01
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	7.41		6.43E-01		6.43E-01	e		6.43E-01
3,3',4,4',5-Pentachlorobiphenyl	no cas #	No data		No data		1.00E+01		Default	1.00E+01
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	No data		No data		1.00E+01		Default	1.00E+01
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	No data		No data		1.00E+01		Default	1.00E+01
Phthalates									
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	5.20	4.04E-03			4.02E-03	e		4.04E-03
Butylbenzyl phthalate	85-68-7	4.41		6.51E-04		6.51E-04	e		6.51E-04
Dibutyl phthalate	84-74-2	4.72		1.32E-03		1.32E-03	e		1.32E-03
Diethyl phthalate	84-66-2	4.44		6.86E-04		6.86E-04	e		6.86E-04
Dimethylphthalate	131-11-3	1.63		1.08E-06		1.08E-06	e		1.08E-06
n-Dioctyl phthalate	117-84-0	9.33	5.36E+01			5.38E+01	e		5.36E+01
Light Polycyclic Aromatic Hydrocarbons (molecular weight <200 g/mole)									
2-Chloronapthalene	91-58-7	4.07		2.94E-04		2.94E-04	e		2.94E-04
2-Methyl naphthalene	91-57-6	3.86		1.82E-04		1.82E-04	e		1.82E-04
5-Nitroacenaphthene	602-87-9	No data		No data		1.00E+01		Default	1.00E+01
Acenaphthene	83-32-9	3.96		2.32E-04		2.32E-04	e		2.32E-04
Acenaphthylene	208-96-8	4.07		2.95E-04		2.95E-04	e		2.95E-04
Anthracene	120-12-7	4.47		7.41E-04		7.41E-04	e		7.41E-04
Fluorene	86-73-7	4.18		3.80E-04		3.80E-04	e		3.80E-04

Table C2-5. Terrestrial Animal-to-Animal Transfer Factors (Ba) for Ecological Receptors (mg/kg tissue/mg ingested/day)

Constituent of Potential Concern	CAS Registry Number	k	EPA (1999) Ba, Mammals[b]	Ba Calculated by EPA (1999) Methods[c]	Ecology Guidance Preferred Ba	SAIC Compiled Ba	Notes	Comments	Recommended Mammal Ba[d]
Indene	95-13-6	No data		No data		1.00E+01		Default	1.00E+01
Naphthalene	91-20-3	3.37		5.93E-05		5.93E-05	e		5.93E-05
Phenanthrene	85-01-8	4.55		8.92E-04		8.92E-04	e		8.92E-04
Pyrene	129-00-0	5.00		2.51E-03		2.51E-03	e		2.51E-03
Heavy Polycyclic Aromatic Hydrocarbons (molecular weight >200 g/mole)									
3-Methylcholanthrene	56-49-5	7.11		3.24E-01		3.24E-01	e		3.24E-01
5-Methylchrysene	3697-24-3	No data		No data		1.00E+01		Default	1.00E+01
Benzo(a)anthracene	56-55-3	5.68	1.20E-02			1.20E-02	e		1.20E-02
Benzo(a)pyrene	50-32-8	6.13	3.38E-02			3.39E-02	e		3.38E-02
Benzo(b)fluoranthene	205-99-2	6.20	4.01E-02			3.99E-02	e		4.01E-02
Benzo(e)pyrene	192-97-2	7.40		6.31E-01		6.31E-01	e		6.31E-01
Benzo(g,h,i)perylene	191-24-2	7.10		3.16E-01		3.16E-01	e		3.16E-01
Benzo(j)fluoranthene	205-82-3	6.44		6.92E-02		6.92E-02	e		6.92E-02
Benzo(k)fluoranthene	207-08-9	6.20	3.97E-02			4.00E-02	e		3.97E-02
Benzo[a,i]pyrene	191-30-0	No data		No data		1.00E+01		Default	1.00E+01
Chrysene	218-01-9	5.74	1.38E-02			1.38E-02	e		1.38E-02
Dibenz(a,h)anthracene	53-70-3	6.55	8.88E-02			8.87E-02	e		8.88E-02
Dibenz[a,h]acridine	226-36-8	No data		No data		1.00E+01		Default	1.00E+01
Dibenz[a,j]acridine	224-42-0	No data		No data		1.00E+01		Default	1.00E+01
Dibenzo(a,e)fluoranthene	5385-75-1	No data		No data		1.00E+01		Default	1.00E+01
Dibenzo(a,h)fluoranthene	no cas #	No data		No data		1.00E+01		Default	1.00E+01
Dibenzo[a,e]pyrene	192-65-4	No data		No data		1.00E+01		Default	1.00E+01
Dibenzo[a,h]pyrene	189-64-0	No data		No data		1.00E+01		Default	1.00E+01
Dibenzo[a,i]pyrene	189-55-9	7.29		4.90E-01		4.90E-01	e		4.90E-01
Fluoranthene	206-44-0	5.08		3.04E-03		3.04E-03	e		3.04E-03
Hexachloronaphthalene	1335-87-1	7.59		9.77E-01		9.77E-01	e		9.77E-01
Indeno(1,2,3-cd)pyrene	193-39-5	6.91	2.07E-01			2.06E-01	e		2.07E-01
Octachloronaphthalene	2234-13-1	6.42		6.61E-02		6.61E-02	e		6.61E-02
Pentachloronaphthalene	1321-64-8	No data		No data		1.00E+01		Default	1.00E+01

Table C2-5. Terrestrial Animal-to-Animal Transfer Factors (Ba) for Ecological Receptors (mg/kg tissue/mg ingested/day)

Constituent of Potential Concern	CAS Registry Number	k	EPA (1999) Ba, Mammals[b]	Ba Calculated by EPA (1999) Methods[c]	Ecology Guidance Preferred Ba	SAIC Compiled Ba	Notes	Comments	Recommended Mammal Ba[d]
Tetrachloronaphthalene	1335-88-2	No data		No data		1.00E+01		Default	1.00E+01
Trichloronaphthalene	1321-65-9	No data		No data		1.00E+01		Default	1.00E+01
Light Substituted Benzene Compounds (MW <200 g/mole)									
1,2,3-Trichlorobenzene	87-61-6	4.05		2.79E-04		2.79E-04	e		2.79E-04
1,2,4-Trichlorobenzene	120-82-1	3.99		2.44E-04		2.44E-04	e		2.44E-04
1,2,4-Trimethyl benzene	95-63-6	3.65		1.12E-04		1.12E-04	e		1.12E-04
1,2-Dichlorobenzene	95-50-1	3.45		7.01E-05		7.01E-05	e		7.01E-05
1,3,5-Trimethyl benzene	108-67-8	3.42		6.61E-05		6.61E-05	e		6.61E-05
1,3-Dichlorobenzene	541-73-1	3.53		8.52E-05		8.52E-05	e		8.52E-05
1,3-Dinitrobenzene	99-65-0	1.49	7.79E-07			7.79E-07	e		7.79E-07
1,4-Dichlorobenzene	106-46-7	3.41		6.48E-05		6.48E-05	e		6.48E-05
1,4-Dinitrobenzene	100-25-4	1.50		7.94E-07		7.94E-07	e		7.94E-07
2,4,5-Trichlorophenol	95-95-4	3.87		1.86E-04		1.86E-04	e		1.86E-04
2,4,6-Trichlorophenol	88-06-2	3.71		1.29E-04		1.29E-04	e		1.29E-04
2,4-Dichlorophenol	120-83-2	3.04		2.74E-05		2.74E-05	e		2.74E-05
2,4-Dimethylphenol	105-67-9	2.36		5.75E-06		5.75E-06	e		5.75E-06
2,4-Dinitrophenol	51-28-5	1.52		8.29E-07		8.29E-07	e		8.29E-07
2,4-Dinitrotoluene	121-14-2	2.00	2.49E-06			2.49E-06	e		2.49E-06
2,6-Dinitrotoluene	606-20-2	1.89	1.93E-06			1.93E-06	e		1.93E-06
2-Chlorophenol	95-57-8	2.16		3.64E-06		3.64E-06	e		3.64E-06
2-Chlorotoluene	95-49-8	3.54		8.64E-05		8.64E-05	e		8.64E-05
2-Nitrophenol	88-75-5	1.79		1.55E-06		1.55E-06	e		1.55E-06
4,6-Dinitro-o-cresol	534-52-1	2.85		1.78E-05		1.78E-05	e		1.78E-05
4-Chlorotoluene	106-43-4	3.33		5.37E-05		5.37E-05	e		5.37E-05
4-Nitrophenol	100-02-7	1.91		2.04E-06		2.04E-06	e		2.04E-06
alpha-Methylstyrene	98-83-9	3.46		7.30E-05		7.30E-05	e		7.30E-05
Aniline	62-53-3	0.98		2.40E-07		2.40E-07	e		2.40E-07
Benzotrichloride	98-07-7	2.92		2.09E-05		2.09E-05	e		2.09E-05
Benzyl chloride	100-44-7	0.36		5.78E-08		5.78E-08	e		5.78E-08

Table C2-5. Terrestrial Animal-to-Animal Transfer Factors (Ba) for Ecological Receptors (mg/kg tissue/mg ingested/day)

Constituent of Potential Concern	CAS Registry Number	k	EPA (1999) Ba, Mammals[b]	Ba Calculated by EPA (1999) Methods[c]	Ecology Guidance Preferred Ba	SAIC Compiled Ba	Notes	Comments	Recommended Mammal Ba[d]
Bromobenzene	108-86-1	2.99		2.45E-05		2.45E-05	e		2.45E-05
Chlorobenzene	108-90-7	2.79		1.55E-05		1.55E-05	e		1.55E-05
Cumene	98-82-8	3.61		1.03E-04		1.03E-04	e		1.03E-04
m-Cresol	108-39-4	1.96		2.29E-06		2.29E-06	e		2.29E-06
n-Butyl benzene	104-51-8	4.28		4.79E-04		4.79E-04	e		4.79E-04
Nitrobenzene	98-95-3	1.83	1.71E-06			1.71E-06	e		1.71E-06
n-Propyl benzene	103-65-1	3.69		1.23E-04		1.23E-04	e		1.23E-04
o-Cresol	95-48-7	2.02		2.64E-06		2.64E-06	e		2.64E-06
o-Dinitrobenzene	528-29-0	1.60		1.00E-06		1.00E-06	e		1.00E-06
o-Nitroaniline	88-74-4	1.85		1.78E-06		1.78E-06	e		1.78E-06
o-Toluidine	95-53-4	1.34		5.50E-07		5.50E-07	e		5.50E-07
p-Chloroaniline	106-47-8	1.87		1.86E-06		1.86E-06	e		1.86E-06
p-Cresol	106-44-5	1.94		2.19E-06		2.19E-06	e		2.19E-06
Phenol	108-95-2	1.48		7.54E-07		7.54E-07	e		7.54E-07
p-Nitrochlorobenzene	100-00-5	2.39		6.17E-06		6.17E-06	e		6.17E-06
p-Toluidine	106-49-0	1.40		6.31E-07		6.31E-07	e		6.31E-07
sec-Butyl benzene	135-98-8	4.57		9.33E-04		9.33E-04	e		9.33E-04
tert-Butyl benzene	98-06-6	4.11		3.24E-04		3.24E-04	e		3.24E-04
Toluene-2,6-diamine	823-40-5	1.45		7.16E-07		7.16E-07	e		7.16E-07
Trimethyl benzene	25551-13-7	3.40		6.31E-05		6.31E-05	e		6.31E-05
Other Light Semivolatile Compounds (molecular weight <200 g/mole)									
1,1'-Biphenyl	92-52-4	3.90		2.00E-04		2.00E-04	e		2.00E-04
1,1-Dimethylhydrazine	57-14-7	0.60		9.89E-08		9.89E-08	e		9.89E-08
1,2-Dimethylhydrazine	540-73-8	-1.37		1.08E-09		1.08E-09	e		1.08E-09
1,2-Diphenylhydrazine	122-66-7	2.94		2.19E-05		2.19E-05	e		2.19E-05
1,3-Propane sultone	1120-71-4	-0.52		7.53E-09		7.53E-09	e		7.53E-09
2,4-Toluene diisocyante	584-84-9	No data		No data		NA		Decomposes rapidly in water	NA

Table C2-5. Terrestrial Animal-to-Animal Transfer Factors (Ba) for Ecological Receptors (mg/kg tissue/mg ingested/day)

Constituent of Potential Concern	CAS Registry Number	k	EPA (1999) Ba, Mammals[b]	Ba Calculated by EPA (1999) Methods[c]	Ecology Guidance Preferred Ba	SAIC Compiled Ba	Notes	Comments	Recommended Mammal Ba[d]
2-Chloroacetophenone	532-27-4	2.59		9.88E-06		9.88E-06	e		9.88E-06
2-Propenoic acid	79-10-7	0.33		5.37E-08		5.37E-08	e		5.37E-08
4,4-Methylenedianiline	101-77-9	3.38		6.02E-05		6.02E-05	e		6.02E-05
Acetophenone	98-86-2	1.64		1.10E-06		1.10E-06	e		1.10E-06
Benzoic acid	65-85-0	1.86		1.82E-06		1.82E-06	e		1.82E-06
bis(2-Chloroethoxy)methane	111-91-1	7.59		9.77E-01		9.77E-01	e		9.77E-01
bis(2-Chloroethyl) ether	111-44-4	1.30		5.02E-07		5.02E-07	e		5.02E-07
Chlorocyclopentadiene	41851-50-7	2.43		6.76E-06		6.76E-06	e		6.76E-06
Cyclohexanol	108-93-0	1.23		4.27E-07		4.27E-07	e		4.27E-07
Dichloroisopropyl ether	108-60-1	2.58		9.55E-06		9.55E-06	e		9.55E-06
Dichloromethyl ether	542-88-1	-0.38		1.05E-08		1.05E-08	e		1.05E-08
Dichloropentadiene	no cas #	No data		No data		1.00E+01		Default	1.00E+01
Dimethyl sulfate	77-78-1	0.32		5.23E-08		5.23E-08	e		5.23E-08
Dimethylaniline	121-69-7	-0.88		3.31E-09		3.31E-09	e		3.31E-09
Di-n-propylnitrosamine	621-64-7	1.38		6.03E-07		6.03E-07	e		6.03E-07
Diphenyl ether	101-84-8	4.21		4.07E-04		4.07E-04	e		4.07E-04
Epichlorohydrin	106-89-8	0.25		4.47E-08		4.47E-08	e		4.47E-08
Ethyl carbamate (urethane)	51-79-6	-0.15		1.78E-08		1.78E-08	e		1.78E-08
Ethyl methanesulfonate	62-50-0	0.05		2.81E-08		2.81E-08	e		2.81E-08
Ethylene dibromide	106-93-4	1.75		1.41E-06		1.41E-06	e		1.41E-06
Ethylene glycol	107-21-1	-0.91		3.06E-09		3.06E-09	e		3.06E-09
Ethylene glycol monobutyl ether	111-76-2	1.55		9.00E-07		9.00E-07	e		9.00E-07
Ethylene glycol monoethyl ether acetate	111-15-9	0.62		1.05E-07		1.05E-07	e		1.05E-07
Ethylene thiourea	96-45-7	-0.64		5.73E-09		5.73E-09	e		5.73E-09
Furfural	98-01-1	0.96		2.29E-07		2.29E-07	e		2.29E-07
Maleic hydrazide	123-33-1	-0.74		4.57E-09		4.57E-09	e		4.57E-09
Malononitrile	109-77-3	0.04		2.75E-08		2.75E-08	e		2.75E-08
Methyl styrene (mixed isomers)	25013-15-4	3.35		5.62E-05		5.62E-05	e		5.62E-05
Methylhydrazine	60-34-4	-1.06		2.19E-09		2.19E-09	e		2.19E-09

Table C2-5. Terrestrial Animal-to-Animal Transfer Factors (Ba) for Ecological Receptors (mg/kg tissue/mg ingested/day)

Constituent of Potential Concern	CAS Registry Number	k	EPA (1999) Ba, Mammals[b]	Ba Calculated by EPA (1999) Methods[c]	Ecology Guidance Preferred Ba	SAIC Compiled Ba	Notes	Comments	Recommended Mammal Ba[d]
N,N-Diphenylamine	122-39-4	3.50		7.94E-05		7.94E-05	e		7.94E-05
Nitric acid, propyl ester	627-13-4	No data		No data		1.00E+01		Default	1.00E+01
N-Nitrosodi-n-butylamine	924-16-3	2.41		6.46E-06		6.46E-06	e		6.46E-06
N-Nitrosomorpholine	59-89-2	0.98		2.40E-07		2.40E-07	e		2.40E-07
N-Nitroso-N,N-dimethylamine	62-75-9	-0.47		8.51E-09		8.51E-09	e		8.51E-09
o-Anisidine	90-04-0	1.18		3.80E-07		3.80E-07	e		3.80E-07
Oxalic acid	144-62-7	No data		No data		1.00E+01		Default	1.00E+01
Phthalic anhydride	85-44-9	-0.60		6.28E-09		6.28E-09	e		6.28E-09
p-Phthalic acid	100-21-0	0.82		1.67E-07		1.67E-07	e		1.67E-07
Pyridine	110-86-1	0.67		1.18E-07		1.18E-07	e		1.18E-07
Quinoline	91-22-5	2.03		2.69E-06		2.69E-06	e		2.69E-06
Quinone	106-51-4	0.20		3.98E-08		3.98E-08	e		3.98E-08
Safrole	94-59-7	2.66		1.15E-05		1.15E-05	e		1.15E-05
Tetrahydrofuran	109-99-9	0.45		7.03E-08		7.03E-08	e		7.03E-08
Other Heavy Semivolatile Compounds (molecular weight >200 g/mole)									
1,2,4,5-Tetrachlorobenzene	95-94-3	4.64		1.10E-03		1.10E-03	e		1.10E-03
1,3,5-Trinitrobenzene	99-35-4	1.18		3.79E-07		3.79E-07	e		3.79E-07
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	4.17		3.72E-04		3.72E-04	e		3.72E-04
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	-2.70		5.01E-11		5.01E-11	e		5.01E-11
2-sec-Butyl-4,6-dinitrophenol	88-85-7	3.14		3.47E-05		3.47E-05	e		3.47E-05
3,3'-Dimethoxybenzidine	119-90-4	1.81		1.62E-06		1.62E-06	e		1.62E-06
3,3-Dichlorobenzidine	91-94-1	3.58		9.44E-05		9.44E-05	e		9.44E-05
4-Bromophenylphenyl ether	101-55-3	5.00		2.51E-03		2.51E-03	e		2.51E-03
Ammonium perfluorooctanoate	3825-26-1	No data		No data		1.00E+01		Default	1.00E+01
Azobenzene	103-33-3	3.82		1.66E-04		1.66E-04	e		1.66E-04
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	No data		No data		1.00E+01		Default	1.00E+01
Captan	133-06-2	2.35		5.62E-06		5.62E-06	e		5.62E-06
Chlorobenzilate	510-15-6	4.38		6.03E-04		6.03E-04	e		6.03E-04
Dibutylphosphate	107-66-4	No data		No data		1.00E+01		Default	1.00E+01

Table C2-5. Terrestrial Animal-to-Animal Transfer Factors (Ba) for Ecological Receptors (mg/kg tissue/mg ingested/day)

Constituent of Potential Concern	CAS Registry Number	k	EPA (1999) Ba, Mammals[b]	Ba Calculated by EPA (1999) Methods[c]	Ecology Guidance Preferred Ba	SAIC Compiled Ba	Notes	Comments	Recommended Mammal Ba[d]
Dimethyl aminoazobenzene	60-11-7	4.58		9.55E-04		9.55E-04	e		9.55E-04
Hexachlorobenzene	118-74-1	5.50	7.98E-03			7.99E-03	e		7.98E-03
Hexachlorobutadiene	87-68-3	4.73	1.35E-03			1.35E-03	e		1.35E-03
Hexachlorocyclopentadiene	77-47-4	5.04	2.03E-03			2.75E-03	e		2.03E-03
Hexachloroethane	67-72-1	3.98		2.43E-04		2.43E-04	e		2.43E-04
Hexachlorophene	70-30-4	7.54	8.73E-01			8.72E-01	e		8.73E-01
Hexamethylene-1,5-diisocyanate	822-06-0	1.27		4.71E-07		4.71E-07	e		4.71E-07
Mirex	2385-85-5	6.89		1.95E-01		1.95E-01	e		1.95E-01
Nitrofen	1836-75-5	5.53		8.51E-03		8.51E-03	e		8.51E-03
Pentachlorobenzene	608-93-5	5.09	3.07E-03			3.06E-03	e		3.07E-03
Pentachloronitrobenzene	82-68-8	4.64	1.10E-03			1.10E-03	e		1.10E-03
Pentachlorophenol	87-86-5	5.08	3.02E-03			3.01E-03	e		3.02E-03
Picric acid	88-89-1	2.03		2.69E-06		2.69E-06	e		2.69E-06
Pronamide	23950-58-5	3.51		8.14E-05		8.14E-05	e		8.14E-05
Strychnine	57-24-9	1.93		2.14E-06		2.14E-06	e		2.14E-06
Terphenyls	26140-60-3	No data		No data		1.00E+01		Default	1.00E+01
Tributyl phosphate	126-73-8	4.00		2.51E-04		2.51E-04	e		2.51E-04
Trifluralin	1582-09-8	5.34		5.50E-03		5.50E-03	e		5.50E-03
Triphenylamine	603-34-9	No data		No data		1.00E+01		Default	1.00E+01
Herbicides and Organochlorinated Pesticides									
2,4,5-T	93-76-5	3.36		5.75E-05		5.75E-05	e		5.75E-05
2,4-D and esters	94-75-7	2.81		1.62E-05		1.62E-05	e		1.62E-05
4,4-DDD	72-54-8	6.20		3.98E-02		3.98E-02	e		3.98E-02
4,4-DDE	72-55-9	6.26	4.53E-02			4.52E-02	e		4.53E-02
4,4-DDT	50-29-3	6.00		2.51E-02		2.51E-02	e		2.51E-02
Aldrin	309-00-2	6.18		3.79E-02		3.79E-02	e		3.79E-02
alpha-BHC	319-84-6	3.80		1.58E-04		1.58E-04	e		1.58E-04
beta-BHC	319-85-7	3.83		1.71E-04		1.71E-04	e		1.71E-04
Chlordane	57-74-9	5.94		2.18E-02		2.18E-02	e		2.18E-02

Table C2-5. Terrestrial Animal-to-Animal Transfer Factors (Ba) for Ecological Receptors (mg/kg tissue/mg ingested/day)

Constituent of Potential Concern	CAS Registry Number	k	EPA (1999) Ba, Mammals[b]	Ba Calculated by EPA (1999) Methods[c]	Ecology Guidance Preferred Ba	SAIC Compiled Ba	Notes	Comments	Recommended Mammal Ba[d]
Delta-BHC	319-86-8	4.14		3.47E-04		3.47E-04	e		3.47E-04
Dieldrin	60-57-1	5.27		4.67E-03		4.67E-03	e		4.67E-03
Endothall	145-73-3	-0.87		3.39E-09		3.39E-09	e		3.39E-09
Endrin	72-20-8	4.89		1.96E-03		1.96E-03	e		1.96E-03
gamma-BHC (Lindane)	58-89-9	3.72		1.32E-04		1.32E-04	e		1.32E-04
Heptachlor	76-44-8	5.02	2.60E-03			2.61E-03	e		2.60E-03
Isodrin	465-73-6	3.55		8.91E-05		8.91E-05	e		8.91E-05
Methoxychlor	72-43-5	4.53		8.44E-04		8.44E-04	e		8.44E-04
Silvex (2,4,5-TP)	93-72-1	4.07		2.97E-04		2.97E-04	e		2.97E-04
Toxaphene	8001-35-2	5.50		7.94E-03		7.94E-03	e		7.94E-03
Inorganic Chemicals and Compounds									
Metals									
Aluminum	7429-90-5	NA		1.50E-03		1.50E-03	g		1.50E-03
Antimony	7440-36-0	NA	1.00E-03			1.00E-03	g		1.00E-03
Arsenic	7440-38-2	NA	2.00E-03			2.00E-03	g		2.00E-03
Barium	7440-39-3	NA	1.50E-04			1.50E-04	g		1.50E-04
Beryllium	7440-41-7	NA	1.00E-03			1.00E-03	g		1.00E-03
Bismuth	7440-69-9	NA		4.00E-04		4.00E-04	g		4.00E-04
Boron	7440-42-8	NA		8.00E-04		8.00E-04	g		8.00E-04
Cadmium	7440-43-9	NA	1.20E-04			5.50E-04	g		1.20E-04
Calcium	7440-70-2	NA		7.00E-04		7.00E-04	g		7.00E-04
Chromium (and VI)	18540-29-9	NA	5.51E-03			5.50E-03	g		5.51E-03
Cobalt	7440-48-4	NA		2.00E-02		2.00E-02	g		2.00E-02
Copper	7440-50-8	NA		1.00E-02		1.00E-02	g		1.00E-02
Iron	7439-89-6	NA		2.00E-02		2.00E-02	g		2.00E-02
Lead	7439-92-1	NA	3.00E-04			3.00E-04	g		3.00E-04
Lithium	7439-93-2	NA		1.00E-02		1.00E-02	g		1.00E-02
Magnesium	7439-95-4	NA		5.00E-03		5.00E-03	g		5.00E-03
Manganese	7439-96-5	NA		4.00E-04		4.00E-04	g		4.00E-04

Table C2-5. Terrestrial Animal-to-Animal Transfer Factors (Ba) for Ecological Receptors (mg/kg tissue/mg ingested/day)

Constituent of Potential Concern	CAS Registry Number	k	EPA (1999) Ba, Mammals[b]	Ba Calculated by EPA (1999) Methods[c]	Ecology Guidance Preferred Ba	SAIC Compiled Ba	Notes	Comments	Recommended Mammal Ba[d]
Mercury	7439-97-6	NA	5.21E-03			2.50E-01	g		5.21E-03
Molybdenum	7439-98-7	NA		6.00E-03		6.00E-03	g		6.00E-03
Nickel	7440-02-0	NA	5.99E-03			6.00E-03	g		5.99E-03
Potassium	7440-09-7	NA		2.00E-02		2.00E-02	g		2.00E-02
Rhodium	7440-16-6	NA		2.00E-03		2.00E-03	g		2.00E-03
Selenium	7782-49-2	NA	2.27E-03			1.50E-02	g		2.27E-03
Silicon	7440-21-3	NA		4.00E-05		4.00E-05	g		4.00E-05
Silver	7440-22-4	NA	3.00E-03			3.00E-03	g		3.00E-03
Sodium	7440-23-5	NA		5.50E-02		5.50E-02	g		5.50E-02
Strontium	7440-24-6	NA		3.00E-04		3.00E-04	g		3.00E-04
Tantalum	7440-25-7	NA		6.00E-04		6.00E-04	g		6.00E-04
Thallium	7440-28-0	NA	4.01E-02			4.00E-02	g		4.01E-02
Tin	7440-31-5	NA		8.00E-02		8.00E-02	g		8.00E-02
Tungsten	7440-33-7	NA		4.50E-02		4.50E-02	g		4.50E-02
Uranium	7440-61-1	NA		2.00E-04		2.00E-04	g		2.00E-04
Vanadium	7440-62-2	NA		2.50E-03		2.50E-03	g		2.50E-03
Yttrium	7440-65-5	NA		3.00E-04		3.00E-04	g		3.00E-04
Zinc	7440-66-6	NA	8.99E-05			1.00E-01	g		8.99E-05
Zirconium	7440-67-7	NA		5.50E-03		5.50E-03	g		5.50E-03
Non-metals and Anions									
Ammonia/Ammonium	7664-41-7	NA				1.00E+00			1.00E+00
Bromide	24959-67-9	NA		2.50E-02		2.50E-02	g		2.50E-02
Chloride	16887-00-6	NA		8.00E-02		8.00E-02	g		8.00E-02
Cyanide	57-12-5	NA				1.00E+00			1.00E+00
Fluoride	16984-48-8	NA		1.50E-01		1.50E-01	g		1.50E-01
Hydroxide	14280-30-9	NA		NA				Depends on pH	NA
Iodine	7553-56-2	NA		7.00E-03		7.00E-03	g		7.00E-03
Nitrate	14797-55-8	NA		7.50E-02		7.50E-02	g		7.50E-02

Table C2-5. Terrestrial Animal-to-Animal Transfer Factors (Ba) for Ecological Receptors (mg/kg tissue/mg ingested/day)

Constituent of Potential Concern	CAS Registry Number	k	EPA (1999) Ba, Mammals[b]	Ba Calculated by EPA (1999) Methods[c]	Ecology Guidance Preferred Ba	SAIC Compiled Ba	Notes	Comments	Recommended Mammal Ba[d]
Nitrite	14797-65-0	NA		7.50E-02		7.50E-02	g		7.50E-02
Phosphate	14265-44-2	NA		5.50E-02		5.50E-02	g		5.50E-02
Phosphorus	7723-14-0	NA		5.50E-02		5.50E-02	g		5.50E-02
Sulfate	14808-79-8	NA		1.00E-01		1.00E-01	g		1.00E-01
Total Sulfur	63705-05-5	NA		1.00E-01		1.00E-01	g		1.00E-01
Priority Pollutants									
Carbon Dioxide	124-38-9	NA				1.00E+00			1.00E+00
Nitrogen Dioxide	10102-44-0	NA				1.00E+00			1.00E+00
Ozone	10028-15-6	NA				1.00E+00			1.00E+00
Particulate Matter	No CAS #	NA		NA		NA			NA
Sulfur Dioxode	7446-09-5	NA				1.00E+00			1.00E+00
Radionuclides									
Americium-241	1596-10-2	NA		3.50E-06		3.50E-06	g		3.50E-06
Antimony-125	14234-35-6	NA		1.00E-03		1.00E-03	g		1.00E-03
Barium-137	13981-97-0	NA		1.50E-04		1.50E-04	g		1.50E-04
Cadmium-113	None	NA	1.20E-04			5.50E-04	g		1.20E-04
Cesium-134	13967-70-9	NA		2.00E-02		2.00E-02	g		2.00E-02
Cesium-137	10045-97-3	NA		2.00E-02		2.00E-02	g		2.00E-02
Europium-154	15585-10-1	NA		5.00E-03		5.00E-03	g		5.00E-03
Europium-155	14391-16-3	NA		5.00E-03		5.00E-03	g		5.00E-03
Nickel-63	13981-37-8	NA		6.00E-03		6.00E-03	g		6.00E-03
Plutonium-239	15117-48-3	NA		5.00E-07		5.00E-07	g		5.00E-07
Plutonium-241	14119-32-5	NA		5.00E-07		5.00E-07	g		5.00E-07
Samarium-151	15715-94-3	NA		5.00E-03		5.00E-03	g		5.00E-03
Strontium-90	10098-97-2	NA		3.00E-04		3.00E-04	g		3.00E-04
Technetium-99	14133-79-7	NA		8.50E-03		8.50E-03	g		8.50E-03
Tritium	10028-17-8	NA							
Yttrium-90	10098-91-6	NA		3.00E-04		3.00E-04	g		3.00E-04
Uranium-232	14158-29-3	NA		2.00E-04		2.00E-04	g		2.00E-04

Table C2-5. Terrestrial Animal-to-Animal Transfer Factors (Ba) for Ecological Receptors (mg/kg tissue/mg ingested/day)

Constituent of Potential Concern	CAS Registry Number	k	EPA (1999) Ba, Mammals[b]	Ba Calculated by EPA (1999) Methods[c]	Ecology Guidance Preferred Ba	SAIC Compiled Ba	Notes	Comments	Recommended Mammal Ba[d]
Uranium-233	13968-55-3	NA		2.00E-04		2.00E-04	g		2.00E-04
Uranium-234	13966-29-5	NA		2.00E-04		2.00E-04	g		2.00E-04
Uranium-235	15117-96-1	NA		2.00E-04		2.00E-04	g		2.00E-04
Uranium-236	13982-70-2	NA		2.00E-04		2.00E-04	g		2.00E-04
Uranium-238	7440-61-1	NA		2.00E-04		2.00E-04	g		2.00E-04
Actinium-227	14952-40-0	NA		2.50E-03		2.50E-03	g		2.50E-03
Americium-243	14993-75-0	NA		3.50E-06		3.50E-06	g		3.50E-06
Carbon-14	14762-75-5	NA							
Cobalt-60	10198-40-0	NA		2.00E-02		2.00E-02	g		2.00E-02
Curium-242	15510-73-3	NA		3.50E-06		3.50E-06	g		3.50E-06
Curium-243	15757-87-6	NA		3.50E-06		3.50E-06	g		3.50E-06
Curium-244	13981-15-2	NA		3.50E-06		3.50E-06	g		3.50E-06
Europium-152	14683-23-9	NA		5.00E-03		5.00E-03	g		5.00E-03
Iodine-129	15046-84-1	NA		7.00E-03		7.00E-03	g		7.00E-03
Neptunium-237	13994-20-2	NA		5.50E-03		5.50E-03	g		5.50E-03
Nickel-59	14336-70-0	NA		6.00E-03		6.00E-03	g		6.00E-03
Niobium-93		NA		2.50E-01		2.50E-01	g		2.50E-01
Plutonium-238	13981-16-3	NA		5.00E-07		5.00E-07	g		5.00E-07
Plutonium-240	14119-33-6	NA		5.00E-07		5.00E-07	g		5.00E-07
Plutonium-242	13982-10-0	NA		5.00E-07		5.00E-07	g		5.00E-07
Protactinium-231	14331-85-2	NA		1.00E-05		1.00E-05	g		1.00E-05
Radium-226	13982-63-3	NA		2.50E-04		2.50E-04	g		2.50E-04
Radium-228	15262-20-1	NA		2.50E-04		2.50E-04	g		2.50E-04
Ruthenium-106	13967-48-1	NA		2.00E-03		2.00E-03	g		2.00E-03
Selenium-79	None	NA		1.50E-02		1.50E-02	g		1.50E-02
Thorium-229	15594-54-4	NA		6.00E-06		6.00E-06	g		6.00E-06
Thorium-232	7440-29-1	NA		6.00E-06		6.00E-06	g		6.00E-06
Tin-126	15832-50-5	NA		8.00E-02		8.00E-02	g		8.00E-02
Zirconium-93	15751-77-6	NA		5.50E-03		5.50E-03	g		5.50E-03

Table C2-5. Terrestrial Animal-to-Animal Transfer Factors (Ba) for Ecological Receptors (mg/kg tissue/mg ingested/day)

Constituent of Potential Concern	CAS Registry Number	k	EPA (1999) Ba, Mammals[b]	Ba Calculated by EPA (1999) Methods[c]	Ecology Guidance Preferred Ba	SAIC Compiled Ba	Notes	Comments	Recommended Mammal Ba[d]

NA = Not applicable

[a] $\log_{10}$ of K_{ow} values in Table 4.1

[b] EPA (1999) calculates receptor-specific BCFs by multiplying the beef Ba [(mg/kg tissue)/(mg COPC ingested/day)] from Baes and othersl. (1984) by the receptor's specific ingestion rate (kg food/kg body wt/day). Receptor-specific values are found in Appendix D, Tables D-1 through D-3 in EPA (1999).
In this table, Ba values are back-calculated by dividing BCFs for muskrats by the ingestion rate of 0.267 kg/kg BW/d EPA (1999), Table 5-1.
EPA (1999) calculates bird Ba values for organics by multiplying mammal Ba values by 0.79, the nominal ratio of body fat in birds to body fat in mammals.

c Appendix D, Sect. D-1.1 EPA (1999) specifies that Ba values for organics other than dioxins and dibenzofurans, log Ba = -7.6 +log K_{ow} (Travis and Arms 1988).
For inorganic constituents, Ba values from Baes and others. (1984) were used EPA (1999), Appendix D, Sect. D-1.1

[d] Selection criteria described in Sect. 8.2.4.3

[e] Calculated by using log K_{ow}: log Ba = log K_{ow} -7.6 (Travis and Arms 1988)

[f] Value for Aroclor 1254 in Appendix D of EPA (1999), Table D-1, was used for PCB mixtures

[g] Baes and others. (1984), Figure 2.25

Table C2-6. Aquatic Water-to-Plant Transfer Factors (WP) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) WP[b]	WP Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance WP	Soil Kd[d]	SAIC Computed WP[e]	Comments	Recommended WP[f]
Organic Compounds										
Aromatic Halogenated Hydrocarbons										
4-Chloro-3-methylphenol	59-50-7	3.10		2.47E+01	g		6.03E+00	7.54E-01		7.54E-01
2,3,4,6-Tetrachlorophenol	58-90-2	4.42		2.98E+02	g		6.61E+01	1.43E+00		1.43E+00
Aromatic Nonhalogenated Hydrocarbons										
2-Nitrotoluene	88-72-2	2.30		5.47E+00	g		4.27E+00	1.55E+00		1.55E+00
4-Nitrobiphenyl	92-93-3	3.77		8.74E+01	g		No Kd	1.00E+00	Default	1.00E+00
Benzaldehyde	100-52-7	1.48		1.16E+00	g		2.01E-01	2.18E-01		2.18E-01
Benzene	71-43-2	-4.99		5.84E-06	g		6.20E+01	3.69E+05		3.69E+05
Benzyl alcohol	100-51-6	1.10		5.69E-01	g		1.02E-01	1.83E-01		1.83E-01
Ethyl benzene	100-41-4	3.12		2.58E+01	g		2.04E+00	2.47E-01		2.47E-01
m-Xylene	108-38-3	3.20		2.99E+01	g		1.96E+00	2.14E-01		2.14E-01
o-Xylene	95-47-6	3.13		2.62E+01	g		2.41E+00	2.90E-01		2.90E-01
p-Xylene	106-42-3	3.17		2.82E+01	g		3.11E+00	3.54E-01		3.54E-01
Styrene	100-42-5	2.93		1.79E+01	g		9.12E+00	1.43E+00		1.43E+00
Toluene	108-88-3	2.67		1.09E+01	g		1.40E+00	3.11E-01		3.11E-01
Non-aromatic Nonhalogenated Hydrocarbons										
1,2-Epoxybutane	106-88-7	1.44		1.08E+00	g		No Kd	1.00E+00	Default	1.00E+00
1,3-Butadiene	106-99-0	1.90		2.57E+00	g		7.41E-01	4.58E-01		4.58E-01
1,4-Dioxane	123-91-1	-0.27	4.00E-02		g		8.76E-03	9.69E-02		9.69E-02
1-Methylpropyl alcohol	78-92-2	0.61		2.26E-01	g		4.00E-02	1.37E-01		1.37E-01
1-Nitropropane	108-03-2	0.87		3.69E-01	g		No Kd	1.00E+00	Default	1.00E+00
2,2,4-Trimethylpentane	540-84-1	5.02		9.23E+02	g		8.64E+02	8.40E+00		8.40E+00
2-Butanone	78-93-3	0.28		1.21E-01	g		2.34E-02	1.25E-01		1.25E-01
2-Butenaldehyde (2-Butenal)	4170-30-3	No data		No data			No Kd	1.00E+00	Default	1.00E+00
2-Ethoxyethanol	110-80-5	-0.10		5.92E-02	g		2.09E-01	1.85E+00		1.85E+00
2-Heptanone	110-43-0	1.98		2.99E+00	g		8.88E-01	4.93E-01		4.93E-01
2-Hexanone	591-78-6	1.38		9.64E-01	g		1.34E+00	1.65E+00		1.65E+00
2-Methoxyethanol	109-86-4	0.25		1.15E-01	g		No Kd	1.00E+00	Default	1.00E+00
2-Methyl-2-propanol	75-65-0	0.35		1.38E-01	g		3.72E-01	1.81E+00		1.81E+00

Table C2-6. Aquatic Water-to-Plant Transfer Factors (WP) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) WP[b]	WP Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance WP	Soil Kd[d]	SAIC Computed WP[e]	Comments	Recommended WP[f]
2-Methyl-2-propenenitrile	126-98-7	0.54		1.98E-01	g		3.74E-02	1.41E-01		1.41E-01
2-Methylaziridine	75-55-8	-0.60		2.29E-02	g		No Kd	1.00E+00	Default	1.00E+00
2-Methylpropyl alcohol	78-83-1	0.76		3.00E-01	g		5.61E-02	1.58E-01		1.58E-01
2-Pentanone	107-87-9	0.91		3.97E-01	g		7.40E-01	1.71E+00		1.71E+00
2-Propanone (Acetone)	67-64-1	-0.22	5.00E-02		g		9.51E-03	9.89E-02		9.89E-02
2-Propene-1-ol	107-18-6	0.17		9.85E-02	g		1.48E-02	9.12E-02		9.12E-02
2-Propyl alcohol	67-63-0	0.05		7.85E-02	g		1.12E-02	8.15E-02		8.15E-02
3-Heptanone	106-35-4	No data		No data			No Kd	1.00E+00	Default	1.00E+00
3-Methyl-1-butanol	123-51-3	No data		No data			No Kd	1.00E+00	Default	1.00E+00
3-Methyl-2-butanone	563-80-4	No data		No data			No Kd	1.00E+00	Default	1.00E+00
3-Pentanone	96-22-0	0.99		4.62E-01	g		1.20E-01	2.49E-01		2.49E-01
4-Heptanone	123-19-3	No data		No data			No Kd	1.00E+00	Default	1.00E+00
4-Methyl-2-pentanone	108-10-1	1.19		6.74E-01	g		1.20E-01	1.91E-01		1.91E-01
4-Methyl-3-penten-2-one	141-79-7	No data		No data			No Kd	1.00E+00	Default	1.00E+00
5-Methyl-2-hexanone	110-12-3	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Acetaldehyde	75-07-0	-0.22		4.72E-02	g		9.53E-03	9.90E-02		9.90E-02
Acetamide	60-35-5	-1.26		6.64E-03	g		2.82E-04	1.17E-02		1.17E-02
Acetic acid	64-19-7	-0.17		5.19E-02	g		1.00E-02	9.71E-02		9.71E-02
Acetic acid ethyl ester	141-78-6	0.73		2.83E-01	g		2.30E-02	6.73E-02		6.73E-02
Acetic acid n-butyl ester	123-86-4	1.73		1.87E+00	g		5.04E-01	3.91E-01		3.91E-01
Acetonitrile	75-05-8	-0.34		3.76E-02	g		7.69E-03	9.37E-02		9.37E-02
Acrolein	107-02-8	-0.01		7.03E-02	g		1.39E-02	1.09E-01		1.09E-01
Acrylonitrile	107-13-1	0.25	1.10E-01		g		2.22E-02	1.23E-01		1.23E-01
Bis(isopropyl)ether	108-20-3	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Butane	106-97-8	2.89		1.66E+01	g		No Kd	1.00E+00	Default	1.00E+00
Carbon disulfide	75-15-0	2.00		3.10E+00	g		5.14E-01	2.78E-01		2.78E-01
Cyanogen	460-19-5	0.81		3.28E-01	g		No Kd	1.00E+00	Default	1.00E+00
Cyclohexane	110-82-7	3.44		4.69E+01	g		4.79E+00	3.81E-01		3.81E-01
Cyclohexanone	108-94-1	0.81		3.29E-01	g		6.28E-02	1.66E-01		1.66E-01
Cyclohexene	110-83-8	2.86		1.57E+01	g		6.51E+00	1.12E+00		1.12E+00

Table C2-6. Aquatic Water-to-Plant Transfer Factors (WP) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) WP[b]	WP Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance WP	Soil Kd[d]	SAIC Computed WP[e]	Comments	Recommended WP[f]
Cyclopentane	287-92-3	3.00		2.05E+01	g		8.93E+00	1.28E+00		1.28E+00
Ethyl alcohol	64-17-5	0.31		1.28E-01	g		2.03E-02	1.04E-01		1.04E-01
Ethyl ether	60-29-7	0.89		3.83E-01	g		7.53E-02	1.78E-01		1.78E-01
Ethyl methacrylate	97-63-2	1.59		1.43E+00	g		2.46E-01	2.30E-01		2.30E-01
Formaldehyde	50-00-0	0.34	1.40E-01		g		2.62E-02	1.29E-01		1.29E-01
Formamide	75-12-7	-1.51		4.14E-03	g		No Kd	1.00E+00	Default	1.00E+00
Formic acid	64-18-6	-0.54		2.59E-02	g		5.39E-02	8.54E-01		8.54E-01
Formic acid, methyl ester	107-31-3	-0.26		4.34E-02	g		No Kd	1.00E+00	Default	1.00E+00
Glycidylaldehyde	765-34-4	-0.73		1.80E-02	g		No Kd	1.00E+00	Default	1.00E+00
Methyl acetate	79-20-9	0.18		1.00E-01	g		4.80E-02	2.92E-01		2.92E-01
Methyl alcohol	67-56-1	-0.71		1.87E-02	g		3.96E-03	7.89E-02		7.89E-02
Methyl isocyanate	624-83-9	No data		No data			No Kd	NA	Reacts with water	NA
Methyl methacrylate	80-62-6	0.79		3.17E-01	g		6.31E-01	1.71E+00		1.71E+00
Methyl tert-butyl ether	1634-04-4	0.94		4.21E-01	g		8.43E-02	1.87E-01		1.87E-01
Methylacetylene	74-99-7	0.94		4.21E-01	g		No Kd	1.00E+00	Default	1.00E+00
Methylcyclohexane	108-87-2	4.10		1.63E+02	g		1.08E+02	3.56E+00		3.56E+00
N,N-Dimethylacetamide	127-19-5	No data		No data			No Kd	1.00E+00	Default	1.00E+00
n-Butyl alcohol	71-36-3	0.88		3.76E-01	g		7.36E-02	1.77E-01		1.77E-01
n-Heptane	142-82-5	4.66		4.68E+02	g		3.83E+02	6.00E+00		6.00E+00
n-Hexane	110-54-3	4.11		1.66E+02	g		1.10E+02	3.59E+00		3.59E+00
Nitromethane	75-52-5	-0.35		3.69E-02	g		No Kd	1.00E+00	Default	1.00E+00
n-Nonane	111-84-2	5.65		3.03E+03	g		3.60E+03	1.51E+01		1.51E+01
n-Octane	111-65-9	4.00		1.35E+02	g		8.59E+01	3.24E+00		3.24E+00
n-Pentane	109-66-0	3.21		3.04E+01	g		1.44E+01	1.55E+00		1.55E+00
n-Propionaldehyde	123-38-6	0.59		2.17E-01	g		3.82E-02	1.35E-01		1.35E-01
n-Propyl alcohol	71-23-8	0.25		1.14E-01	g		No Kd	1.00E+00	Default	1.00E+00
n-Valeraldehyde	110-62-3	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Oxirane	75-21-8	-0.30		4.06E-02	g		8.26E-03	9.54E-02		9.54E-02
p-Cymene	99-87-6	4.10		1.63E+02	g		No Kd	1.00E+00	Default	1.00E+00
Phosgene	75-44-5	No data		No data			No Kd	1.00E+00	Default	1.00E+00

Table C2-6. Aquatic Water-to-Plant Transfer Factors (WP) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) WP[b]	WP Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance WP	Soil Kd[d]	SAIC Computed WP[e]	Comments	Recommended WP[f]
Propargyl alcohol	107-19-7	0.26		1.17E-01	g		No Kd	1.00E+00	Default	1.00E+00
Propionic acid	79-09-4	0.33		1.33E-01	g		2.12E-02	1.06E-01		1.06E-01
Propionitrile	107-12-0	0.16		9.66E-02	g		1.44E-02	9.03E-02		9.03E-02
Propylene gylcol monomethyl ether	107-98-2	-0.18		5.09E-02	g		No Kd	1.00E+00	Default	1.00E+00
p-tert-Butyltoluene	98-51-1	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Triethylamine	121-44-8	0.16		9.66E-02	g		1.44E-02	9.03E-02		9.03E-02
Trimethylamine	75-50-3	0.16		9.66E-02	g		4.00E-02	2.50E-01		2.50E-01
Vinyl acetate	108-05-4	0.70		2.67E-01	g		4.97E-02	1.52E-01		1.52E-01
Non-aromatic Halogenated Hydrocarbons										
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	No data		No data			No Kd	1.00E+00	Default	1.00E+00
1,1,1,2-Tetrachloroethane	630-20-6	2.63		1.02E+01	g		1.59E+00	3.72E-01		3.72E-01
1,1,1-Trichloroethane	71-55-6	2.42		6.88E+00	g		1.35E+03	4.17E+02		4.17E+02
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	3.73		8.11E+01	g		3.16E+00	1.71E-01		1.71E-01
1,1,2,2-Tetrachloroethane	79-34-5	4.64		4.54E+02	g		7.90E-01	1.27E-02		1.27E-02
1,1,2,2-Tetrachloroethene	127-18-4	2.55		8.68E+00	g		2.65E+00	6.94E-01		6.94E-01
1,1,2-Trichloroethane	79-00-5	2.10		3.73E+00	g		7.50E-01	3.57E-01		3.57E-01
1,1,2-Trichloroethylene	79-01-6	2.43		7.02E+00	g		9.40E-01	2.86E-01		2.86E-01
1,1-Dichloroethane	75-34-3	1.79		2.10E+00	g		5.30E-01	3.78E-01		3.78E-01
1,1-Dichloroethene	75-35-4	2.12		3.90E+00	g		6.50E-01	2.99E-01		2.99E-01
1,2,2-Trichloro-1,1,2-trifluoroethane	76-13-1	3.16		2.77E+01	g		2.57E+00	2.97E-01		2.97E-01
1,2,3-Trichloropropane	96-18-4	2.25		4.98E+00	g		8.10E-01	3.14E-01		3.14E-01
1,2-Dibromo-3-chloropropane	96-12-8	2.34		5.90E+00	g		9.47E-01	3.26E-01		3.26E-01
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	2.82		1.46E+01	g		No Kd	1.00E+00	Default	1.00E+00
1,2-Dichloroethane	107-06-2	1.46		1.13E+00	g		1.96E-01	2.17E-01		2.17E-01
1,2-Dichloroethylene	540-59-0	0.48		1.77E-01	g		No Kd	1.00E+00	Default	1.00E+00
1,2-Dichloropropane	78-87-5	2.25		4.98E+00	g		4.70E-01	1.82E-01		1.82E-01
1,3-Dichloropropene	542-75-6	1.75		1.93E+00	g		2.70E-01	2.04E-01		2.04E-01
1,4-Dichloro-2-butene	764-41-0	0.87		3.70E-01	g		No Kd	1.00E+00	Default	1.00E+00
1-Chloroethene	75-01-4	1.15		6.20E-01	g		1.11E-01	1.87E-01		1.87E-01
2,2-Dichloropropionic acid	75-99-0	0.78		3.10E-01	g		No Kd	1.00E+00	Default	1.00E+00

Table C2-6. Aquatic Water-to-Plant Transfer Factors (WP) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) WP[b]	WP Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance WP	Soil Kd[d]	SAIC Computed WP[e]	Comments	Recommended WP[f]
2-Chloropropane	75-29-6	1.90		2.57E+00	g		7.41E-01	4.58E-01		4.58E-01
3-Chloropropene (allyl chloride)	107-05-1	0.95		4.32E-01	g		No Kd	1.00E+00	Default	1.00E+00
Bromochloromethane	74-97-5	1.41		1.02E+00	g		2.44E-01	2.90E-01		2.90E-01
Bromodichloromethane	75-27-4	2.03		3.26E+00	g		5.38E-01	2.81E-01		2.81E-01
Bromoethene	593-60-2	1.07		5.34E-01	g		No Kd	1.00E+00	Default	1.00E+00
Bromoform	75-25-2	2.35		6.01E+00	g		1.26E+00	4.28E-01		4.28E-01
Bromomethane	74-83-9	1.11		5.84E-01	g		9.00E-02	1.58E-01		1.58E-01
Carbon tetrachloride	56-23-5	2.72	3.00E+02				1.52E+00	3.16E-01		3.00E+02
Chlorodibromomethane	124-48-1	2.18		4.33E+00	g		7.05E-01	3.02E-01		3.02E-01
Chlorodifluoromethane	75-45-6	1.08		5.47E-01	g		9.83E-02	1.81E-01		1.81E-01
Chloroethane	75-00-3	3.10		2.47E+01	g		3.71E+00	4.64E-01		4.64E-01
Chloroform	67-66-3	1.95	2.82E+00		g		5.30E+01	3.07E+01		3.07E+01
Chloromethane	74-87-3	0.90		3.92E-01	g		6.00E-02	1.40E-01		1.40E-01
Chloropentafluoroethane	76-15-3	No data		No data			No Kd	1.00E+00	Default	1.00E+00
cis-1,2-Dichloroethene	156-59-2	1.98		3.00E+00	g		4.98E+00	2.76E+00		2.76E+00
cis-1,3-Dichloropropene	10061-01-5	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Cyanogen bromide	506-68-3	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Cyanogen chloride	506-77-4	0.20		1.04E-01	g		No Kd	1.00E+00	Default	1.00E+00
Dichlorodifluoromethane	75-71-8	2.16		4.19E+00	g		6.85E-01	3.00E-01		3.00E-01
Dichlorofluoromethane	75-43-4	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Dichloromethane	75-09-2	1.26		7.62E-01	g		1.00E-01	1.46E-01		1.46E-01
Difluorodibromomethane	75-61-6	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Hexafluoroacetone	684-16-2	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Iodomethane	74-88-4	1.69		1.73E+00	g		4.61E-01	3.76E-01		3.76E-01
Methylene bromide	74-95-3	1.62		1.52E+00	g		2.60E-01	2.33E-01		2.33E-01
Pentachloroethane	76-01-7	3.05		2.25E+01	g		1.00E+01	1.34E+00		1.34E+00
trans-1,2-Dichloroethene	156-60-5	1.98		3.00E+00	g		3.80E-01	2.10E-01		2.10E-01
trans-1,3-Dichloropropene	10061-02-6	2.06		3.48E+00	g		No Kd	1.00E+00	Default	1.00E+00
Trichloroacetic acid	76-03-9	1.33		8.78E-01	g		2.04E-01	2.69E-01		2.69E-01
Trichlorofluoroethane	27154-33-2	No data		No data			No Kd	1.00E+00	Default	1.00E+00

Table C2-6. Aquatic Water-to-Plant Transfer Factors (WP) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) WP[b]	WP Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance WP	Soil Kd[d]	SAIC Computed WP[e]	Comments	Recommended WP[f]
Trichlorofluoromethane	75-69-4	2.53		8.46E+00	g		1.34E+00	3.57E-01		3.57E-01
Trifluorobromomethane	75-63-8	1.86		2.38E+00	g		No Kd	1.00E+00	Default	1.00E+00
Dioxin and Furan Compounds										
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	35822-46-9	8.20	1.68E+02		h		9.77E+05	5.67E+01		1.68E+02
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	7.92	3.63E+01		h		5.13E+05	6.36E+00		3.63E+01
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	7.92	1.29E+03		h		5.13E+05	2.26E+02		1.29E+03
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	39227-28-6	7.79	1.02E+03		h		3.80E+05	1.29E+02		1.02E+03
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	7.25	2.51E+02		h		1.10E+05	9.46E+00		2.51E+02
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	57653-85-7	7.25	3.96E+02		h		1.10E+05	1.47E+01		3.96E+02
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	7.25	6.27E+02		h		1.10E+05	2.42E+01		6.27E+02
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	19408-74-3	7.25	4.62E+02		h		1.10E+05	1.72E+01		4.62E+02
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	7.25	2.08E+03		h		1.10E+05	7.70E+01		2.08E+03
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	40321-76-4	6.64	3.04E+03		h		2.69E+04	2.80E+01		3.04E+03
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	6.79	7.26E+02		h		3.80E+04	8.36E+00		7.26E+02
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	7.25	2.21E+03		h		1.10E+05	8.36E+01		2.21E+03
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	6.92	5.28E+03		h		5.13E+04	9.23E+01		5.28E+03
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1746-01-6	6.64	3.30E+03				2.69E+04	3.02E+01		3.30E+03
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	6.53	2.64E+03		h		2.09E+04	1.88E+01		2.64E+03
Dibenzofuran	132-64-9	4.33		2.51E+02	g		1.81E+02	4.41E+00		4.41E+00
Octachlorodibenzo(p)dioxin	3268-87-9	8.20	3.96E+01		h		2.24E+01	3.00E-04		3.96E+01
Octachlorodibenzofuran	39001-02-0	8.78	5.28E+01		h		3.72E+06	6.70E+01		5.28E+01
PCBs										
Polychlorinated biphenyls (PCBs)[i]	1336-36-3	7.31	4.77E+05				3.09E+03	6.18E+00		4.77E+05
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	7.08		4.49E+04	g		No Kd	1.00E+00	Default	1.00E+00
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	7.12		4.84E+04	g		No Kd	1.00E+00	Default	1.00E+00
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	No data		No data			No Kd	1.00E+00	Default	1.00E+00
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	No data		No data			No Kd	1.00E+00	Default	1.00E+00
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	No data		No data			No Kd	1.00E+00	Default	1.00E+00
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	No data		No data			No Kd	1.00E+00	Default	1.00E+00
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	No data		No data			No Kd	1.00E+00	Default	1.00E+00

Table C2-6. Aquatic Water-to-Plant Transfer Factors (WP) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) WP[b]	WP Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance WP	Soil Kd[d]	SAIC Computed WP[e]	Comments	Recommended WP[f]
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	No data		No data			No Kd	1.00E+00	Default	1.00E+00
2',3,4,4',5-Pentachlorobiphenyl	no cas #	No data		No data			No Kd	1.00E+00	Default	1.00E+00
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	7.12		4.84E+04	g		No Kd	1.00E+00	Default	1.00E+00
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	7.41		8.34E+04	g		No Kd	1.00E+00	Default	1.00E+00
3,3',4,4',5-Pentachlorobiphenyl	no cas #	No data		No data			No Kd	1.00E+00	Default	1.00E+00
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	No data		No data			No Kd	1.00E+00	Default	1.00E+00
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Phthalates										
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	5.20	9.93E+03				1.11E+03	8.44E+00		9.93E+03
Butylbenzyl phthalate	85-68-7	4.41		2.94E+02	g		1.37E+02	2.98E+00		2.98E+00
Dibutyl phthalate	84-74-2	4.72		5.25E+02	g		1.57E+01	2.27E-01		2.27E-01
Diethyl phthalate	84-66-2	4.44		3.07E+02	g		8.20E-01	1.73E-02		1.73E-02
Dimethylphthalate	131-11-3	1.63		1.56E+00	g		2.66E+00	2.34E+00		2.34E+00
n-Dioctyl phthalate	117-84-0	9.33	2.85E+04				9.03E+06	2.84E+02		2.85E+04
Light Polycyclic Aromatic Hydrocarbons (molecular weight <200 g/mole)										
2-Chloronapthalene	91-58-7	4.07		1.53E+02	g		7.14E+01	2.46E+00		2.46E+00
2-Methyl naphthalene	91-57-6	3.86		1.04E+02	g		6.26E+01	2.85E+00		2.85E+00
5-Nitroacenaphthene	602-87-9	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Acenaphthene	83-32-9	3.96		1.26E+02	g		4.90E+01	1.94E+00		1.94E+00
Acenaphthylene	208-96-8	4.07		1.54E+02	g		6.76E+01	2.33E+00		2.33E+00
Anthracene	120-12-7	4.47		3.27E+02	g		2.35E+02	4.75E+00		4.75E+00
Fluorene	86-73-7	4.18		1.89E+02	g		1.41E+02	4.20E+00		4.20E+00
Indene	95-13-6	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Naphthalene	91-20-3	3.37		4.13E+01	g		1.19E+01	1.04E+00		1.04E+00
Phenanthrene	85-01-8	4.55		3.81E+02	g		5.01E+02	9.10E+00		9.10E+00
Pyrene	129-00-0	5.00		8.89E+02	g		6.80E+02	6.78E+00		6.78E+00
Heavy Polycyclic Aromatic Hydrocarbons (molecular weight >200 g/mole)										
3-Methylcholanthrene	56-49-5	7.11		4.75E+04	g		1.51E+04	9.11E+00		9.11E+00
5-Methylchrysene	3697-24-3	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Benzo(a)anthracene	56-55-3	5.68	5.26E+03		j		2.60E+03	1.05E+01		5.26E+03

Table C2-6. Aquatic Water-to-Plant Transfer Factors (WP) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) WP [b]	WP Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance WP	Soil Kd [d]	SAIC Computed WP [e]	Comments	Recommended WP [f]
Benzo(a)pyrene	50-32-8	6.13	5.26E+03				9.69E+03	2.15E+01		5.26E+03
Benzo(b)fluoranthene	205-99-2	6.20	5.26E+03		j		8.36E+03	1.69E+01		5.26E+03
Benzo(e)pyrene	192-97-2	7.40		8.21E+04	g		1.58E+05	6.48E+01		6.48E+01
Benzo(g,h,i)perylene	191-24-2	7.10		4.67E+04	g		1.82E+04	1.11E+01		1.11E+01
Benzo(j)fluoranthene	205-82-3	6.44		1.34E+04	g		2.15E+04	3.16E+01		3.16E+01
Benzo(k)fluoranthene	207-08-9	6.20	5.26E+03		j		8.32E+03	1.68E+01		5.26E+03
Benzo[a,i]pyrene	191-30-0	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Chrysene	218-01-9	5.74	5.26E+03		j		2.97E+03	1.11E+01		5.26E+03
Dibenz(a,h)anthracene	53-70-3	6.55	5.26E+03		j		1.79E+04	2.29E+01		5.26E+03
Dibenz[a,h]acridine	226-36-8	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Dibenz[a,j]acridine	224-42-0	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Dibenzo(a,e)fluoranthene	5385-75-1	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Dibenzo(a,h)fluoranthene	no cas #	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Dibenzo[a,e]pyrene	192-65-4	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Dibenzo[a,h]pyrene	189-64-0	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Dibenzo[a,i]pyrene	189-55-9	7.29		6.68E+04	g		No Kd	1.00E+00	Default	1.00E+00
Fluoranthene	206-44-0	5.08		1.04E+03	g		4.91E+02	4.39E+00		4.39E+00
Hexachloronaphthalene	1335-87-1	7.59		1.18E+05	g		No Kd	1.00E+00	Default	1.00E+00
Indeno(1,2,3-cd)pyrene	193-39-5	6.91	5.26E+03		j		4.11E+04	3.21E+01		5.26E+03
Octachloronaphthalene	2234-13-1	6.42		1.29E+04	g		No Kd	1.00E+00	Default	1.00E+00
Pentachloronaphthalene	1321-64-8	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Tetrachloronaphthalene	1335-88-2	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Trichloronaphthalene	1321-65-9	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Light Substituted Benzene Compounds (MW <200 g/mole)										
1,2,3-Trichlorobenzene	87-61-6	4.05		1.47E+02	g		2.02E+02	7.18E+00		7.18E+00
1,2,4-Trichlorobenzene	120-82-1	3.99		1.32E+02	g		1.66E+01	6.37E-01		6.37E-01
1,2,4-Trimethyl benzene	95-63-6	3.65		6.97E+01	g		3.89E+01	2.34E+00		2.34E+00
1,2-Dichlorobenzene	95-50-1	3.45		4.74E+01	g		3.79E+00	2.99E-01		2.99E-01
1,3,5-Trimethyl benzene	108-67-8	3.42		4.52E+01	g		1.67E+01	1.36E+00		1.36E+00
1,3-Dichlorobenzene	541-73-1	3.53		5.56E+01	g		8.03E+01	5.67E+00		5.67E+00

Table C2-6. Aquatic Water-to-Plant Transfer Factors (WP) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) WP[b]	WP Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance WP	Soil Kd[d]	SAIC Computed WP[e]	Comments	Recommended WP[f]
1,3-Dinitrobenzene	99-65-0	1.49	2.51E+03		k		2.06E-01	2.19E-01		2.51E+03
1,4-Dichlorobenzene	106-46-7	3.41		4.45E+01	g		6.16E+00	5.09E-01		5.09E-01
1,4-Dinitrobenzene	100-25-4	1.50		1.21E+00	g		No Kd	1.00E+00	Default	1.00E+00
2,4,5-Trichlorophenol	95-95-4	3.87		1.06E+02	g		1.13E+01	5.07E-01		5.07E-01
2,4,6-Trichlorophenol	88-06-2	3.71		7.83E+01	g		2.26E+00	1.25E-01		1.25E-01
2,4-Dichlorophenol	120-83-2	3.04		2.20E+01	g		1.40E+00	1.90E-01		1.90E-01
2,4-Dimethylphenol	105-67-9	2.36		6.12E+00	g		1.26E+00	4.22E-01		4.22E-01
2,4-Dinitrophenol	51-28-5	1.52		1.25E+00	g		1.00E-04	1.03E-04		1.03E-04
2,4-Dinitrotoluene	121-14-2	2.00	2.51E+03				5.10E-01	2.77E-01		2.51E+03
2,6-Dinitrotoluene	606-20-2	1.89	2.51E+03		k		4.19E-01	2.64E-01		2.51E+03
2-Chlorophenol	95-57-8	2.16		4.21E+00	g		3.87E+00	1.69E+00		1.69E+00
2-Chlorotoluene	95-49-8	3.54		5.63E+01	g		No Kd	1.00E+00	Default	1.00E+00
2-Nitrophenol	88-75-5	1.79		2.09E+00	g		3.53E+00	2.52E+00		2.52E+00
4,6-Dinitro-o-cresol	534-52-1	2.85		1.54E+01	g		No Kd	1.00E+00	Default	1.00E+00
4-Chlorotoluene	106-43-4	3.33		3.81E+01	g		No Kd	1.00E+00	Default	1.00E+00
4-Nitrophenol	100-02-7	1.91		2.62E+00	g		4.37E+02	2.66E+02		2.66E+02
alpha-Methylstyrene	98-83-9	3.46		4.90E+01	g		No Kd	1.00E+00	Default	1.00E+00
Aniline	62-53-3	0.98		4.54E-01	g		8.23E-02	1.73E-01		1.73E-01
Benzotrichloride	98-07-7	2.92		1.76E+01	g		No Kd	1.00E+00	Default	1.00E+00
Benzyl chloride	100-44-7	0.36		1.41E-01	g		2.71E-02	1.30E-01		1.30E-01
Bromobenzene	108-86-1	2.99		2.01E+01	g		4.47E+00	6.47E-01		6.47E-01
Chlorobenzene	108-90-7	2.79		1.38E+01	g		2.24E+00	4.24E-01		4.24E-01
Cumene	98-82-8	3.61		6.50E+01	g		9.13E+01	5.77E+00		5.77E+00
m-Cresol	108-39-4	1.96		2.87E+00	g		4.78E-01	2.73E-01		2.73E-01
n-Butyl benzene	104-51-8	4.28		2.29E+02	g		2.51E+01	6.53E-01		6.53E-01
Nitrobenzene	98-95-3	1.83	2.40E+01				1.19E+00	8.04E-01		2.40E+01
n-Propyl benzene	103-65-1	3.69		7.52E+01	g		7.24E+00	4.13E-01		4.13E-01
o-Cresol	95-48-7	2.02		3.23E+00	g		5.34E-01	2.81E-01		2.81E-01
o-Dinitrobenzene	528-29-0	1.60		1.46E+00	g		No Kd	1.00E+00	Default	1.00E+00
o-Nitroaniline	88-74-4	1.85		2.34E+00	g		3.93E+00	2.59E+00		2.59E+00

Table C2-6. Aquatic Water-to-Plant Transfer Factors (WP) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) WP[b]	WP Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance WP	Soil Kd[d]	SAIC Computed WP[e]	Comments	Recommended WP[f]
o-Toluidine	95-53-4	1.34		8.95E-01	g		1.57E-01	2.04E-01		2.04E-01
p-Chloroaniline	106-47-8	1.87		2.43E+00	g		4.06E-01	2.61E-01		2.61E-01
p-Cresol	106-44-5	1.94		2.77E+00	g		4.61E-01	2.70E-01		2.70E-01
Phenol	108-95-2	1.48		1.16E+00	g		2.20E-01	2.39E-01		2.39E-01
p-Nitrochlorobenzene	100-00-5	2.39		6.48E+00	g		No Kd	1.00E+00	Default	1.00E+00
p-Toluidine	106-49-0	1.40		1.00E+00	g		2.39E-01	2.87E-01		2.87E-01
sec-Butyl benzene	135-98-8	4.57		3.95E+02	g		No Kd	1.00E+00	Default	1.00E+00
tert-Butyl benzene	98-06-6	4.11		1.66E+02	g		1.10E+02	3.59E+00		3.59E+00
Toluene-2,6-diamine	823-40-5	1.45		1.11E+00	g		No Kd	1.00E+00	Default	1.00E+00
Trimethyl benzene	25551-13-7	3.40		4.35E+01	g		No Kd	1.00E+00	Default	1.00E+00
Other Light Semivolatile Compounds (molecular weight <200 g/mole)										
1,1'-Biphenyl	92-52-4	3.90		1.12E+02	g		2.51E+01	1.08E+00		1.08E+00
1,1-Dimethylhydrazine	57-14-7	0.60		2.20E-01	g		No Kd	1.00E+00	Default	1.00E+00
1,2-Dimethylhydrazine	540-73-8	-1.37		5.42E-03	g		No Kd	1.00E+00	Default	1.00E+00
1,2-Diphenylhydrazine	122-66-7	2.94		1.83E+01	g		2.78E+00	4.30E-01		4.30E-01
1,3-Propane sultone	1120-71-4	-0.52		2.66E-02	g		No Kd	1.00E+00	Default	1.00E+00
2,4-Toluene diisocyante	584-84-9	No data		No data			No Kd	NA	Decomposes rapidly in water	NA
2-Chloroacetophenone	532-27-4	2.59		9.53E+00	g		No Kd	1.00E+00	Default	1.00E+00
2-Propenoic acid	79-10-7	0.33		1.33E-01	g		2.12E-02	1.06E-01		1.06E-01
4,4-Methylenedianiline	101-77-9	3.38		4.19E+01	g		No Kd	1.00E+00	Default	1.00E+00
Acetophenone	98-86-2	1.64		1.58E+00	g		2.69E-01	2.35E-01		2.35E-01
Benzoic acid	65-85-0	1.86		2.38E+00	g		5.50E-03	3.58E-03		3.58E-03
bis(2-Chloroethoxy)methane	111-91-1	7.59		1.18E+05	g		2.40E+05	7.63E+01		7.63E+01
bis(2-Chloroethyl) ether	111-44-4	1.30		8.31E-01	g		7.60E-01	1.04E+00		1.04E+00
Chlorocyclopentadiene	41851-50-7	2.43		6.99E+00	g		No Kd	1.00E+00	Default	1.00E+00
Cyclohexanol	108-93-0	1.23		7.27E-01	g		1.30E-01	1.96E-01		1.96E-01
Dichloroisopropyl ether	108-60-1	2.58		9.27E+00	g		6.10E-01	1.52E-01		1.52E-01
Dichloromethyl ether	542-88-1	-0.38		3.49E-02	g		7.94E-01	1.02E+01		1.02E+01
Dichloropentadiene	no cas #	No data		No data			No Kd	1.00E+00	Default	1.00E+00

Table C2-6. Aquatic Water-to-Plant Transfer Factors (WP) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) WP[b]	WP Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance WP	Soil Kd[d]	SAIC Computed WP[e]	Comments	Recommended WP[f]
Dimethyl sulfate	77-78-1	0.32		1.30E-01	g		No Kd	1.00E+00	Default	1.00E+00
Dimethylaniline	121-69-7	-0.88		1.36E-02	g		No Kd	1.00E+00	Default	1.00E+00
Di-n-propylnitrosamine	621-64-7	1.38		9.65E-01	g		1.70E-01	2.10E-01		2.10E-01
Diphenyl ether	101-84-8	4.21		2.00E+02	g		1.38E+02	3.95E+00		3.95E+00
Epichlorohydrin	106-89-8	0.25		1.15E-01	g		2.22E-02	1.23E-01		1.23E-01
Ethyl carbamate (urethane)	51-79-6	-0.15		5.38E-02	g		No Kd	1.00E+00	Default	1.00E+00
Ethyl methanesulfonate	62-50-0	0.05		7.84E-02	g		1.55E-02	1.12E-01		1.12E-01
Ethylene dibromide	106-93-4	1.75		1.94E+00	g		3.28E-01	2.47E-01		2.47E-01
Ethylene glycol	107-21-1	-0.91		1.27E-02	g		No Kd	1.00E+00	Default	1.00E+00
Ethylene glycol monobutyl ether	111-76-2	1.55		1.34E+00	g		No Kd	1.00E+00	Default	1.00E+00
Ethylene glycol monoethyl ether acetate	111-15-9	0.62		2.30E-01	g		No Kd	1.00E+00	Default	1.00E+00
Ethylene thiourea	96-45-7	-0.64		2.13E-02	g		No Kd	1.00E+00	Default	1.00E+00
Furfural	98-01-1	0.96		4.37E-01	g		8.82E-02	1.90E-01		1.90E-01
Maleic hydrazide	123-33-1	-0.74		1.77E-02	g		No Kd	1.00E+00	Default	1.00E+00
Malononitrile	109-77-3	0.04		7.70E-02	g		No Kd	1.00E+00	Default	1.00E+00
Methyl styrene (mixed isomers)	25013-15-4	3.35		3.96E+01	g		No Kd	1.00E+00	Default	1.00E+00
Methylhydrazine	60-34-4	-1.06		9.68E-03	g		No Kd	1.00E+00	Default	1.00E+00
N,N-Diphenylamine	122-39-4	3.50		5.25E+01	g		3.47E+00	2.55E-01		2.55E-01
Nitric acid, propyl ester	627-13-4	No data		No data			No Kd	1.00E+00	Default	1.00E+00
N-Nitrosodi-n-butylamine	924-16-3	2.41		6.73E+00	g		1.07E+00	3.35E-01		3.35E-01
N-Nitrosomorpholine	59-89-2	0.98		4.54E-01	g		No Kd	1.00E+00	Default	1.00E+00
N-Nitroso-N,N-dimethylamine	62-75-9	-0.47		2.94E-02	g		No Kd	1.00E+00	Default	1.00E+00
o-Anisidine	90-04-0	1.18		6.61E-01	g		No Kd	1.00E+00	Default	1.00E+00
Oxalic acid	144-62-7	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Phthalic anhydride	85-44-9	-0.60		2.30E-02	g		4.80E-03	8.28E-02		8.28E-02
p-Phthalic acid	100-21-0	0.82		3.37E-01	g		No Kd	1.00E+00	Default	1.00E+00
Pyridine	110-86-1	0.67		2.53E-01	g		4.72E-02	1.50E-01		1.50E-01
Quinoline	91-22-5	2.03		3.29E+00	g		No Kd	1.00E+00	Default	1.00E+00
Quinone	106-51-4	0.20		1.04E-01	g		No Kd	1.00E+00	Default	1.00E+00
Safrole	94-59-7	2.66		1.08E+01	g		1.68E+00	3.77E-01		3.77E-01

Table C2-6. Aquatic Water-to-Plant Transfer Factors (WP) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) WP[b]	WP Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance WP	Soil Kd[d]	SAIC Computed WP[e]	Comments	Recommended WP[f]
Tetrahydrofuran	109-99-9	0.45		1.66E-01	g		3.16E-02	1.35E-01		1.35E-01
Other Heavy Semivolatile Compounds (molecular weight >200 g/mole)										
1,2,4,5-Tetrachlorobenzene	95-94-3	4.64		4.51E+02	g		5.89E+01	9.50E-01		9.50E-01
1,3,5-Trinitrobenzene	99-35-4	1.18		6.60E-01	g		1.18E-01	1.90E-01		1.90E-01
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	4.17		1.86E+02	g		No Kd	1.00E+00	Default	1.00E+00
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	-2.70		4.39E-04	g		No Kd	1.00E+00	Default	1.00E+00
2-sec-Butyl-4,6-dinitrophenol	88-85-7	3.14		2.66E+01	g		No Kd	1.00E+00	Default	1.00E+00
3,3'-Dimethoxybenzidine	119-90-4	1.81		2.17E+00	g		3.65E-01	2.54E-01		2.54E-01
3,3-Dichlorobenzidine	91-94-1	3.58		6.05E+01	g		8.70E+00	5.78E-01		5.78E-01
4-Bromophenylphenyl ether	101-55-3	5.00		8.89E+02	g		No Kd	1.00E+00	Default	1.00E+00
Ammonium perfluorooctanoate	3825-26-1	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Azobenzene	103-33-3	3.82		9.61E+01	g		No Kd	1.00E+00	Default	1.00E+00
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Captan	133-06-2	2.35		6.01E+00	g		2.00E+00	6.77E-01		6.77E-01
Chlorobenzilate	510-15-6	4.38		2.76E+02	g		3.69E+01	8.40E-01		8.40E-01
Dibutylphosphate	107-66-4	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Dimethyl aminoazobenzene	60-11-7	4.58		4.03E+02	g		No Kd	1.00E+00	Default	1.00E+00
Hexachlorobenzene	118-74-1	5.50	1.11E+04				8.00E+02	4.08E+00		1.11E+04
Hexachlorobutadiene	87-68-3	4.73	1.60E+02				6.94E+01	9.91E-01		1.60E+02
Hexachlorocyclopentadiene	77-47-4	5.04	6.10E+02				9.51E+01	1.07E+00		6.10E+02
Hexachloroethane	67-72-1	3.98		1.31E+02	g		1.82E+01	7.01E-01		7.01E-01
Hexachlorophene	70-30-4	7.54		1.07E+05	g		1.08E+04	3.67E+00		3.67E+00
Hexamethylene-1,5-diisocyanate	822-06-0	1.27		7.89E-01	g		No Kd	1.00E+00	Default	1.00E+00
Mirex	2385-85-5	6.89		3.14E+04	g		1.00E+04	8.07E+00		8.07E+00
Nitrofen	1836-75-5	5.53		2.42E+03	g		No Kd	1.00E+00	Default	1.00E+00
Pentachlorobenzene	608-93-5	5.09	4.00E+03				3.21E+02	2.82E+00		4.00E+03
Pentachloronitrobenzene	82-68-8	4.64	4.74E+03				5.89E+01	9.48E-01		4.74E+03
Pentachlorophenol	87-86-5	5.08	1.71E+03				1.99E+02	1.79E+00		1.71E+03
Picric acid	88-89-1	2.03		3.29E+00	g		No Kd	1.00E+00	Default	1.00E+00
Pronamide	23950-58-5	3.51		5.36E+01	g		7.74E+00	5.61E-01		5.61E-01

Table C2-6. Aquatic Water-to-Plant Transfer Factors (WP) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) WP[b]	WP Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance WP	Soil Kd[d]	SAIC Computed WP[e]	Comments	Recommended WP[f]
Strychnine	57-24-9	1.93		2.72E+00	g		4.53E-01	2.69E-01		2.69E-01
Terphenyls	26140-60-3	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Tributyl phosphate	126-73-8	4.00		1.35E+02	g		No Kd	1.00E+00	Default	1.00E+00
Trifluralin	1582-09-8	5.34		1.69E+03	g		6.03E+01	3.82E-01		3.82E-01
Triphenylamine	603-34-9	No data		No data			No Kd	1.00E+00	Default	1.00E+00
Herbicides and Organochlorinated Pesticides										
2,4,5-T	93-76-5	3.36		4.03E+01	g		5.25E-01	4.64E-02		4.64E-02
2,4-D and esters	94-75-7	2.81		1.43E+01	g		2.00E-01	3.67E-02		3.67E-02
4,4-DDD	72-54-8	6.20		8.55E+03	g		1.00E+03	2.02E+00		2.02E+00
4,4-DDE	72-55-9	6.26	1.13E+04				8.64E+06	1.62E+04		1.13E+04
4,4-DDT	50-29-3	6.00		5.86E+03	g		2.40E+03	6.32E+00		6.32E+00
Aldrin	309-00-2	6.18		8.21E+03	g		4.87E+02	1.01E+00		1.01E+00
alpha-BHC	319-84-6	3.80		9.24E+01	g		1.76E+01	8.68E-01		8.68E-01
beta-BHC	319-85-7	3.83		9.85E+01	g		2.14E+01	1.01E+00		1.01E+00
Chlordane	57-74-9	5.94		5.21E+03	g		5.13E+02	1.47E+00		1.47E+00
Delta-BHC	319-86-8	4.14		1.76E+02	g		6.61E+00	2.07E-01		2.07E-01
Dieldrin	60-57-1	5.27		1.48E+03	g		2.55E+02	1.78E+00		1.78E+00
Endothall	145-73-3	-0.87		1.39E-02	g		1.40E-03	3.46E-02		3.46E-02
Endrin	72-20-8	4.89		7.25E+02	g		1.08E+02	1.24E+00		1.24E+00
gamma-BHC (Lindane)	58-89-9	3.72		7.96E+01	g		1.07E+01	5.87E-01		5.87E-01
Heptachlor	76-44-8	5.02	2.10E+04				9.53E+01	9.32E-01		2.10E+04
Isodrin	465-73-6	3.55		5.77E+01	g		No Kd	1.00E+00	Default	1.00E+00
Methoxychlor	72-43-5	4.53		3.64E+02	g		8.00E+02	1.50E+01		1.50E+01
Silvex (2,4,5-TP)	93-72-1	4.07		1.55E+02	g		No Kd	1.00E+00	Default	1.00E+00
Toxaphene	8001-35-2	5.50		2.28E+03	g		1.00E+03	5.13E+00		5.13E+00
Inorganic Chemicals and Compounds										
Metals										
Aluminum	7429-90-5	NA	8.33E+02				1.50E+03	1.20E+00		8.33E+02
Antimony	7440-36-0	NA	1.48E+03				4.50E+01	1.80E+00		1.48E+03
Arsenic	7440-38-2	NA	2.93E+02				2.50E+01	1.80E-01		2.99E+03

Table C2-6. Aquatic Water-to-Plant Transfer Factors (WP) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) WP[b]	WP Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance WP	Soil Kd[d]	SAIC Computed WP[e]	Comments	Recommended WP[f]
Barium	7440-39-3	NA	2.60E+02				1.10E+01	3.30E-01		2.60E+02
Beryllium	7440-41-7	NA	1.41E+02				2.30E+01	4.60E-02		1.41E+02
Bismuth	7440-69-9	NA					2.00E+02	1.40E+00		1.40E+00
Boron	7440-42-8	NA					3.00E+00	2.40E+00		2.40E+00
Cadmium	7440-43-9	NA	7.82E+02				1.50E+01	1.09E+00		7.82E+02
Calcium	7440-70-2	NA					4.00E+00	2.80E+00		2.80E+00
Chromium (and VI)	18540-29-9	NA	4.41E+03				1.20E+03	1.80E+00		4.41E+03
Cobalt	7440-48-4	NA					4.50E+01	1.80E-01		1.80E-01
Copper	7440-50-8	NA	5.41E+02				3.50E+01	2.80E+00		5.41E+02
Iron	7439-89-6	NA					2.50E+01	2.00E-02		2.00E-02
Lead	7439-92-1	NA	1.71E+03				9.00E+02	8.10E+00		1.71E+03
Lithium	7439-93-2	NA					3.00E+02	1.50E+00		1.50E+00
Magnesium	7439-95-4	NA					4.50E+00	9.00E-01		9.00E-01
Manganese	7439-96-5	NA					6.50E+01	3.25E+00		3.25E+00
Mercury	7439-97-6	NA	2.48E+04				1.00E+03	7.50E+00		2.48E+04
Molybdenum	7439-98-7	NA					2.00E+01	1.00E+00		1.00E+00
Nickel	7440-02-0	NA	6.10E+01				1.60E+01	1.02E-01		6.10E+01
Potassium	7440-09-7	NA					5.50E+00	1.10E+00		1.10E+00
Rhodium	7440-16-6	NA					6.00E+01	1.80E+00		1.80E+00
Selenium	7782-49-2	NA	1.85E+03				1.80E+01	5.76E-02		1.85E+03
Silicon	7440-21-3	NA					3.00E+01	2.10E+00		2.10E+00
Silver	7440-22-4	NA	1.07E+04				1.00E-01	8.00E-03		1.07E+04
Sodium	7440-23-5	NA					1.00E+02	1.50E+00		1.50E+00
Strontium	7440-24-6	NA					3.50E+01	1.75E+01		1.75E+01
Tantalum	7440-25-7	NA					6.50E+02	1.30E+00		1.30E+00
Thallium	7440-28-0	NA	1.50E+04				4.40E+01	3.52E-02		1.50E+04
Tin	7440-31-5	NA					2.50E+02	1.50E+00		1.50E+00
Tungsten	7440-33-7	NA					1.50E+02	1.35E+00		1.35E+00
Uranium	7440-61-1	NA					4.50E+02	7.65E-01		7.65E-01
Vanadium	7440-62-2	NA					1.00E+03	1.10E+00		1.10E+00

Table C2-6. Aquatic Water-to-Plant Transfer Factors (WP) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) WP [b]	WP Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance WP	Soil Kd [d]	SAIC Computed WP [e]	Comments	Recommended WP [f]
Yttrium	7440-65-5	NA					5.00E+02	1.50E+00		1.50E+00
Zinc	7440-66-6	NA	2.18E+03				6.20E+01	1.49E-11		2.18E+03
Zirconium	7440-67-7	NA					3.00E+03	1.20E+00		1.20E+00
Non-metals and Anions										
Ammonia/Ammonium	7664-41-7	NA					No Kd	1.00E+00	Default	1.00E+00
Bromide	24959-67-9	NA					7.50E+00	2.25E+00		2.25E+00
Chloride	16887-00-6	NA					2.50E-01	3.50E+00		3.50E+00
Cyanide	57-12-5	NA	2.20E+01				No Kd	1.00E+00	Default	1.00E+00
Fluoride	16984-48-8	NA					1.50E+02	1.80E+00		1.80E+00
Hydroxide	14280-30-9	NA	NA				No Kd	NA	Depends on pH	NA
Iodine	7553-56-2	NA					6.00E+01	1.80E+00		1.80E+00
Nitrate	14797-55-8	NA					No Kd	1.00E+00	Default	1.00E+00
Nitrite	14797-65-0	NA					No Kd	1.00E+00	Default	1.00E+00
Phosphate	14265-44-2	NA					3.50E+00	2.45E+00		2.45E+00
Phosphorus	7723-14-0	NA					3.50E+00	2.45E+00		2.45E+00
Sulfate	14808-79-8	NA					No Kd	1.00E+00	Default	1.00E+00
Total Sulfur	63705-05-5	NA					7.50E+00	2.25E+00		2.25E+00
Priority Pollutants										
Carbon Dioxide	124-38-9	NA					No Kd	1.00E+00	Default	1.00E+00
Nitrogen Dioxide	10102-44-0	NA					No Kd	1.00E+00	Default	1.00E+00
Ozone	10028-15-6	NA					No Kd	1.00E+00	Default	1.00E+00
Particulate Matter	No CAS #	NA	NA				No Kd	1.00E+00	Default	1.00E+00
Sulfur Dioxode	7446-09-5	NA					No Kd	1.00E+00	Default	1.00E+00
Radionuclides										
Americium-241	1596-10-2	NA					7.00E+02	7.70E-01		7.70E-01
Antimony-125	14234-35-6	NA	1.48E+03				4.50E+01	1.80E+00		1.48E+03
Barium-137	13981-97-0	NA	2.60E+02				6.00E+01	1.80E+00		2.60E+02
Cadmium-113	None	NA	7.82E+02				6.50E+00	7.15E-01		7.82E+02
Cesium-134	13967-70-9	NA					1.00E+03	1.60E+01		1.60E+01
Cesium-137	10045-97-3	NA					1.00E+03	1.60E+01		1.60E+01

Table C2-6. Aquatic Water-to-Plant Transfer Factors (WP) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) WP[b]	WP Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance WP	Soil Kd[d]	SAIC Computed WP[e]	Comments	Recommended WP[f]
Europium-154	15585-10-1	NA					6.50E+02	1.30E+00		1.30E+00
Europium-155	14391-16-3	NA					6.50E+02	1.30E+00		1.30E+00
Nickel-63	13981-37-8	NA	6.10E+01				1.50E+02	1.80E+00		6.10E+01
Plutonium-239	15117-48-3	NA					4.50E+03	4.05E-01		4.05E-01
Plutonium-241	14119-32-5	NA					4.50E+03	4.05E-01		4.05E-01
Samarium-151	15715-94-3	NA					6.50E+02	1.30E+00		1.30E+00
Strontium-90	10098-97-2	NA					3.50E+01	1.75E+01		1.75E+01
Technetium-99	14133-79-7	NA					1.50E+00	2.85E+00		2.85E+00
Tritium	10028-17-8	NA						1.00E+00	Default	1.00E+00
Yttrium-90	10098-91-6	NA					5.00E+02	1.50E+00		1.50E+00
Uranium-232	14158-29-3	NA					4.50E+02	7.65E-01		7.65E-01
Uranium-233	13968-55-3	NA					4.50E+02	7.65E-01		7.65E-01
Uranium-234	13966-29-5	NA					4.50E+02	7.65E-01		7.65E-01
Uranium-235	15117-96-1	NA					4.50E+02	7.65E-01		7.65E-01
Uranium-236	13982-70-2	NA					4.50E+02	7.65E-01		7.65E-01
Uranium-238	7440-61-1	NA					4.50E+02	7.65E-01		7.65E-01
Actinium-227	14952-40-0	NA					1.50E+03	1.05E+00		1.05E+00
Americium-243	14993-75-0	NA					7.00E+02	7.70E-01		7.70E-01
Carbon-14	14762-75-5	NA						1.00E+00	Default	1.00E+00
Cobalt-60	10198-40-0	NA					4.50E+01	1.80E-01		1.80E-01
Curium-242	15510-73-3	NA					2.00E+03	3.40E-01		3.40E-01
Curium-243	15757-87-6	NA					2.00E+03	3.40E-01		3.40E-01
Curium-244	13981-15-2	NA					2.00E+03	3.40E-01		3.40E-01
Europium-152	14683-23-9	NA					6.50E+02	1.30E+00		1.30E+00
Iodine-129	15046-84-1	NA					6.00E+01	1.80E+00		1.80E+00
Neptunium-237	13994-20-2	NA					3.00E+01	6.00E-01		6.00E-01
Nickel-59	14336-70-0	NA	6.10E+01				1.50E+02	9.60E-01		6.10E+01
Niobium-93	None	NA					3.50E+02	1.40E+00		1.40E+00
Plutonium-238	13981-16-3	NA					4.50E+03	4.05E-01		4.05E-01
Plutonium-240	14119-33-6	NA					4.50E+03	4.05E-01		4.05E-01

Table C2-6. Aquatic Water-to-Plant Transfer Factors (WP) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) WP[b]	WP Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance WP	Soil Kd[d]	SAIC Computed WP[e]	Comments	Recommended WP[f]
Plutonium-242	13982-10-0	NA					4.50E+07	4.05E+03		4.05E+03
Protactinium-231	14331-85-2	NA					2.50E+03	1.25E+00		1.25E+00
Radium-226	13982-63-3	NA					4.50E+02	1.35E+00		1.35E+00
Radium-228	15262-20-1	NA					4.50E+02	1.35E+00		1.35E+00
Ruthenium-106	13967-48-1	NA					3.50E+02	5.25E+00		5.25E+00
Selenium-79	None	NA	1.85E+03				3.00E+02	1.50E+00		1.85E+03
Thorium-229	15594-54-4	NA					1.50E+05	2.55E+01		2.55E+01
Thorium-232	7440-29-1	NA					1.50E+05	2.55E+01		2.55E+01
Tin-126	15832-50-5	NA					2.50E+02	1.50E+00		1.50E+00
Zirconium-93	15751-77-6	NA					3.00E+03	1.20E+00		1.20E+00

NA = Not applicable

[a] $\log_{10}$ of K_{ow} values in Table 4.1

[b] Data published in Appendix C, Table C-4 of EPA (1999)

[c] Calculated or chosen as described in Appendix C, Sect. C-1.4 of EPA (1999)

[d] Soil Kd values from Appendix B Table B-1-1 or from Baes and others. (1984) Fig. 2.31 for radionuclides and other elements not shown in Appendix B Table B-1-1

[e] Calculated by using the Kd and the recommended SPv values for plants in soil (Table C.2-1): WP = SPv x Kd

[f] Selection criteria described in Sect. 8.2.4.4

[g] Calculated by using log K_{ow}: log WP = 0.819 x log K_{ow} -1.146 (Southworth, Beauchamp, and Schmieder 1978)

[h] Calculated by using the WP for TCDD from Appendix C, Table C-4 of EPA (1999) and BEFs for other congeners

[i] Value for Aroclor 1254 from Appendix C of EPA (1999), Table C-4, was used for PCB mixtures

[j] Benzo(a)pyrene value from Appendix C of EPA (1999), Table C-4, was used as a surrogate value

[k] 2,4-dinitrotoluene value from Appendix C of EPA (1999), Table C-4, was used as a surrogate value

Table C2-7. Aquatic Sediment-to-Plant Transfer Factors (SP) for Ecological Receptors (mg/kg tissue wet weight / mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SP[b] (per dry weight of plants)	Calculated from EPA (1999) SP[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SP	Soil Kd[d]	Sediment Kd[e]	SAIC Computed SP[f]	Comments	Recommended SP[g]
Organic Compounds											
Aromatic Halogenated Hydrocarbons											
4-Chloro-3-methylphenol	59-50-7	3.10		1.25E-01	h		6.03E+00	2.41E+01	3.13E-02		3.13E-02
2,3,4,6-Tetrachlorophenol	58-90-2	4.42		2.16E-02	h		6.61E+01	2.64E+02	5.40E-03		5.40E-03
Aromatic Nonhalogenated Hydrocarbons											
2-Nitrotoluene	88-72-2	2.30		3.63E-01	h		4.27E+00	1.71E+01	9.07E-02		9.07E-02
4-Nitrobiphenyl	92-93-3	3.77		5.13E-02	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Benzaldehyde	100-52-7	1.48		1.08E+00	h		2.01E-01	8.04E-01	2.71E-01		2.71E-01
Benzene	71-43-2	-4.99		5.94E+03	h		6.20E+01	4.65E+00	7.92E+04		7.92E+04
Benzyl alcohol	100-51-6	1.10		1.79E+00	h		1.02E-01	4.09E-01	4.47E-01		4.47E-01
Ethyl benzene	100-41-4	3.12		1.21E-01	h		2.04E+00	8.16E+00	3.03E-02		3.03E-02
m-Xylene	108-38-3	3.20		1.09E-01	h		1.96E+00	7.84E+00	2.73E-02		2.73E-02
o-Xylene	95-47-6	3.13		1.20E-01	h		2.41E+00	9.64E+00	3.00E-02		3.00E-02
p-Xylene	106-42-3	3.17		1.14E-01	h		3.11E+00	1.24E+01	2.86E-02		2.86E-02
Styrene	100-42-5	2.93		1.57E-01	h		9.12E+00	3.65E+01	3.92E-02		3.92E-02
Toluene	108-88-3	2.67		2.22E-01	h		1.40E+00	5.60E+00	5.56E-02		5.56E-02
Non-aromatic Nonhalogenated Hydrocarbons											
1,2-Epoxybutane	106-88-7	1.44		1.14E+00	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
1,3-Butadiene	106-99-0	1.90		6.18E-01	h		7.41E-01	2.96E+00	1.54E-01		1.54E-01
1,4-Dioxane	123-91-1	-0.27	5.53E+01	1.11E+01	k		8.76E-03	3.50E-02	2.77E+00		2.77E+00
1-Methylpropyl alcohol	78-92-2	0.61		3.44E+00	h		4.00E-02	1.60E-01	8.60E-01		8.60E-01
1-Nitropropane	108-03-2	0.87		2.43E+00	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
2,2,4-Trimethylpentane	540-84-1	5.02		9.72E-03	h		8.64E+02	3.46E+03	2.43E-03		2.43E-03
2-Butanone	78-93-3	0.28		5.33E+00	h		2.34E-02	9.36E-02	1.33E+00		1.33E+00
2-Butenaldehyde (2-Butenal)	4170-30-3	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
2-Ethoxyethanol	110-80-5	-0.10		8.85E+00	h		2.09E-01	8.36E-01	2.21E+00		2.21E+00
2-Heptanone	110-43-0	1.98		5.55E-01	h		8.88E-01	3.55E+00	1.39E-01		1.39E-01
2-Hexanone	591-78-6	1.38		1.23E+00	h		1.34E+00	5.36E+00	3.09E-01		3.09E-01

Table C2-7. Aquatic Sediment-to-Plant Transfer Factors (SP) for Ecological Receptors (mg/kg tissue wet weight / mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SP[b] (per dry weight of plants)	Calculated from EPA (1999) SP[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SP	Soil Kd[d]	Sediment Kd[e]	SAIC Computed SP[f]	Comments	Recommended SP[g]
2-Methoxyethanol	109-86-4	0.25		5.54E+00	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
2-Methyl-2-propanol	75-65-0	0.35		4.86E+00	h		3.72E-01	1.49E+00	1.22E+00		1.22E+00
2-Methyl-2-propenenitrile	126-98-7	0.54		3.77E+00	h		3.74E-02	1.49E-01	9.47E-01		9.47E-01
2-Methylaziridine	75-55-8	-0.60		1.73E+01	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
2-Methylpropyl alcohol	78-83-1	0.76		2.82E+00	h		5.61E-02	2.24E-01	7.04E-01		7.04E-01
2-Pentanone	107-87-9	0.91		2.31E+00	h		7.40E-01	2.96E+00	5.77E-01		5.77E-01
2-Propanone (Acetone)	67-64-1	-0.22	5.20E+01	1.04E+01	k		9.51E-03	3.80E-02	2.60E+00		2.60E+00
2-Propene-1-ol	107-18-6	0.17		6.18E+00	h		1.48E-02	5.90E-02	1.54E+00		1.54E+00
2-Propyl alcohol	67-63-0	0.05		7.25E+00	h		1.12E-02	4.50E-02	1.81E+00		1.81E+00
3-Heptanone	106-35-4	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
3-Methyl-1-butanol	123-51-3	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
3-Methyl-2-butanone	563-80-4	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
3-Pentanone	96-22-0	0.99		2.07E+00	h		1.20E-01	4.81E-01	5.19E-01		5.19E-01
4-Heptanone	123-19-3	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
4-Methyl-2-pentanone	108-10-1	1.19		1.59E+00	h		1.20E-01	4.80E-01	3.97E-01		3.97E-01
4-Methyl-3-penten-2-one	141-79-7	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
5-Methyl-2-hexanone	110-12-3	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
Acetaldehyde	75-07-0	-0.22		1.04E+01	h		9.53E-03	3.81E-02	2.60E+00		2.60E+00
Acetamide	60-35-5	-1.26		4.14E+01	h		2.82E-04	1.13E-03	1.04E+01		1.04E+01
Acetic acid	64-19-7	-0.17		9.71E+00	h		1.00E-02	4.00E-02	2.43E+00		2.43E+00
Acetic acid ethyl ester	141-78-6	0.73		2.93E+00	h		2.30E-02	9.18E-02	7.33E-01		7.33E-01
Acetic acid n-butyl ester	123-86-4	1.73		7.75E-01	h		5.04E-01	2.02E+00	1.94E-01		1.94E-01
Acetonitrile	75-05-8	-0.34		1.22E+01	h		7.69E-03	3.07E-02	3.05E+00		3.05E+00
Acrolein	107-02-8	-0.01		7.84E+00	h		1.39E-02	5.57E-02	1.96E+00		1.96E+00
Acrylonitrile	107-13-1	0.25	2.78E+01	5.55E+00	k		2.22E-02	8.88E-02	1.39E+00		1.39E+00
Bis(isopropyl)ether	108-20-3	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
Butane	106-97-8	2.89		1.65E-01	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Carbon disulfide	75-15-0	2.00		5.41E-01	h		5.14E-01	2.06E+00	1.35E-01		1.35E-01

Table C2-7. Aquatic Sediment-to-Plant Transfer Factors (SP) for Ecological Receptors (mg/kg tissue wet weight / mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SP[b] (per dry weight of plants)	Calculated from EPA (1999) SP[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SP	Soil Kd[d]	Sediment Kd[e]	SAIC Computed SP[f]	Comments	Recommended SP[g]
Cyanogen	460-19-5	0.81		2.64E+00	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Cyclohexane	110-82-7	3.44		7.96E-02	h		4.79E+00	1.91E+01	1.99E-02		1.99E-02
Cyclohexanone	108-94-1	0.81		2.64E+00	h		6.28E-02	2.51E-01	6.59E-01		6.59E-01
Cyclohexene	110-83-8	2.86		1.72E-01	h		6.51E+00	2.60E+01	4.30E-02		4.30E-02
Cyclopentane	287-92-3	3.00		1.43E-01	h		8.93E+00	3.57E+01	3.57E-02		3.57E-02
Ethyl alcohol	64-17-5	0.31		5.13E+00	h		2.03E-02	8.11E-02	1.28E+00		1.28E+00
Ethyl ether	60-29-7	0.89		2.37E+00	h		7.53E-02	3.01E-01	5.92E-01		5.92E-01
Ethyl methacrylate	97-63-2	1.59		9.33E-01	h		2.46E-01	9.80E-01	2.34E-01		2.34E-01
Formaldehyde	50-00-0	0.34	2.46E+01	4.91E+00	k		2.62E-02	1.05E-01	1.23E+00		1.23E+00
Formamide	75-12-7	-1.51		5.78E+01	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Formic acid	64-18-6	-0.54		1.58E+01	h		5.39E-02	2.16E-01	3.95E+00		3.95E+00
Formic acid, methyl ester	107-31-3	-0.26		1.10E+01	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Glycidylaldehyde	765-34-4	-0.73		2.05E+01	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Methyl acetate	79-20-9	0.18		6.10E+00	h		4.80E-02	1.92E-01	1.52E+00		1.52E+00
Methyl alcohol	67-56-1	-0.71		1.99E+01	h		3.96E-03	1.58E-02	4.99E+00		4.99E+00
Methyl isocyanate	624-83-9	No data		No data			No Kd	No Kd	NA	Reacts with water	NA
Methyl methacrylate	80-62-6	0.79		2.71E+00	h		6.31E-01	2.52E+00	6.77E-01		6.77E-01
Methyl tert-butyl ether	1634-04-4	0.94		2.22E+00	h		8.43E-02	3.37E-01	5.54E-01		5.54E-01
Methylacetylene	74-99-7	0.94		2.22E+00	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Methylcyclohexane	108-87-2	4.10		3.31E-02	h		1.08E+02	4.31E+02	8.26E-03		8.26E-03
N,N-Dimethylacetamide	127-19-5	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
n-Butyl alcohol	71-36-3	0.88		2.40E+00	h		7.36E-02	2.95E-01	6.00E-01		6.00E-01
n-Heptane	142-82-5	4.66		1.57E-02	h		3.83E+02	1.53E+03	3.92E-03		3.92E-03
n-Hexane	110-54-3	4.11		3.26E-02	h		1.10E+02	4.41E+02	8.15E-03		8.15E-03
Nitromethane	75-52-5	-0.35		1.23E+01	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
n-Nonane	111-84-2	5.65		4.20E-03	h		3.60E+03	1.44E+04	1.05E-03		1.05E-03
n-Octane	111-65-9	4.00		3.78E-02	h		8.59E+01	3.44E+02	9.44E-03		9.44E-03
n-Pentane	109-66-0	3.21		1.08E-01	h		1.44E+01	5.75E+01	2.70E-02		2.70E-02

Table C2-7. Aquatic Sediment-to-Plant Transfer Factors (SP) for Ecological Receptors (mg/kg tissue wet weight / mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{oc}[a]	EPA (1999) SP[b] (per dry weight of plants)	Calculated from EPA (1999) SP[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SP	Soil Kd[d]	Sediment Kd[e]	SAIC Computed SP[f]	Comments	Recommended SP[g]
n-Propionaldehyde	123-38-6	0.59		3.53E+00	h		3.82E-02	1.53E-01	8.83E-01		8.83E-01
n-Propyl alcohol	71-23-8	0.25		5.55E+00	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
n-Valeraldehyde	110-62-3	No data		No data	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Oxirane	75-21-8	-0.30		1.15E+01	h		8.26E-03	3.30E-02	2.89E+00		2.89E+00
p-Cymene	99-87-6	4.10		3.31E-02	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Phosgene	75-44-5	No data		No data	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Propargyl alcohol	107-19-7	0.26		5.48E+00	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Propionic acid	79-09-4	0.33		4.99E+00	h		2.12E-02	8.48E-02	1.25E+00		1.25E+00
Propionitrile	107-12-0	0.16		6.26E+00	h		1.44E-02	5.77E-02	1.56E+00		1.56E+00
Propylene gylcol monomethyl ether	107-98-2	-0.18		9.84E+00	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
p-tert-Butyltoluene	98-51-1	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
Triethylamine	121-44-8	0.16		6.26E+00	h		1.44E-02	5.77E-02	1.56E+00		1.56E+00
Trimethylamine	75-50-3	0.16		6.26E+00	h		4.00E-02	1.60E-01	1.56E+00		1.56E+00
Vinyl acetate	108-05-4	0.70		3.06E+00	h		4.97E-02	1.99E-01	7.63E-01		7.63E-01
Non-aromatic Halogenated Hydrocarbons											
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
1,1,1,2-Tetrachloroethane	630-20-6	2.63		2.34E-01	h		1.59E+00	6.37E+00	5.83E-02		5.83E-02
1,1,1-Trichloroethane	71-55-6	2.42		3.09E-01	h		1.35E+03	5.40E+03	7.71E-02		7.71E-02
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	3.73		5.41E-02	h		3.16E+00	1.26E+01	1.35E-02		1.35E-02
1,1,2,2-Tetrachloroethane	79-34-5	4.64		1.60E-02	h		7.90E-01	3.16E+00	4.01E-03		4.01E-03
1,1,2,2-Tetrachloroethene	127-18-4	2.55		2.62E-01	h		2.65E+00	1.06E+01	6.54E-02		6.54E-02
1,1,2-Trichloroethane	79-00-5	2.10		4.75E-01	h		7.50E-01	3.00E+00	1.19E-01		1.19E-01
1,1,2-Trichloroethylene	79-01-6	2.43		3.04E-01	h		9.40E-01	3.76E+00	7.60E-02		7.60E-02
1,1-Dichloroethane	75-34-3	1.79		7.13E-01	h		5.30E-01	2.12E+00	1.78E-01		1.78E-01
1,1-Dichloroethene	75-35-4	2.12		4.61E-01	h		6.50E-01	2.60E+00	1.15E-01		1.15E-01
1,2,2-Trichloro-1,1,2-trifluoroethane	76-13-1	3.16		1.15E-01	h		2.57E+00	1.03E+01	2.89E-02		2.89E-02
1,2,3-Trichloropropane	96-18-4	2.25		3.88E-01	h		8.10E-01	3.22E+00	9.75E-02		9.75E-02
1,2-Dibromo-3-chloropropane	96-12-8	2.34		3.44E-01	h		9.47E-01	3.79E+00	8.59E-02		8.59E-02

Table C2-7. Aquatic Sediment-to-Plant Transfer Factors (SP) for Ecological Receptors (mg/kg tissue wet weight / mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SP[b] (per dry weight of plants)	Calculated from EPA (1999) SP[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SP	Soil Kd[d]	Sediment Kd[e]	SAIC Computed SP[f]	Comments	Recommended SP[g]
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	2.82		1.82E-01	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
1,2-Dichloroethane	107-06-2	1.46		1.11E+00	h		1.96E-01	7.83E-01	2.77E-01		2.77E-01
1,2-Dichloroethylene	540-59-0	0.48		4.09E+00	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
1,2-Dichloropropane	78-87-5	2.25		3.88E-01	h		4.70E-01	1.88E+00	9.69E-02		9.69E-02
1,3-Dichloropropene	542-75-6	1.75		7.56E-01	h		2.70E-01	1.08E+00	1.89E-01		1.89E-01
1,4-Dichloro-2-butene	764-41-0	0.87		2.42E+00	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
1-Chloroethene	75-01-4	1.15		1.68E+00	h		1.11E-01	4.44E-01	4.21E-01		4.21E-01
2,2-Dichloropropionic acid	75-99-0	0.78		2.75E+00	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
2-Chloropropane	75-29-6	1.90		6.18E-01	h		7.41E-01	2.96E+00	1.54E-01		1.54E-01
3-Chloropropene (allyl chloride)	107-05-1	0.95		2.17E+00	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Bromochloromethane	74-97-5	1.41		1.19E+00	h		2.44E-01	9.77E-01	2.96E-01		2.96E-01
Bromodichloromethane	75-27-4	2.03		5.23E-01	h		5.38E-01	2.15E+00	1.31E-01		1.31E-01
Bromoethene	593-60-2	1.07		1.87E+00	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Bromoform	75-25-2	2.35		3.39E-01	h		1.26E+00	5.04E+00	8.48E-02		8.48E-02
Bromomethane	74-83-9	1.11		1.76E+00	h		9.00E-02	3.60E-01	4.40E-01		4.40E-01
Carbon tetrachloride	56-23-5	2.72	1.04E+00	2.08E-01	k		1.52E+00	6.08E+00	5.20E-02		5.20E-02
Chlorodibromomethane	124-48-1	2.18		4.28E-01	h		7.05E-01	2.82E+00	1.07E-01		1.07E-01
Chlorodifluoromethane	75-45-6	1.08		1.84E+00	h		9.83E-02	3.93E-01	4.61E-01		4.61E-01
Chloroethane	75-00-3	3.10		1.25E-01	h		3.71E+00	1.48E+01	3.13E-02		3.13E-02
Chloroform	67-66-3	1.95	2.90E+00	5.80E-01	k		5.30E+01	2.12E+00	1.45E+01		1.45E+01
Chloromethane	74-87-3	0.90		2.33E+00	h		6.00E-02	2.40E-01	5.82E-01		5.82E-01
Chloropentafluoroethane	76-15-3	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
cis-1,2-Dichloroethene	156-59-2	1.98		5.54E-01	h		4.98E+00	1.99E+01	1.39E-01		1.39E-01
cis-1,3-Dichloropropene	10061-01-5	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
Cyanogen bromide	506-68-3	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
Cyanogen chloride	506-77-4	0.20		5.94E+00	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Dichlorodifluoromethane	75-71-8	2.16		4.38E-01	h		6.85E-01	2.74E+00	1.10E-01		1.10E-01
Dichlorofluoromethane	75-43-4	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00

Table C2-7. Aquatic Sediment-to-Plant Transfer Factors (SP) for Ecological Receptors (mg/kg tissue wet weight / mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SP[b] (per dry weight of plants)	Calculated from EPA (1999) SP[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SP	Soil Kd[d]	Sediment Kd[e]	SAIC Computed SP[f]	Comments	Recommended SP[g]
Dichloromethane	75-09-2	1.26		1.46E+00	h		1.00E-01	4.00E-01	3.64E-01		3.64E-01
Difluorodibromomethane	75-61-6	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
Hexafluoroacetone	684-16-2	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
Iodomethane	74-88-4	1.69		8.17E-01	h		4.61E-01	1.84E+00	2.04E-01		2.04E-01
Methylene bromide	74-95-3	1.62		8.97E-01	h		2.60E-01	1.04E+00	2.24E-01		2.24E-01
Pentachloroethane	76-01-7	3.05		1.34E-01	h		1.00E+01	4.00E+01	3.34E-02		3.34E-02
trans-1,2-Dichloroethene	156-60-5	1.98		5.54E-01	h		3.80E-01	1.52E+00	1.38E-01		1.38E-01
trans-1,3-Dichloropropene	10061-02-6	2.06		4.99E-01	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Trichloroacetic acid	76-03-9	1.33		1.32E+00	h		2.04E-01	8.16E-01	3.30E-01		3.30E-01
Trichlorofluoroethane	27154-33-2	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
Trichlorofluoromethane	75-69-4	2.53		2.67E-01	h		1.34E+00	5.34E+00	6.69E-02		6.69E-02
Trifluorobromomethane	75-63-8	1.86		6.52E-01	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Dioxin and Furan Compounds											
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	35822-46-9	8.20	2.90E-04	5.80E-05	i		9.77E+05	3.91E+06	1.45E-05	h	1.45E-05
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	7.92	6.20E-05	1.24E-05	i		5.13E+05	2.05E+06	3.10E-06	h	3.10E-06
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	7.92	2.20E-03	4.40E-04	i		5.13E+05	2.05E+06	1.10E-04	h	1.10E-04
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	39227-28-6	7.79	1.70E-03	3.40E-04	i		3.80E+05	1.52E+06	8.50E-05	h	8.50E-05
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	7.25	4.30E-04	8.60E-05	i		1.10E+05	4.39E+05	2.15E-05	h	2.15E-05
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	57653-85-7	7.25	6.70E-04	1.34E-04	i		1.10E+05	4.39E+05	3.36E-05	h	3.36E-05
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	7.25	1.10E-03	2.20E-04	i		1.10E+05	4.39E+05	5.51E-05	h	5.51E-05
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	19408-74-3	7.25	7.80E-04	1.56E-04	i		1.10E+05	4.39E+05	3.91E-05	h	3.91E-05
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	7.25	3.50E-03	7.00E-04	i		1.10E+05	4.39E+05	1.75E-04	h	1.75E-04
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	40321-76-4	6.64	5.20E-03	1.04E-03	i		2.69E+04	1.08E+05	2.59E-04	h	2.59E-04
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	6.79	1.10E-03	2.20E-04	i		3.80E+04	1.52E+05	5.50E-05	h	5.50E-05
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	7.25	3.80E-03	7.60E-04	i		1.10E+05	4.39E+05	1.90E-04	h	1.90E-04
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	6.92	9.00E-03	1.80E-03	i		5.13E+04	2.05E+05	4.50E-04	h	4.50E-04
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1746-01-6	6.64	5.62E-03	1.12E-03	h		2.69E+04	1.08E+05	2.80E-04		2.80E-04
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	6.53	4.50E-03	9.00E-04	i		2.09E+04	8.36E+04	2.25E-04	h	2.25E-04

Table C2-7. Aquatic Sediment-to-Plant Transfer Factors (SP) for Ecological Receptors (mg/kg tissue wet weight / mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SP[b] (per dry weight of plants)	Calculated from EPA (1999) SP[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SP	Soil Kd[d]	Sediment Kd[e]	SAIC Computed SP[f]	Comments	Recommended SP[g]
Dibenzofuran	132-64-9	4.33		2.43E-02	h		1.81E+02	7.25E+02	6.08E-03		6.08E-03
Octachlorodibenzo(p)dioxin	3268-87-9	8.20	6.70E-05	1.34E-05	i		2.24E+01	8.95E+01	3.35E-06	h	3.35E-06
Octachlorodibenzofuran	39001-02-0	8.78	9.00E-05	1.80E-05	i		3.72E+06	1.49E+07	4.49E-06	h	4.49E-06
PCBs											
Polychlorinated biphenyls (PCBs)[j]	1336-36-3	7.31	1.00E-02	2.00E-03	h		3.09E+03	1.24E+04	5.00E-04		5.00E-04
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	7.08		6.26E-04	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	7.12		5.94E-04	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	No data		No data	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	No data		No data	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	No data		No data	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	No data		No data	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	No data		No data	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	No data		No data	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
2',3,4,4',5-Pentachlorobiphenyl	no cas #	No data		No data	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	7.12		5.94E-04	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	7.41		4.05E-04	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
3,3',4,4',5-Pentachlorobiphenyl	no cas #	No data		No data	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	No data		No data	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	No data		No data	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Phthalates											
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	5.20	3.80E-02	7.60E-03	h		1.11E+03	4.44E+03	1.90E-03		1.90E-03
Butylbenzyl phthalate	85-68-7	4.41		2.18E-02	h		1.37E+02	5.50E+02	5.43E-03		5.43E-03
Dibutyl phthalate	84-74-2	4.72		1.45E-02	h		1.57E+01	6.27E+01	3.63E-03		3.63E-03
Diethyl phthalate	84-66-2	4.44		2.11E-02	h		8.20E-01	3.28E+00	5.28E-03		5.28E-03
Dimethylphthalate	131-11-3	1.63		8.81E-01	h		2.66E+00	1.06E+01	2.21E-01		2.21E-01
n-Dioctyl phthalate	117-84-0	9.33	1.57E-04	3.14E-05	h		9.03E+06	3.61E+07	7.85E-06		7.85E-06
Light Polycyclic Aromatic Hydrocarbons (molecular weight <200 g/mole)											
2-Chloronapthalene	91-58-7	4.07		3.45E-02	h		7.14E+01	2.86E+02	8.61E-03		8.61E-03

Table C2-7. Aquatic Sediment-to-Plant Transfer Factors (SP) for Ecological Receptors (mg/kg tissue wet weight / mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SP[b] (per dry weight of plants)	Calculated from EPA (1999) SP[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SP	Soil Kd[d]	Sediment Kd[e]	SAIC Computed SP[f]	Comments	Recommended SP[g]
2-Methyl naphthalene	91-57-6	3.86		4.55E-02	h		6.26E+01	2.50E+02	1.14E-02		1.14E-02
5-Nitroacenaphthene	602-87-9	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
Acenaphthene	83-32-9	3.96		3.96E-02	h		4.90E+01	1.96E+02	9.89E-03		9.89E-03
Acenaphthylene	208-96-8	4.07		3.44E-02	h		6.76E+01	2.70E+02	8.60E-03		8.60E-03
Anthracene	120-12-7	4.47		2.02E-02	h		2.35E+02	9.40E+02	5.05E-03		5.05E-03
Fluorene	86-73-7	4.18		2.97E-02	h		1.41E+02	5.65E+02	7.43E-03		7.43E-03
Indene	95-13-6	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
Naphthalene	91-20-3	3.37		8.70E-02	h		1.19E+01	4.76E+01	2.17E-02		2.17E-02
Phenanthrene	85-01-8	4.55		1.82E-02	h		5.01E+02	2.01E+03	4.53E-03		4.53E-03
Pyrene	129-00-0	5.00		9.98E-03	h		6.80E+02	2.72E+03	2.49E-03		2.49E-03
Heavy Polycyclic Aromatic Hydrocarbons (molecular weight >200 g/mole)											
3-Methylcholanthrene	56-49-5	7.11		6.02E-04	h		1.51E+04	6.05E+04	1.50E-04		1.50E-04
5-Methylchrysene	3697-24-3	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
Benzo(a)anthracene	56-55-3	5.68	2.02E-02	4.04E-03	h		2.60E+03	1.04E+04	1.01E-03		1.01E-03
Benzo(a)pyrene	50-32-8	6.13	1.11E-02	2.22E-03	h		9.69E+03	3.87E+04	5.56E-04		5.56E-04
Benzo(b)fluoranthene	205-99-2	6.20	1.01E-02	2.02E-03	h		8.36E+03	3.34E+04	5.06E-04		5.06E-04
Benzo(e)pyrene	192-97-2	7.40		4.09E-04	h		1.58E+05	6.34E+05	1.02E-04		1.02E-04
Benzo(g,h,i)perylene	191-24-2	7.10		6.10E-04	h		1.82E+04	7.28E+04	1.52E-04		1.52E-04
Benzo(j)fluoranthene	205-82-3	6.44		1.47E-03	h		2.15E+04	8.60E+04	3.67E-04		3.67E-04
Benzo(k)fluoranthene	207-08-9	6.20	1.01E-02	2.02E-03	h		8.32E+03	3.33E+04	5.05E-04		5.05E-04
Benzo[a,i]pyrene	191-30-0	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
Chrysene	218-01-9	5.74	1.87E-02	3.74E-03	h		2.97E+03	1.19E+04	9.33E-04		9.33E-04
Dibenz(a,h)anthracene	53-70-3	6.55	6.40E-03	1.28E-03	h		1.79E+04	7.16E+04	3.20E-04		3.20E-04
Dibenz[a,h]acridine	226-36-8	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
Dibenz[a,j]acridine	224-42-0	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
Dibenzo(a,e)fluoranthene	5385-75-1	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
Dibenzo(a,h)fluoranthene	no cas #	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
Dibenzo[a,e]pyrene	192-65-4	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00

Table C2-7. Aquatic Sediment-to-Plant Transfer Factors (SP) for Ecological Receptors (mg/kg tissue wet weight / mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SP[b] (per dry weight of plants)	Calculated from EPA (1999) SP[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SP	Soil Kd[d]	Sediment Kd[e]	SAIC Computed SP[f]	Comments	Recommended SP[g]
Dibenzo[a,h]pyrene	189-64-0	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
Dibenzo[a,i]pyrene	189-55-9	7.29		4.74E-04	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Fluoranthene	206-44-0	5.08		8.94E-03	h		4.91E+02	1.96E+03	2.24E-03		2.24E-03
Hexachloronaphthalene	1335-87-1	7.59		3.18E-04	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Indeno(1,2,3-cd)pyrene	193-39-5	6.91	3.90E-03	7.80E-04	h		4.11E+04	1.64E+05	1.95E-04		1.95E-04
Octachloronaphthalene	2234-13-1	6.42		1.51E-03	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Pentachloronaphthalene	1321-64-8	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
Tetrachloronaphthalene	1335-88-2	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
Trichloronaphthalene	1321-65-9	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
Light Substituted Benzene Compounds (MW <200 g/mole)											
1,2,3-Trichlorobenzene	87-61-6	4.05		3.55E-02	h		2.02E+02	8.10E+02	8.87E-03		8.87E-03
1,2,4-Trichlorobenzene	120-82-1	3.99		3.84E-02	h		1.66E+01	6.64E+01	9.59E-03		9.59E-03
1,2,4-Trimethyl benzene	95-63-6	3.65		6.02E-02	h		3.89E+01	1.56E+02	1.50E-02		1.50E-02
1,2-Dichlorobenzene	95-50-1	3.45		7.90E-02	h		3.79E+00	1.52E+01	1.97E-02		1.97E-02
1,3,5-Trimethyl benzene	108-67-8	3.42		8.17E-02	h		1.67E+01	6.69E+01	2.04E-02		2.04E-02
1,3-Dichlorobenzene	541-73-1	3.53		7.06E-02	h		8.03E+01	3.21E+02	1.77E-02		1.77E-02
1,3-Dinitrobenzene	99-65-0	1.49	5.32E+00	1.06E+00	h		2.06E-01	8.25E-01	2.66E-01		2.66E-01
1,4-Dichlorobenzene	106-46-7	3.41		8.26E-02	h		6.16E+00	2.46E+01	2.07E-02		2.07E-02
1,4-Dinitrobenzene	100-25-4	1.50		1.05E+00	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
2,4,5-Trichlorophenol	95-95-4	3.87		4.49E-02	h		1.13E+01	4.51E+01	1.13E-02		1.13E-02
2,4,6-Trichlorophenol	88-06-2	3.71		5.54E-02	h		2.26E+00	9.05E+00	1.38E-02		1.38E-02
2,4-Dichlorophenol	120-83-2	3.04		1.36E-01	h		1.40E+00	5.58E+00	3.41E-02		3.41E-02
2,4-Dimethylphenol	105-67-9	2.36		3.35E-01	h		1.26E+00	5.04E+00	8.38E-02		8.38E-02
2,4-Dinitrophenol	51-28-5	1.52		1.03E+00	h		1.00E-04	4.00E-04	2.57E-01		2.57E-01
2,4-Dinitrotoluene	121-14-2	2.00	2.72E+00	5.44E-01	h		5.10E-01	2.04E+00	1.36E-01		1.36E-01
2,6-Dinitrotoluene	606-20-2	1.89	3.15E+00	6.30E-01	h		4.19E-01	1.68E+00	1.57E-01		1.57E-01
2-Chlorophenol	95-57-8	2.16		4.36E-01	h		3.87E+00	1.55E+01	1.09E-01		1.09E-01
2-Chlorotoluene	95-49-8	3.54		7.00E-02	h		No Kd	No Kd	1.00E+00	Default	1.00E+00

Table C2-7. Aquatic Sediment-to-Plant Transfer Factors (SP) for Ecological Receptors (mg/kg tissue wet weight / mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) SP [b] (per dry weight of plants)	Calculated from EPA (1999) SP [c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SP	Soil Kd [d]	Sediment Kd [e]	SAIC Computed SP [f]	Comments	Recommended SP [g]
2-Nitrophenol	88-75-5	1.79		7.15E-01	h		3.53E+00	1.41E+01	1.79E-01		1.79E-01
4,6-Dinitro-o-cresol	534-52-1	2.85		1.74E-01	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
4-Chlorotoluene	106-43-4	3.33		9.21E-02	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
4-Nitrophenol	100-02-7	1.91		6.10E-01	h		4.37E+02	1.75E+01	1.52E+01		1.52E+01
alpha-Methylstyrene	98-83-9	3.46		7.71E-02	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Aniline	62-53-3	0.98		2.10E+00	h		8.23E-02	3.29E-01	5.26E-01		5.26E-01
Benzotrichloride	98-07-7	2.92		1.59E-01	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Benzyl chloride	100-44-7	0.36		4.79E+00	h		2.71E-02	1.08E-01	1.20E+00		1.20E+00
Bromobenzene	108-86-1	2.99		1.45E-01	h		4.47E+00	1.79E+01	3.62E-02		3.62E-02
Chlorobenzene	108-90-7	2.79		1.89E-01	h		2.24E+00	8.96E+00	4.73E-02		4.73E-02
Cumene	98-82-8	3.61		6.32E-02	h		9.13E+01	3.72E+02	1.55E-02		1.55E-02
m-Cresol	108-39-4	1.96		5.71E-01	h		4.78E-01	1.91E+00	1.43E-01		1.43E-01
n-Butyl benzene	104-51-8	4.28		2.60E-02	h		2.51E+01	1.00E+02	6.50E-03		6.50E-03
Nitrobenzene	98-95-3	1.83	3.38E+00	6.76E-01	h		1.19E+00	4.76E+04	1.69E-05		1.69E-05
n-Propyl benzene	103-65-1	3.69		5.70E-02	h		7.24E+00	2.90E+01	1.43E-02		1.43E-02
o-Cresol	95-48-7	2.02		5.26E-01	h		5.34E-01	2.14E+00	1.31E-01		1.31E-01
o-Dinitrobenzene	528-29-0	1.60		9.21E-01	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
o-Nitroaniline	88-74-4	1.85		6.60E-01	h		3.93E+00	1.57E+01	1.65E-01		1.65E-01
o-Toluidine	95-53-4	1.34		1.30E+00	h		1.57E-01	6.28E-01	3.25E-01		3.25E-01
p-Chloroaniline	106-47-8	1.87		6.44E-01	h		4.06E-01	1.63E+00	1.60E-01		1.60E-01
p-Cresol	106-44-5	1.94		5.86E-01	h		4.61E-01	1.84E+00	1.47E-01		1.47E-01
Phenol	108-95-2	1.48		1.08E+00	h		2.20E-01	8.79E-01	2.71E-01		2.71E-01
p-Nitrochlorobenzene	100-00-5	2.39		3.22E-01	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
p-Toluidine	106-49-0	1.40		1.20E+00	h		2.39E-01	9.56E-01	3.00E-01		3.00E-01
sec-Butyl benzene	135-98-8	4.57		1.77E-02	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
tert-Butyl benzene	98-06-6	4.11		3.26E-02	h		1.10E+02	4.41E+02	8.15E-03		8.15E-03
Toluene-2,6-diamine	823-40-5	1.45		1.12E+00	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Trimethyl benzene	25551-13-7	3.40		8.39E-02	h		No Kd	No Kd	1.00E+00	Default	1.00E+00

Table C2-7. Aquatic Sediment-to-Plant Transfer Factors (SP) for Ecological Receptors (mg/kg tissue wet weight / mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SP[b] (per dry weight of plants)	Calculated from EPA (1999) SP[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SP	Soil Kd[d]	Sediment Kd[e]	SAIC Computed SP[f]	Comments	Recommended SP[g]
Other Light Semivolatile Compounds (molecular weight <200 g/mole)											
1,1'-Biphenyl	92-52-4	3.90		4.31E-02	h		2.51E+01	1.00E+02	1.08E-02		1.08E-02
1,1-Dimethylhydrazine	57-14-7	0.60		3.51E+00	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
1,2-Dimethylhydrazine	540-73-8	-1.37		4.78E+01	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
1,2-Diphenylhydrazine	122-66-7	2.94		1.55E-01	h		2.78E+00	1.11E+01	3.88E-02		3.88E-02
1,3-Propane sultone	1120-71-4	-0.52		1.55E+01	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
2,4-Toluene diisocyante	584-84-9	No data		No data			No Kd	No Kd	NA	Decomposes rapidly in water	NA
2-Chloroacetophenone	532-27-4	2.59		2.45E-01	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
2-Propenoic acid	79-10-7	0.33		4.99E+00	h		2.12E-02	8.48E-02	1.25E+00		1.25E+00
4,4-Methylenedianiline	101-77-9	3.38		8.62E-02	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Acetophenone	98-86-2	1.64		8.73E-01	h		2.69E-01	1.08E+00	2.17E-01		2.17E-01
Benzoic acid	65-85-0	1.86		6.52E-01	h		5.50E-03	2.20E-02	1.63E-01		1.63E-01
bis(2-Chloroethoxy)methane	111-91-1	7.59		3.18E-04	h		2.40E+05	9.60E+05	7.94E-05		7.94E-05
bis(2-Chloroethyl) ether	111-44-4	1.30		1.37E+00	h		7.60E-01	3.04E+00	3.43E-01		3.43E-01
Chlorocyclopentadiene	41851-50-7	2.43		3.05E-01	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Cyclohexanol	108-93-0	1.23		1.51E+00	h		1.30E-01	5.20E-01	3.77E-01		3.77E-01
Dichloroisopropyl ether	108-60-1	2.58		2.50E-01	h		6.10E-01	2.44E+00	6.25E-02		6.25E-02
Dichloromethyl ether	542-88-1	-0.38		1.28E+01	h		7.94E-01	3.18E+00	3.21E+00		3.21E+00
Dichloropentadiene	no cas #	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
Dimethyl sulfate	77-78-1	0.32		5.07E+00	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Dimethylaniline	121-69-7	-0.88		2.50E+01	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Di-n-propylnitrosamine	621-64-7	1.38		1.23E+00	h		1.70E-01	6.80E-01	3.08E-01		3.08E-01
Diphenyl ether	101-84-8	4.21		2.86E-02	h		1.38E+02	5.53E+02	7.14E-03		7.14E-03
Epichlorohydrin	106-89-8	0.25		5.55E+00	h		2.22E-02	8.88E-02	1.39E+00		1.39E+00
Ethyl carbamate (urethane)	51-79-6	-0.15		9.46E+00	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Ethyl methanesulfonate	62-50-0	0.05		7.25E+00	h		1.55E-02	6.19E-02	1.82E+00		1.82E+00
Ethylene dibromide	106-93-4	1.75		7.55E-01	h		3.28E-01	1.31E+00	1.89E-01		1.89E-01
Ethylene glycol	107-21-1	-0.91		2.62E+01	h		No Kd	No Kd	1.00E+00	Default	1.00E+00

Table C2-7. Aquatic Sediment-to-Plant Transfer Factors (SP) for Ecological Receptors (mg/kg tissue wet weight / mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SP[b] (per dry weight of plants)	Calculated from EPA (1999) SP[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SP	Soil Kd[d]	Sediment Kd[e]	SAIC Computed SP[f]	Comments	Recommended SP[g]
Ethylene glycol monobutyl ether	111-76-2	1.55		9.78E-01	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Ethylene glycol monoethyl ether acetate	111-15-9	0.62		3.40E+00	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Ethylene thiourea	96-45-7	-0.64		1.32E+01	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Furfural	98-01-1	0.96		2.16E+00	h		8.82E-02	3.53E-01	5.40E-01		5.40E-01
Maleic hydrazide	123-33-1	-0.74		2.07E+01	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Malononitrile	109-77-3	0.04		7.34E+00	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Methyl styrene (mixed isomers)	25013-15-4	3.35		8.97E-02	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Methylhydrazine	60-34-4	-1.06		3.17E+01	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
N,N-Diphenylamine	122-39-4	3.50		7.35E-02	h		3.47E+00	1.39E+01	1.84E-02		1.84E-02
Nitric acid, propyl ester	627-13-4	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
N-Nitrosodi-n-butylamine	924-16-3	2.41		3.13E-01	h		1.07E+00	4.29E+00	7.82E-02		7.82E-02
N-Nitrosomorpholine	59-89-2	0.98		2.10E+00	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
N-Nitroso-N,N-dimethylamine	62-75-9	-0.47		1.45E+01	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
o-Anisidine	90-04-0	1.18		1.61E+00	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Oxalic acid	144-62-7	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
Phthalic anhydride	85-44-9	-0.60		1.73E+01	h		4.80E-03	1.92E-02	4.31E+00		4.31E+00
p-Phthalic acid	100-21-0	0.82		2.59E+00	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Pyridine	110-86-1	0.67		3.17E+00	h		4.72E-02	1.89E-01	7.93E-01		7.93E-01
Quinoline	91-22-5	2.03		5.20E-01	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Quinone	106-51-4	0.20		5.94E+00	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Safrole	94-59-7	2.66		2.25E-01	h		1.68E+00	6.73E+00	5.61E-02		5.61E-02
Tetrahydrofuran	109-99-9	0.45		4.27E+00	h		3.16E-02	1.26E-01	1.07E+00		1.07E+00
Other Heavy Semivolatile Compounds (molecular weight >200 g/mole)											
1,2,4,5-Tetrachlorobenzene	95-94-3	4.64		1.61E-02	h		5.89E+01	2.36E+02	4.02E-03		4.02E-03
1,3,5-Trinitrobenzene	99-35-4	1.18		1.61E+00	h		1.18E-01	4.72E-01	4.03E-01		4.03E-01
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	4.17		3.01E-02	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	-2.70		2.82E+02	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
2-sec-Butyl-4,6-dinitrophenol	88-85-7	3.14		1.19E-01	h		No Kd	No Kd	1.00E+00	Default	1.00E+00

Table C2-7. Aquatic Sediment-to-Plant Transfer Factors (SP) for Ecological Receptors (mg/kg tissue wet weight / mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SP[b] (per dry weight of plants)	Calculated from EPA (1999) SP[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SP	Soil Kd[d]	Sediment Kd[e]	SAIC Computed SP[f]	Comments	Recommended SP[g]
3,3'-Dimethoxybenzidine	119-90-4	1.81		6.96E-01	h		3.65E-01	1.46E+00	1.74E-01		1.74E-01
3,3-Dichlorobenzidine	91-94-1	3.58		6.65E-02	h		8.70E+00	3.48E+01	1.66E-02		1.66E-02
4-Bromophenylphenyl ether	101-55-3	5.00		9.98E-03	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Ammonium perfluorooctanoate	3825-26-1	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
Azobenzene	103-33-3	3.82		4.80E-02	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
Captan	133-06-2	2.35		3.39E-01	h		2.00E+00	7.98E+00	8.49E-02		8.49E-02
Chlorobenzilate	510-15-6	4.38		2.28E-02	h		3.69E+01	1.48E+02	5.68E-03		5.68E-03
Dibutylphosphate	107-66-4	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
Dimethyl aminoazobenzene	60-11-7	4.58		1.74E-02	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Hexachlorobenzene	118-74-1	5.50	2.55E-02	5.10E-03	k		8.00E+02	3.20E+03	1.28E-03		1.28E-03
Hexachlorobutadiene	87-68-3	4.73	7.14E-02	1.43E-02	k		6.94E+01	2.77E+02	3.58E-03		3.58E-03
Hexachlorocyclopentadiene	77-47-4	5.04	5.65E-02	1.13E-02	k		9.51E+01	3.80E+02	2.83E-03		2.83E-03
Hexachloroethane	67-72-1	3.98		3.85E-02	h		1.82E+01	7.27E+01	9.64E-03		9.64E-03
Hexachlorophene	70-30-4	7.54		3.39E-04	h		1.08E+04	4.31E+04	8.50E-05		8.50E-05
Hexamethylene-1,5-diisocyanate	822-06-0	1.27		1.42E+00	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Mirex	2385-85-5	6.89		8.07E-04	h		1.00E+04	4.00E+04	2.02E-04		2.02E-04
Nitrofen	1836-75-5	5.53		4.93E-03	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Pentachlorobenzene	608-93-5	5.09	4.40E-02	8.80E-03	k		3.21E+02	1.29E+03	2.19E-03		2.19E-03
Pentachloronitrobenzene	82-68-8	4.64	8.00E-02	1.61E-02	k		5.89E+01	2.36E+02	4.02E-03		4.02E-03
Pentachlorophenol	87-86-5	5.08	4.49E-02	8.98E-03	k		1.99E+02	7.97E+02	2.25E-03		2.25E-03
Picric acid	88-89-1	2.03		5.20E-01	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Pronamide	23950-58-5	3.51		7.24E-02	h		7.74E+00	3.10E+01	1.81E-02		1.81E-02
Strychnine	57-24-9	1.93		5.94E-01	h		4.53E-01	1.81E+00	1.49E-01		1.49E-01
Terphenyls	26140-60-3	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00
Tributyl phosphate	126-73-8	4.00		3.78E-02	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Trifluralin	1582-09-8	5.34		6.35E-03	h		6.03E+01	2.41E+02	1.59E-03		1.59E-03
Triphenylamine	603-34-9	No data		No data			No Kd	No Kd	1.00E+00	Default	1.00E+00

Table C2-7. Aquatic Sediment-to-Plant Transfer Factors (SP) for Ecological Receptors (mg/kg tissue wet weight / mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SP[b] (per dry weight of plants)	Calculated from EPA (1999) SP[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SP	Soil Kd[d]	Sediment Kd[e]	SAIC Computed SP[f]	Comments	Recommended SP[g]
Herbicides and Organochlorinated Pesticides											
2,4,5-T	93-76-5	3.36		8.85E-02	h		5.25E-01	2.10E+00	2.21E-02		2.21E-02
2,4-D and esters	94-75-7	2.81		1.84E-01	h		2.00E-01	7.98E-01	4.60E-02		4.60E-02
4,4-DDD	72-54-8	6.20		2.02E-03	h		1.00E+03	4.00E+03	5.05E-04		5.05E-04
4,4-DDE	72-55-9	6.26	9.37E-03	1.87E-03	k		8.64E+06	3.46E+03	4.68E+00		4.68E+00
4,4-DDT	50-29-3	6.00		2.64E-03	h		2.40E+03	9.60E+03	6.59E-04		6.59E-04
Aldrin	309-00-2	6.18		2.08E-03	h		4.87E+02	1.95E+03	5.19E-04		5.19E-04
alpha-BHC	319-84-6	3.80		4.93E-02	h		1.76E+01	7.05E+01	1.23E-02		1.23E-02
beta-BHC	319-85-7	3.83		4.71E-02	h		2.14E+01	8.56E+01	1.18E-02		1.18E-02
Chlordane	57-74-9	5.94		2.87E-03	h		5.13E+02	2.05E+03	7.17E-04		7.17E-04
Delta-BHC	319-86-8	4.14		3.13E-02	h		6.61E+00	2.64E+01	7.84E-03		7.84E-03
Dieldrin	60-57-1	5.27		6.97E-03	h		2.55E+02	1.02E+03	1.74E-03		1.74E-03
Endothall	145-73-3	-0.87		2.47E+01	h		1.40E-03	5.61E-03	6.16E+00		6.16E+00
Endrin	72-20-8	4.89		1.15E-02	h		1.08E+02	4.32E+02	2.88E-03		2.88E-03
gamma-BHC (Lindane)	58-89-9	3.72		5.48E-02	h		1.07E+01	4.29E+01	1.37E-02		1.37E-02
Heptachlor	76-44-8	5.02	4.89E-02	9.78E-03	k		9.53E+01	3.81E+02	2.45E-03		2.45E-03
Isodrin	465-73-6	3.55		6.87E-02	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Methoxychlor	72-43-5	4.53		1.87E-02	h		8.00E+02	3.20E+03	4.69E-03		4.69E-03
Silvex (2,4,5-TP)	93-72-1	4.07		3.43E-02	h		No Kd	No Kd	1.00E+00	Default	1.00E+00
Toxaphene	8001-35-2	5.50		5.13E-03	h		1.00E+03	4.00E+03	1.28E-03		1.28E-03
Inorganic Chemicals and Compounds											
Metals											
Aluminum	7429-90-5	NA	4.00E-03	8.00E-04	k		1.50E+03	1.50E+03	8.00E-04		8.00E-04
Antimony	7440-36-0	NA	2.00E-01	4.00E-02	k		4.50E+01	4.50E+01	4.00E-02		4.00E-02
Arsenic	7440-38-2	NA	3.60E-02	7.20E-03	k		2.50E+01	2.50E+01	7.20E-03		7.20E-03
Barium	7440-39-3	NA	1.50E-01	3.00E-02	k		1.10E+01	1.10E+01	3.00E-02		3.00E-02
Beryllium	7440-41-7	NA	1.00E-02	2.00E-03	k		2.30E+01	2.30E+01	2.00E-03		2.00E-03
Bismuth	7440-69-9	NA		7.00E-03	l		2.00E+02	2.00E+02	7.00E-03		7.00E-03

Table C2-7. Aquatic Sediment-to-Plant Transfer Factors (SP) for Ecological Receptors (mg/kg tissue wet weight / mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SP[b] (per dry weight of plants)	Calculated from EPA (1999) SP[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SP	Soil Kd[d]	Sediment Kd[e]	SAIC Computed SP[f]	Comments	Recommended SP[g]
Boron	7440-42-8	NA		8.00E-01	l		3.00E+00	3.00E+00	8.00E-01		8.00E-01
Cadmium	7440-43-9	NA	3.64E-01	7.28E-02	k		1.50E+01	1.50E+01	7.28E-02		7.28E-02
Calcium	7440-70-2	NA		7.00E-01	l		4.00E+00	4.00E+00	7.00E-01		7.00E-01
Chromium (and VI)	18540-29-9	NA	7.50E-03	1.50E-03	k		1.20E+03	1.20E+03	1.50E-03		1.50E-03
Cobalt	7440-48-4	NA		4.00E-03	l		4.50E+01	4.50E+01	4.00E-03		4.00E-03
Copper	7440-50-8	NA	4.00E-01	8.00E-02	k		3.50E+01	3.50E+01	8.00E-02		8.00E-02
Iron	7439-89-6	NA		8.00E-04	l		2.50E+01	2.50E+01	8.00E-04		8.00E-04
Lead	7439-92-1	NA	4.50E-02	9.00E-03	k		9.00E+02	9.00E+02	9.00E-03		9.00E-03
Lithium	7439-93-2	NA		5.00E-03	l		3.00E+02	3.00E+02	5.00E-03		5.00E-03
Magnesium	7439-95-4	NA		2.00E-01	l		4.50E+00	4.50E+00	2.00E-01		2.00E-01
Manganese	7439-96-5	NA		5.00E-02	l		6.50E+01	6.50E+01	5.00E-02		5.00E-02
Mercury	7439-97-6	NA	3.75E-02	7.50E-03	k		1.00E+03	1.00E+03	7.50E-03		7.50E-03
Molybdenum	7439-98-7	NA		5.00E-02	l		2.00E+01	2.00E+01	5.00E-02		5.00E-02
Nickel	7440-02-0	NA	3.20E-02	6.40E-03	k		1.60E+01	1.60E+01	6.40E-03		6.40E-03
Potassium	7440-09-7	NA		2.00E-01	l		5.50E+00	5.50E+00	2.00E-01		2.00E-01
Rhodium	7440-16-6	NA		3.00E-02	l		6.00E+01	6.00E+01	3.00E-02		3.00E-02
Selenium	7782-49-2	NA	1.60E-02	3.20E-03	k		1.80E+01	1.80E+01	3.20E-03		3.20E-03
Silicon	7440-21-3	NA		7.00E-02	l		3.00E+01	3.00E+01	7.00E-02		7.00E-02
Silver	7440-22-4	NA	4.00E-01	8.00E-02	k		1.00E-01	1.00E-01	8.00E-02		8.00E-02
Sodium	7440-23-5	NA		1.50E-02	l		1.00E+02	1.00E+02	1.50E-02		1.50E-02
Strontium	7440-24-6	NA		5.00E-01	l		3.50E+01	3.50E+01	5.00E-01		5.00E-01
Tantalum	7440-25-7	NA		2.00E-03	l		6.50E+02	6.50E+02	2.00E-03		2.00E-03
Thallium	7440-28-0	NA	4.00E-03	8.00E-04	k		4.40E+01	4.40E+01	8.00E-04		8.00E-04
Tin	7440-31-5	NA		6.00E-03	l		2.50E+02	2.50E+02	6.00E-03		6.00E-03
Tungsten	7440-33-7	NA		9.00E-03	l		1.50E+02	1.50E+02	9.00E-03		9.00E-03
Uranium	7440-61-1	NA		1.70E-03	l		4.50E+02	4.50E+02	1.70E-03		1.70E-03
Vanadium	7440-62-2	NA		1.10E-03	l		1.00E+03	1.00E+03	1.10E-03		1.10E-03
Yttrium	7440-65-5	NA		3.00E-03	l		5.00E+02	5.00E+02	3.00E-03		3.00E-03

Table C2-7. Aquatic Sediment-to-Plant Transfer Factors (SP) for Ecological Receptors (mg/kg tissue wet weight / mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SP[b] (per dry weight of plants)	Calculated from EPA (1999) SP[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SP	Soil Kd[d]	Sediment Kd[e]	SAIC Computed SP[f]	Comments	Recommended SP[g]
Zinc	7440-66-6	NA	1.20E-12	2.40E-13	k		6.20E+01	6.20E+01	2.40E-13		2.40E-13
Zirconium	7440-67-7	NA		4.00E-04	l		3.00E+03	3.00E+03	4.00E-04		4.00E-04
Non-metals and Anions											
Ammonia/Ammonium	7664-41-7	NA		4.37E-01	m		No Kd	No Kd	1.00E+00	Default	4.37E-01
Bromide	24959-67-9	NA		3.00E-01	l		7.50E+00	7.50E+00	3.00E-01		3.00E-01
Chloride	16887-00-6	NA		1.40E+01	l		2.50E-01	2.50E-01	1.40E+01		1.40E+01
Cyanide	57-12-5	NA		4.37E-01	m		No Kd	No Kd	1.00E+00	Default	4.37E-01
Fluoride	16984-48-8	NA		1.20E-02	l		1.50E+02	1.50E+02	1.20E-02		1.20E-02
Hydroxide	14280-30-9	NA		NA			No Kd	No Kd	NA		NA
Iodine	7553-56-2	NA		3.00E-02	l		6.00E+01	6.00E+01	3.00E-02		3.00E-02
Nitrate	14797-55-8	NA		4.37E-01	m		No Kd	No Kd	1.00E+00	Default	4.37E-01
Nitrite	14797-65-0	NA		4.37E-01	m		No Kd	No Kd	1.00E+00	Default	4.37E-01
Phosphate	14265-44-2	NA		7.00E-01	l		3.50E+00	3.50E+00	7.00E-01		7.00E-01
Phosphorus	7723-14-0	NA		7.00E-01	l		3.50E+00	3.50E+00	7.00E-01		7.00E-01
Sulfate	14808-79-8	NA		3.00E-01	l		No Kd	No Kd	1.00E+00	Default	3.00E-01
Total Sulfur	63705-05-5	NA		3.00E-01	l		7.50E+00	7.50E+00	3.00E-01		3.00E-01
Priority Pollutants											
Carbon Dioxide	124-38-9	NA		4.37E-01	m		No Kd	No Kd	1.00E+00	Default	4.37E-01
Nitrogen Dioxide	10102-44-0	NA		4.37E-01	m		No Kd	No Kd	1.00E+00	Default	4.37E-01
Ozone	10028-15-6	NA		4.37E-01	m		No Kd	No Kd	1.00E+00	Default	4.37E-01
Particulate Matter	No CAS #	NA		NA			No Kd	No Kd	NA		NA
Sulfur Dioxode	7446-09-5	NA		4.37E-01	m		No Kd	No Kd	1.00E+00	Default	4.37E-01
Radionuclides											
Americium-241	1596-10-2	NA		1.10E-03	l		7.00E+02	7.00E+02	1.10E-03		1.10E-03
Antimony-125	14234-35-6	NA		4.00E-02	l		4.50E+01	4.50E+01	4.00E-02		4.00E-02
Barium-137	13981-97-0	NA		3.00E-02	l		6.00E+01	6.00E+01	3.00E-02		3.00E-02
Cadmium-113	None	NA		1.10E-01	l		6.50E+00	6.50E+00	1.10E-01		1.10E-01
Cesium-134	13967-70-9	NA		1.60E-02	l		1.00E+03	1.00E+03	1.60E-02		1.60E-02

Table C2-7. Aquatic Sediment-to-Plant Transfer Factors (SP) for Ecological Receptors (mg/kg tissue wet weight / mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SP[b] (per dry weight of plants)	Calculated from EPA (1999) SP[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SP	Soil Kd[d]	Sediment Kd[e]	SAIC Computed SP[f]	Comments	Recommended SP[g]
Cesium-137	10045-97-3	NA		1.60E-02	l		1.00E+03	1.00E+03	1.60E-02		1.60E-02
Europium-154	15585-10-1	NA		2.00E-03	l		6.50E+02	6.50E+02	2.00E-03		2.00E-03
Europium-155	14391-16-3	NA		2.00E-03	l		6.50E+02	6.50E+02	2.00E-03		2.00E-03
Nickel-63	13981-37-8	NA		1.20E-02	l		1.50E+02	1.50E+02	1.20E-02		1.20E-02
Plutonium-239	15117-48-3	NA		9.00E-05	l		4.50E+03	4.50E+03	9.00E-05		9.00E-05
Plutonium-241	14119-32-5	NA		9.00E-05	l		4.50E+03	4.50E+03	9.00E-05		9.00E-05
Samarium-151	15715-94-3	NA		2.00E-03	l		6.50E+02	6.50E+02	2.00E-03		2.00E-03
Strontium-90	10098-97-2	NA		5.00E-01	l		3.50E+01	3.50E+01	5.00E-01		5.00E-01
Technetium-99	14133-79-7	NA		1.90E+00	l		1.50E+00	1.50E+00	1.90E+00		1.90E+00
Tritium	10028-17-8	NA		4.27E-01	m		No Kd	No Kd	1.00E+00	Default	4.27E-01
Yttrium-90	10098-91-6	NA		3.00E-03	l		5.00E+02	5.00E+02	3.00E-03		3.00E-03
Uranium-232	14158-29-3	NA		1.70E-03	l		4.50E+02	4.50E+02	1.70E-03		1.70E-03
Uranium-233	13968-55-3	NA		1.70E-03	l		4.50E+02	4.50E+02	1.70E-03		1.70E-03
Uranium-234	13966-29-5	NA		1.70E-03	l		4.50E+02	4.50E+02	1.70E-03		1.70E-03
Uranium-235	15117-96-1	NA		1.70E-03	l		4.50E+02	4.50E+02	1.70E-03		1.70E-03
Uranium-236	13982-70-2	NA		1.70E-03	l		4.50E+02	4.50E+02	1.70E-03		1.70E-03
Uranium-238	7440-61-1	NA		1.70E-03	l		4.50E+02	4.50E+02	1.70E-03		1.70E-03
Actinium-227	14952-40-0	NA		7.00E-04	l		1.50E+03	1.50E+03	7.00E-04		7.00E-04
Americium-243	14993-75-0	NA		1.10E-03	l		7.00E+02	7.00E+02	1.10E-03		1.10E-03
Carbon-14	14762-75-5	NA		4.27E-01	m		No Kd	No Kd	1.00E+00	Default	4.27E-01
Cobalt-60	10198-40-0	NA		4.00E-03	l		4.50E+01	4.50E+01	4.00E-03		4.00E-03
Curium-242	15510-73-3	NA		1.70E-04	l		2.00E+03	2.00E+03	1.70E-04		1.70E-04
Curium-243	15757-87-6	NA		1.70E-04	l		2.00E+03	2.00E+03	1.70E-04		1.70E-04
Curium-244	13981-15-2	NA		1.70E-04	l		2.00E+03	2.00E+03	1.70E-04		1.70E-04
Europium-152	14683-23-9	NA		2.00E-03	l		6.50E+02	6.50E+02	2.00E-03		2.00E-03
Iodine-129	15046-84-1	NA		3.00E-02	l		6.00E+01	6.00E+01	3.00E-02		3.00E-02
Neptunium-237	13994-20-2	NA		2.00E-02	l		3.00E+01	3.00E+01	2.00E-02		2.00E-02
Nickel-59	14336-70-0	NA	3.20E-02	6.40E-03	k		1.50E+02	1.50E+02	6.40E-03		6.40E-03

Table C2-7. Aquatic Sediment-to-Plant Transfer Factors (SP) for Ecological Receptors (mg/kg tissue wet weight / mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) SP[b] (per dry weight of plants)	Calculated from EPA (1999) SP[c] (adjusted to wet weight of plants)	Notes	Ecology Guidance SP	Soil Kd[d]	Sediment Kd[e]	SAIC Computed SP[f]	Comments	Recommended SP[g]
Niobium-93	None	NA		4.00E-03	l		3.50E+02	3.50E+02	4.00E-03		4.00E-03
Plutonium-238	13981-16-3	NA		9.00E-05	l		4.50E+03	4.50E+03	9.00E-05		9.00E-05
Plutonium-240	14119-33-6	NA		9.00E-05	l		4.50E+03	4.50E+03	9.00E-05		9.00E-05
Plutonium-242	13982-10-0	NA		9.00E-05	l		4.50E+07	4.50E+07	9.00E-05		9.00E-05
Protactinium-231	14331-85-2	NA		5.00E-04	l		2.50E+03	2.50E+03	5.00E-04		5.00E-04
Radium-226	13982-63-3	NA		3.00E-03	l		4.50E+02	4.50E+02	3.00E-03		3.00E-03
Radium-228	15262-20-1	NA		3.00E-03	l		4.50E+02	4.50E+02	3.00E-03		3.00E-03
Ruthenium-106	13967-48-1	NA		1.50E-02	l		3.50E+02	3.50E+02	1.50E-02		1.50E-02
Selenium-79	None	NA		5.00E-03	l		3.00E+02	3.00E+02	5.00E-03		5.00E-03
Thorium-229	15594-54-4	NA		1.70E-04	l		1.50E+05	1.50E+05	1.70E-04		1.70E-04
Thorium-232	7440-29-1	NA		1.70E-04	l		1.50E+05	1.50E+05	1.70E-04		1.70E-04
Tin-126	15832-50-5	NA		6.00E-03	l		2.50E+02	2.50E+02	6.00E-03		6.00E-03
Zirconium-93	15751-77-6	NA		4.00E-04	l		3.00E+03	3.00E+03	4.00E-04		4.00E-04

NA = Not applicable

[a] $\log_{10}$ of K_{ow} values in Table 4.1

[b] Published in Appendix C of EPA (1999), Table C-2

[c] Calculated or chosen as described in Appendix C, Sect. C-1.2 of EPA (1999), then multiplied by 0.2 to adjust to wet weight of plants

[d] Soil Kd values from Appendix B Table B-1-1 or from Baes and others. (1984) Fig. 2.31 for radionuclides and other elements not shown in Appendix B Table B-1-1

[e] Sediment Kd values from Appendix B Table B-1-1 or soil Kds from Baes and others. (1984) Fig. 2.31 for radionuclides and other elements not shown in Appendix B Table B-1-1

[f] Calculated by using the Kds and the recommended SPv values for plants in soil (Table C.2-1): SP = SPv x Kd(soil)/Kd(sediment)

[g] Selection criteria described in Sect. 8.2.4.4

[h] Wet weight values calculated by using log K_{ow}: log SPv = 1.588 - 0.578 x log K_{ow} (Travis and Arms 1988) for dry weight and multiplying by 0.2 to adjust to wet weight

[i] Calculated from value for TCDD in Appendix C of EPA (1999), Table C-2 by multiplying by BEF, then by 0.2 to adjust to wet weight of plants

[j] Value for Aroclor 1254 from Appendix C of EPA (1999), Table C-2, was used for PCB mixtures and was multiplied by 0.2 to adjust to wet weight of plants

[k] EPA (1999) values published in Table C-2 multiplied by 0.2 to adjust to wet weight of plants

[l] Baes et al. (1984) (Figure 2.1) values multiplied by 0.2 to adjust to wet weight of plants

[m] Average of values for other inorganics as published in Appendix C of EPA (1999), Table C-2

Table C2-8. Aquatic Water-to-Invertebrate Transfer Factors (BCF_{inv}) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{inv} [b]	BCF_{inv} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{inv}	SAIC Compiled BCF_{inv}	Comments	Recommended BCF_{inv} [d]
Organic Compounds									
Aromatic Halogenated Hydrocarbons									
4-Chloro-3-methylphenol	59-50-7	3.10		2.47E+01	e				2.47E+01
2,3,4,6-Tetrachlorophenol	58-90-2	4.42		2.98E+02	e				2.98E+02
Aromatic Nonhalogenated Hydrocarbons									
2-Nitrotoluene	88-72-2	2.30		5.47E+00	e				5.47E+00
4-Nitrobiphenyl	92-93-3	3.77		8.74E+01	e				8.74E+01
Benzaldehyde	100-52-7	1.48		1.16E+00	e				1.16E+00
Benzene	71-43-2	-4.99		5.84E-06	e				5.84E-06
Benzyl alcohol	100-51-6	1.10		5.69E-01	e				5.69E-01
Ethyl benzene	100-41-4	3.12		2.58E+01	e				2.58E+01
m-Xylene	108-38-3	3.20		2.99E+01	e				2.99E+01
o-Xylene	95-47-6	3.13		2.62E+01	e				2.62E+01
p-Xylene	106-42-3	3.17		2.82E+01	e				2.82E+01
Styrene	100-42-5	2.93		1.79E+01	e				1.79E+01
Toluene	108-88-3	2.67		1.09E+01	e				1.09E+01
Non-aromatic Nonhalogenated Hydrocarbons									
1,2-Epoxybutane	106-88-7	1.44		1.08E+00	e				1.08E+00
1,3-Butadiene	106-99-0	1.90		2.57E+00	e				2.57E+00
1,4-Dioxane	123-91-1	-0.27	4.31E-02		e				4.31E-02
1-Methylpropyl alcohol	78-92-2	0.61		2.26E-01	e				2.26E-01
1-Nitropropane	108-03-2	0.87		3.69E-01	e				3.69E-01
2,2,4-Trimethylpentane	540-84-1	5.02		9.23E+02	e				9.23E+02
2-Butanone	78-93-3	0.28		1.21E-01	e				1.21E-01
2-Butenaldehyde (2-Butenal)	4170-30-3	No data		No data			1.00E+05	Default	1.00E+05
2-Ethoxyethanol	110-80-5	-0.10		5.92E-02	e				5.92E-02
2-Heptanone	110-43-0	1.98		2.99E+00	e				2.99E+00
2-Hexanone	591-78-6	1.38		9.64E-01	e				9.64E-01
2-Methoxyethanol	109-86-4	0.25		1.15E-01	e				1.15E-01

Table C2-8. Aquatic Water-to-Invertebrate Transfer Factors (BCF_{inv}) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{inv} [b]	BCF_{inv} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{inv}	SAIC Compiled BCF_{inv}	Comments	Recommended BCF_{inv} [d]
2-Methyl-2-propanol	75-65-0	0.35		1.38E-01	e				1.38E-01
2-Methyl-2-propenenitrile	126-98-7	0.54		1.98E-01	e				1.98E-01
2-Methylaziridine	75-55-8	-0.60		2.29E-02	e				2.29E-02
2-Methylpropyl alcohol	78-83-1	0.76		3.00E-01	e				3.00E-01
2-Pentanone	107-87-9	0.91		3.97E-01	e				3.97E-01
2-Propanone (Acetone)	67-64-1	-0.22	5.00E-02		e				5.00E-02
2-Propene-1-ol	107-18-6	0.17		9.85E-02	e				9.85E-02
2-Propyl alcohol	67-63-0	0.05		7.85E-02	e				7.85E-02
3-Heptanone	106-35-4	No data		No data			1.00E+05	Default	1.00E+05
3-Methyl-1-butanol	123-51-3	No data		No data			1.00E+05	Default	1.00E+05
3-Methyl-2-butanone	563-80-4	No data		No data			1.00E+05	Default	1.00E+05
3-Pentanone	96-22-0	0.99		4.62E-01	e				4.62E-01
4-Heptanone	123-19-3	No data		No data			1.00E+05	Default	1.00E+05
4-Methyl-2-pentanone	108-10-1	1.19		6.74E-01	e				6.74E-01
4-Methyl-3-penten-2-one	141-79-7	No data		No data			1.00E+05	Default	1.00E+05
5-Methyl-2-hexanone	110-12-3	No data		No data			1.00E+05	Default	1.00E+05
Acetaldehyde	75-07-0	-0.22		4.72E-02	e				4.72E-02
Acetamide	60-35-5	-1.26		6.64E-03	e				6.64E-03
Acetic acid	64-19-7	-0.17		5.19E-02	e				5.19E-02
Acetic acid ethyl ester	141-78-6	0.73		2.83E-01	e				2.83E-01
Acetic acid n-butyl ester	123-86-4	1.73		1.87E+00	e				1.87E+00
Acetonitrile	75-05-8	-0.34		3.76E-02	e				3.76E-02
Acrolein	107-02-8	-0.01		7.03E-02	e				7.03E-02
Acrylonitrile	107-13-1	0.25	1.15E-01		e				1.15E-01
Bis(isopropyl)ether	108-20-3	No data		No data			1.00E+05	Default	1.00E+05
Butane	106-97-8	2.89		1.66E+01	e				1.66E+01
Carbon disulfide	75-15-0	2.00		3.10E+00	e				3.10E+00
Cyanogen	460-19-5	0.81		3.28E-01	e				3.28E-01
Cyclohexane	110-82-7	3.44		4.69E+01	e				4.69E+01

Table C2-8. Aquatic Water-to-Invertebrate Transfer Factors (BCF_{inv}) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{inv} [b]	BCF_{inv} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{inv}	SAIC Compiled BCF_{inv}	Comments	Recommended BCF_{inv} [d]
Cyclohexanone	108-94-1	0.81		3.29E-01	e				3.29E-01
Cyclohexene	110-83-8	2.86		1.57E+01	e				1.57E+01
Cyclopentane	287-92-3	3.00		2.05E+01	e				2.05E+01
Ethyl alcohol	64-17-5	0.31		1.28E-01	e				1.28E-01
Ethyl ether	60-29-7	0.89		3.83E-01	e				3.83E-01
Ethyl methacrylate	97-63-2	1.59		1.43E+00	e				1.43E+00
Formaldehyde	50-00-0	0.34	1.36E-01		e				1.36E-01
Formamide	75-12-7	-1.51		4.14E-03	e				4.14E-03
Formic acid	64-18-6	-0.54		2.59E-02	e				2.59E-02
Formic acid, methyl ester	107-31-3	-0.26		4.34E-02	e				4.34E-02
Glycidylaldehyde	765-34-4	-0.73		1.80E-02	e				1.80E-02
Methyl acetate	79-20-9	0.18		1.00E-01	e				1.00E-01
Methyl alcohol	67-56-1	-0.71		1.87E-02	e				1.87E-02
Methyl isocyanate	624-83-9	No data		No data				Reacts with water	NA
Methyl methacrylate	80-62-6	0.79		3.17E-01	e				3.17E-01
Methyl tert-butyl ether	1634-04-4	0.94		4.21E-01	e				4.21E-01
Methylacetylene	74-99-7	0.94		4.21E-01	e				4.21E-01
Methylcyclohexane	108-87-2	4.10		1.63E+02	e				1.63E+02
N,N-Dimethylacetamide	127-19-5	No data		No data			1.00E+05	Default	1.00E+05
n-Butyl alcohol	71-36-3	0.88		3.76E-01	e				3.76E-01
n-Heptane	142-82-5	4.66		4.68E+02	e				4.68E+02
n-Hexane	110-54-3	4.11		1.66E+02	e				1.66E+02
Nitromethane	75-52-5	-0.35		3.69E-02	e				3.69E-02
n-Nonane	111-84-2	5.65		3.03E+03	e				3.03E+03
n-Octane	111-65-9	4.00		1.35E+02	e				1.35E+02
n-Pentane	109-66-0	3.21		3.04E+01	e				3.04E+01
n-Propionaldehyde	123-38-6	0.59		2.17E-01	e				2.17E-01
n-Propyl alcohol	71-23-8	0.25		1.14E-01	e				1.14E-01
n-Valeraldehyde	110-62-3	No data		No data			1.00E+05	Default	1.00E+05

Table C2-8. Aquatic Water-to-Invertebrate Transfer Factors (BCF_{inv}) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{inv} [b]	BCF_{inv} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{inv}	SAIC Compiled BCF_{inv}	Comments	Recommended BCF_{inv} [d]
Oxirane	75-21-8	-0.30		4.06E-02	e				4.06E-02
p-Cymene	99-87-6	4.10		1.63E+02	e				1.63E+02
Phosgene	75-44-5	No data		No data			1.00E+05	Default	1.00E+05
Propargyl alcohol	107-19-7	0.26		1.17E-01	e				1.17E-01
Propionic acid	79-09-4	0.33		1.33E-01	e				1.33E-01
Propionitrile	107-12-0	0.16		9.66E-02	e				9.66E-02
Propylene gylcol monomethyl ether	107-98-2	-0.18		5.09E-02	e				5.09E-02
p-tert-Butyltoluene	98-51-1	No data		No data			1.00E+05	Default	1.00E+05
Triethylamine	121-44-8	0.16		9.66E-02	e				9.66E-02
Trimethylamine	75-50-3	0.16		9.66E-02	e				9.66E-02
Vinyl acetate	108-05-4	0.70		2.67E-01	e				2.67E-01
Non-aromatic Halogenated Hydrocarbons									
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	No data		No data			1.00E+05	Default	1.00E+05
1,1,1,2-Tetrachloroethane	630-20-6	2.63		1.02E+01	e				1.02E+01
1,1,1-Trichloroethane	71-55-6	2.42		6.88E+00	e				6.88E+00
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	3.73		8.11E+01	e				8.11E+01
1,1,2,2-Tetrachloroethane	79-34-5	4.64		4.54E+02	e				4.54E+02
1,1,2,2-Tetrachloroethene	127-18-4	2.55		8.68E+00	e				8.68E+00
1,1,2-Trichloroethane	79-00-5	2.10		3.73E+00	e				3.73E+00
1,1,2-Trichloroethylene	79-01-6	2.43		7.02E+00	e				7.02E+00
1,1-Dichloroethane	75-34-3	1.79		2.10E+00	e				2.10E+00
1,1-Dichloroethene	75-35-4	2.12		3.90E+00	e				3.90E+00
1,2,2-Trichloro-1,1,2-trifluoroethane	76-13-1	3.16		2.77E+01	e				2.77E+01
1,2,3-Trichloropropane	96-18-4	2.25		4.98E+00	e				4.98E+00
1,2-Dibromo-3-chloropropane	96-12-8	2.34		5.90E+00	e				5.90E+00
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	2.82		1.46E+01	e				1.46E+01
1,2-Dichloroethane	107-06-2	1.46		1.13E+00	e				1.13E+00
1,2-Dichloroethylene	540-59-0	0.48		1.77E-01	e				1.77E-01
1,2-Dichloropropane	78-87-5	2.25		4.98E+00	e				4.98E+00

Table C2-8. Aquatic Water-to-Invertebrate Transfer Factors (BCF_{inv}) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{inv}[a]	EPA (1999) BCF_{inv}[b]	BCF_{inv} Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance BCF_{inv}	SAIC Compiled BCF_{inv}	Comments	Recommended BCF_{inv}[d]
1,3-Dichloropropene	542-75-6	1.75		1.93E+00	e				1.93E+00
1,4-Dichloro-2-butene	764-41-0	0.87		3.70E-01	e				3.70E-01
1-Chloroethene	75-01-4	1.15		6.20E-01	e				6.20E-01
2,2-Dichloropropionic acid	75-99-0	0.78		3.10E-01	e				3.10E-01
2-Chloropropane	75-29-6	1.90		2.57E+00	e				2.57E+00
3-Chloropropene (allyl chloride)	107-05-1	0.95		4.32E-01	e				4.32E-01
Bromochloromethane	74-97-5	1.41		1.02E+00	e				1.02E+00
Bromodichloromethane	75-27-4	2.03		3.26E+00	e				3.26E+00
Bromoethene	593-60-2	1.07		5.34E-01	e				5.34E-01
Bromoform	75-25-2	2.35		6.01E+00	e				6.01E+00
Bromomethane	74-83-9	1.11		5.84E-01	e				5.84E-01
Carbon tetrachloride	56-23-5	2.72	1.20E+01		e				1.20E+01
Chlorodibromomethane	124-48-1	2.18		4.33E+00	e				4.33E+00
Chlorodifluoromethane	75-45-6	1.08		5.47E-01	e				5.47E-01
Chloroethane	75-00-3	3.10		2.47E+01	e				2.47E+01
Chloroform	67-66-3	1.95	2.82E+00		e				2.82E+00
Chloromethane	74-87-3	0.90		3.92E-01	e				3.92E-01
Chloropentafluoroethane	76-15-3	No data		No data			1.00E+05	Default	1.00E+05
cis-1,2-Dichloroethene	156-59-2	1.98		3.00E+00	e				3.00E+00
cis-1,3-Dichloropropene	10061-01-5	No data		No data			1.00E+05	Default	1.00E+05
Cyanogen bromide	506-68-3	No data		No data			1.00E+05	Default	1.00E+05
Cyanogen chloride	506-77-4	0.20		1.04E-01	e				1.04E-01
Dichlorodifluoromethane	75-71-8	2.16		4.19E+00	e				4.19E+00
Dichlorofluoromethane	75-43-4	No data		No data			1.00E+05	Default	1.00E+05
Dichloromethane	75-09-2	1.26		7.62E-01	e				7.62E-01
Difluorodibromomethane	75-61-6	No data		No data			1.00E+05	Default	1.00E+05
Hexafluoroacetone	684-16-2	No data		No data			1.00E+05	Default	1.00E+05
Iodomethane	74-88-4	1.69		1.73E+00	e				1.73E+00
Methylene bromide	74-95-3	1.62		1.52E+00	e				1.52E+00

Table C2-8. Aquatic Water-to-Invertebrate Transfer Factors (BCF_{inv}) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{inv} [b]	BCF_{inv} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{inv}	SAIC Compiled BCF_{inv}	Comments	Recommended BCF_{inv} [d]
Pentachloroethane	76-01-7	3.05		2.25E+01	e				2.25E+01
trans-1,2-Dichloroethene	156-60-5	1.98		3.00E+00	e				3.00E+00
trans-1,3-Dichloropropene	10061-02-6	2.06		3.48E+00	e				3.48E+00
Trichloroacetic acid	76-03-9	1.33		8.78E-01	e				8.78E-01
Trichlorofluoroethane	27154-33-2	No data		No data			1.00E+05	Default	1.00E+05
Trichlorofluoromethane	75-69-4	2.53		8.46E+00	e				8.46E+00
Trifluorobromomethane	75-63-8	1.86		2.38E+00	e				2.38E+00
Dioxin and Furan Compounds									
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	35822-46-9	8.20	7.96E+01		f				7.96E+01
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	7.92	1.72E+01		f				1.72E+01
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	7.92	6.08E+02		f				6.08E+02
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	39227-28-6	7.79	4.84E+02		f				4.84E+02
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	7.25	1.19E+02		f				1.19E+02
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	57653-85-7	7.25	1.87E+02		f				1.87E+02
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	7.25	2.96E+02		f				2.96E+02
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	19408-74-3	7.25	2.18E+02		f				2.18E+02
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	7.25	9.83E+02		f				9.83E+02
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	40321-76-4	6.64	1.44E+03		f				1.44E+03
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	6.79	3.43E+02		f				3.43E+02
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	7.25	1.05E+03		f				1.05E+03
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	6.92	2.50E+03		f				2.50E+03
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1746-01-6	6.64	1.56E+03						1.56E+03
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	6.53	1.25E+03		f				1.25E+03
Dibenzofuran	132-64-9	4.33		2.51E+02	e				2.51E+02
Octachlorodibenzo(p)dioxin	3268-87-9	8.20	1.87E+01		f				1.87E+01
Octachlorodibenzofuran	39001-02-0	8.78	2.50E+01		f				2.50E+01
PCBs									
Polychlorinated biphenyls (PCBs)[g]	1336-36-3	7.31	5.54E+03						5.54E+03
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	7.08		4.49E+04	e				4.49E+04

Table C2-8. Aquatic Water-to-Invertebrate Transfer Factors (BCF_{inv}) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{inv} [b]	BCF_{inv} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{inv}	SAIC Compiled BCF_{inv}	Comments	Recommended BCF_{inv} [d]
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	7.12		4.84E+04	e				4.84E+04
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	No data		No data			1.00E+05	Default	1.00E+05
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	No data		No data			1.00E+05	Default	1.00E+05
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	No data		No data			1.00E+05	Default	1.00E+05
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	No data		No data			1.00E+05	Default	1.00E+05
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	No data		No data			1.00E+05	Default	1.00E+05
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	No data		No data			1.00E+05	Default	1.00E+05
2',3,4,4',5-Pentachlorobiphenyl	no cas #	No data		No data			1.00E+05	Default	1.00E+05
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	7.12		4.84E+04	e				4.84E+04
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	7.41		8.34E+04	e				8.34E+04
3,3',4,4',5-Pentachlorobiphenyl	no cas #	No data		No data			1.00E+05	Default	1.00E+05
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	No data		No data			1.00E+05	Default	1.00E+05
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	No data		No data			1.00E+05	Default	1.00E+05
Phthalates									
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	5.20	3.18E+02						3.18E+02
Butylbenzyl phthalate	85-68-7	4.41		2.94E+02	e				2.94E+02
Dibutyl phthalate	84-74-2	4.72		5.25E+02	e				5.25E+02
Diethyl phthalate	84-66-2	4.44		3.07E+02	e				3.07E+02
Dimethylphthalate	131-11-3	1.63		1.56E+00	e				1.56E+00
n-Dioctyl phthalate	117-84-0	9.33	5.95E+03						5.95E+03
Light Polycyclic Aromatic Hydrocarbons (molecular weight <200 g/mole)									
2-Chloronapthalene	91-58-7	4.07		1.53E+02	e				1.53E+02
2-Methyl naphthalene	91-57-6	3.86		1.04E+02	e				1.04E+02
5-Nitroacenaphthene	602-87-9	No data		No data			1.00E+05	Default	1.00E+05
Acenaphthene	83-32-9	3.96		1.26E+02	e				1.26E+02
Acenaphthylene	208-96-8	4.07		1.54E+02	e				1.54E+02
Anthracene	120-12-7	4.47		3.27E+02	e				3.27E+02
Fluorene	86-73-7	4.18		1.89E+02	e				1.89E+02
Indene	95-13-6	No data		No data			1.00E+05	Default	1.00E+05

Table C2-8. Aquatic Water-to-Invertebrate Transfer Factors (BCF_{inv}) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{inv} [b]	BCF_{inv} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{inv}	SAIC Compiled BCF_{inv}	Comments	Recommended BCF_{inv} [d]
Naphthalene	91-20-3	3.37		4.13E+01	e				4.13E+01
Phenanthrene	85-01-8	4.55		3.81E+02	e				3.81E+02
Pyrene	129-00-0	5.00		8.89E+02	e				8.89E+02
Heavy Polycyclic Aromatic Hydrocarbons (molecular weight >200 g/mole)									
3-Methylcholanthrene	56-49-5	7.11		4.75E+04	e				4.75E+04
5-Methylchrysene	3697-24-3	No data		No data			1.00E+05	Default	1.00E+05
Benzo(a)anthracene	56-55-3	5.68	1.23E+04						1.23E+04
Benzo(a)pyrene	50-32-8	6.13	4.70E+03						4.70E+03
Benzo(b)fluoranthene	205-99-2	6.20	4.70E+03		h				4.70E+03
Benzo(e)pyrene	192-97-2	7.40		8.21E+04	e				8.21E+04
Benzo(g,h,i)perylene	191-24-2	7.10		4.67E+04	e				4.67E+04
Benzo(j)fluoranthene	205-82-3	6.44		1.34E+04	e				1.34E+04
Benzo(k)fluoranthene	207-08-9	6.20	1.32E+04						1.32E+04
Benzo[a,i]pyrene	191-30-0	No data		No data			1.00E+05	Default	1.00E+05
Chrysene	218-01-9	5.74	9.80E+02						9.80E+02
Dibenz(a,h)anthracene	53-70-3	6.55	7.10E+02						7.10E+02
Dibenz[a,h]acridine	226-36-8	No data		No data			1.00E+05	Default	1.00E+05
Dibenz[a,j]acridine	224-42-0	No data		No data			1.00E+05	Default	1.00E+05
Dibenzo(a,e)fluoranthene	5385-75-1	No data		No data			1.00E+05	Default	1.00E+05
Dibenzo(a,h)fluoranthene	no cas #	No data		No data			1.00E+05	Default	1.00E+05
Dibenzo[a,e]pyrene	192-65-4	No data		No data			1.00E+05	Default	1.00E+05
Dibenzo[a,h]pyrene	189-64-0	No data		No data			1.00E+05	Default	1.00E+05
Dibenzo[a,i]pyrene	189-55-9	7.29		6.68E+04	e				6.68E+04
Fluoranthene	206-44-0	5.08		1.04E+03	e				1.04E+03
Hexachloronaphthalene	1335-87-1	7.59		1.18E+05	e				1.18E+05
Indeno(1,2,3-cd)pyrene	193-39-5	6.91	4.70E+03		h				4.70E+03
Octachloronaphthalene	2234-13-1	6.42		1.29E+04	e				1.29E+04
Pentachloronaphthalene	1321-64-8	No data		No data			1.00E+05	Default	1.00E+05
Tetrachloronaphthalene	1335-88-2	No data		No data			1.00E+05	Default	1.00E+05

Table C2-8. Aquatic Water-to-Invertebrate Transfer Factors (BCF_{inv}) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) BCF_{inv}[b]	BCF_{inv} Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance BCF_{inv}	SAIC Compiled BCF_{inv}	Comments	Recommended BCF_{inv}[d]
Trichloronaphthalene	1321-65-9	No data		No data			1.00E+05	Default	1.00E+05
Light Substituted Benzene Compounds (MW <200 g/mole)									
1,2,3-Trichlorobenzene	87-61-6	4.05		1.47E+02	e				1.47E+02
1,2,4-Trichlorobenzene	120-82-1	3.99		1.32E+02	e				1.32E+02
1,2,4-Trimethyl benzene	95-63-6	3.65		6.97E+01	e				6.97E+01
1,2-Dichlorobenzene	95-50-1	3.45		4.74E+01	e				4.74E+01
1,3,5-Trimethyl benzene	108-67-8	3.42		4.52E+01	e				4.52E+01
1,3-Dichlorobenzene	541-73-1	3.53		5.56E+01	e				5.56E+01
1,3-Dinitrobenzene	99-65-0	1.49	1.30E+01		i				1.30E+01
1,4-Dichlorobenzene	106-46-7	3.41		4.45E+01	e				4.45E+01
1,4-Dinitrobenzene	100-25-4	1.50		1.21E+00	e				1.21E+00
2,4,5-Trichlorophenol	95-95-4	3.87		1.06E+02	e				1.06E+02
2,4,6-Trichlorophenol	88-06-2	3.71		7.83E+01	e				7.83E+01
2,4-Dichlorophenol	120-83-2	3.04		2.20E+01	e				2.20E+01
2,4-Dimethylphenol	105-67-9	2.36		6.12E+00	e				6.12E+00
2,4-Dinitrophenol	51-28-5	1.52		1.25E+00	e				1.25E+00
2,4-Dinitrotoluene	121-14-2	2.00	1.30E+01						1.30E+01
2,6-Dinitrotoluene	606-20-2	1.89	1.30E+01		i				1.30E+01
2-Chlorophenol	95-57-8	2.16		4.21E+00	e				4.21E+00
2-Chlorotoluene	95-49-8	3.54		5.63E+01	e				5.63E+01
2-Nitrophenol	88-75-5	1.79		2.09E+00	e				2.09E+00
4,6-Dinitro-o-cresol	534-52-1	2.85		1.54E+01	e				1.54E+01
4-Chlorotoluene	106-43-4	3.33		3.81E+01	e				3.81E+01
4-Nitrophenol	100-02-7	1.91		2.62E+00	e				2.62E+00
alpha-Methylstyrene	98-83-9	3.46		4.90E+01	e				4.90E+01
Aniline	62-53-3	0.98		4.54E-01	e				4.54E-01
Benzotrichloride	98-07-7	2.92		1.76E+01	e				1.76E+01
Benzyl chloride	100-44-7	0.36		1.41E-01	e				1.41E-01
Bromobenzene	108-86-1	2.99		2.01E+01	e				2.01E+01

Table C2-8. Aquatic Water-to-Invertebrate Transfer Factors (BCF_{inv}) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) BCF_{inv}[b]	BCF_{inv} Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance BCF_{inv}	SAIC Compiled BCF_{inv}	Comments	Recommended BCF_{inv}[d]
Chlorobenzene	108-90-7	2.79		1.38E+01	e				1.38E+01
Cumene	98-82-8	3.61		6.50E+01	e				6.50E+01
m-Cresol	108-39-4	1.96		2.87E+00	e				2.87E+00
n-Butyl benzene	104-51-8	4.28		2.29E+02	e				2.29E+02
Nitrobenzene	98-95-3	1.83	1.30E+01		i				1.30E+01
n-Propyl benzene	103-65-1	3.69		7.52E+01	e				7.52E+01
o-Cresol	95-48-7	2.02		3.23E+00	e				3.23E+00
o-Dinitrobenzene	528-29-0	1.60		1.46E+00	e				1.46E+00
o-Nitroaniline	88-74-4	1.85		2.34E+00	e				2.34E+00
o-Toluidine	95-53-4	1.34		8.95E-01	e				8.95E-01
p-Chloroaniline	106-47-8	1.87		2.43E+00	e				2.43E+00
p-Cresol	106-44-5	1.94		2.77E+00	e				2.77E+00
Phenol	108-95-2	1.48		1.16E+00	e				1.16E+00
p-Nitrochlorobenzene	100-00-5	2.39		6.48E+00	e				6.48E+00
p-Toluidine	106-49-0	1.40		1.00E+00	e				1.00E+00
sec-Butyl benzene	135-98-8	4.57		3.95E+02	e				3.95E+02
tert-Butyl benzene	98-06-6	4.11		1.66E+02	e				1.66E+02
Toluene-2,6-diamine	823-40-5	1.45		1.11E+00	e				1.11E+00
Trimethyl benzene	25551-13-7	3.40		4.35E+01	e				4.35E+01
Other Light Semivolatile Compounds (molecular weight <200 g/mole)									
1,1'-Biphenyl	92-52-4	3.90		1.12E+02	e				1.12E+02
1,1-Dimethylhydrazine	57-14-7	0.60		2.20E-01	e				2.20E-01
1,2-Dimethylhydrazine	540-73-8	-1.37		5.42E-03	e				5.42E-03
1,2-Diphenylhydrazine	122-66-7	2.94		1.83E+01	e				1.83E+01
1,3-Propane sultone	1120-71-4	-0.52		2.66E-02	e				2.66E-02
2,4-Toluene diisocyante	584-84-9	No data		No data				Decomposes rapidly in water	NA
2-Chloroacetophenone	532-27-4	2.59		9.53E+00	e				9.53E+00
2-Propenoic acid	79-10-7	0.33		1.33E-01	e				1.33E-01
4,4-Methylenedianiline	101-77-9	3.38		4.19E+01	e				4.19E+01

Table C2-8. Aquatic Water-to-Invertebrate Transfer Factors (BCF_{inv}) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{inv} [b]	BCF_{inv} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{inv}	SAIC Compiled BCF_{inv}	Comments	Recommended BCF_{inv} [d]
Acetophenone	98-86-2	1.64		1.58E+00	e				1.58E+00
Benzoic acid	65-85-0	1.86		2.38E+00	e				2.38E+00
bis(2-Chloroethoxy)methane	111-91-1	7.59		1.18E+05	e				1.18E+05
bis(2-Chloroethyl) ether	111-44-4	1.30		8.31E-01	e				8.31E-01
Chlorocyclopentadiene	41851-50-7	2.43		6.99E+00	e				6.99E+00
Cyclohexanol	108-93-0	1.23		7.27E-01	e				7.27E-01
Dichloroisopropyl ether	108-60-1	2.58		9.27E+00	e				9.27E+00
Dichloromethyl ether	542-88-1	-0.38		3.49E-02	e				3.49E-02
Dichloropentadiene	no cas #	No data		No data			1.00E+05	Default	1.00E+05
Dimethyl sulfate	77-78-1	0.32		1.30E-01	e				1.30E-01
Dimethylaniline	121-69-7	-0.88		1.36E-02	e				1.36E-02
Di-n-propylnitrosamine	621-64-7	1.38		9.65E-01	e				9.65E-01
Diphenyl ether	101-84-8	4.21		2.00E+02	e				2.00E+02
Epichlorohydrin	106-89-8	0.25		1.15E-01	e				1.15E-01
Ethyl carbamate (urethane)	51-79-6	-0.15		5.38E-02	e				5.38E-02
Ethyl methanesulfonate	62-50-0	0.05		7.84E-02	e				7.84E-02
Ethylene dibromide	106-93-4	1.75		1.94E+00	e				1.94E+00
Ethylene glycol	107-21-1	-0.91		1.27E-02	e				1.27E-02
Ethylene glycol monobutyl ether	111-76-2	1.55		1.34E+00	e				1.34E+00
Ethylene glycol monoethyl ether acetate	111-15-9	0.62		2.30E-01	e				2.30E-01
Ethylene thiourea	96-45-7	-0.64		2.13E-02	e				2.13E-02
Furfural	98-01-1	0.96		4.37E-01	e				4.37E-01
Maleic hydrazide	123-33-1	-0.74		1.77E-02	e				1.77E-02
Malononitrile	109-77-3	0.04		7.70E-02	e				7.70E-02
Methyl styrene (mixed isomers)	25013-15-4	3.35		3.96E+01	e				3.96E+01
Methylhydrazine	60-34-4	-1.06		9.68E-03	e				9.68E-03
N,N-Diphenylamine	122-39-4	3.50		5.25E+01	e				5.25E+01
Nitric acid, propyl ester	627-13-4	No data		No data			1.00E+05	Default	1.00E+05
N-Nitrosodi-n-butylamine	924-16-3	2.41		6.73E+00	e				6.73E+00

Table C2-8. Aquatic Water-to-Invertebrate Transfer Factors (BCF_{inv}) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{inv} [b]	BCF_{inv} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{inv}	SAIC Compiled BCF_{inv}	Comments	Recommended BCF_{inv} [d]
N-Nitrosomorpholine	59-89-2	0.98		4.54E-01	e				4.54E-01
N-Nitroso-N,N-dimethylamine	62-75-9	-0.47		2.94E-02	e				2.94E-02
o-Anisidine	90-04-0	1.18		6.61E-01	e				6.61E-01
Oxalic acid	144-62-7	No data		No data			1.00E+05	Default	1.00E+05
Phthalic anhydride	85-44-9	-0.60		2.30E-02	e				2.30E-02
p-Phthalic acid	100-21-0	0.82		3.37E-01	e				3.37E-01
Pyridine	110-86-1	0.67		2.53E-01	e				2.53E-01
Quinoline	91-22-5	2.03		3.29E+00	e				3.29E+00
Quinone	106-51-4	0.20		1.04E-01	e				1.04E-01
Safrole	94-59-7	2.66		1.08E+01	e				1.08E+01
Tetrahydrofuran	109-99-9	0.45		1.66E-01	e				1.66E-01
Other Heavy Semivolatile Compounds (molecular weight >200 g/mole)									
1,2,4,5-Tetrachlorobenzene	95-94-3	4.64		4.51E+02	e				4.51E+02
1,3,5-Trinitrobenzene	99-35-4	1.18		6.60E-01	e				6.60E-01
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	4.17		1.86E+02	e				1.86E+02
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	-2.70		4.39E-04	e				4.39E-04
2-sec-Butyl-4,6-dinitrophenol	88-85-7	3.14		2.66E+01	e				2.66E+01
3,3'-Dimethoxybenzidine	119-90-4	1.81		2.17E+00	e				2.17E+00
3,3-Dichlorobenzidine	91-94-1	3.58		6.05E+01	e				6.05E+01
4-Bromophenylphenyl ether	101-55-3	5.00		8.89E+02	e				8.89E+02
Ammonium perfluorooctanoate	3825-26-1	No data		No data			1.00E+05	Default	1.00E+05
Azobenzene	103-33-3	3.82		9.61E+01	e				9.61E+01
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	No data		No data			1.00E+05	Default	1.00E+05
Captan	133-06-2	2.35		6.01E+00	e				6.01E+00
Chlorobenzilate	510-15-6	4.38		2.76E+02	e				2.76E+02
Dibutylphosphate	107-66-4	No data		No data			1.00E+05	Default	1.00E+05
Dimethyl aminoazobenzene	60-11-7	4.58		4.03E+02	e				4.03E+02
Hexachlorobenzene	118-74-1	5.50	2.60E+03						2.60E+03
Hexachlorobutadiene	87-68-3	4.73	1.05E+01						1.05E+01

Table C2-8. Aquatic Water-to-Invertebrate Transfer Factors (BCF_{inv}) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{inv} [b]	BCF_{inv} Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance BCF_{inv}	SAIC Compiled BCF_{inv}	Comments	Recommended BCF_{inv} [d]
Hexachlorocyclopentadiene	77-47-4	5.04	1.23E+03						1.23E+03
Hexachloroethane	67-72-1	3.98		1.31E+02	e				1.31E+02
Hexachlorophene	70-30-4	7.54		1.07E+05	e				1.07E+05
Hexamethylene-1,5-diisocyanate	822-06-0	1.27		7.89E-01	e				7.89E-01
Mirex	2385-85-5	6.89		3.14E+04	e				3.14E+04
Nitrofen	1836-75-5	5.53		2.42E+03	e				2.42E+03
Pentachlorobenzene	608-93-5	5.09	2.60E+03		j				2.60E+03
Pentachloronitrobenzene	82-68-8	4.64	1.30E+01		i				#REF!
Pentachlorophenol	87-86-5	5.08	5.20E+01						5.20E+01
Picric acid	88-89-1	2.03		3.29E+00	e				3.29E+00
Pronamide	23950-58-5	3.51		5.36E+01	e				5.36E+01
Strychnine	57-24-9	1.93		2.72E+00	e				2.72E+00
Terphenyls	26140-60-3	No data		No data			1.00E+05	Default	1.00E+05
Tributyl phosphate	126-73-8	4.00		1.35E+02	e				1.35E+02
Trifluralin	1582-09-8	5.34		1.69E+03	e				1.69E+03
Triphenylamine	603-34-9	No data		No data			1.00E+05	Default	1.00E+05
Herbicides and Organochlorinated Pesticides									
2,4,5-T	93-76-5	3.36		4.03E+01	e				4.03E+01
2,4-D and esters	94-75-7	2.81		1.43E+01	e				1.43E+01
4,4-DDD	72-54-8	6.20		8.55E+03	e				8.55E+03
4,4-DDE	72-55-9	6.26	1.19E+04						1.19E+04
4,4-DDT	50-29-3	6.00		5.86E+03	e				5.86E+03
Aldrin	309-00-2	6.18		8.21E+03	e				8.21E+03
alpha-BHC	319-84-6	3.80		9.24E+01	e				9.24E+01
beta-BHC	319-85-7	3.83		9.85E+01	e				9.85E+01
Chlordane	57-74-9	5.94		5.21E+03	e				5.21E+03
Delta-BHC	319-86-8	4.14		1.76E+02	e				1.76E+02
Dieldrin	60-57-1	5.27		1.48E+03	e				1.48E+03
Endothall	145-73-3	-0.87		1.39E-02	e				1.39E-02

Table C2-8. Aquatic Water-to-Invertebrate Transfer Factors (BCF_{inv}) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{inv} [b]	BCF_{inv} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{inv}	SAIC Compiled BCF_{inv}	Comments	Recommended BCF_{inv} [d]
Endrin	72-20-8	4.89		7.25E+02	e				7.25E+02
gamma-BHC (Lindane)	58-89-9	3.72		7.96E+01	e				7.96E+01
Heptachlor	76-44-8	5.02	3.81E+03						3.81E+03
Isodrin	465-73-6	3.55		5.77E+01	e				5.77E+01
Methoxychlor	72-43-5	4.53		3.64E+02	e				3.64E+02
Silvex (2,4,5-TP)	93-72-1	4.07		1.55E+02	e				1.55E+02
Toxaphene	8001-35-2	5.50		2.28E+03	e				2.28E+03
Inorganic Chemicals and Compounds									
Metals									
Aluminum	7429-90-5	NA	4.07E+03		k				4.07E+03
Antimony	7440-36-0	NA	7.00E+00						7.00E+00
Arsenic	7440-38-2	NA	7.30E+01						7.30E+01
Barium	7440-39-3	NA	2.00E+02						2.00E+02
Beryllium	7440-41-7	NA	4.50E+01						4.50E+01
Bismuth	7440-69-9	NA		4.07E+03	k				4.07E+03
Boron	7440-42-8	NA		4.07E+03	k				4.07E+03
Cadmium	7440-43-9	NA	3.46E+03						3.46E+03
Calcium	7440-70-2	NA		4.07E+03	k				4.07E+03
Chromium (and VI)	18540-29-9	NA	3.00E+03						3.00E+03
Cobalt	7440-48-4	NA		4.07E+03	k				4.07E+03
Copper	7440-50-8	NA	3.72E+03						3.72E+03
Iron	7439-89-6	NA		4.07E+03	k				4.07E+03
Lead	7439-92-1	NA	5.06E+03						5.06E+03
Lithium	7439-93-2	NA		4.07E+03	k				4.07E+03
Magnesium	7439-95-4	NA		4.07E+03	k				4.07E+03
Manganese	7439-96-5	NA		4.07E+03	k				4.07E+03
Mercury	7439-97-6	NA	2.02E+04						2.02E+04
Molybdenum	7439-98-7	NA		4.07E+03	k				4.07E+03
Nickel	7440-02-0	NA	2.80E+01						2.80E+01

Table C2-8. Aquatic Water-to-Invertebrate Transfer Factors (BCF_{inv}) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{inv} [b]	BCF_{inv} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{inv}	SAIC Compiled BCF_{inv}	Comments	Recommended BCF_{inv} [d]
Potassium	7440-09-7	NA		4.07E+03	k				4.07E+03
Rhodium	7440-16-6	NA		4.07E+03	k				4.07E+03
Selenium	7782-49-2	NA	1.26E+03						1.26E+03
Silicon	7440-21-3	NA		4.07E+03	k				4.07E+03
Silver	7440-22-4	NA	2.98E+02						2.98E+02
Sodium	7440-23-5	NA		4.07E+03	k				4.07E+03
Strontium	7440-24-6	NA		4.07E+03	k				4.07E+03
Tantalum	7440-25-7	NA		4.07E+03	k				4.07E+03
Thallium	7440-28-0	NA	1.50E+04						1.50E+04
Tin	7440-31-5	NA		4.07E+03	k				4.07E+03
Tungsten	7440-33-7	NA		4.07E+03	k				4.07E+03
Uranium	7440-61-1	NA		4.07E+03	k				4.07E+03
Vanadium	7440-62-2	NA		4.07E+03	k				4.07E+03
Yttrium	7440-65-5	NA		4.07E+03	k				4.07E+03
Zinc	7440-66-6	NA	4.58E+03						4.58E+03
Zirconium	7440-67-7	NA		4.07E+03	k				4.07E+03
Non-metals and Anions									
Ammonia/Ammonium	7664-41-7	NA					5.00E+02	Default	5.00E+02
Bromide	24959-67-9	NA		4.07E+03	k				4.07E+03
Chloride	16887-00-6	NA		4.07E+03	k				4.07E+03
Cyanide	57-12-5	NA	4.07E+03		k				4.07E+03
Fluoride	16984-48-8	NA		4.07E+03	k				4.07E+03
Hydroxide	14280-30-9	NA		NA				Depends on pH	NA
Iodine	7553-56-2	NA		4.07E+03	k				4.07E+03
Nitrate	14797-55-8	NA		4.07E+03	k				4.07E+03
Nitrite	14797-65-0	NA		4.07E+03	k				4.07E+03
Phosphate	14265-44-2	NA		4.07E+03	k				4.07E+03
Phosphorus	7723-14-0	NA		4.07E+03	k				4.07E+03
Sulfate	14808-79-8	NA		4.07E+03	k				4.07E+03

Table C2-8. Aquatic Water-to-Invertebrate Transfer Factors (BCF_{inv}) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{inv} [b]	BCF_{inv} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{inv}	SAIC Compiled BCF_{inv}	Comments	Recommended BCF_{inv} [d]
Total Sulfur	63705-05-5	NA		4.07E+03	k				4.07E+03
Priority Pollutants									
Carbon Dioxide	124-38-9	NA		NA					NA
Nitrogen Dioxide	10102-44-0	NA					5.00E+02	Default	5.00E+02
Ozone	10028-15-6	NA					5.00E+02	Default	5.00E+02
Particulate Matter	No CAS #	NA		NA					NA
Sulfur Dioxode	7446-09-5	NA					5.00E+02	Default	5.00E+02
Radionuclides									
Americium-241	1596-10-2	NA		4.07E+03	k				4.07E+03
Antimony-125	14234-35-6	NA	7.00E+00						7.00E+00
Barium-137	13981-97-0	NA	2.00E+02						2.00E+02
Cadmium-113	None	NA	3.46E+03						3.46E+03
Cesium-134	13967-70-9	NA		4.07E+03	k				4.07E+03
Cesium-137	10045-97-3	NA		4.07E+03	k				4.07E+03
Europium-154	15585-10-1	NA		4.07E+03	k				4.07E+03
Europium-155	14391-16-3	NA		4.07E+03	k				4.07E+03
Nickel-63	13981-37-8	NA	2.80E+01						2.80E+01
Plutonium-239	15117-48-3	NA		4.07E+03	k				4.07E+03
Plutonium-241	14119-32-5	NA		4.07E+03	k				4.07E+03
Samarium-151	15715-94-3	NA		4.07E+03	k				4.07E+03
Strontium-90	10098-97-2	NA		4.07E+03	k				4.07E+03
Technetium-99	14133-79-7	NA		4.07E+03	k				4.07E+03
Tritium	10028-17-8	NA		4.07E+03	k				4.07E+03
Yttrium-90	10098-91-6	NA		4.07E+03	k				4.07E+03
Uranium-232	14158-29-3	NA		4.07E+03	k				4.07E+03
Uranium-233	13968-55-3	NA		4.07E+03	k				4.07E+03
Uranium-234	13966-29-5	NA		4.07E+03	k				4.07E+03
Uranium-235	15117-96-1	NA		4.07E+03	k				4.07E+03
Uranium-236	13982-70-2	NA		4.07E+03	k				4.07E+03

Table C2-8. Aquatic Water-to-Invertebrate Transfer Factors (BCF_{inv}) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{inv} [b]	BCF_{inv} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{inv}	SAIC Compiled BCF_{inv}	Comments	Recommended BCF_{inv} [d]
Uranium-238	7440-61-1	NA		4.07E+03	k				4.07E+03
Actinium-227	14952-40-0	NA		4.07E+03	k				4.07E+03
Americium-243	14993-75-0	NA		4.07E+03	k				4.07E+03
Carbon-14	14762-75-5	NA		4.07E+03	k				4.07E+03
Cobalt-60	10198-40-0	NA		4.07E+03	k				4.07E+03
Curium-242	15510-73-3	NA		4.07E+03	k				4.07E+03
Curium-243	15757-87-6	NA		4.07E+03	k				4.07E+03
Curium-244	13981-15-2	NA		4.07E+03	k				4.07E+03
Europium-152	14683-23-9	NA		4.07E+03	k				4.07E+03
Iodine-129	15046-84-1	NA		4.07E+03	k				4.07E+03
Neptunium-237	13994-20-2	NA		4.07E+03	k				4.07E+03
Nickel-59	14336-70-0	NA	2.80E+01						2.80E+01
Niobium-93	None	NA		4.07E+03	k				4.07E+03
Plutonium-238	13981-16-3	NA		4.07E+03	k				4.07E+03
Plutonium-240	14119-33-6	NA		4.07E+03	k				4.07E+03
Plutonium-242	13982-10-0	NA		4.07E+03	k				4.07E+03
Protactinium-231	14331-85-2	NA		4.07E+03	k				4.07E+03
Radium-226	13982-63-3	NA		4.07E+03	k				4.07E+03
Radium-228	15262-20-1	NA		4.07E+03	k				4.07E+03
Ruthenium-106	13967-48-1	NA		4.07E+03	k				4.07E+03
Selenium-79	None	NA	1.26E+03						1.26E+03
Thorium-229	15594-54-4	NA		4.07E+03	k				4.07E+03
Thorium-232	7440-29-1	NA		4.07E+03	k				4.07E+03
Tin-126	15832-50-5	NA		4.07E+03	k				4.07E+03
Zirconium-93	15751-77-6	NA		4.07E+03	k				4.07E+03

NA = Not applicable

[a] $\log_{10}$ of K_{ow} values in Table 4.1

[b] Published in Appendix C of EPA (1999), Table C-3

[c] Calculated or chosen as described in Appendix C, Sect. C-1.3 of EPA (1999)

Table C2-8. Aquatic Water-to-Invertebrate Transfer Factors (BCF_{inv}) for Ecological Receptors (mg/kg tissue wet weight / mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{inv} [b]	BCF_{inv} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{inv}	SAIC Compiled BCF_{inv}	Comments	Recommended BCF_{inv} [d]

[d] Selection criteria described in Sect. 8.2.4.4

[c] Calculated by using log K_{ow}: log BCF-I = 0.819 x log K_{ow} -1.146 (Southworth, Beauchamp, and Schmieder 1978)

[f] Calculated by using the BCF_{inv} for TCDD from Appendix C, Table C-3 of EPA (1999) and BEFs for other congeners

[g] Value for Aroclor 1254 from Appendix C of EPA (1999), Table C-3, was used for PCB mixtures

[h] Benzo(a)pyrene value from Appendix C of EPA (1999), Table C-3, was used as a surrogate value

[i] 2,4-dinitrotoluene value from Appendix C of EPA (1999), Table C-3, was used as a surrogate value

[j] Pentachlorobenzene value from Appendix C of EPA (1999), Table C-3, was used as a surrogate value

[k] Average of values for other inorganics as published in Appendix C of EPA (1999), Table C-3

Table C2-9. Aquatic Water-to-Fish Transfer Factors (BCF_{fish}) for Ecological Receptors (mg/kg tissue wet weight/mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{fish} [b]	BCF_{fish} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{fish}	SAIC compiled BCF_{fish}	Notes	Comments	Recommended BCF_{fish} [d]
Organic Compounds										
Aromatic Halogenated Hydrocarbons										
4-Chloro-3-methylphenol	59-50-7	3.10		1.08E+02	e					1.08E+02
2,3,4,6-Tetrachlorophenol	58-90-2	4.42		1.66E+03	e					1.66E+03
Aromatic Nonhalogenated Hydrocarbons										
2-Nitrotoluene	88-72-2	2.30		2.03E+01	e					2.03E+01
4-Nitrobiphenyl	92-93-3	3.77		4.38E+02	e					4.38E+02
Benzaldehyde	100-52-7	1.48		3.62E+00	e					3.62E+00
Benzene	71-43-2	-4.99		4.70E-06	e		3.20E+01	f		4.70E-06
Benzyl alcohol	100-51-6	1.10		1.64E+00	e					1.64E+00
Ethyl benzene	100-41-4	3.12		1.14E+02	e		2.90E+02	f		1.14E+02
m-Xylene	108-38-3	3.20		1.34E+02	e		1.70E+01	f		1.34E+02
o-Xylene	95-47-6	3.13		1.15E+02	e		1.70E+01	f		1.15E+02
p-Xylene	106-42-3	3.17		1.25E+02	e		1.70E+01	f		1.25E+02
Styrene	100-42-5	2.93		7.57E+01	e					7.57E+01
Toluene	108-88-3	2.67		4.38E+01	e		8.30E+01	f		4.38E+01
Non-aromatic Nonhalogenated Hydrocarbons										
1,2-Epoxybutane	106-88-7	1.44		3.35E+00	e					3.35E+00
1,3-Butadiene	106-99-0	1.90		8.77E+00	e					8.77E+00
1,4-Dioxane	123-91-1	-0.27		9.34E-02	e					9.34E-02
1-Methylpropyl alcohol	78-92-2	0.61		5.88E-01	e					5.88E-01
1-Nitropropane	108-03-2	0.87		1.01E+00	e					1.01E+00
2,2,4-Trimethylpentane	540-84-1	5.02		5.29E+03	e					5.29E+03
2-Butanone	78-93-3	0.28		2.95E-01	e		6.00E-01	f		2.95E-01
2-Butenaldehyde (2-Butenal)	4170-30-3	No data		No data			1.00E+05		Default	1.00E+05
2-Ethoxyethanol	110-80-5	-0.10		1.33E-01	e					1.33E-01
2-Heptanone	110-43-0	1.98		1.04E+01	e					1.04E+01
2-Hexanone	591-78-6	1.38		2.95E+00	e					2.95E+00
2-Methoxyethanol	109-86-4	0.25		2.78E-01	e					2.78E-01

Table C2-9. Aquatic Water-to-Fish Transfer Factors (BCF_{fish}) for Ecological Receptors (mg/kg tissue wet weight/mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{fish} [b]	BCF_{fish} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{fish}	SAIC compiled BCF_{fish}	Notes	Comments	Recommended BCF_{fish} [d]
2-Methyl-2-propanol	75-65-0	0.35		3.41E-01	e					3.41E-01
2-Methyl-2-propenenitrile	126-98-7	0.54		5.08E-01	e					5.08E-01
2-Methylaziridine	75-55-8	-0.60		4.63E-02	e					4.63E-02
2-Methylpropyl alcohol	78-83-1	0.76		8.05E-01	e					8.05E-01
2-Pentanone	107-87-9	0.91		1.10E+00	e					1.10E+00
2-Propanone (Acetone)	67-64-1	-0.22	1.03E-01		e		2.00E-01	f		1.03E-01
2-Propene-1-ol	107-18-6	0.17		2.34E-01	e					2.34E-01
2-Propyl alcohol	67-63-0	0.05		1.82E-01	e					1.82E-01
3-Heptanone	106-35-4	No data		No data			1.00E+05		Default	1.00E+05
3-Methyl-1-butanol	123-51-3	No data		No data			1.00E+05		Default	1.00E+05
3-Methyl-2-butanone	563-80-4	No data		No data			1.00E+05		Default	1.00E+05
3-Pentanone	96-22-0	0.99		1.30E+00	e					1.30E+00
4-Heptanone	123-19-3	No data		No data			1.00E+05		Default	1.00E+05
4-Methyl-2-pentanone	108-10-1	1.19		1.98E+00	e		6.00E+00	f		1.98E+00
4-Methyl-3-penten-2-one	141-79-7	No data		No data			1.00E+05		Default	1.00E+05
5-Methyl-2-hexanone	110-12-3	No data		No data			1.00E+05		Default	1.00E+05
Acetaldehyde	75-07-0	-0.22		1.03E-01	e					1.03E-01
Acetamide	60-35-5	-1.26		1.17E-02	e					1.17E-02
Acetic acid	64-19-7	-0.17		1.15E-01	e					1.15E-01
Acetic acid ethyl ester	141-78-6	0.73		7.56E-01	e					7.56E-01
Acetic acid n-butyl ester	123-86-4	1.73		6.14E+00	e					6.14E+00
Acetonitrile	75-05-8	-0.34		8.03E-02	e					8.03E-02
Acrolein	107-02-8	-0.01		1.61E-01	e					1.61E-01
Acrylonitrile	107-13-1	0.25	4.80E+01							4.80E+01
Bis(isopropyl)ether	108-20-3	No data		No data			1.00E+05		Default	1.00E+05
Butane	106-97-8	2.89		6.97E+01	e					6.97E+01
Carbon disulfide	75-15-0	2.00		1.08E+01	e					1.08E+01
Cyanogen	460-19-5	0.81		8.89E-01	e					8.89E-01
Cyclohexane	110-82-7	3.44		2.20E+02	e					2.20E+02

Table C2-9. Aquatic Water-to-Fish Transfer Factors (BCF_{fish}) for Ecological Receptors (mg/kg tissue wet weight/mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{fish} [b]	BCF_{fish} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{fish}	SAIC compiled BCF_{fish}	Notes	Comments	Recommended BCF_{fish} [d]
Cyclohexanone	108-94-1	0.81		8.94E-01	e					8.94E-01
Cyclohexene	110-83-8	2.86		6.55E+01	e					6.55E+01
Cyclopentane	287-92-3	3.00		8.78E+01	e					8.78E+01
Ethyl alcohol	64-17-5	0.31		3.13E-01	e					3.13E-01
Ethyl ether	60-29-7	0.89		1.06E+00	e					1.06E+00
Ethyl methacrylate	97-63-2	1.59		4.58E+00	e					4.58E+00
Formaldehyde	50-00-0	0.34	3.35E-01		e					3.35E-01
Formamide	75-12-7	-1.51		6.92E-03	e					6.92E-03
Formic acid	64-18-6	-0.54		5.31E-02	e					5.31E-02
Formic acid, methyl ester	107-31-3	-0.26		9.41E-02	e					9.41E-02
Glycidylaldehyde	765-34-4	-0.73		3.55E-02	e					3.55E-02
Methyl acetate	79-20-9	0.18		2.39E-01	e					2.39E-01
Methyl alcohol	67-56-1	-0.71		3.70E-02	e					3.70E-02
Methyl isocyanate	624-83-9	No data		No data			1.00E+05		Default	1.00E+05
Methyl methacrylate	80-62-6	0.79		8.57E-01	e					8.57E-01
Methyl tert-butyl ether	1634-04-4	0.94		1.17E+00	e					1.17E+00
Methylacetylene	74-99-7	0.94		1.17E+00	e					1.17E+00
Methylcyclohexane	108-87-2	4.10		8.66E+02	e					8.66E+02
N,N-Dimethylacetamide	127-19-5	No data		No data			1.00E+05		Default	1.00E+05
n-Butyl alcohol	71-36-3	0.88		1.03E+00	e					1.03E+00
n-Heptane	142-82-5	4.66		2.68E+03	e					2.68E+03
n-Hexane	110-54-3	4.11		8.84E+02	e					8.84E+02
Nitromethane	75-52-5	-0.35		7.86E-02	e					7.86E-02
n-Nonane	111-84-2	5.65		1.34E+04	e					1.34E+04
n-Octane	111-65-9	4.00		7.05E+02	e					7.05E+02
n-Pentane	109-66-0	3.21		1.36E+02	e					1.36E+02
n-Propionaldehyde	123-38-6	0.59		5.64E-01	e					5.64E-01
n-Propyl alcohol	71-23-8	0.25		2.76E-01	e					2.76E-01
n-Valeraldehyde	110-62-3	No data		No data			1.00E+05		Default	1.00E+05

Table C2-9. Aquatic Water-to-Fish Transfer Factors (BCF_{fish}) for Ecological Receptors (mg/kg tissue wet weight/mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{fish} [b]	BCF_{fish} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{fish}	SAIC compiled BCF_{fish}	Notes	Comments	Recommended BCF_{fish} [d]
Oxirane	75-21-8	-0.30		8.73E-02	e					8.73E-02
p-Cymene	99-87-6	4.10		8.66E+02	e					8.66E+02
Phosgene	75-44-5	No data		No data			1.00E+05		Default	1.00E+05
Propargyl alcohol	107-19-7	0.26		2.82E-01	e					2.82E-01
Propionic acid	79-09-4	0.33		3.27E-01	e					3.27E-01
Propionitrile	107-12-0	0.16		2.29E-01	e					2.29E-01
Propylene gylcol monomethyl ether	107-98-2	-0.18		1.12E-01	e					1.12E-01
p-tert-Butyltoluene	98-51-1	No data		No data			1.00E+05		Default	1.00E+05
Triethylamine	121-44-8	0.16		2.29E-01	e					2.29E-01
Trimethylamine	75-50-3	0.16		2.29E-01	e					2.29E-01
Vinyl acetate	108-05-4	0.70		7.08E-01	e					7.08E-01
Non-aromatic Halogenated Hydrocarbons										
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	No data		No data			1.00E+05		Default	1.00E+05
1,1,1,2-Tetrachloroethane	630-20-6	2.63		4.05E+01	e					4.05E+01
1,1,1-Trichloroethane	71-55-6	2.42		2.62E+01	e					2.62E+01
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	3.73		4.03E+02	e					4.03E+02
1,1,2,2-Tetrachloroethane	79-34-5	4.64		2.60E+03	e					2.60E+03
1,1,2,2-Tetrachloroethene	127-18-4	2.55		3.39E+01	e		4.40E+01	f		3.39E+01
1,1,2-Trichloroethane	79-00-5	2.10		1.32E+01	e					1.32E+01
1,1,2-Trichloroethylene	79-01-6	2.43		2.68E+01	e		1.70E+01	f		2.68E+01
1,1-Dichloroethane	75-34-3	1.79		7.00E+00	e					7.00E+00
1,1-Dichloroethene	75-35-4	2.12		1.39E+01	e					1.39E+01
1,2,2-Trichloro-1,1,2-trifluoroethane	76-13-1	3.16		1.23E+02	e					1.23E+02
1,2,3-Trichloropropane	96-18-4	2.25		1.83E+01	e					1.83E+01
1,2-Dibromo-3-chloropropane	96-12-8	2.34		2.21E+01	e					2.21E+01
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	2.82		6.02E+01	e					6.02E+01
1,2-Dichloroethane	107-06-2	1.46		3.51E+00	e		2.00E+00	f		3.51E+00
1,2-Dichloroethylene	540-59-0	0.48		4.48E-01	e					4.48E-01
1,2-Dichloropropane	78-87-5	2.25		1.83E+01	e					1.83E+01

Table C2-9. Aquatic Water-to-Fish Transfer Factors (BCF_{fish}) for Ecological Receptors (mg/kg tissue wet weight/mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{fish} [b]	BCF_{fish} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{fish}	SAIC compiled BCF_{fish}	Notes	Comments	Recommended BCF_{fish} [d]
1,3-Dichloropropene	542-75-6	1.75		6.38E+00	e					6.38E+00
1,4-Dichloro-2-butene	764-41-0	0.87		1.02E+00	e					1.02E+00
1-Chloroethene	75-01-4	1.15		1.81E+00	e					1.81E+00
2,2-Dichloropropionic acid	75-99-0	0.78		8.36E-01	e					8.36E-01
2-Chloropropane	75-29-6	1.90		8.77E+00	e					8.77E+00
3-Chloropropene (allyl chloride)	107-05-1	0.95		1.21E+00	e					1.21E+00
Bromochloromethane	74-97-5	1.41		3.14E+00	e					3.14E+00
Bromodichloromethane	75-27-4	2.03		1.14E+01	e					1.14E+01
Bromoethene	593-60-2	1.07		1.53E+00	e					1.53E+00
Bromoform	75-25-2	2.35		2.25E+01	e					2.25E+01
Bromomethane	74-83-9	1.11		1.69E+00	e					1.69E+00
Carbon tetrachloride	56-23-5	2.72	3.00E+01							3.00E+01
Chlorodibromomethane	124-48-1	2.18		1.56E+01	e					1.56E+01
Chlorodifluoromethane	75-45-6	1.08		1.57E+00	e					1.57E+00
Chloroethane	75-00-3	3.10		1.08E+02	e					1.08E+02
Chloroform	67-66-3	1.95	3.59E+00				6.00E+00	f		3.59E+00
Chloromethane	74-87-3	0.90		1.09E+00	e					1.09E+00
Chloropentafluoroethane	76-15-3	No data		No data			1.00E+05		Default	1.00E+05
cis-1,2-Dichloroethene	156-59-2	1.98		1.04E+01	e		8.60E-01	f		1.04E+01
cis-1,3-Dichloropropene	10061-01-5	No data		No data			1.00E+05		Default	1.00E+05
Cyanogen bromide	506-68-3	No data		No data			1.00E+05		Default	1.00E+05
Cyanogen chloride	506-77-4	0.20		2.49E-01	e					2.49E-01
Dichlorodifluoromethane	75-71-8	2.16		1.51E+01	e					1.51E+01
Dichlorofluoromethane	75-43-4	No data		No data			1.00E+05		Default	1.00E+05
Dichloromethane	75-09-2	1.26		2.27E+00	e					2.27E+00
Difluorodibromomethane	75-61-6	No data		No data			1.00E+05		Default	1.00E+05
Hexafluoroacetone	684-16-2	No data		No data			1.00E+05		Default	1.00E+05
Iodomethane	74-88-4	1.69		5.65E+00	e					5.65E+00
Methylene bromide	74-95-3	1.62		4.88E+00	e					4.88E+00

Table C2-9. Aquatic Water-to-Fish Transfer Factors (BCF_{fish}) for Ecological Receptors (mg/kg tissue wet weight/mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{fish} [b]	BCF_{fish} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{fish}	SAIC compiled BCF_{fish}	Notes	Comments	Recommended BCF_{fish} [d]
Pentachloroethane	76-01-7	3.05		9.75E+01	e					9.75E+01
trans-1,2-Dichloroethene	156-60-5	1.98		1.04E+01	e		8.60E-01	f		1.04E+01
trans-1,3-Dichloropropene	10061-02-6	2.06		1.23E+01	e					1.23E+01
Trichloroacetic acid	76-03-9	1.33		2.66E+00	e					2.66E+00
Trichlorofluoroethane	27154-33-2	No data		No data			1.00E+05		Default	1.00E+05
Trichlorofluoromethane	75-69-4	2.53		3.29E+01	e					3.29E+01
Trifluorobromomethane	75-63-8	1.86		8.06E+00	e					8.06E+00
Dioxin and Furan Compounds										
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	35822-46-9	8.20	2.16E+02		g					2.16E+02
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	7.92	4.66E+01		g					4.66E+01
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	7.92	1.65E+03		g					1.65E+03
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	39227-28-6	7.79	1.31E+03		g					1.31E+03
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	7.25	3.22E+02		g					3.22E+02
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	57653-85-7	7.25	5.08E+02		g					5.08E+02
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	7.25	8.05E+02		g					8.05E+02
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	19408-74-3	7.25	5.93E+02		g					5.93E+02
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	7.25	2.67E+03		g					2.67E+03
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	40321-76-4	6.64	3.90E+03		g					3.90E+03
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	6.79	9.32E+02		g					9.32E+02
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	7.25	2.84E+03		g					2.84E+03
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	6.92	6.78E+03		g					6.78E+03
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1746-01-6	6.64	4.24E+03							4.24E+03
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	6.53	3.39E+03		g					3.39E+03
Dibenzofuran	132-64-9	4.33		1.39E+03	e		6.90E+02	f		1.39E+03
Octachlorodibenzo(p)dioxin	3268-87-9	8.20	5.08E+01		g					5.08E+01
Octachlorodibenzofuran	39001-02-0	8.78	6.78E+01		g					6.78E+01
PCBs										
Polychlorinated biphenyls (PCBs) [h]	1336-36-3	7.31	2.30E+05							2.30E+05
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	7.08		5.70E+03	e					5.70E+03

Table C2-9. Aquatic Water-to-Fish Transfer Factors (BCF_{fish}) for Ecological Receptors (mg/kg tissue wet weight/mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{fish} [b]	BCF_{fish} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{fish}	SAIC compiled BCF_{fish}	Notes	Comments	Recommended BCF_{fish} [d]
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	7.12		5.26E+03	e					5.26E+03
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	No data		No data			1.00E+05		Default	1.00E+05
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	No data		No data			1.00E+05		Default	1.00E+05
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	No data		No data			1.00E+05		Default	1.00E+05
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	No data		No data			1.00E+05		Default	1.00E+05
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	No data		No data			1.00E+05		Default	1.00E+05
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	No data		No data			1.00E+05		Default	1.00E+05
2',3,4,4',5-Pentachlorobiphenyl	no cas #	No data		No data			1.00E+05		Default	1.00E+05
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	7.12		5.26E+03	e					5.26E+03
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	7.41		2.87E+03	e					2.87E+03
3,3',4,4',5-Pentachlorobiphenyl	no cas #	No data		No data			1.00E+05		Default	1.00E+05
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	No data		No data			1.00E+05		Default	1.00E+05
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	No data		No data			1.00E+05		Default	1.00E+05
Phthalates										
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	5.20	7.00E+01				3.10E+02	f		7.00E+01
Butylbenzyl phthalate	85-68-7	4.41		1.64E+03	e		6.60E+02	f		1.64E+03
Dibutyl phthalate	84-74-2	4.72		3.01E+03	e		5.10E+03	f		3.01E+03
Diethyl phthalate	84-66-2	4.44		1.72E+03	e		1.20E+02	f		1.72E+03
Dimethylphthalate	131-11-3	1.63		5.02E+00	e					5.02E+00
n-Dioctyl phthalate	117-84-0	9.33	9.40E+03				9.30E+03	f		9.40E+03
Light Polycyclic Aromatic Hydrocarbons (molecular weight <200 g/mole)										
2-Chloronapthalene	91-58-7	4.07		8.11E+02	e					8.11E+02
2-Methyl naphthalene	91-57-6	3.86		5.28E+02	e		4.30E+02	f		5.28E+02
5-Nitroacenaphthene	602-87-9	No data		No data			1.00E+05		Default	1.00E+05
Acenaphthene	83-32-9	3.96		6.55E+02	e					6.55E+02
Acenaphthylene	208-96-8	4.07		8.14E+02	e					8.14E+02
Anthracene	120-12-7	4.47		1.84E+03	e		1.40E+03	f		1.84E+03
Fluorene	86-73-7	4.18		1.02E+03	e					1.02E+03
Indene	95-13-6	No data		No data			1.00E+05		Default	1.00E+05

Table C2-9. Aquatic Water-to-Fish Transfer Factors (BCF_{fish}) for Ecological Receptors (mg/kg tissue wet weight/mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{fish} [b]	BCF_{fish} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{fish}	SAIC compiled BCF_{fish}	Notes	Comments	Recommended BCF_{fish} [d]
Naphthalene	91-20-3	3.37		1.91E+02	e		4.30E+02	f		1.91E+02
Phenanthrene	85-01-8	4.55		2.16E+03	e		1.40E+03	f		2.16E+03
Pyrene	129-00-0	5.00		5.10E+03	e		6.10E+03	f		5.10E+03
Heavy Polycyclic Aromatic Hydrocarbons (molecular weight >200 g/mole)										
3-Methylcholanthrene	56-49-5	7.11		5.37E+03	e					5.37E+03
5-Methylchrysene	3697-24-3	No data		No data			1.00E+05		Default	1.00E+05
Benzo(a)anthracene	56-55-3	5.68	5.00E+02		i		1.30E+04	f		5.00E+02
Benzo(a)pyrene	50-32-8	6.13	5.00E+02				3.00E+01	f		5.00E+02
Benzo(b)fluoranthene	205-99-2	6.20	5.00E+02		i		2.60E+04	f		5.00E+02
Benzo(e)pyrene	192-97-2	7.40		2.92E+03	e					2.92E+03
Benzo(g,h,i)perylene	191-24-2	7.10		5.48E+03	e		6.50E+04	f		5.48E+03
Benzo(j)fluoranthene	205-82-3	6.44		1.48E+04	e					1.48E+04
Benzo(k)fluoranthene	207-08-9	6.20	5.00E+02		i		2.60E+04	f		5.00E+02
Benzo[a,i]pyrene	191-30-0	No data		No data			1.00E+05		Default	1.00E+05
Chrysene	218-01-9	5.74	5.00E+02		i		1.30E+04	f		5.00E+02
Dibenz(a,h)anthracene	53-70-3	6.55	5.00E+02		i		5.40E+04	f		5.00E+02
Dibenz[a,h]acridine	226-36-8	No data		No data			1.00E+05		Default	1.00E+05
Dibenz[a,j]acridine	224-42-0	No data		No data			1.00E+05		Default	1.00E+05
Dibenzo(a,e)fluoranthene	5385-75-1	No data		No data			1.00E+05		Default	1.00E+05
Dibenzo(a,h)fluoranthene	no cas #	No data		No data			1.00E+05		Default	1.00E+05
Dibenzo[a,e]pyrene	192-65-4	No data		No data			1.00E+05		Default	1.00E+05
Dibenzo[a,h]pyrene	189-64-0	No data		No data			1.00E+05		Default	1.00E+05
Dibenzo[a,i]pyrene	189-55-9	7.29		3.70E+03	e					3.70E+03
Fluoranthene	206-44-0	5.08		5.91E+03	e		3.20E+03	f		5.91E+03
Hexachloronaphthalene	1335-87-1	7.59		1.90E+03	e					1.90E+03
Indeno(1,2,3-cd)pyrene	193-39-5	6.91	5.00E+02		i		6.50E+04	f		5.00E+02
Octachloronaphthalene	2234-13-1	6.42		1.50E+04	e					1.50E+04
Pentachloronaphthalene	1321-64-8	No data		No data			1.00E+05		Default	1.00E+05
Tetrachloronaphthalene	1335-88-2	No data		No data			1.00E+05		Default	1.00E+05

Table C2-9. Aquatic Water-to-Fish Transfer Factors (BCF_{fish}) for Ecological Receptors (mg/kg tissue wet weight/mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{fish} [b]	BCF_{fish} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{fish}	SAIC compiled BCF_{fish}	Notes	Comments	Recommended BCF_{fish} [d]
Trichloronaphthalene	1321-65-9	No data		No data			1.00E+05		Default	1.00E+05
Light Substituted Benzene Compounds (MW <200 g/mole)										
1,2,3-Trichlorobenzene	87-61-6	4.05		7.74E+02	e					7.74E+02
1,2,4-Trichlorobenzene	120-82-1	3.99		6.88E+02	e					6.88E+02
1,2,4-Trimethyl benzene	95-63-6	3.65		3.41E+02	e					3.41E+02
1,2-Dichlorobenzene	95-50-1	3.45		2.23E+02	e					2.23E+02
1,3,5-Trimethyl benzene	108-67-8	3.42		2.11E+02	e					2.11E+02
1,3-Dichlorobenzene	541-73-1	3.53		2.66E+02	e					2.66E+02
1,3-Dinitrobenzene	99-65-0	1.49	7.40E+01							7.40E+01
1,4-Dichlorobenzene	106-46-7	3.41		2.08E+02	e					2.08E+02
1,4-Dinitrobenzene	100-25-4	1.50		3.79E+00	e					3.79E+00
2,4,5-Trichlorophenol	95-95-4	3.87		5.39E+02	e					5.39E+02
2,4,6-Trichlorophenol	88-06-2	3.71		3.88E+02	e					3.88E+02
2,4-Dichlorophenol	120-83-2	3.04		9.49E+01	e					9.49E+01
2,4-Dimethylphenol	105-67-9	2.36		2.30E+01	e					2.30E+01
2,4-Dinitrophenol	51-28-5	1.52		3.94E+00	e					3.94E+00
2,4-Dinitrotoluene	121-14-2	2.00	2.10E+01		j					2.10E+01
2,6-Dinitrotoluene	606-20-2	1.89	2.10E+01		j		2.60E+01	f		2.10E+01
2-Chlorophenol	95-57-8	2.16		1.52E+01	e					1.52E+01
2-Chlorotoluene	95-49-8	3.54		2.71E+02	e					2.71E+02
2-Nitrophenol	88-75-5	1.79		6.97E+00	e					6.97E+00
4,6-Dinitro-o-cresol	534-52-1	2.85		6.41E+01	e					6.41E+01
4-Chlorotoluene	106-43-4	3.33		1.75E+02	e					1.75E+02
4-Nitrophenol	100-02-7	1.91		8.96E+00	e		1.30E+01	f		8.96E+00
alpha-Methylstyrene	98-83-9	3.46		2.31E+02	e					2.31E+02
Aniline	62-53-3	0.98		1.28E+00	e					1.28E+00
Benzotrichloride	98-07-7	2.92		7.43E+01	e					7.43E+01
Benzyl chloride	100-44-7	0.36		3.49E-01	e					3.49E-01
Bromobenzene	108-86-1	2.99		8.60E+01	e					8.60E+01

Table C2-9. Aquatic Water-to-Fish Transfer Factors (BCF_{fish}) for Ecological Receptors (mg/kg tissue wet weight/mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{fish} [b]	BCF_{fish} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{fish}	SAIC compiled BCF_{fish}	Notes	Comments	Recommended BCF_{fish} [d]
Chlorobenzene	108-90-7	2.79		5.65E+01	e		4.50E+02	f		5.65E+01
Cumene	98-82-8	3.61		3.16E+02	e					3.16E+02
m-Cresol	108-39-4	1.96		9.92E+00	e					9.92E+00
n-Butyl benzene	104-51-8	4.28		1.25E+03	e					1.25E+03
Nitrobenzene	98-95-3	1.83	2.10E+01							2.10E+01
n-Propyl benzene	103-65-1	3.69		3.71E+02	e					3.71E+02
o-Cresol	95-48-7	2.02		1.13E+01	e					1.13E+01
o-Dinitrobenzene	528-29-0	1.60		4.68E+00	e					4.68E+00
o-Nitroaniline	88-74-4	1.85		7.90E+00	e					7.90E+00
o-Toluidine	95-53-4	1.34		2.72E+00	e					2.72E+00
p-Chloroaniline	106-47-8	1.87		8.22E+00	e					8.22E+00
p-Cresol	106-44-5	1.94		9.53E+00	e					9.53E+00
Phenol	108-95-2	1.48		3.62E+00	e		7.80E+02	f		3.62E+00
p-Nitrochlorobenzene	100-00-5	2.39		2.45E+01	e					2.45E+01
p-Toluidine	106-49-0	1.40		3.08E+00	e					3.08E+00
sec-Butyl benzene	135-98-8	4.57		2.25E+03	e					2.25E+03
tert-Butyl benzene	98-06-6	4.11		8.84E+02	e					8.84E+02
Toluene-2,6-diamine	823-40-5	1.45		3.45E+00	e					3.45E+00
Trimethyl benzene	25551-13-7	3.40		2.03E+02	e					2.03E+02
Other Light Semivolatile Compounds (molecular weight <200 g/mole)										
1,1'-Biphenyl	92-52-4	3.90		5.73E+02	e					5.73E+02
1,1-Dimethylhydrazine	57-14-7	0.60		5.75E-01	e					5.75E-01
1,2-Dimethylhydrazine	540-73-8	-1.37		9.31E-03	e					9.31E-03
1,2-Diphenylhydrazine	122-66-7	2.94		7.74E+01	e					7.74E+01
1,3-Propane sultone	1120-71-4	-0.52		5.47E-02	e					5.47E-02
2,4-Toluene diisocyante	584-84-9	No data		No data					Decomposes rapidly in water	NA
2-Chloroacetophenone	532-27-4	2.59		3.76E+01	e					3.76E+01
2-Propenoic acid	79-10-7	0.33		3.27E-01	e					3.27E-01
4,4-Methylenedianiline	101-77-9	3.38		1.94E+02	e					1.94E+02

Table C2-9. Aquatic Water-to-Fish Transfer Factors (BCF_{fish}) for Ecological Receptors (mg/kg tissue wet weight/mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{fish} [b]	BCF_{fish} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{fish}	SAIC compiled BCF_{fish}	Notes	Comments	Recommended BCF_{fish} [d]
Acetophenone	98-86-2	1.64		5.09E+00	e					5.09E+00
Benzoic acid	65-85-0	1.86		8.06E+00	e					8.06E+00
bis(2-Chloroethoxy)methane	111-91-1	7.59		1.90E+03	e					1.90E+03
bis(2-Chloroethyl) ether	111-44-4	1.30		2.50E+00	e					2.50E+00
Chlorocyclopentadiene	41851-50-7	2.43		2.66E+01	e					2.66E+01
Cyclohexanol	108-93-0	1.23		2.15E+00	e					2.15E+00
Dichloroisopropyl ether	108-60-1	2.58		3.64E+01	e					3.64E+01
Dichloromethyl ether	542-88-1	-0.38		7.38E-02	e					7.38E-02
Dichloropentadiene	no cas #	No data		No data			1.00E+05		Default	1.00E+05
Dimethyl sulfate	77-78-1	0.32		3.19E-01	e					3.19E-01
Dimethylaniline	121-69-7	-0.88		2.59E-02	e					2.59E-02
Di-n-propylnitrosamine	621-64-7	1.38		2.95E+00	e					2.95E+00
Diphenyl ether	101-84-8	4.21		1.09E+03	e					1.09E+03
Epichlorohydrin	106-89-8	0.25		2.77E-01	e					2.77E-01
Ethyl carbamate (urethane)	51-79-6	-0.15		1.20E-01	e					1.20E-01
Ethyl methanesulfonate	62-50-0	0.05		1.81E-01	e					1.81E-01
Ethylene dibromide	106-93-4	1.75		6.40E+00	e					6.40E+00
Ethylene glycol	107-21-1	-0.91		2.41E-02	e					2.41E-02
Ethylene glycol monobutyl ether	111-76-2	1.55		4.25E+00	e					4.25E+00
Ethylene glycol monoethyl ether acetate	111-15-9	0.62		5.99E-01	e					5.99E-01
Ethylene thiourea	96-45-7	-0.64		4.27E-02	e					4.27E-02
Furfural	98-01-1	0.96		1.22E+00	e					1.22E+00
Maleic hydrazide	123-33-1	-0.74		3.47E-02	e					3.47E-02
Malononitrile	109-77-3	0.04		1.78E-01	e					1.78E-01
Methyl styrene (mixed isomers)	25013-15-4	3.35		1.82E+02	e					1.82E+02
Methylhydrazine	60-34-4	-1.06		1.78E-02	e					1.78E-02
N,N-Diphenylamine	122-39-4	3.50		2.50E+02	e					2.50E+02
Nitric acid, propyl ester	627-13-4	No data		No data			1.00E+05		Default	1.00E+05
N-Nitrosodi-n-butylamine	924-16-3	2.41		2.55E+01	e					2.55E+01

Table C2-9. Aquatic Water-to-Fish Transfer Factors (BCF_{fish}) for Ecological Receptors (mg/kg tissue wet weight/mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{fish} [b]	BCF_{fish} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{fish}	SAIC compiled BCF_{fish}	Notes	Comments	Recommended BCF_{fish} [d]
N-Nitrosomorpholine	59-89-2	0.98		1.28E+00	e					1.28E+00
N-Nitroso-N,N-dimethylamine	62-75-9	-0.47		6.11E-02	e					6.11E-02
o-Anisidine	90-04-0	1.18		1.94E+00	e					1.94E+00
Oxalic acid	144-62-7	No data		No data			1.00E+05		Default	1.00E+05
Phthalic anhydride	85-44-9	-0.60		4.64E-02	e					4.64E-02
p-Phthalic acid	100-21-0	0.82		9.17E-01	e					9.17E-01
Pyridine	110-86-1	0.67		6.67E-01	e					6.67E-01
Quinoline	91-22-5	2.03		1.15E+01	e					1.15E+01
Quinone	106-51-4	0.20		2.49E-01	e					2.49E-01
Safrole	94-59-7	2.66		4.31E+01	e					4.31E+01
Tetrahydrofuran	109-99-9	0.45		4.18E-01	e					4.18E-01
Other Heavy Semivolatile Compounds (molecular weight >200 g/mole)										
1,2,4,5-Tetrachlorobenzene	95-94-3	4.64		2.58E+03						2.58E+03
1,3,5-Trinitrobenzene	99-35-4	1.18		1.94E+00						1.94E+00
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	4.17		1.00E+03						1.00E+03
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	-2.70		5.71E-04						5.71E-04
2-sec-Butyl-4,6-dinitrophenol	88-85-7	3.14		1.18E+02						1.18E+02
3,3'-Dimethoxybenzidine	119-90-4	1.81		7.27E+00						7.27E+00
3,3-Dichlorobenzidine	91-94-1	3.58		2.92E+02						2.92E+02
4-Bromophenylphenyl ether	101-55-3	5.00		5.10E+03						5.10E+03
Ammonium perfluorooctanoate	3825-26-1	No data		No data			1.00E+05		Default	1.00E+05
Azobenzene	103-33-3	3.82		4.86E+02						4.86E+02
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	No data		No data			1.00E+05		Default	1.00E+05
Captan	133-06-2	2.35		2.25E+01	e					2.25E+01
Chlorobenzilate	510-15-6	4.38		1.53E+03						1.53E+03
Dibutylphosphate	107-66-4	No data		No data			1.00E+05		Default	1.00E+05
Dimethyl aminoazobenzene	60-11-7	4.58		2.29E+03	e					2.29E+03
Hexachlorobenzene	118-74-1	5.50	2.53E+02							2.53E+02
Hexachlorobutadiene	87-68-3	4.73	7.83E+02							7.83E+02

Table C2-9. Aquatic Water-to-Fish Transfer Factors (BCF_{fish}) for Ecological Receptors (mg/kg tissue wet weight/mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{fish} [b]	BCF_{fish} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{fish}	SAIC compiled BCF_{fish}	Notes	Comments	Recommended BCF_{fish} [d]
Hexachlorocyclopentadiene	77-47-4	5.04	1.65E+02							1.65E+02
Hexachloroethane	67-72-1	3.98		6.83E+02	e					6.83E+02
Hexachlorophene	70-30-4	7.54		2.13E+03	e					2.13E+03
Hexamethylene-1,5-diisocyanate	822-06-0	1.27		2.36E+00	e					2.36E+00
Mirex	2385-85-5	6.89		8.09E+03	e					8.09E+03
Nitrofen	1836-75-5	5.53		1.17E+04	e					1.17E+04
Pentachlorobenzene	608-93-5	5.09	1.27E+04							1.27E+04
Pentachloronitrobenzene	82-68-8	4.64	2.14E+02							2.14E+02
Pentachlorophenol	87-86-5	5.08	1.09E+02							1.09E+02
Picric acid	88-89-1	2.03		1.15E+01	e					1.15E+01
Pronamide	23950-58-5	3.51		2.55E+02	e					2.55E+02
Strychnine	57-24-9	1.93		9.34E+00	e					9.34E+00
Terphenyls	26140-60-3	No data		No data			1.00E+05		Default	1.00E+05
Tributyl phosphate	126-73-8	4.00		7.05E+02	e					7.05E+02
Trifluralin	1582-09-8	5.34		9.00E+03	e					9.00E+03
Triphenylamine	603-34-9	No data		No data			1.00E+05		Default	1.00E+05
Herbicides and Organochlorinated Pesticides										
2,4,5-T	93-76-5	3.36		1.86E+02	e					1.86E+02
2,4-D and esters	94-75-7	2.81		5.90E+01	e					5.90E+01
4,4-DDD	72-54-8	6.20		1.69E+04	e					1.69E+04
4,4-DDE	72-55-9	6.26	2.55E+04							2.55E+04
4,4-DDT	50-29-3	6.00		1.69E+04	e					1.69E+04
Aldrin	309-00-2	6.18		1.70E+04	e		1.10E+04	f		1.70E+04
alpha-BHC	319-84-6	3.80		4.65E+02	e					4.65E+02
beta-BHC	319-85-7	3.83		4.99E+02	e					4.99E+02
Chlordane	57-74-9	5.94		1.66E+04	e		1.40E+06	f		1.66E+04
Delta-BHC	319-86-8	4.14		9.41E+02	e					9.41E+02
Dieldrin	60-57-1	5.27		8.07E+03	e		1.40E+04	f		8.07E+03
Endothall	145-73-3	-0.87		2.64E-02	e					2.64E-02

Table C2-9. Aquatic Water-to-Fish Transfer Factors (BCF_{fish}) for Ecological Receptors (mg/kg tissue wet weight/mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{fish} [b]	BCF_{fish} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{fish}	SAIC compiled BCF_{fish}	Notes	Comments	Recommended BCF_{fish} [d]
Endrin	72-20-8	4.89		4.18E+03	e		2.60E+03	f		4.18E+03
gamma-BHC (Lindane)	58-89-9	3.72		3.95E+02	e		1.00E+03	f		3.95E+02
Heptachlor	76-44-8	5.02	5.52E+03				1.40E+04	f		5.52E+03
Isodrin	465-73-6	3.55		2.77E+02	e					2.77E+02
Methoxychlor	72-43-5	4.53		2.06E+03	e					2.06E+03
Silvex (2,4,5-TP)	93-72-1	4.07		8.19E+02	e					8.19E+02
Toxaphene	8001-35-2	5.50		1.13E+04	e					1.13E+04
Inorganic Chemicals and Compounds										
Metals										
Aluminum	7429-90-5	NA	2.70E+00				1.00E+01	f		2.70E+00
Antimony	7440-36-0	NA	4.00E+01				1.00E+00	f		4.00E+01
Arsenic	7440-38-2	NA	1.14E+02				2.80E+02	f		1.14E+02
Barium	7440-39-3	NA	6.33E+02		l		4.00E+00	f		6.33E+02
Beryllium	7440-41-7	NA	6.20E+01				2.00E+00	f		6.20E+01
Bismuth	7440-69-9	NA		6.33E+02	l					6.33E+02
Boron	7440-42-8	NA		6.33E+02	l					6.33E+02
Cadmium	7440-43-9	NA	9.07E+02				5.00E+01	f		9.07E+02
Calcium	7440-70-2	NA		6.33E+02	l		4.00E+01	k		6.33E+02
Chromium (and VI)	18540-29-9	NA	1.90E+01				2.00E+02	f		1.90E+01
Cobalt	7440-48-4	NA		6.33E+02	l		3.00E+02	f		6.33E+02
Copper	7440-50-8	NA	7.10E+02				2.10E+02	f		7.10E+02
Iron	7439-89-6	NA		6.33E+02	l		2.00E+03	k		6.33E+02
Lead	7439-92-1	NA	9.00E-02				3.00E+02	f		9.00E-02
Lithium	7439-93-2	NA		6.33E+02	l					6.33E+02
Magnesium	7439-95-4	NA		6.33E+02	l					6.33E+02
Manganese	7439-96-5	NA		6.33E+02	l		4.00E+02	f		6.33E+02
Mercury	7439-97-6	NA	3.53E+03				6.30E+04	f		3.53E+03
Molybdenum	7439-98-7	NA		6.33E+02	l					6.33E+02
Nickel	7440-02-0	NA	7.80E+01				1.00E+02	f		7.80E+01

Table C2-9. Aquatic Water-to-Fish Transfer Factors (BCF_{fish}) for Ecological Receptors (mg/kg tissue wet weight/mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{fish} [b]	BCF_{fish} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{fish}	SAIC compiled BCF_{fish}	Notes	Comments	Recommended BCF_{fish} [d]
Potassium	7440-09-7	NA		6.33E+02	l		1.00E+03	k		6.33E+02
Rhodium	7440-16-6	NA		6.33E+02	l					6.33E+02
Selenium	7782-49-2	NA	1.29E+02				8.00E+00	f		1.29E+02
Silicon	7440-21-3	NA		6.33E+02	l					6.33E+02
Silver	7440-22-4	NA	8.77E+01				2.00E+00	f		8.77E+01
Sodium	7440-23-5	NA		6.33E+02	l		1.00E+02	k		6.33E+02
Strontium	7440-24-6	NA		6.33E+02	l					6.33E+02
Tantalum	7440-25-7	NA		6.33E+02	l					6.33E+02
Thallium	7440-28-0	NA	1.00E+04							1.00E+04
Tin	7440-31-5	NA		6.33E+02	l					6.33E+02
Tungsten	7440-33-7	NA		6.33E+02	l					6.33E+02
Uranium	7440-61-1	NA		6.33E+02	l					6.33E+02
Vanadium	7440-62-2	NA		6.33E+02	l		1.00E-02	f		6.33E+02
Yttrium	7440-65-5	NA		6.33E+02	l					6.33E+02
Zinc	7440-66-6	NA	2.06E+03				1.00E+03	f		2.06E+03
Zirconium	7440-67-7	NA		6.33E+02	l					6.33E+02
Non-metals and Anions										
Ammonia/Ammonium	7664-41-7	NA					5.00E+02		Default	5.00E+02
Bromide	24959-67-9	NA		6.33E+02	l					6.33E+02
Chloride	16887-00-6	NA		6.33E+02	l					6.33E+02
Cyanide	57-12-5	NA	6.33E+02		l		0.00E+00	f		0.00E+00
Fluoride	16984-48-8	NA		6.33E+02	l					6.33E+02
Hydroxide	14280-30-9	NA	NA						Depends on pH	NA
Iodine	7553-56-2	NA		6.33E+02	l					6.33E+02
Nitrate	14797-55-8	NA		6.33E+02	l					6.33E+02
Nitrite	14797-65-0	NA		6.33E+02	l					6.33E+02
Phosphate	14265-44-2	NA		6.33E+02	l					6.33E+02
Phosphorus	7723-14-0	NA		6.33E+02	l					6.33E+02
Sulfate	14808-79-8	NA		6.33E+02	l					6.33E+02

Table C2-9. Aquatic Water-to-Fish Transfer Factors (BCF_{fish}) for Ecological Receptors (mg/kg tissue wet weight/mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{fish} [b]	BCF_{fish} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{fish}	SAIC compiled BCF_{fish}	Notes	Comments	Recommended BCF_{fish} [d]
Total Sulfur	63705-05-5	NA		6.33E+02	l					6.33E+02
Priority Pollutants										
Carbon Dioxide	124-38-9	NA					5.00E+02		Default	5.00E+02
Nitrogen Dioxide	10102-44-0	NA					5.00E+02		Default	5.00E+02
Ozone	10028-15-6	NA					5.00E+02		Default	5.00E+02
Particulate Matter	No CAS #	NA	NA							NA
Sulfur Dioxode	7446-09-5	NA					5.00E+02		Default	5.00E+02
Radionuclides										
Americium-241	1596-10-2	NA		6.33E+02	l		1.00E+02	k		6.33E+02
Antimony-125	14234-35-6	NA	4.00E+01				2.00E+02	k		4.00E+01
Barium-137	13981-97-0	NA		6.33E+02	l		2.00E+02	k		6.33E+02
Cadmium-113	None	NA	9.07E+02				2.00E+02	k		9.07E+02
Cesium-134	13967-70-9	NA		6.33E+02	l		2.00E+03	k		6.33E+02
Cesium-137	10045-97-3	NA		6.33E+02	l		2.00E+03	k		6.33E+02
Europium-154	15585-10-1	NA		6.33E+02	l		3.00E+02	k		6.33E+02
Europium-155	14391-16-3	NA		6.33E+02	l		3.00E+02	k		6.33E+02
Nickel-63	13981-37-8	NA	7.80E+01				1.00E+02	k		7.80E+01
Plutonium-239	15117-48-3	NA		6.33E+02	l		2.50E+02	k		6.33E+02
Plutonium-241	14119-32-5	NA		6.33E+02	l		2.50E+02	k		6.33E+02
Samarium-151	15715-94-3	NA		6.33E+02	l		3.00E+02	k		6.33E+02
Strontium-90	10098-97-2	NA		6.33E+02	l		5.00E+01	k		6.33E+02
Technetium-99	14133-79-7	NA		6.33E+02	l		1.50E+01	k		6.33E+02
Tritium	10028-17-8	NA		6.33E+02	l		1.00E+00	k		6.33E+02
Yttrium-90	10098-91-6	NA		6.33E+02	l		2.50E+01	k		6.33E+02
Uranium-232	14158-29-3	NA		6.33E+02	l		5.00E+01	k		6.33E+02
Uranium-233	13968-55-3	NA		6.33E+02	l		5.00E+01	k		6.33E+02
Uranium-234	13966-29-5	NA		6.33E+02	l		5.00E+01	k		6.33E+02
Uranium-235	15117-96-1	NA		6.33E+02	l		5.00E+01	k		6.33E+02
Uranium-236	13982-70-2	NA		6.33E+02	l		5.00E+01	k		6.33E+02

Table C2-9. Aquatic Water-to-Fish Transfer Factors (BCF_{fish}) for Ecological Receptors (mg/kg tissue wet weight/mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{fish} [b]	BCF_{fish} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{fish}	SAIC compiled BCF_{fish}	Notes	Comments	Recommended BCF_{fish} [d]
Uranium-238	7440-61-1	NA		6.33E+02	l		5.00E+01	k		6.33E+02
Actinium-227	14952-40-0	NA		6.33E+02	l		3.30E+02	k		6.33E+02
Americium-243	14993-75-0	NA		6.33E+02	l		1.00E+02	k		6.33E+02
Carbon-14	14762-75-5	NA		6.33E+02	l		9.00E+03	k		6.33E+02
Cobalt-60	10198-40-0	NA		6.33E+02	l		3.30E+02	k		6.33E+02
Curium-242	15510-73-3	NA		6.33E+02	l		3.00E+01	k		6.33E+02
Curium-243	15757-87-6	NA		6.33E+02	l		3.00E+01	k		6.33E+02
Curium-244	13981-15-2	NA		6.33E+02	l		3.00E+01	k		6.33E+02
Europium-152	14683-23-9	NA		6.33E+02	l		3.00E+02	k		6.33E+02
Iodine-129	15046-84-1	NA		6.33E+02	l		5.00E+01	k		6.33E+02
Neptunium-237	13994-20-2	NA		6.33E+02	l		2.50E+03	k		6.33E+02
Nickel-59	14336-70-0	NA	7.80E+01				1.00E+02	k		7.80E+01
Niobium-93		NA		6.33E+02	l		1.00E+02	k		1.00E+02
Plutonium-238	13981-16-3	NA		6.33E+02	l		2.50E+02	k		2.50E+02
Plutonium-240	14119-33-6	NA		6.33E+02	l		2.50E+02	k		2.50E+02
Plutonium-242	13982-10-0	NA		6.33E+02	l		2.50E+02	k		2.50E+02
Protactinium-231	14331-85-2	NA		6.33E+02	l		3.00E+01	k		3.00E+01
Radium-226	13982-63-3	NA		6.33E+02	l		5.00E+01	k		5.00E+01
Radium-228	15262-20-1	NA		6.33E+02	l		5.00E+01	k		5.00E+01
Ruthenium-106	13967-48-1	NA		6.33E+02	l		1.00E+02	k		1.00E+02
Selenium-79	None	NA	1.29E+02				1.00E+03	k		1.29E+02
Thorium-229	15594-54-4	NA		6.33E+02	l		1.00E+02	k		1.00E+02
Thorium-232	7440-29-1	NA		6.33E+02	l		1.00E+02	k		1.00E+02
Tin-126	15832-50-5	NA		6.33E+02	l		1.00E+03	k		1.00E+03
Zirconium-93	15751-77-6	NA		6.33E+02	l		2.00E+02	k		2.00E+02

NA = Not applicable

[a] $\log_{10}$ of K_{ow} values in Table 4.1

[b] Published in Appendix C of EPA (1999), Table C-5

[c] Calculated or chosen as described in Appendix C, Sect. C-1.5 of EPA (1999)

Table C2-9. Aquatic Water-to-Fish Transfer Factors (BCF_{fish}) for Ecological Receptors (mg/kg tissue wet weight/mg/L)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BCF_{fish} [b]	BCF_{fish} Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BCF_{fish}	SAIC compiled BCF_{fish}	Notes	Comments	Recommended BCF_{fish} [d]
[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]

[d] Selection criteria described in Sect. 8.2.4.4

[c] Calculated by using log K_{ow}: log BCF = 0.91 x log K_{ow} - 1.975 x log(0.00000068 x K_{ow}+1)-0.786 (Bintein, Devillers, and Karcher 1993)

[f] HAZWRAP (1994), Table 4.4

[g] Calculated by using the BCF_{fish} for TCDD from Appendix C, Table C-5 of EPA (1999) and BEFs for other congeners

[h] Value for Aroclor 1254 in Appendix C of EPA (1999), Table C-5, was used for PCB mixtures

[i] Benzo(a)pyrene value from Appendix C of EPA (1999), Table C-5, was used as a surrogate value

[j] Nitrobenzene value from Appendix C of EPA (1999), Table C-5, was used as a surrogate value

[k] NRC (1992), Table 6.19

[l] Average of values for other inorganics as published in Appendix C of EPA (1999), Table C-5

Table C2-10. Aquatic Sediment-to-Animal Transfer Factors (BASF) for Ecological Receptors (mg/kg tissue wet weight/mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BASF [b]	BASF Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BASF	Kd [d]	BCF_{inv} [e]	SAIC computed BASF [f]	Comments	Preferred BASF [g]
Organic Compounds											
Aromatic Halogenated Hydrocarbons											
4-Chloro-3-methylphenol	59-50-7	3.10		2.47E+01	h		2.41E+01	2.47E+01	1.03E+00		1.03E+00
2,3,4,6-Tetrachlorophenol	58-90-2	4.42		2.98E+02	h		2.64E+02	2.98E+02	1.13E+00		1.13E+00
Aromatic Nonhalogenated Hydrocarbons											
2-Nitrotoluene	88-72-2	2.30		5.47E+00	h		1.71E+01	5.47E+00	3.20E-01		3.20E-01
4-Nitrobiphenyl	92-93-3	3.77		8.74E+01	h		No Kd	8.74E+01	1.00E+01	Default	1.00E+01
Benzaldehyde	100-52-7	1.48		1.16E+00	h		8.04E-01	1.16E+00	1.44E+00		1.44E+00
Benzene	71-43-2	-4.99		5.84E-06	h		4.65E+00	5.84E-06	1.25E-06		1.25E-06
Benzyl alcohol	100-51-6	1.10		5.69E-01	h		4.09E-01	5.69E-01	1.39E+00		1.39E+00
Ethyl benzene	100-41-4	3.12		2.58E+01	h		8.16E+00	2.58E+01	3.17E+00		3.17E+00
m-Xylene	108-38-3	3.20		2.99E+01	h		7.84E+00	2.99E+01	3.82E+00		3.82E+00
o-Xylene	95-47-6	3.13		2.62E+01	h		9.64E+00	2.62E+01	2.71E+00		2.71E+00
p-Xylene	106-42-3	3.17		2.82E+01	h		1.24E+01	2.82E+01	2.28E+00		2.28E+00
Styrene	100-42-5	2.93		1.79E+01	h		3.65E+01	1.79E+01	4.90E-01		4.90E-01
Toluene	108-88-3	2.67		1.09E+01	h		5.60E+00	1.09E+01	1.95E+00		1.95E+00
Non-aromatic Nonhalogenated Hydrocarbons											
1,2-Epoxybutane	106-88-7	1.44		1.08E+00	h		No Kd	1.08E+00	1.00E+01	Default	1.00E+01
1,3-Butadiene	106-99-0	1.90		2.57E+00	h		2.96E+00	2.57E+00	8.68E-01		8.68E-01
1,4-Dioxane	123-91-1	-0.27	4.31E-02		h		3.50E-02	4.31E-02	1.23E+00		1.23E+00
1-Methylpropyl alcohol	78-92-2	0.61		2.26E-01	h		1.60E-01	2.26E-01	1.41E+00		1.41E+00
1-Nitropropane	108-03-2	0.87		3.69E-01	h		No Kd	3.69E-01	1.00E+01	Default	1.00E+01
2,2,4-Trimethylpentane	540-84-1	5.02		9.23E+02	h		3.46E+03	9.23E+02	2.67E-01		2.67E-01
2-Butanone	78-93-3	0.28		1.21E-01	h		9.36E-02	1.21E-01	1.30E+00		1.30E+00
2-Butenaldehyde (2-Butenal)	4170-30-3	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
2-Ethoxyethanol	110-80-5	-0.10		5.92E-02	h		8.36E-01	5.92E-02	7.08E-02		7.08E-02
2-Heptanone	110-43-0	1.98		2.99E+00	h		3.55E+00	2.99E+00	8.42E-01		8.42E-01
2-Hexanone	591-78-6	1.38		9.64E-01	h		5.36E+00	9.64E-01	1.80E-01		1.80E-01
2-Methoxyethanol	109-86-4	0.25		1.15E-01	h		No Kd	1.15E-01	1.00E+01	Default	1.00E+01

Table C2-10. Aquatic Sediment-to-Animal Transfer Factors (BASF) for Ecological Receptors (mg/kg tissue wet weight/mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BASF [b]	BASF Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BASF	Kd [d]	BCF_{fish} [e]	SAIC computed BASF [f]	Comments	Preferred BASF [g]
2-Methyl-2-propanol	75-65-0	0.35		1.38E-01	h		1.49E+00	1.38E-01	9.30E-02		9.30E-02
2-Methyl-2-propenenitrile	126-98-7	0.54		1.98E-01	h		1.49E-01	1.98E-01	1.33E+00		1.33E+00
2-Methylaziridine	75-55-8	-0.60		2.29E-02	h		No Kd	2.29E-02	1.00E+01	Default	1.00E+01
2-Methylpropyl alcohol	78-83-1	0.76		3.00E-01	h		2.24E-01	3.00E-01	1.33E+00		1.33E+00
2-Pentanone	107-87-9	0.91		3.97E-01	h		2.96E+00	3.97E-01	1.34E-01		1.34E-01
2-Propanone (Acetone)	67-64-1	-0.22	5.00E-02		h		3.80E-02	5.00E-02	1.32E+00		1.32E+00
2-Propene-1-ol	107-18-6	0.17		9.85E-02	h		5.90E-02	9.85E-02	1.67E+00		1.67E+00
2-Propyl alcohol	67-63-0	0.05		7.85E-02	h		4.50E-02	7.85E-02	1.74E+00		1.74E+00
3-Heptanone	106-35-4	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
3-Methyl-1-butanol	123-51-3	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
3-Methyl-2-butanone	563-80-4	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
3-Pentanone	96-22-0	0.99		4.62E-01	h		4.81E-01	4.62E-01	9.61E-01		9.61E-01
4-Heptanone	123-19-3	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
4-Methyl-2-pentanone	108-10-1	1.19		6.74E-01	h		4.80E-01	6.74E-01	1.40E+00		1.40E+00
4-Methyl-3-penten-2-one	141-79-7	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
5-Methyl-2-hexanone	110-12-3	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Acetaldehyde	75-07-0	-0.22		4.72E-02	h		3.81E-02	4.72E-02	1.24E+00		1.24E+00
Acetamide	60-35-5	-1.26		6.64E-03	h		1.13E-03	6.64E-03	5.89E+00		5.89E+00
Acetic acid	64-19-7	-0.17		5.19E-02	h		4.00E-02	5.19E-02	1.30E+00		1.30E+00
Acetic acid ethyl ester	141-78-6	0.73		2.83E-01	h		9.18E-02	2.83E-01	3.08E+00		3.08E+00
Acetic acid n-butyl ester	123-86-4	1.73		1.87E+00	h		2.02E+00	1.87E+00	9.25E-01		9.25E-01
Acetonitrile	75-05-8	-0.34		3.76E-02	h		3.07E-02	3.76E-02	1.23E+00		1.23E+00
Acrolein	107-02-8	-0.01		7.03E-02	h		5.57E-02	7.03E-02	1.26E+00		1.26E+00
Acrylonitrile	107-13-1	0.25	1.15E-01		h		8.88E-02	1.15E-01	1.29E+00		1.29E+00
Bis(isopropyl)ether	108-20-3	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Butane	106-97-8	2.89		1.66E+01	h		No Kd	1.66E+01	1.00E+01	Default	1.00E+01
Carbon disulfide	75-15-0	2.00		3.10E+00	h		2.06E+00	3.10E+00	1.51E+00		1.51E+00
Cyanogen	460-19-5	0.81		3.28E-01	h		No Kd	3.28E-01	1.00E+01	Default	1.00E+01
Cyclohexane	110-82-7	3.44		4.69E+01	h		1.91E+01	4.69E+01	2.45E+00		2.45E+00

Table C2-10. Aquatic Sediment-to-Animal Transfer Factors (BASF) for Ecological Receptors (mg/kg tissue wet weight/mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BASF [b]	BASF Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BASF	Kd [d]	BCF_{inv} [e]	SAIC computed BASF [f]	Comments	Preferred BASF [g]
Cyclohexanone	108-94-1	0.81		3.29E-01	h		2.51E-01	3.29E-01	1.31E+00		1.31E+00
Cyclohexene	110-83-8	2.86		1.57E+01	h		2.60E+01	1.57E+01	6.04E-01		6.04E-01
Cyclopentane	287-92-3	3.00		2.05E+01	h		3.57E+01	2.05E+01	5.73E-01		5.73E-01
Ethyl alcohol	64-17-5	0.31		1.28E-01	h		8.11E-02	1.28E-01	1.58E+00		1.58E+00
Ethyl ether	60-29-7	0.89		3.83E-01	h		3.01E-01	3.83E-01	1.27E+00		1.27E+00
Ethyl methacrylate	97-63-2	1.59		1.43E+00	h		9.80E-01	1.43E+00	1.46E+00		1.46E+00
Formaldehyde	50-00-0	0.34	1.36E-01		h		1.05E-01	1.36E-01	1.30E+00		1.30E+00
Formamide	75-12-7	-1.51		4.14E-03	h		No Kd	4.14E-03	1.00E+01	Default	1.00E+01
Formic acid	64-18-6	-0.54		2.59E-02	h		2.16E-01	2.59E-02	1.20E-01		1.20E-01
Formic acid, methyl ester	107-31-3	-0.26		4.34E-02	h		No Kd	4.34E-02	1.00E+01	Default	1.00E+01
Glycidylaldehyde	765-34-4	-0.73		1.80E-02	h		No Kd	1.80E-02	1.00E+01	Default	1.00E+01
Methyl acetate	79-20-9	0.18		1.00E-01	h		1.92E-01	1.00E-01	5.23E-01		5.23E-01
Methyl alcohol	67-56-1	-0.71		1.87E-02	h		1.58E-02	1.87E-02	1.19E+00		1.19E+00
Methyl isocyanate	624-83-9	No data		No data			No Kd	NA	1.00E+01	Default	1.00E+01
Methyl methacrylate	80-62-6	0.79		3.17E-01	h		2.52E+00	3.17E-01	1.26E-01		1.26E-01
Methyl tert-butyl ether	1634-04-4	0.94		4.21E-01	h		3.37E-01	4.21E-01	1.25E+00		1.25E+00
Methylacetylene	74-99-7	0.94		4.21E-01	h		No Kd	4.21E-01	1.00E+01	Default	1.00E+01
Methylcyclohexane	108-87-2	4.10		1.63E+02	h		4.31E+02	1.63E+02	3.78E-01		3.78E-01
N,N-Dimethylacetamide	127-19-5	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
n-Butyl alcohol	71-36-3	0.88		3.76E-01	h		2.95E-01	3.76E-01	1.28E+00		1.28E+00
n-Heptane	142-82-5	4.66		4.68E+02	h		1.53E+03	4.68E+02	3.06E-01		3.06E-01
n-Hexane	110-54-3	4.11		1.66E+02	h		4.41E+02	1.66E+02	3.77E-01		3.77E-01
Nitromethane	75-52-5	-0.35		3.69E-02	h		No Kd	3.69E-02	1.00E+01	Default	1.00E+01
n-Nonane	111-84-2	5.65		3.03E+03	h		1.44E+04	3.03E+03	2.11E-01		2.11E-01
n-Octane	111-65-9	4.00		1.35E+02	h		3.44E+02	1.35E+02	3.93E-01		3.93E-01
n-Pentane	109-66-0	3.21		3.04E+01	h		5.75E+01	3.04E+01	5.29E-01		5.29E-01
n-Propionaldehyde	123-38-6	0.59		2.17E-01	h		1.53E-01	2.17E-01	1.42E+00		1.42E+00
n-Propyl alcohol	71-23-8	0.25		1.14E-01	h		No Kd	1.14E-01	1.00E+01	Default	1.00E+01
n-Valeraldehyde	110-62-3	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01

Table C2-10. Aquatic Sediment-to-Animal Transfer Factors (BASF) for Ecological Receptors (mg/kg tissue wet weight/mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) BASF[b]	BASF Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance BASF	Kd[d]	BCF_{inv}[e]	SAIC computed BASF[f]	Comments	Preferred BASF[g]
Oxirane	75-21-8	-0.30		4.06E-02	h		3.30E-02	4.06E-02	1.23E+00		1.23E+00
p-Cymene	99-87-6	4.10		1.63E+02	h		No Kd	1.63E+02	1.00E+01	Default	1.00E+01
Phosgene	75-44-5	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Propargyl alcohol	107-19-7	0.26		1.17E-01	h		No Kd	1.17E-01	1.00E+01	Default	1.00E+01
Propionic acid	79-09-4	0.33		1.33E-01	h		8.48E-02	1.33E-01	1.57E+00		1.57E+00
Propionitrile	107-12-0	0.16		9.66E-02	h		5.77E-02	9.66E-02	1.67E+00		1.67E+00
Propylene gylcol monomethyl ether	107-98-2	-0.18		5.09E-02	h		No Kd	5.09E-02	1.00E+01	Default	1.00E+01
p-tert-Butyltoluene	98-51-1	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Triethylamine	121-44-8	0.16		9.66E-02	h		5.77E-02	9.66E-02	1.67E+00		1.67E+00
Trimethylamine	75-50-3	0.16		9.66E-02	h		1.60E-01	9.66E-02	6.04E-01		6.04E-01
Vinyl acetate	108-05-4	0.70		2.67E-01	h		1.99E-01	2.67E-01	1.34E+00		1.34E+00
Non-aromatic Halogenated Hydrocarbons											
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
1,1,1,2-Tetrachloroethane	630-20-6	2.63		1.02E+01	h		6.37E+00	1.02E+01	1.60E+00		1.60E+00
1,1,1-Trichloroethane	71-55-6	2.42		6.88E+00	h		5.40E+03	6.88E+00	1.27E-03		1.27E-03
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	3.73		8.11E+01	h		1.26E+01	8.11E+01	6.41E+00		6.41E+00
1,1,2,2-Tetrachloroethane	79-34-5	4.64		4.54E+02	h		3.16E+00	4.54E+02	1.44E+02		1.44E+02
1,1,2,2-Tetrachloroethene	127-18-4	2.55		8.68E+00	h		1.06E+01	8.68E+00	8.19E-01		8.19E-01
1,1,2-Trichloroethane	79-00-5	2.10		3.73E+00	h		3.00E+00	3.73E+00	1.24E+00		1.24E+00
1,1,2-Trichloroethylene	79-01-6	2.43		7.02E+00	h		3.76E+00	7.02E+00	1.87E+00		1.87E+00
1,1-Dichloroethane	75-34-3	1.79		2.10E+00	h		2.12E+00	2.10E+00	9.90E-01		9.90E-01
1,1-Dichloroethene	75-35-4	2.12		3.90E+00	h		2.60E+00	3.90E+00	1.50E+00		1.50E+00
1,2,2-Trichloro-1,1,2-trifluoroethane	76-13-1	3.16		2.77E+01	h		1.03E+01	2.77E+01	2.69E+00		2.69E+00
1,2,3-Trichloropropane	96-18-4	2.25		4.98E+00	h		3.22E+00	4.98E+00	1.55E+00		1.55E+00
1,2-Dibromo-3-chloropropane	96-12-8	2.34		5.90E+00	h		3.79E+00	5.90E+00	1.56E+00		1.56E+00
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	2.82		1.46E+01	h		No Kd	1.46E+01	1.00E+01	Default	1.00E+01
1,2-Dichloroethane	107-06-2	1.46		1.13E+00	h		7.83E-01	1.13E+00	1.44E+00		1.44E+00
1,2-Dichloroethylene	540-59-0	0.48		1.77E-01	h		No Kd	1.77E-01	1.00E+01	Default	1.00E+01
1,2-Dichloropropane	78-87-5	2.25		4.98E+00	h		1.88E+00	4.98E+00	2.65E+00		2.65E+00

Table C2-10. Aquatic Sediment-to-Animal Transfer Factors (BASF) for Ecological Receptors (mg/kg tissue wet weight/mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) BASF[b]	BASF Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance BASF	Kd[d]	BCF_{inv}[e]	SAIC computed BASF[f]	Comments	Preferred BASF[g]
1,3-Dichloropropene	542-75-6	1.75		1.93E+00	h		1.08E+00	1.93E+00	1.79E+00		1.79E+00
1,4-Dichloro-2-butene	764-41-0	0.87		3.70E-01	h		No Kd	3.70E-01	1.00E+01	Default	1.00E+01
1-Chloroethene	75-01-4	1.15		6.20E-01	h		4.44E-01	6.20E-01	1.40E+00		1.40E+00
2,2-Dichloropropionic acid	75-99-0	0.78		3.10E-01	h		No Kd	3.10E-01	1.00E+01	Default	1.00E+01
2-Chloropropane	75-29-6	1.90		2.57E+00	h		2.96E+00	2.57E+00	8.68E-01		8.68E-01
3-Chloropropene (allyl chloride)	107-05-1	0.95		4.32E-01	h		No Kd	4.32E-01	1.00E+01	Default	1.00E+01
Bromochloromethane	74-97-5	1.41		1.02E+00	h		9.77E-01	1.02E+00	1.04E+00		1.04E+00
Bromodichloromethane	75-27-4	2.03		3.26E+00	h		2.15E+00	3.26E+00	1.51E+00		1.51E+00
Bromoethene	593-60-2	1.07		5.34E-01	h		No Kd	5.34E-01	1.00E+01	Default	1.00E+01
Bromoform	75-25-2	2.35		6.01E+00	h		5.04E+00	6.01E+00	1.19E+00		1.19E+00
Bromomethane	74-83-9	1.11		5.84E-01	h		3.60E-01	5.84E-01	1.62E+00		1.62E+00
Carbon tetrachloride	56-23-5	2.72	1.20E+01		h		6.08E+00	1.20E+01	1.97E+00		1.97E+00
Chlorodibromomethane	124-48-1	2.18		4.33E+00	h		2.82E+00	4.33E+00	1.53E+00		1.53E+00
Chlorodifluoromethane	75-45-6	1.08		5.47E-01	h		3.93E-01	5.47E-01	1.39E+00		1.39E+00
Chloroethane	75-00-3	3.10		2.47E+01	h		1.48E+01	2.47E+01	1.67E+00		1.67E+00
Chloroform	67-66-3	1.95	2.82E+00		h		2.12E+00	2.82E+00	1.33E+00		1.33E+00
Chloromethane	74-87-3	0.90		3.92E-01	h		2.40E-01	3.92E-01	1.63E+00		1.63E+00
Chloropentafluoroethane	76-15-3	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
cis-1,2-Dichloroethene	156-59-2	1.98		3.00E+00	h		1.99E+01	3.00E+00	1.51E-01		1.51E-01
cis-1,3-Dichloropropene	10061-01-5	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Cyanogen bromide	506-68-3	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Cyanogen chloride	506-77-4	0.20		1.04E-01	h		No Kd	1.04E-01	1.00E+01	Default	1.00E+01
Dichlorodifluoromethane	75-71-8	2.16		4.19E+00	h		2.74E+00	4.19E+00	1.53E+00		1.53E+00
Dichlorofluoromethane	75-43-4	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Dichloromethane	75-09-2	1.26		7.62E-01	h		4.00E-01	7.62E-01	1.91E+00		1.91E+00
Difluorodibromomethane	75-61-6	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Hexafluoroacetone	684-16-2	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Iodomethane	74-88-4	1.69		1.73E+00	h		1.84E+00	1.73E+00	9.39E-01		9.39E-01
Methylene bromide	74-95-3	1.62		1.52E+00	h		1.04E+00	1.52E+00	1.46E+00		1.46E+00

Table C2-10. Aquatic Sediment-to-Animal Transfer Factors (BASF) for Ecological Receptors (mg/kg tissue wet weight/mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) BASF[b]	BASF Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance BASF	Kd[d]	BCF_{inv}[e]	SAIC computed BASF[f]	Comments	Preferred BASF[g]
Pentachloroethane	76-01-7	3.05		2.25E+01	h		4.00E+01	2.25E+01	5.62E-01		5.62E-01
trans-1,2-Dichloroethene	156-60-5	1.98		3.00E+00	h		1.52E+00	3.00E+00	1.98E+00		1.98E+00
trans-1,3-Dichloropropene	10061-02-6	2.06		3.48E+00	h		No Kd	3.48E+00	1.00E+01	Default	1.00E+01
Trichloroacetic acid	76-03-9	1.33		8.78E-01	h		8.16E-01	8.78E-01	1.08E+00		1.08E+00
Trichlorofluoroethane	27154-33-2	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Trichlorofluoromethane	75-69-4	2.53		8.46E+00	h		5.34E+00	8.46E+00	1.58E+00		1.58E+00
Trifluorobromomethane	75-63-8	1.86		2.38E+00	h		No Kd	2.38E+00	1.00E+01	Default	1.00E+01
Dioxin and Furan Compounds											
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	35822-46-9	8.20	9.94E+01		i		3.91E+06	7.96E+01	2.04E-05		2.04E-05
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	7.92	2.16E+02		i		2.05E+06	1.72E+01	8.39E-06		8.39E-06
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	7.92	7.64E+03		i		2.05E+06	6.08E+02	2.97E-04		2.97E-04
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	39227-28-6	7.79	6.08E+03		i		1.52E+06	4.84E+02	3.18E-04		3.18E-04
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	7.25	1.49E+03		i		4.39E+05	1.19E+02	2.70E-04		2.70E-04
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	57653-85-7	7.25	2.35E+03		i		4.39E+05	1.87E+02	4.26E-04		4.26E-04
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	7.25	3.72E+03		i		4.39E+05	2.96E+02	6.75E-04		6.75E-04
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	19408-74-3	7.25	2.74E+03		i		4.39E+05	2.18E+02	4.97E-04		4.97E-04
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	7.25	1.23E+04		i		4.39E+05	9.83E+02	2.24E-03		2.24E-03
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	40321-76-4	6.64	1.80E+04		i		1.08E+05	1.44E+03	1.33E-02		1.33E-02
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	6.79	4.31E+03		i		1.52E+05	3.43E+02	2.26E-03		2.26E-03
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	7.25	1.31E+04		i		4.39E+05	1.05E+03	2.38E-03		2.38E-03
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	6.92	3.14E+04		i		2.05E+05	2.50E+03	1.22E-02		1.22E-02
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1746-01-6	6.64	1.96E+04		h		1.08E+05	1.56E+03	1.44E-02		1.44E-02
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	6.53	2.64E+03		i		8.36E+04	1.25E+03	1.49E-02		1.49E-02
Dibenzofuran	132-64-9	4.33		2.51E+02	h		7.25E+02	2.51E+02	3.47E-01		3.47E-01
Octachlorodibenzo(p)dioxin	3268-87-9	8.20	2.35E+01		i		8.95E+01	1.87E+01	2.09E-01		2.09E-01
Octachlorodibenzofuran	39001-02-0	8.78	3.14E+02		i		1.49E+07	2.50E+01	1.68E-06		1.68E-06
PCBs											
Polychlorinated biphenyls (PCBs)[j]	1336-36-3	7.31	5.30E-01				1.24E+04	5.54E+03	4.48E-01		4.48E-01
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	7.08		4.49E+04	h		No Kd	4.49E+04	1.00E+01	Default	1.00E+01

Table C2-10. Aquatic Sediment-to-Animal Transfer Factors (BASF) for Ecological Receptors (mg/kg tissue wet weight/mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) BASF[b]	BASF Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance BASF	Kd[d]	BCF_{inv}[e]	SAIC computed BASF[f]	Comments	Preferred BASF[g]
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	7.12		4.84E+04	h		No Kd	4.84E+04	1.00E+01	Default	1.00E+01
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
2',3,4,4',5-Pentachlorobiphenyl	no cas #	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	7.12		4.84E+04	h		No Kd	4.84E+04	1.00E+01	Default	1.00E+01
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	7.41		8.34E+04	h		No Kd	8.34E+04	1.00E+01	Default	1.00E+01
3,3',4,4',5-Pentachlorobiphenyl	no cas #	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Phthalates											
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	5.20	1.31E+03		h		4.44E+03	3.18E+02	7.16E-02		7.16E-02
Butylbenzyl phthalate	85-68-7	4.41		2.94E+02	h		5.50E+02	2.94E+02	5.35E-01		5.35E-01
Dibutyl phthalate	84-74-2	4.72		5.25E+02	h		6.27E+01	5.25E+02	8.37E+00		8.37E+00
Diethyl phthalate	84-66-2	4.44		3.07E+02	h		3.28E+00	3.07E+02	9.36E+01		9.36E+01
Dimethylphthalate	131-11-3	1.63		1.56E+00	h		1.06E+01	1.56E+00	1.47E-01		1.47E-01
n-Dioctyl phthalate	117-84-0	9.33	3.13E+06		h		3.61E+07	5.95E+03	1.65E-04		1.65E-04
Light Polycyclic Aromatic Hydrocarbons (molecular weight <200 g/mole)											
2-Chloronapthalene	91-58-7	4.07		1.53E+02	h		2.86E+02	1.53E+02	5.36E-01		5.36E-01
2-Methyl naphthalene	91-57-6	3.86		1.04E+02	h		2.50E+02	1.04E+02	4.14E-01		4.14E-01
5-Nitroacenaphthene	602-87-9	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Acenaphthene	83-32-9	3.96		1.26E+02	h		1.96E+02	1.26E+02	6.44E-01		6.44E-01
Acenaphthylene	208-96-8	4.07		1.54E+02	h		2.70E+02	1.54E+02	5.69E-01		5.69E-01
Anthracene	120-12-7	4.47		3.27E+02	h		9.40E+02	3.27E+02	3.48E-01		3.48E-01
Fluorene	86-73-7	4.18		1.89E+02	h		5.65E+02	1.89E+02	3.35E-01		3.35E-01
Indene	95-13-6	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01

Table C2-10. Aquatic Sediment-to-Animal Transfer Factors (BASF) for Ecological Receptors (mg/kg tissue wet weight/mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BASF[b]	BASF Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance BASF	Kd[d]	BCF_{inv} [e]	SAIC computed BASF[f]	Comments	Preferred BASF[g]
Naphthalene	91-20-3	3.37		4.13E+01	h		4.76E+01	4.13E+01	8.69E-01		8.69E-01
Phenanthrene	85-01-8	4.55		3.81E+02	h		2.01E+03	3.81E+02	1.89E-01		1.89E-01
Pyrene	129-00-0	5.00		8.89E+02	h		2.72E+03	8.89E+02	3.27E-01		3.27E-01
Heavy Polycyclic Aromatic Hydrocarbons (molecular weight >200 g/mole)											
3-Methylcholanthrene	56-49-5	7.11		4.75E+04	h		6.05E+04	4.75E+04	7.85E-01		7.85E-01
5-Methylchrysene	3697-24-3	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Benzo(a)anthracene	56-55-3	5.68	1.45E+00		k		1.04E+04	1.23E+04	1.18E+00		1.18E+00
Benzo(a)pyrene	50-32-8	6.13	1.59E+00				3.87E+04	4.70E+03	1.21E-01		1.21E-01
Benzo(b)fluoranthene	205-99-2	6.20	1.61E+00		k		3.34E+04	4.70E+03	1.41E-01		1.41E-01
Benzo(e)pyrene	192-97-2	7.40		8.21E+04	h		6.34E+05	8.21E+04	1.30E-01		1.30E-01
Benzo(g,h,i)perylene	191-24-2	7.10		4.67E+04	h		7.28E+04	4.67E+04	6.41E-01		6.41E-01
Benzo(j)fluoranthene	205-82-3	6.44		1.34E+04	h		8.60E+04	1.34E+04	1.56E-01		1.56E-01
Benzo(k)fluoranthene	207-08-9	6.20	1.61E+00	1.61E+00	k		3.33E+04	1.32E+04	3.97E-01		3.97E-01
Benzo[a,i]pyrene	191-30-0	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Chrysene	218-01-9	5.74	1.38E+00				1.19E+04	9.80E+02	8.24E-02		8.24E-02
Dibenz(a,h)anthracene	53-70-3	6.55	1.61E+00		k		7.16E+04	7.10E+02	9.92E-03		9.92E-03
Dibenz[a,h]acridine	226-36-8	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Dibenz[a,j]acridine	224-42-0	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Dibenzo(a,e)fluoranthene	5385-75-1	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Dibenzo(a,h)fluoranthene	no cas #	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Dibenzo[a,e]pyrene	192-65-4	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Dibenzo[a,h]pyrene	189-64-0	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Dibenzo[a,i]pyrene	189-55-9	7.29		6.68E+04	h		No Kd	6.68E+04	1.00E+01	Default	1.00E+01
Fluoranthene	206-44-0	5.08		1.04E+03	h		1.96E+03	1.04E+03	5.30E-01		5.30E-01
Hexachloronaphthalene	1335-87-1	7.59		1.18E+05	h		No Kd	1.18E+05	1.00E+01	Default	1.00E+01
Indeno(1,2,3-cd)pyrene	193-39-5	6.91	1.61E+00		k		1.64E+05	4.70E+03	2.86E-02		1.61E+00
Octachloronaphthalene	2234-13-1	6.42		1.29E+04	h		No Kd	1.29E+04	1.00E+01	Default	1.00E+01
Pentachloronaphthalene	1321-64-8	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Tetrachloronaphthalene	1335-88-2	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01

Table C2-10. Aquatic Sediment-to-Animal Transfer Factors (BASF) for Ecological Receptors (mg/kg tissue wet weight/mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BASF [b]	BASF Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BASF	Kd [d]	BCF_{inv} [e]	SAIC computed BASF [f]	Comments	Preferred BASF [g]
Trichloronaphthalene	1321-65-9	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Light Substituted Benzene Compounds (MW <200 g/mole)											
1,2,3-Trichlorobenzene	87-61-6	4.05		1.47E+02	h		8.10E+02	1.47E+02	1.81E-01		1.81E-01
1,2,4-Trichlorobenzene	120-82-1	3.99		1.32E+02	h		6.64E+01	1.32E+02	1.99E+00		1.99E+00
1,2,4-Trimethyl benzene	95-63-6	3.65		6.97E+01	h		1.56E+02	6.97E+01	4.48E-01		4.48E-01
1,2-Dichlorobenzene	95-50-1	3.45		4.74E+01	h		1.52E+01	4.74E+01	3.12E+00		3.12E+00
1,3,5-Trimethyl benzene	108-67-8	3.42		4.52E+01	h		6.69E+01	4.52E+01	6.75E-01		6.75E-01
1,3-Dichlorobenzene	541-73-1	3.53		5.56E+01	h		3.21E+02	5.56E+01	1.73E-01		1.73E-01
1,3-Dinitrobenzene	99-65-0	1.49	1.19E+00		h		8.25E-01	1.30E+01	1.58E+01		1.58E+01
1,4-Dichlorobenzene	106-46-7	3.41		4.45E+01	h		2.46E+01	4.45E+01	1.81E+00		1.81E+00
1,4-Dinitrobenzene	100-25-4	1.50		1.21E+00	h		No Kd	1.21E+00	1.00E+01	Default	1.00E+01
2,4,5-Trichlorophenol	95-95-4	3.87		1.06E+02	h		4.51E+01	1.06E+02	2.34E+00		2.34E+00
2,4,6-Trichlorophenol	88-06-2	3.71		7.83E+01	h		9.05E+00	7.83E+01	8.66E+00		8.66E+00
2,4-Dichlorophenol	120-83-2	3.04		2.20E+01	h		5.58E+00	2.20E+01	3.94E+00		3.94E+00
2,4-Dimethylphenol	105-67-9	2.36		6.12E+00	h		5.04E+00	6.12E+00	1.21E+00		1.21E+00
2,4-Dinitrophenol	51-28-5	1.52		1.25E+00	h		4.00E-04	1.25E+00	3.13E+03		3.13E+03
2,4-Dinitrotoluene	121-14-2	2.00	5.80E+01				2.04E+00	1.30E+01	6.37E+00		5.80E+01
2,6-Dinitrotoluene	606-20-2	1.89	2.51E+00		h		1.68E+00	1.30E+01	7.74E+00		7.74E+00
2-Chlorophenol	95-57-8	2.16		4.21E+00	h		1.55E+01	4.21E+00	2.72E-01		2.72E-01
2-Chlorotoluene	95-49-8	3.54		5.63E+01	h		No Kd	5.63E+01	1.00E+01	Default	1.00E+01
2-Nitrophenol	88-75-5	1.79		2.09E+00	h		1.41E+01	2.09E+00	1.48E-01		1.48E-01
4,6-Dinitro-o-cresol	534-52-1	2.85		1.54E+01	h		No Kd	1.54E+01	1.00E+01	Default	1.00E+01
4-Chlorotoluene	106-43-4	3.33		3.81E+01	h		No Kd	3.81E+01	1.00E+01	Default	1.00E+01
4-Nitrophenol	100-02-7	1.91		2.62E+00	h		1.75E+01	2.62E+00	1.50E-01		1.50E-01
alpha-Methylstyrene	98-83-9	3.46		4.90E+01	h		No Kd	4.90E+01	1.00E+01	Default	1.00E+01
Aniline	62-53-3	0.98		4.54E-01	h		3.29E-01	4.54E-01	1.38E+00		1.38E+00
Benzotrichloride	98-07-7	2.92		1.76E+01	h		No Kd	1.76E+01	1.00E+01	Default	1.00E+01
Benzyl chloride	100-44-7	0.36		1.41E-01	h		1.08E-01	1.41E-01	1.31E+00		1.31E+00
Bromobenzene	108-86-1	2.99		2.01E+01	h		1.79E+01	2.01E+01	1.12E+00		1.12E+00

Table C2-10. Aquatic Sediment-to-Animal Transfer Factors (BASF) for Ecological Receptors (mg/kg tissue wet weight/mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) BASF[b]	BASF Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance BASF	Kd[d]	BCF_{inv}[e]	SAIC computed BASF[f]	Comments	Preferred BASF[g]
Chlorobenzene	108-90-7	2.79		1.38E+01	h		8.96E+00	1.38E+01	1.54E+00		1.54E+00
Cumene	98-82-8	3.61		6.50E+01	h		3.72E+02	6.50E+01	1.75E-01		1.75E-01
m-Cresol	108-39-4	1.96		2.87E+00	h		1.91E+00	2.87E+00	1.50E+00		1.50E+00
n-Butyl benzene	104-51-8	4.28		2.29E+02	h		1.00E+02	2.29E+02	2.28E+00		2.28E+00
Nitrobenzene	98-95-3	1.83	2.26E+00		h		4.76E+04	1.30E+01	2.73E-04		2.73E-04
n-Propyl benzene	103-65-1	3.69		7.52E+01	h		2.90E+01	7.52E+01	2.59E+00		2.59E+00
o-Cresol	95-48-7	2.02		3.23E+00	h		2.14E+00	3.23E+00	1.51E+00		1.51E+00
o-Dinitrobenzene	528-29-0	1.60		1.46E+00	h		No Kd	1.46E+00	1.00E+01	Default	1.00E+01
o-Nitroaniline	88-74-4	1.85		2.34E+00	h		1.57E+01	2.34E+00	1.49E-01		1.49E-01
o-Toluidine	95-53-4	1.34		8.95E-01	h		6.28E-01	8.95E-01	1.43E+00		1.43E+00
p-Chloroaniline	106-47-8	1.87		2.43E+00	h		1.63E+00	2.43E+00	1.49E+00		1.49E+00
p-Cresol	106-44-5	1.94		2.77E+00	h		1.84E+00	2.77E+00	1.51E+00		1.51E+00
Phenol	108-95-2	1.48		1.16E+00	h		8.79E-01	1.16E+00	1.32E+00		1.32E+00
p-Nitrochlorobenzene	100-00-5	2.39		6.48E+00	h		No Kd	6.48E+00	1.00E+01	Default	1.00E+01
p-Toluidine	106-49-0	1.40		1.00E+00	h		9.56E-01	1.00E+00	1.05E+00		1.05E+00
sec-Butyl benzene	135-98-8	4.57		3.95E+02	h		No Kd	3.95E+02	1.00E+01	Default	1.00E+01
tert-Butyl benzene	98-06-6	4.11		1.66E+02	h		4.41E+02	1.66E+02	3.77E-01		3.77E-01
Toluene-2,6-diamine	823-40-5	1.45		1.11E+00	h		No Kd	1.11E+00	1.00E+01	Default	1.00E+01
Trimethyl benzene	25551-13-7	3.40		4.35E+01	h		No Kd	4.35E+01	1.00E+01	Default	1.00E+01
Other Light Semivolatile Compounds (molecular weight <200 g/mole)											
1,1'-Biphenyl	92-52-4	3.90		1.12E+02	h		1.00E+02	1.12E+02	1.11E+00		1.11E+00
1,1-Dimethylhydrazine	57-14-7	0.60		2.20E-01	h		No Kd	2.20E-01	1.00E+01	Default	1.00E+01
1,2-Dimethylhydrazine	540-73-8	-1.37		5.42E-03	h		No Kd	5.42E-03	1.00E+01	Default	1.00E+01
1,2-Diphenylhydrazine	122-66-7	2.94		1.83E+01	h		1.11E+01	1.83E+01	1.65E+00		1.65E+00
1,3-Propane sultone	1120-71-4	-0.52		2.66E-02	h		No Kd	2.66E-02	1.00E+01	Default	1.00E+01
2,4-Toluene diisocyante	584-84-9	No data		No data			No Kd	NA	1.00E+01	Default	1.00E+01
2-Chloroacetophenone	532-27-4	2.59		9.45E+00	h		No Kd	9.53E+00	1.00E+01	Default	1.00E+01
2-Propenoic acid	79-10-7	0.33		1.33E-01	h		8.48E-02	1.33E-01	1.57E+00		1.57E+00
4,4-Methylenedianiline	101-77-9	3.38		4.19E+01	h		No Kd	4.19E+01	1.00E+01	Default	1.00E+01

Table C2-10. Aquatic Sediment-to-Animal Transfer Factors (BASF) for Ecological Receptors (mg/kg tissue wet weight/mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) BASF[b]	BASF Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance BASF	Kd[d]	BCF_{inv}[e]	SAIC computed BASF[f]	Comments	Preferred BASF[g]
Acetophenone	98-86-2	1.64		1.58E+00	h		1.08E+00	1.58E+00	1.46E+00		1.46E+00
Benzoic acid	65-85-0	1.86		2.38E+00	h		2.20E-02	2.38E+00	1.08E+02		1.08E+02
bis(2-Chloroethoxy)methane	111-91-1	7.59		1.18E+05	h		9.60E+05	1.18E+05	1.22E-01		1.22E-01
bis(2-Chloroethyl) ether	111-44-4	1.30		8.31E-01	h		3.04E+00	8.31E-01	2.73E-01		2.73E-01
Chlorocyclopentadiene	41851-50-7	2.43		6.99E+00	h		No Kd	6.99E+00	1.00E+01	Default	1.00E+01
Cyclohexanol	108-93-0	1.23		7.27E-01	h		5.20E-01	7.27E-01	1.40E+00		1.40E+00
Dichloroisopropyl ether	108-60-1	2.58		9.27E+00	h		2.44E+00	9.27E+00	3.80E+00		3.80E+00
Dichloromethyl ether	542-88-1	-0.38		3.49E-02	h		3.18E+00	3.49E-02	1.10E-02		1.10E-02
Dichloropentadiene	no cas #	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Dimethyl sulfate	77-78-1	0.32		1.30E-01	h		No Kd	1.30E-01	1.00E+01	Default	1.00E+01
Dimethylaniline	121-69-7	-0.88		1.36E-02	h		No Kd	1.36E-02	1.00E+01	Default	1.00E+01
Di-n-propylnitrosamine	621-64-7	1.38		9.65E-01	h		6.80E-01	9.65E-01	1.42E+00		1.42E+00
Diphenyl ether	101-84-8	4.21		2.00E+02	h		5.53E+02	2.00E+02	3.63E-01		3.63E-01
Epichlorohydrin	106-89-8	0.25		1.15E-01	h		8.88E-02	1.15E-01	1.29E+00		1.29E+00
Ethyl carbamate (urethane)	51-79-6	-0.15		5.38E-02	h		No Kd	5.38E-02	1.00E+01	Default	1.00E+01
Ethyl methanesulfonate	62-50-0	0.05		7.84E-02	h		6.19E-02	7.84E-02	1.27E+00		1.27E+00
Ethylene dibromide	106-93-4	1.75		1.94E+00	h		1.31E+00	1.94E+00	1.48E+00		1.48E+00
Ethylene glycol	107-21-1	-0.91		1.27E-02	h		No Kd	1.27E-02	1.00E+01	Default	1.00E+01
Ethylene glycol monobutyl ether	111-76-2	1.55		1.34E+00	h		No Kd	1.34E+00	1.00E+01	Default	1.00E+01
Ethylene glycol monoethyl ether acetate	111-15-9	0.62		2.30E-01	h		No Kd	2.30E-01	1.00E+01	Default	1.00E+01
Ethylene thiourea	96-45-7	-0.64		2.13E-02	h		No Kd	2.13E-02	1.00E+01	Default	1.00E+01
Furfural	98-01-1	0.96		4.37E-01	h		3.53E-01	4.37E-01	1.24E+00		1.24E+00
Maleic hydrazide	123-33-1	-0.74		1.77E-02	h		No Kd	1.77E-02	1.00E+01	Default	1.00E+01
Malononitrile	109-77-3	0.04		7.70E-02	h		No Kd	7.70E-02	1.00E+01	Default	1.00E+01
Methyl styrene (mixed isomers)	25013-15-4	3.35		3.96E+01	h		No Kd	3.96E+01	1.00E+01	Default	1.00E+01
Methylhydrazine	60-34-4	-1.06		9.68E-03	h		No Kd	9.68E-03	1.00E+01	Default	1.00E+01
N,N-Diphenylamine	122-39-4	3.50		5.25E+01	h		1.39E+01	5.25E+01	3.79E+00		3.79E+00
Nitric acid, propyl ester	627-13-4	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
N-Nitrosodi-n-butylamine	924-16-3	2.41		6.73E+00	h		4.29E+00	6.73E+00	1.57E+00		1.57E+00

Table C2-10. Aquatic Sediment-to-Animal Transfer Factors (BASF) for Ecological Receptors (mg/kg tissue wet weight/mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) BASF[b]	BASF Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance BASF	Kd[d]	BCF_{inv}[e]	SAIC computed BASF[f]	Comments	Preferred BASF[g]
N-Nitrosomorpholine	59-89-2	0.98		4.54E-01	h		No Kd	4.54E-01	1.00E+01	Default	1.00E+01
N-Nitroso-N,N-dimethylamine	62-75-9	-0.47		2.94E-02	h		No Kd	2.94E-02	1.00E+01	Default	1.00E+01
o-Anisidine	90-04-0	1.18		6.61E-01	h		No Kd	6.61E-01	1.00E+01	Default	1.00E+01
Oxalic acid	144-62-7	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Phthalic anhydride	85-44-9	-0.60		2.30E-02	h		1.92E-02	2.30E-02	1.20E+00		1.20E+00
p-Phthalic acid	100-21-0	0.82		3.37E-01	h		No Kd	3.37E-01	1.00E+01	Default	1.00E+01
Pyridine	110-86-1	0.67		2.53E-01	h		1.89E-01	2.53E-01	1.34E+00		1.34E+00
Quinoline	91-22-5	2.03		3.29E+00	h		No Kd	3.29E+00	1.00E+01	Default	1.00E+01
Quinone	106-51-4	0.20		1.04E-01	h		No Kd	1.04E-01	1.00E+01	Default	1.00E+01
Safrole	94-59-7	2.66		1.08E+01	h		6.73E+00	1.08E+01	1.60E+00		1.60E+00
Tetrahydrofuran	109-99-9	0.45		1.66E-01	h		1.26E-01	1.66E-01	1.32E+00		1.32E+00
Other Heavy Semivolatile Compounds (molecular weight >200 g/mole)											
1,2,4,5-Tetrachlorobenzene	95-94-3	4.64		4.51E+02	h		2.36E+02	4.51E+02	1.91E+00		1.91E+00
1,3,5-Trinitrobenzene	99-35-4	1.18		6.60E-01	h		4.72E-01	6.60E-01	1.40E+00		1.40E+00
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	4.17		1.86E+02	h		No Kd	1.86E+02	1.00E+01	Default	1.00E+01
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	-2.70		4.39E-04	h		No Kd	4.39E-04	1.00E+01	Default	1.00E+01
2-sec-Butyl-4,6-dinitrophenol	88-85-7	3.14		2.66E+01	h		No Kd	2.66E+01	1.00E+01	Default	1.00E+01
3,3'-Dimethoxybenzidine	119-90-4	1.81		2.17E+00	h		1.46E+00	2.17E+00	1.49E+00		1.49E+00
3,3-Dichlorobenzidine	91-94-1	3.58		6.05E+01	h		3.48E+01	6.05E+01	1.74E+00		1.74E+00
4-Bromophenylphenyl ether	101-55-3	5.00		8.89E+02	h		No Kd	8.89E+02	1.00E+01	Default	1.00E+01
Ammonium perfluorooctanoate	3825-26-1	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Azobenzene	103-33-3	3.82		9.61E+01	h		No Kd	9.61E+01	1.00E+01	Default	1.00E+01
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Captan	133-06-2	2.35		6.01E+00	h		7.98E+00	6.01E+00	7.53E-01		7.53E-01
Chlorobenzilate	510-15-6	4.38		2.76E+02	h		1.48E+02	2.76E+02	1.87E+00		1.87E+00
Dibutylphosphate	107-66-4	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Dimethyl aminoazobenzene	60-11-7	4.58		4.03E+02	h		No Kd	4.03E+02	1.00E+01	Default	1.00E+01
Hexachlorobenzene	118-74-1	5.50	2.29E+03		h		3.20E+03	2.60E+03	8.11E-01		8.11E-01
Hexachlorobutadiene	87-68-3	4.73	4.40E-01				2.77E+02	1.05E+01	3.79E-02		4.40E-01

Table C2-10. Aquatic Sediment-to-Animal Transfer Factors (BASF) for Ecological Receptors (mg/kg tissue wet weight/mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BASF [b]	BASF Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BASF	Kd [d]	BCF_{inv} [e]	SAIC computed BASF [f]	Comments	Preferred BASF [g]
Hexachlorocyclopentadiene	77-47-4	5.04	7.46E+02		h		3.80E+02	1.23E+03	3.24E+00		3.24E+00
Hexachloroethane	67-72-1	3.98		1.31E+02	h		7.27E+01	1.31E+02	1.80E+00		1.80E+00
Hexachlorophene	70-30-4	7.54		1.07E+05	h		4.31E+04	1.07E+05	2.48E+00		2.48E+00
Hexamethylene-1,5-diisocyanate	822-06-0	1.27		7.89E-01	h		No Kd	7.89E-01	1.00E+01	Default	1.00E+01
Mirex	2385-85-5	6.89		3.14E+04	h		4.00E+04	3.14E+04	7.85E-01		7.85E-01
Nitrofen	1836-75-5	5.53		2.42E+03	h		No Kd	2.42E+03	1.00E+01	Default	1.00E+01
Pentachlorobenzene	608-93-5	5.09	3.20E-01				1.29E+03	2.60E+03	2.01E+00		3.20E-01
Pentachloronitrobenzene	82-68-8	4.64	4.51E+02		h		2.36E+02	#REF!	FALSE		5.51E-02
Pentachlorophenol	87-86-5	5.08	1.03E+03		h		7.97E+02	5.20E+01	6.53E-02		6.53E-02
Picric acid	88-89-1	2.03		3.29E+00	h		No Kd	3.29E+00	1.00E+01	Default	1.00E+01
Pronamide	23950-58-5	3.51		5.36E+01	h		3.10E+01	5.36E+01	1.73E+00		1.73E+00
Strychnine	57-24-9	1.93		2.72E+00	h		1.81E+00	2.72E+00	1.50E+00		1.50E+00
Terphenyls	26140-60-3	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Tributyl phosphate	126-73-8	4.00		1.35E+02	h		No Kd	1.35E+02	1.00E+01	Default	1.00E+01
Trifluralin	1582-09-8	5.34		1.69E+03	h		2.41E+02	1.69E+03	7.00E+00		7.00E+00
Triphenylamine	603-34-9	No data		No data			No Kd	1.00E+05	1.00E+01	Default	1.00E+01
Herbicides and Organochlorinated Pesticides											
2,4,5-T	93-76-5	3.36		4.03E+01	h		2.10E+00	4.03E+01	1.92E+01		1.92E+01
2,4-D and esters	94-75-7	2.81		1.43E+01	h		7.98E-01	1.43E+01	1.79E+01		1.79E+01
4,4-DDD	72-54-8	6.20		8.55E+03	h		4.00E+03	8.55E+03	2.14E+00		2.14E+00
4,4-DDE	72-55-9	6.26	9.50E-01				3.46E+03	1.19E+04	3.45E+00		9.50E-01
4,4-DDT	50-29-3	6.00		5.86E+03	h		9.60E+03	5.86E+03	6.11E-01		6.11E-01
Aldrin	309-00-2	6.18		8.21E+03	h		1.95E+03	8.21E+03	4.21E+00		4.21E+00
alpha-BHC	319-84-6	3.80		9.24E+01	h		7.05E+01	9.24E+01	1.31E+00		1.31E+00
beta-BHC	319-85-7	3.83		9.85E+01	h		8.56E+01	9.85E+01	1.15E+00		1.15E+00
Chlordane	57-74-9	5.94		5.21E+03	h		2.05E+03	5.21E+03	2.54E+00		2.54E+00
Delta-BHC	319-86-8	4.14		1.76E+02	h		2.64E+01	1.76E+02	6.65E+00		6.65E+00
Dieldrin	60-57-1	5.27		1.48E+03	h		1.02E+03	1.48E+03	1.45E+00		1.45E+00
Endothall	145-73-3	-0.87		1.39E-02	h		5.61E-03	1.39E-02	2.47E+00		2.47E+00

Table C2-10. Aquatic Sediment-to-Animal Transfer Factors (BASF) for Ecological Receptors (mg/kg tissue wet weight/mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BASF [b]	BASF Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BASF	Kd [d]	BCF_{inv} [e]	SAIC computed BASF [f]	Comments	Preferred BASF [g]
Endrin	72-20-8	4.89		7.25E+02	h		4.32E+02	7.25E+02	1.68E+00		1.68E+00
gamma-BHC (Lindane)	58-89-9	3.72		7.96E+01	h		4.29E+01	7.96E+01	1.86E+00		1.86E+00
Heptachlor	76-44-8	5.02	1.67E+00				3.81E+02	3.81E+03	9.99E+00		1.67E+00
Isodrin	465-73-6	3.55		5.77E+01	h		No Kd	5.77E+01	1.00E+01	Default	1.00E+01
Methoxychlor	72-43-5	4.53		3.64E+02	h		3.20E+03	3.64E+02	1.14E-01		1.14E-01
Silvex (2,4,5-TP)	93-72-1	4.07		1.55E+02	h		No Kd	1.55E+02	1.00E+01	Default	1.00E+01
Toxaphene	8001-35-2	5.50		2.28E+03	h		4.00E+03	2.28E+03	5.71E-01		5.71E-01
Inorganic Chemicals and Compounds											
Metals											
Aluminum	7429-90-5	NA	9.00E-01		l		1.50E+03	4.07E+03	2.71E+00		2.71E+00
Antimony	7440-36-0	NA	9.00E-01		l		4.50E+01	7.00E+00	1.56E-01		1.56E-01
Arsenic	7440-38-2	NA	9.00E-01		l		2.50E+01	7.30E+01	2.92E+00		2.92E+00
Barium	7440-39-3	NA	9.00E-01		l		1.10E+01	2.00E+02	1.82E+01		1.82E+01
Beryllium	7440-41-7	NA	9.00E-01		l		2.30E+01	4.50E+01	1.96E+00		1.96E+00
Bismuth	7440-69-9	NA		9.00E-01	l		2.00E+02	4.07E+03	2.03E+01		2.03E+01
Boron	7440-42-8	NA		9.00E-01	l		3.00E+00	4.07E+03	1.36E+03		1.36E+03
Cadmium	7440-43-9	NA	3.40E+00				1.50E+01	3.46E+03	2.31E+02		2.31E+02
Calcium	7440-70-2	NA		9.00E-01	l		4.00E+00	4.07E+03	1.02E+03		1.02E+03
Chromium (and VI)	18540-29-9	NA	3.90E-01				1.20E+03	3.00E+03	2.50E+00		2.50E+00
Cobalt	7440-48-4	NA		9.00E-01	l		4.50E+01	4.07E+03	9.04E+01		9.04E+01
Copper	7440-50-8	NA	3.00E-01				3.50E+01	3.72E+03	1.06E+02		1.06E+02
Iron	7439-89-6	NA		9.00E-01	l		2.50E+01	4.07E+03	1.63E+02		1.63E+02
Lead	7439-92-1	NA	6.30E-01				9.00E+02	5.06E+03	5.62E+00		5.62E+00
Lithium	7439-93-2	NA		9.00E-01	l		3.00E+02	4.07E+03	1.36E+01		1.36E+01
Magnesium	7439-95-4	NA		9.00E-01	l		4.50E+00	4.07E+03	9.04E+02		9.04E+02
Manganese	7439-96-5	NA		9.00E-01	l		6.50E+01	4.07E+03	6.26E+01		6.26E+01
Mercury	7439-97-6	NA	6.80E-02				1.00E+03	2.02E+04	2.02E+01		2.02E+01
Molybdenum	7439-98-7	NA		9.00E-01	l		2.00E+01	4.07E+03	2.03E+02		2.03E+02
Nickel	7440-02-0	NA	9.00E-01		l		1.60E+01	2.80E+01	1.75E+00		1.75E+00

Table C2-10. Aquatic Sediment-to-Animal Transfer Factors (BASF) for Ecological Receptors (mg/kg tissue wet weight/mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BASF [b]	BASF Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BASF	Kd [d]	BCF_{inv} [e]	SAIC computed BASF [f]	Comments	Preferred BASF [g]
Potassium	7440-09-7	NA		9.00E-01	1		5.50E+00	4.07E+03	7.39E+02		7.39E+02
Rhodium	7440-16-6	NA		9.00E-01	1		6.00E+01	4.07E+03	6.78E+01		6.78E+01
Selenium	7782-49-2	NA	9.00E-01		1		1.80E+01	1.26E+03	7.01E+01		7.01E+01
Silicon	7440-21-3	NA		9.00E-01	1		3.00E+01	4.07E+03	1.36E+02		1.36E+02
Silver	7440-22-4	NA	9.00E-01		1		1.00E-01	2.98E+02	2.98E+03		2.98E+03
Sodium	7440-23-5	NA		9.00E-01	1		1.00E+02	4.07E+03	4.07E+01		4.07E+01
Strontium	7440-24-6	NA		9.00E-01	1		No Kd	4.07E+03	1.00E+00	Default	1.00E+00
Tantalum	7440-25-7	NA		9.00E-01	1		6.50E+02	4.07E+03	6.26E+00		6.26E+00
Thallium	7440-28-0	NA	9.00E-01		1		4.40E+01	1.50E+04	3.41E+02		3.41E+02
Tin	7440-31-5	NA		9.00E-01	1		2.50E+02	4.07E+03	1.63E+01		1.63E+01
Tungsten	7440-33-7	NA		9.00E-01	1		1.50E+02	4.07E+03	2.71E+01		2.71E+01
Uranium	7440-61-1	NA		9.00E-01	1		4.50E+02	4.07E+03	9.04E+00		9.04E+00
Vanadium	7440-62-2	NA		9.00E-01	1		1.00E+03	4.07E+03	4.07E+00		4.07E+00
Yttrium	7440-65-5	NA		9.00E-01	1		5.00E+02	4.07E+03	8.13E+00		8.13E+00
Zinc	7440-66-6	NA	5.70E-01				6.20E+01	4.58E+03	7.38E+01		7.38E+01
Zirconium	7440-67-7	NA		9.00E-01	1		3.00E+03	4.07E+03	1.36E+00		1.36E+00
Non-metals and Anions											
Ammonia/Ammonium	7664-41-7	NA					No Kd	5.00E+02	1.00E+00	Default	1.00E+00
Bromide	24959-67-9	NA		9.00E-01	1		No Kd	4.07E+03	1.00E+00	Default	1.00E+00
Chloride	16887-00-6	NA		9.00E-01	1		No Kd	4.07E+03	1.00E+00	Default	1.00E+00
Cyanide	57-12-5	NA	9.00E-01		1		No Kd	4.07E+03	1.00E+00	Default	1.00E+00
Fluoride	16984-48-8	NA		9.00E-01	1		No Kd	4.07E+03	1.00E+00	Default	1.00E+00
Hydroxide	14280-30-9	NA	NA				No Kd	NA	NA		NA
Iodine	7553-56-2	NA		9.00E-01	1		No Kd	4.07E+03	1.00E+00	Default	1.00E+00
Nitrate	14797-55-8	NA		9.00E-01	1		No Kd	4.07E+03	1.00E+00	Default	1.00E+00
Nitrite	14797-65-0	NA		9.00E-01	1		No Kd	4.07E+03	1.00E+00	Default	1.00E+00
Phosphate	14265-44-2	NA		9.00E-01	1		No Kd	4.07E+03	1.00E+00	Default	1.00E+00
Phosphorus	7723-14-0	NA		9.00E-01	1		No Kd	4.07E+03	1.00E+00	Default	1.00E+00
Sulfate	14808-79-8	NA		9.00E-01	1		No Kd	4.07E+03	1.00E+00	Default	1.00E+00

Table C2-10. Aquatic Sediment-to-Animal Transfer Factors (BASF) for Ecological Receptors (mg/kg tissue wet weight/mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) BASF[b]	BASF Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance BASF	Kd[d]	BCF_{inv}[e]	SAIC computed BASF[f]	Comments	Preferred BASF[g]
Total Sulfur	63705-05-5	NA		9.00E-01	1		No Kd	4.07E+03	1.00E+00	Default	1.00E+00
Priority Pollutants											
Carbon Dioxide	124-38-9	NA					No Kd	NA	NA		NA
Nitrogen Dioxide	10102-44-0	NA					No Kd	5.00E+02	1.00E+00	Default	1.00E+00
Ozone	10028-15-6	NA					No Kd	5.00E+02	1.00E+00	Default	1.00E+00
Particulate Matter	No CAS #	NA	NA				No Kd	NA	NA		NA
Sulfur Dioxode	7446-09-5	NA					No Kd	5.00E+02	1.00E+00	Default	1.00E+00
Radionuclides											
Americium-241	1596-10-2	NA		9.00E-01	1		7.00E+02	4.07E+03	5.81E+00		5.81E+00
Antimony-125	14234-35-6	NA	9.00E-01		1		4.50E+01	7.00E+00	1.56E-01		1.56E-01
Barium-137	13981-97-0	NA	9.00E-01		1		6.00E+01	2.00E+02	3.33E+00		3.33E+00
Cadmium-113	None	NA	3.40E+00				6.50E+00	3.46E+03	5.32E+02		5.32E+02
Cesium-134	13967-70-9	NA		9.00E-01	1		1.00E+03	4.07E+03	4.07E+00		4.07E+00
Cesium-137	10045-97-3	NA		9.00E-01	1		1.00E+03	4.07E+03	4.07E+00		4.07E+00
Europium-154	15585-10-1	NA		9.00E-01	1		6.50E+02	4.07E+03	6.26E+00		6.26E+00
Europium-155	14391-16-3	NA		9.00E-01	1		6.50E+02	4.07E+03	6.26E+00		6.26E+00
Nickel-63	13981-37-8	NA	9.00E-01		1		1.50E+02	2.80E+01	1.87E-01		1.87E-01
Plutonium-239	15117-48-3	NA		9.00E-01	1		4.50E+03	4.07E+03	9.04E-01		9.04E-01
Plutonium-241	14119-32-5	NA		9.00E-01	1		4.50E+03	4.07E+03	9.04E-01		9.04E-01
Samarium-151	15715-94-3	NA		9.00E-01	1		6.50E+02	4.07E+03	6.26E+00		6.26E+00
Strontium-90	10098-97-2	NA		9.00E-01	1		3.50E+01	4.07E+03	1.16E+02		1.16E+02
Technetium-99	14133-79-7	NA		9.00E-01	1		1.50E+00	4.07E+03	2.71E+03		2.71E+03
Tritium	10028-17-8	NA		9.00E-01	1		No Kd	4.07E+03			9.00E-01
Yttrium-90	10098-91-6	NA		9.00E-01	1		5.00E+02	4.07E+03	8.13E+00		8.13E+00
Uranium-232	14158-29-3	NA		9.00E-01	1		4.50E+02	4.07E+03	9.04E+00		9.04E+00
Uranium-233	13968-55-3	NA		9.00E-01	1		4.50E+02	4.07E+03	9.04E+00		9.04E+00
Uranium-234	13966-29-5	NA		9.00E-01	1		4.50E+02	4.07E+03	9.04E+00		9.04E+00
Uranium-235	15117-96-1	NA		9.00E-01	1		4.50E+02	4.07E+03	9.04E+00		9.04E+00
Uranium-236	13982-70-2	NA		9.00E-01	1		4.50E+02	4.07E+03	9.04E+00		9.04E+00

Table C2-10. Aquatic Sediment-to-Animal Transfer Factors (BASF) for Ecological Receptors (mg/kg tissue wet weight/mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow} [a]	EPA (1999) BASF [b]	BASF Calculated by EPA (1999) Methods [c]	Notes	Ecology Guidance BASF	Kd [d]	BCF_{inv} [e]	SAIC computed BASF [f]	Comments	Preferred BASF [g]
Uranium-238	7440-61-1	NA		9.00E-01	1		4.50E+02	4.07E+03	9.04E+00		9.04E+00
Actinium-227	14952-40-0	NA		9.00E-01	1		1.50E+03	4.07E+03	2.71E+00		2.71E+00
Americium-243	14993-75-0	NA		9.00E-01	1		7.00E+02	4.07E+03	5.81E+00		5.81E+00
Carbon-14	14762-75-5	NA		9.00E-01	1		No Kd	4.07E+03			9.00E-01
Cobalt-60	10198-40-0	NA		9.00E-01	1		4.50E+01	4.07E+03	9.04E+01		9.04E+01
Curium-242	15510-73-3	NA		9.00E-01	1		2.00E+03	4.07E+03	2.03E+00		2.03E+00
Curium-243	15757-87-6	NA		9.00E-01	1		2.00E+03	4.07E+03	2.03E+00		2.03E+00
Curium-244	13981-15-2	NA		9.00E-01	1		2.00E+03	4.07E+03	2.03E+00		2.03E+00
Europium-152	14683-23-9	NA		9.00E-01	1		6.50E+02	4.07E+03	6.26E+00		6.26E+00
Iodine-129	15046-84-1	NA		9.00E-01	1		6.00E+01	4.07E+03	6.78E+01		6.78E+01
Neptunium-237	13994-20-2	NA		9.00E-01	1		3.00E+01	4.07E+03	1.36E+02		1.36E+02
Nickel-59	14336-70-0	NA	9.00E-01		1		1.50E+02	2.80E+01	1.87E-01		1.87E-01
Niobium-93	None	NA		9.00E-01	1		3.50E+02	4.07E+03	1.16E+01		1.16E+01
Plutonium-238	13981-16-3	NA		9.00E-01	1		4.50E+03	4.07E+03	9.04E-01		9.04E-01
Plutonium-240	14119-33-6	NA		9.00E-01	1		4.50E+03	4.07E+03	9.04E-01		9.04E-01
Plutonium-242	13982-10-0	NA		9.00E-01	1		4.50E+07	4.07E+03	9.03E-05		9.03E-05
Protactinium-231	14331-85-2	NA		9.00E-01	1		2.50E+03	4.07E+03	1.63E+00		1.63E+00
Radium-226	13982-63-3	NA		9.00E-01	1		4.50E+02	4.07E+03	9.04E+00		9.04E+00
Radium-228	15262-20-1	NA		9.00E-01	1		4.50E+02	4.07E+03	9.04E+00		9.04E+00
Ruthenium-106	13967-48-1	NA		9.00E-01	1		3.50E+02	4.07E+03	1.16E+01		1.16E+01
Selenium-79	None	NA	9.00E-01		1		3.00E+02	1.26E+03	4.21E+00		4.21E+00
Thorium-229	15594-54-4	NA		9.00E-01	1		1.50E+05	4.07E+03	2.71E-02		2.71E-02
Thorium-232	7440-29-1	NA		9.00E-01	1		1.50E+05	4.07E+03	2.71E-02		2.71E-02
Tin-126	15832-50-5	NA		9.00E-01	1		2.50E+02	4.07E+03	1.63E+01		1.63E+01
Zirconium-93	15751-77-6	NA		9.00E-01	1		3.00E+03	4.07E+03	1.36E+00		1.36E+00

NA = Not applicable

[a] $\log_{10}$ of K_{ow} values in Table 4.1

[b] Data published in Appendix C, Table C-6 of EPA (1999)

[c] Calculated or chosen as described in Appendix C, Sect. C-1.6 of EPA (1999)

Table C2-10. Aquatic Sediment-to-Animal Transfer Factors (BASF) for Ecological Receptors (mg/kg tissue wet weight/mg/kg sediment)

Constituent of Potential Concern	CAS Registry Number	Log K_{ow}[a]	EPA (1999) BASF[b]	BASF Calculated by EPA (1999) Methods[c]	Notes	Ecology Guidance BASF	Kd[d]	BCF_{inv}[e]	SAIC computed BASF[f]	Comments	Preferred BASF[g]

[d] Sediment Kd values from Appendix B Table B-1-1 or soil Kds from Baes and others. (1984) Fig. 2.31 for radionuclides and other elements not shown in Appendix B Table B-1-1

[e] BCF_{inv} values from Table C2-8

[f] Calculated by using BCF and Kd: BASF = BCF-I/Kd

[g] Selection criteria described in Sect. 8.2.4.4

[h] Calculated by using log K_{ow}: log BASF = 0.819 x log K_{ow} -1.146 (Southworth, Beauchamp, and Schmieder 1978)

[i] Calculated by using the BASF for TCDD from Appendix C, Table C-6 of EPA (1999) and BEFs for other congeners

[j] Value for Aroclor 1254 from Appendix C of EPA (1999), Table C-6, was used for PCB mixtures

[k] Benzo(a)pyrene value from Appendix C of EPA (1999), Table C-6, was used as a surrogate value

[l] Average of values for other inorganics as published in Appendix C of EPA (1999), Table C-6

References for Appendix C-2

Baes CF, Sharp RD, Sjoreen AL, and Shor RW. 1984. *A Review and Analysis of Parameters for Assessing Transport of Environmentally Released Radionuclides through Agriculture*, ORNL-5786, September 1984. Martin Marietta Energy Systems, Oak Ridge, Tennessee, USA.

Corp N and Morgan AJ. 1991. *Accumulation of heavy metals from polluted soils by the earthworm, Lumbricus rubellus: Can laboratory exposure of "control" worms reduce biomonitoring problems?*, Environmental Pollution, volume 74, p 39-52.

HAZWRAP. 1994. *Loring Air Force Base Ecological Risk Methodology*, August 1994, Hazardous Waste Remedial Actions Program. Lockheed Martin Energy Systems, Inc., Oak Ridge, Tennessee, USA.

NRC. 1992. *Residual Radioactive Contamination from Decommissioning, Technical Basis for Translating Contamination Levels to Annual Total Effective Dose Equivalent*, Final Report, NUREG/CR-5512, PNL-7794, Volume 1. US Nuclear Regulatory Commission, Washington, D.C, USA.

Appendix C-3

Toxicity Reference Values for Plants, Earthworms, Birds, Mammals, Sediment-Dwelling Biota, and Surface Water Organisms

Table C3-1. Toxicity Reference Values (TRVs) for Plants

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997a[b] (mg/kg)	Recommended TRV[c] (mg/kg)
Organic Compounds						
Aromatic Halogenated Hydrocarbons						
4-Chloro-3-methylphenol	59-50-7	No data			No data	
2,3,4,6-Tetrachlorophenol	58-90-2	No data			No data	
Aromatic Nonhalogenated Hydrocarbons						
2-Nitrotoluene	88-72-2	No data			No data	
4-Nitrobiphenyl	92-93-3	No data			No data	
Benzaldehyde	100-52-7	No data			No data	
Benzene	71-43-2	No data			No data	
Benzyl alcohol	100-51-6	No data			No data	
Ethyl benzene	100-41-4	No data			No data	
m-Xylene	108-38-3	No data			No data	
o-Xylene	95-47-6	No data			No data	
p-Xylene	106-42-3	No data			No data	
Styrene	100-42-5	No data			3.00E+02	3.00E+02
Toluene	108-88-3	No data			2.00E+02	2.00E+02
Non-aromatic Nonhalogenated Hydrocarbons						
1,2-Epoxybutane	106-88-7	No data			No data	
1,3-Butadiene	106-99-0	No data			No data	
1,4-Dioxane	123-91-1	No data			No data	
1-Methylpropyl alcohol	78-92-2	No data			No data	
1-Nitropropane	108-03-2	No data			No data	
2,2,4-Trimethylpentane	540-84-1	No data			No data	
2-Butanone	78-93-3	No data			No data	
2-Butenaldehyde (2-Butenal)	4170-30-3	No data			No data	
2-Ethoxyethanol	110-80-5	No data			No data	
2-Heptanone	110-43-0	No data			No data	
2-Hexanone	591-78-6	No data			No data	
2-Methoxyethanol	109-86-4	No data			No data	
2-Methyl-2-propanol	75-65-0	No data			No data	

Table C3-1. Toxicity Reference Values (TRVs) for Plants

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997a[b] (mg/kg)	Recommended TRV[c] (mg/kg)
2-Methyl-2-propenenitrile	126-98-7	No data			No data	
2-Methylaziridine	75-55-8	No data			No data	
2-Methylpropyl alcohol	78-83-1	No data			No data	
2-Pentanone	107-87-9	No data			No data	
2-Propanone (Acetone)	67-64-1	No data			No data	
2-Propene-1-ol	107-18-6	No data			No data	
2-Propyl alcohol	67-63-0	No data			No data	
3-Heptanone	106-35-4	No data			No data	
3-Methyl-1-butanol	123-51-3	No data			No data	
3-Methyl-2-butanone	563-80-4	No data			No data	
3-Pentanone	96-22-0	No data			No data	
4-Heptanone	123-19-3	No data			No data	
4-Methyl-2-pentanone	108-10-1	No data			No data	
4-Methyl-3-penten-2-one	141-79-7	No data			No data	
5-Methyl-2-hexanone	110-12-3	No data			No data	
Acetaldehyde	75-07-0	No data			No data	
Acetamide	60-35-5	No data			No data	
Acetic acid	64-19-7	No data			No data	
Acetic acid ethyl ester	141-78-6	No data			No data	
Acetic acid n-butyl ester	123-86-4	No data			No data	
Acetonitrile	75-05-8	No data			No data	
Acrolein	107-02-8	No data			No data	
Acrylonitrile	107-13-1	No data			No data	
Bis(isopropyl)ether	108-20-3	No data			No data	
Butane	106-97-8	No data			No data	
Carbon disulfide	75-15-0	No data			No data	
Cyanogen	460-19-5	No data			No data	
Cyclohexane	110-82-7	No data			No data	
Cyclohexanone	108-94-1	No data			No data	
Cyclohexene	110-83-8	No data			No data	
Cyclopentane	287-92-3	No data			No data	

Table C3-1. Toxicity Reference Values (TRVs) for Plants

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997a[b] (mg/kg)	Recommended TRV[c] (mg/kg)
Ethyl alcohol	64-17-5	No data			No data	
Ethyl ether	60-29-7	No data			No data	
Ethyl methacrylate	97-63-2	No data			No data	
Formaldehyde	50-00-0	No data			No data	
Formamide	75-12-7	No data			No data	
Formic acid	64-18-6	No data			No data	
Formic acid, methyl ester	107-31-3	No data			No data	
Glycidylaldehyde	765-34-4	No data			No data	
Methyl acetate	79-20-9	No data			No data	
Methyl alcohol	67-56-1	No data			No data	
Methyl isocyanate	624-83-9	No data			No data	
Methyl methacrylate	80-62-6	No data			No data	
Methyl tert-butyl ether	1634-04-4	No data			No data	
Methylacetylene	74-99-7	No data			No data	
Methylcyclohexane	108-87-2	No data			No data	
N,N-Dimethylacetamide	127-19-5	No data			No data	
n-Butyl alcohol	71-36-3	No data			No data	
n-Heptane	142-82-5	No data			No data	
n-Hexane	110-54-3	No data			No data	
Nitromethane	75-52-5	No data			No data	
n-Nonane	111-84-2	No data			No data	
n-Octane	111-65-9	No data			No data	
n-Pentane	109-66-0	No data			No data	
n-Propionaldehyde	123-38-6	No data			No data	
n-Propyl alcohol	71-23-8	No data			No data	
n-Valeraldehyde	110-62-3	No data			No data	
Oxirane	75-21-8	No data			No data	
p-Cymene	99-87-6	No data			No data	
Phosgene	75-44-5	No data			No data	
Propargyl alcohol	107-19-7	No data			No data	
Propionic acid	79-09-4	No data			No data	

Table C3-1. Toxicity Reference Values (TRVs) for Plants

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997a[b] (mg/kg)	Recommended TRV[c] (mg/kg)
Propionitrile	107-12-0	No data			No data	
Propylene gylcol monomethyl ether	107-98-2	No data			No data	
p-tert-Butyltoluene	98-51-1	No data			No data	
Triethylamine	121-44-8	No data			No data	
Trimethylamine	75-50-3	No data			No data	
Vinyl acetate	108-05-4	No data			No data	
Non-aromatic Halogenated Hydrocarbons						
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	No data			No data	
1,1,1,2-Tetrachloroethane	630-20-6	No data			No data	
1,1,1-Trichloroethane	71-55-6	No data			No data	
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	No data			No data	
1,1,2,2-Tetrachloroethane	79-34-5	No data			No data	
1,1,2,2-Tetrachloroethene	127-18-4	No data			No data	
1,1,2-Trichloroethane	79-00-5	No data			No data	
1,1,2-Trichloroethylene	79-01-6	No data			No data	
1,1-Dichloroethane	75-34-3	No data			No data	
1,1-Dichloroethene	75-35-4	No data			No data	
1,2,2-Trichloro-1,1,2-trifluoroethane	76-13-1	No data			No data	
1,2,3-Trichloropropane	96-18-4	No data			No data	
1,2-Dibromo-3-chloropropane	96-12-8	No data			No data	
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	No data			No data	
1,2-Dichloroethane	107-06-2	No data			No data	
1,2-Dichloroethylene	540-59-0	No data			No data	
1,2-Dichloropropane	78-87-5	No data			No data	
1,3-Dichloropropene	542-75-6	No data			No data	
1,4-Dichloro-2-butene	764-41-0	No data			No data	
1-Chloroethene	75-01-4	No data			No data	
2,2-Dichloropropionic acid	75-99-0	No data			No data	
2-Chloropropane	75-29-6	No data			No data	
3-Chloropropene (allyl chloride)	107-05-1	No data			No data	
Bromochloromethane	74-97-5	No data			No data	

Table C3-1. Toxicity Reference Values (TRVs) for Plants

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997a[b] (mg/kg)	Recommended TRV[c] (mg/kg)
Bromodichloromethane	75-27-4	No data			No data	
Bromoethene	593-60-2	No data			No data	
Bromoform	75-25-2	No data			No data	
Bromomethane	74-83-9	No data			No data	
Carbon tetrachloride	56-23-5	No data			No data	
Chlorodibromomethane	124-48-1	No data			No data	
Chlorodifluoromethane	75-45-6	No data			No data	
Chloroethane	75-00-3	No data			No data	
Chloroform	67-66-3	No data			No data	
Chloromethane	74-87-3	No data			No data	
Chloropentafluoroethane	76-15-3	No data			No data	
cis-1,2-Dichloroethene	156-59-2	No data			No data	
cis-1,3-Dichloropropene	10061-01-5	No data			No data	
Cyanogen bromide	506-68-3	No data			No data	
Cyanogen chloride	506-77-4	No data			No data	
Dichlorodifluoromethane	75-71-8	No data			No data	
Dichlorofluoromethane	75-43-4	No data			No data	
Dichloromethane	75-09-2	No data			No data	
Difluorodibromomethane	75-61-6	No data			No data	
Hexafluoroacetone	684-16-2	No data			No data	
Iodomethane	74-88-4	No data			No data	
Methylene bromide	74-95-3	No data			No data	
Pentachloroethane	76-01-7	No data			No data	
trans-1,2-Dichloroethene	156-60-5	No data			No data	
trans-1,3-Dichloropropene	10061-02-6	No data			No data	
Trichloroacetic acid	76-03-9	No data			No data	
Trichlorofluoroethane	27154-33-2	No data			No data	
Trichlorofluoromethane	75-69-4	No data			No data	
Trifluorobromomethane	75-63-8	No data			No data	
Dioxin and Furan Compounds						
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	35822-46-9	No data			No data	

Table C3-1. Toxicity Reference Values (TRVs) for Plants

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997a[b] (mg/kg)	Recommended TRV[c] (mg/kg)
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	No data			No data	
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	No data			No data	
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	39227-28-6	No data			No data	
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	No data			No data	
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	57653-85-7	No data			No data	
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	No data			No data	
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	19408-74-3	No data			No data	
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	No data			No data	
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	40321-76-4	No data			No data	
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	No data			No data	
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	No data			No data	
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	No data			No data	
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1746-01-6	No data			No data	
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	No data			No data	
Dibenzofuran	132-64-9	No data			No data	
Octachlorodibenzo(p)dioxin	3268-87-9	No data			No data	
Octachlorodibenzofuran	39001-02-0	No data			No data	
PCBs						
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	No data			No data	
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	No data			No data	
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	No data			No data	
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	No data			No data	
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	No data			No data	
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	No data			No data	
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	No data			No data	
2',3,4,4',5-Pentachlorobiphenyl	no cas #	No data			No data	
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	No data			No data	
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	No data			No data	
3,3',4,4',5-Pentachlorobiphenyl	no cas #	No data			No data	
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	No data			No data	
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	No data			No data	

Table C3-1. Toxicity Reference Values (TRVs) for Plants

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997a[b] (mg/kg)	Recommended TRV[c] (mg/kg)
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	No data			No data	
Polychlorinated biphenyls (PCBs)	1336-36-3	1.00E+01			4.00E+01	1.00E+01
Phthalates						
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	No data			No data	
Butylbenzyl phthalate	85-68-7	No data			No data	
Dibutyl phthalate	84-74-2	No data			2.00E+02	2.00E+02
Diethyl phthalate	84-66-2	No data			1.00E+02	1.00E+02
Dimethylphthalate	131-11-3	No data			No data	
n-Dioctyl phthalate	117-84-0	No data			No data	
Light Polycyclic Aromatic Hydrocarbons (molecular weight <200 g/mole)						
2-Chloronapthalene	91-58-7	No data			No data	
2-Methyl naphthalene	91-57-6	No data			No data	
5-Nitroacenaphthene	602-87-9	No data			No data	
Acenaphthene	83-32-9	No data			2.00E+01	2.00E+01
Acenaphthylene	208-96-8	No data			No data	
Anthracene	120-12-7	No data			No data	
Fluorene	86-73-7	No data			No data	
Indene	95-13-6	No data			No data	
Naphthalene	91-20-3	No data			No data	
Phenanthrene	85-01-8	No data			No data	
Pyrene	129-00-0	No data			No data	
Heavy Polycyclic Aromatic Hydrocarbons (molecular weight >200 g/mole)						
3-Methylcholanthrene	56-49-5	No data			No data	
5-Methylchrysene	3697-24-3	No data			No data	
Benzo(a)anthracene	56-55-3	1.20E+00	d		No data	1.20E+00
Benzo(a)pyrene	50-32-8	1.20E+00			No data	1.20E+00
Benzo(b)fluoranthene	205-99-2	1.20E+00			No data	1.20E+00
Benzo(e)pyrene	192-97-2	No data			No data	
Benzo(g,h,i)perylene	191-24-2	No data			No data	
Benzo(j)fluoranthene	205-82-3	No data			No data	

Table C3-1. Toxicity Reference Values (TRVs) for Plants

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997a[b] (mg/kg)	Recommended TRV[c] (mg/kg)
Benzo(k)fluoranthene	207-08-9	1.20E+00	d		No data	1.20E+00
Benzo[a,i]pyrene	191-30-0	No data			No data	
Chrysene	218-01-9	1.20E+00	d		No data	1.20E+00
Dibenz(a,h)anthracene	53-70-3	1.20E+00	d		No data	1.20E+00
Dibenz[a,h]acridine	226-36-8	No data			No data	
Dibenz[a,j]acridine	224-42-0	No data			No data	
Dibenzo(a,e)fluoranthene	5385-75-1	No data			No data	
Dibenzo(a,h)fluoranthene	no cas #	No data			No data	
Dibenzo[a,e]pyrene	192-65-4	No data			No data	
Dibenzo[a,h]pyrene	189-64-0	No data			No data	
Dibenzo[a,i]pyrene	189-55-9	No data			No data	
Fluoranthene	206-44-0	No data			No data	
Hexachloronaphthalene	1335-87-1	No data			No data	
Indeno(1,2,3-cd)pyrene	193-39-5	1.20E+00	d		No data	1.20E+00
Octachloronaphthalene	2234-13-1	No data			No data	
Pentachloronaphthalene	1321-64-8	No data			No data	
Tetrachloronaphthalene	1335-88-2	No data			No data	
Trichloronaphthalene	1321-65-9	No data			No data	
Light Substituted Benzene Compounds (MW <200 g/mole)						
1,2,3-Trichlorobenzene	87-61-6	No data			No data	
1,2,4-Trichlorobenzene	120-82-1	No data			No data	
1,2,4-Trimethyl benzene	95-63-6	No data			No data	
1,2-Dichlorobenzene	95-50-1	No data			No data	
1,3,5-Trimethyl benzene	108-67-8	No data			No data	
1,3-Dichlorobenzene	541-73-1	No data			No data	
1,3-Dinitrobenzene	99-65-0	No data			No data	
1,4-Dichlorobenzene	106-46-7	No data			No data	
1,4-Dinitrobenzene	100-25-4	No data			No data	
2,4,5-Trichlorophenol	95-95-4	No data			4.00E+00	4.00E+00
2,4,6-Trichlorophenol	88-06-2	No data			No data	
2,4-Dichlorophenol	120-83-2	No data			No data	

Table C3-1. Toxicity Reference Values (TRVs) for Plants

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997a[b] (mg/kg)	Recommended TRV[c] (mg/kg)
2,4-Dimethylphenol	105-67-9	No data			No data	
2,4-Dinitrophenol	51-28-5	No data			2.00E+01	2.00E+01
2,4-Dinitrotoluene	121-14-2	No data			No data	
2,6-Dinitrotoluene	606-20-2	No data			No data	
2-Chlorophenol	95-57-8	No data			No data	
2-Chlorotoluene	95-49-8	No data			No data	
2-Nitrophenol	88-75-5	No data			No data	
4,6-Dinitro-o-cresol	534-52-1	No data			No data	
4-Chlorotoluene	106-43-4	No data			No data	
4-Nitrophenol	100-02-7	No data			No data	
alpha-Methylstyrene	98-83-9	No data			No data	
Aniline	62-53-3	No data			No data	
Benzotrichloride	98-07-7	No data			No data	
Benzyl chloride	100-44-7	No data			No data	
Bromobenzene	108-86-1	No data			No data	
Chlorobenzene	108-90-7	No data			No data	
Cumene	98-82-8	No data			No data	
m-Cresol	108-39-4	No data			No data	
n-Butyl benzene	104-51-8	No data			No data	
Nitrobenzene	98-95-3	No data			No data	
n-Propyl benzene	103-65-1	No data			No data	
o-Cresol	95-48-7	No data			No data	
o-Dinitrobenzene	528-29-0	No data			No data	
o-Nitroaniline	88-74-4	No data			No data	
o-Toluidine	95-53-4	No data			No data	
p-Chloroaniline	106-47-8	No data			No data	
p-Cresol	106-44-5	No data			No data	
Phenol	108-95-2	No data			7.00E+01	7.00E+01
p-Nitrochlorobenzene	100-00-5	No data			No data	
p-Toluidine	106-49-0	No data			No data	
sec-Butyl benzene	135-98-8	No data			No data	

Table C3-1. Toxicity Reference Values (TRVs) for Plants

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997a[b] (mg/kg)	Recommended TRV[c] (mg/kg)
tert-Butyl benzene	98-06-6	No data			No data	
Toluene-2,6-diamine	823-40-5	No data			No data	
Trimethyl benzene	25551-13-7	No data			No data	
Other Light Semivolatile Compounds (molecular weight <200 g/mole)						
1,1'-Biphenyl	92-52-4	No data			6.00E+01	6.00E+01
1,1-Dimethylhydrazine	57-14-7	No data			No data	
1,2-Dimethylhydrazine	540-73-8	No data			No data	
1,2-Diphenylhydrazine	122-66-7	No data			No data	
1,3-Propane sultone	1120-71-4	No data			No data	
2,4-Toluene diisocyante	584-84-9	No data			No data	
2-Chloroacetophenone	532-27-4	No data			No data	
2-Propenoic acid	79-10-7	No data			No data	
4,4-Methylenedianiline	101-77-9	No data			No data	
Acetophenone	98-86-2	No data			No data	
Benzoic acid	65-85-0	No data			No data	
bis(2-Chloroethoxy)methane	111-91-1	No data			No data	
bis(2-Chloroethyl) ether	111-44-4	No data			No data	
Chlorocyclopentadiene	41851-50-7	No data			No data	
Cyclohexanol	108-93-0	No data			No data	
Dichloroisopropyl ether	108-60-1	No data			No data	
Dichloromethyl ether	542-88-1	No data			No data	
Dichloropentadiene	no cas #	No data			No data	
Dimethyl sulfate	77-78-1	No data			No data	
Dimethylaniline	121-69-7	No data			No data	
Di-n-propylnitrosamine	621-64-7	No data			No data	
Diphenyl ether	101-84-8	No data			No data	
Epichlorohydrin	106-89-8	No data			No data	
Ethyl carbamate (urethane)	51-79-6	No data			No data	
Ethyl methanesulfonate	62-50-0	No data			No data	
Ethylene dibromide	106-93-4	No data			No data	
Ethylene glycol	107-21-1	No data			No data	

Table C3-1. Toxicity Reference Values (TRVs) for Plants

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997a[b] (mg/kg)	Recommended TRV[c] (mg/kg)
Ethylene glycol monobutyl ether	111-76-2	No data			No data	
Ethylene glycol monoethyl ether acetate	111-15-9	No data			No data	
Ethylene thiourea	96-45-7	No data			No data	
Furfural	98-01-1	No data			No data	
Maleic hydrazide	123-33-1	No data			No data	
Malononitrile	109-77-3	No data			No data	
Methyl styrene (mixed isomers)	25013-15-4	No data			No data	
Methylhydrazine	60-34-4	No data			No data	
N,N-Diphenylamine	122-39-4	No data			No data	
Nitric acid, propyl ester	627-13-4	No data			No data	
N-Nitrosodi-n-butylamine	924-16-3	No data			No data	
N-Nitrosomorpholine	59-89-2	No data			No data	
N-Nitroso-N,N-dimethylamine	62-75-9	No data			No data	
o-Anisidine	90-04-0	No data			No data	
Oxalic acid	144-62-7	No data			No data	
Phthalic anhydride	85-44-9	No data			No data	
p-Phthalic acid	100-21-0	No data			No data	
Pyridine	110-86-1	No data			No data	
Quinoline	91-22-5	No data			No data	
Quinone	106-51-4	No data			No data	
Safrole	94-59-7	No data			No data	
Tetrahydrofuran	109-99-9	No data			No data	
Other Heavy Semivolatile Compounds (molecular weight >200 g/mole)						
1,2,4,5-Tetrachlorobenzene	95-94-3	No data			No data	
1,3,5-Trinitrobenzene	99-35-4	No data			No data	
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	No data			No data	
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	No data			No data	
2-sec-Butyl-4,6-dinitrophenol	88-85-7	No data			No data	
3,3-Dichlorobenzidine	91-94-1	No data			No data	
3,3'-Dimethoxybenzidine	119-90-4	No data			No data	
4-Bromophenylphenyl ether	101-55-3	No data			No data	

Table C3-1. Toxicity Reference Values (TRVs) for Plants

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997a[b] (mg/kg)	Recommended TRV[c] (mg/kg)
Ammonium perfluorooctanoate	3825-26-1	No data			No data	
Azobenzene	103-33-3	No data			No data	
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	No data			No data	
Captan	133-06-2	No data			No data	
Chlorobenzilate	510-15-6	No data			No data	
Dibutylphosphate	107-66-4	No data			No data	
Dimethyl aminoazobenzene	60-11-7	No data			No data	
Hexachlorobenzene	118-74-1	No data			No data	
Hexachlorobutadiene	87-68-3	No data			No data	
Hexachlorocyclopentadiene	77-47-4	1.00E-01			1.00E+01	1.00E-01
Hexachloroethane	67-72-1	No data			No data	
Hexachlorophene	70-30-4	No data			No data	
Hexamethylene-1,5-diisocyanate	822-06-0	No data			No data	
Mirex	2385-85-5	No data			No data	
Nitrofen	1836-75-5	No data			No data	
Pentachlorobenzene	608-93-5	No data			No data	
Pentachloronitrobenzene	82-68-8	No data			No data	
Pentachlorophenol	87-86-5	1.73E+00			3.00E+00	1.73E+00
Picric acid	88-89-1	No data			No data	
Pronamide	23950-58-5	No data			No data	
Strychnine	57-24-9	No data			No data	
Terphenyls	26140-60-3	No data			No data	
Tributyl phosphate	126-73-8	No data			No data	
Trifluralin	1582-09-8	No data			No data	
Triphenylamine	603-34-9	No data			No data	
Herbicides and Organochlorinated Pesticides						
2,4,5-T	93-76-5	No data			No data	
2,4-D and esters	94-75-7	No data			No data	
4,4-DDD	72-54-8	No data			No data	
4,4-DDE	72-55-9	No data			No data	
4,4-DDT	50-29-3	No data			No data	

Table C3-1. Toxicity Reference Values (TRVs) for Plants

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997a[b] (mg/kg)	Recommended TRV[c] (mg/kg)
Aldrin	309-00-2	No data			No data	
alpha-BHC	319-84-6	No data			No data	
beta-BHC	319-85-7	No data			No data	
Chlordane	57-74-9	No data			No data	
Delta-BHC	319-86-8	No data			No data	
Dieldrin	60-57-1	No data			No data	
Endothall	145-73-3	No data			No data	
Endrin	72-20-8	No data			No data	
gamma-BHC (Lindane)	58-89-9	No data			No data	
Heptachlor	76-44-8	1.00E+00			No data	1.00E+00
Isodrin	465-73-6	No data			No data	
Methoxychlor	72-43-5	No data			No data	
Silvex (2,4,5-TP)	93-72-1	No data			No data	
Toxaphene	8001-35-2	No data			No data	
Inorganic Chemicals and Compounds						
Metals						
Aluminum	7429-90-5	5.00E+00			5.00E+01	5.00E+00
Antimony	7440-36-0	5.00E-01			5.00E+00	5.00E-01
Arsenic	7440-38-2	1.00E+00			1.00E+01	1.00E+00
Barium	7440-39-3	5.00E+00			5.00E+02	5.00E+00
Beryllium	7440-41-7	1.00E-01			1.00E+01	1.00E-01
Bismuth	7440-69-9	No data			No data	
Boron	7440-42-8	No data			5.00E-01	5.00E-01
Cadmium	7440-43-9	2.00E-01			4.00E+00	2.00E-01
Calcium	7440-70-2	No data			No data	
Chromium (and VI)	18540-29-9	1.80E-02			1.00E+00	1.80E-02
Cobalt	7440-48-4	No data			2.00E+01	2.00E+01
Copper	7440-50-8	1.00E+00			1.00E+02	1.00E+00
Iron	7439-89-6	No data			No data	
Lead	7439-92-1	4.60E+00			5.00E+01	4.60E+00

Table C3-1. Toxicity Reference Values (TRVs) for Plants

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997a[b] (mg/kg)	Recommended TRV[c] (mg/kg)
Lithium	7439-93-2	No data			2.00E+00	2.00E+00
Magnesium	7439-95-4	No data			No data	
Manganese	7439-96-5	No data			5.00E+02	5.00E+02
Mercury	7439-97-6	3.50E-01			3.00E-01	3.50E-01
Molybdenum	7439-98-7	No data			2.00E+00	2.00E+00
Nickel	7440-02-0	2.50E+01			3.00E+01	2.50E+01
Potassium	7440-09-7	No data			No data	
Rhodium	7440-16-6	No data			No data	5.00E-02
Selenium	7782-49-2	5.00E-02			1.00E+00	
Silicon	7440-21-3	No data			No data	
Silver	7440-22-4	2.00E-02			2.00E+00	2.00E-02
Sodium	7440-23-5	No data			No data	
Strontium	7440-24-6	No data			No data	
Tantalum	7440-25-7	No data			No data	
Thallium	7440-28-0	1.00E-02			1.00E+00	1.00E-02
Tin	7440-31-5	No data			5.00E+01	5.00E+01
Tungsten	7440-33-7	No data			No data	
Uranium	7440-61-1	No data			5.00E+00	5.00E+00
Vanadium	7440-62-2	No data			2.00E+00	2.00E+00
Yttrium	7440-65-5	No data			No data	
Zinc	7440-66-6	9.00E-01			5.00E+01	9.00E-01
Zirconium	7440-67-7	No data			No data	
Non-metals and Anions						
Ammonia/Ammonium	7664-41-7	No data			No data	
Bromide	24959-67-9	No data			1.00E+01	1.00E+01
Chloride	16887-00-6	No data			No data	
Cyanide	57-12-5	No data			No data	
Fluoride	16984-48-8	No data			2.00E+02	2.00E+02
Hydroxide	14280-30-9	No data			No data	
Iodine	7553-56-2	No data			4.00E+00	4.00E+00
Nitrate	14797-55-8	No data			No data	

Table C3-1. Toxicity Reference Values (TRVs) for Plants

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997a[b] (mg/kg)	Recommended TRV[c] (mg/kg)
Nitrite	14797-65-0	No data			No data	
Phosphate	14265-44-2	No data			No data	
Phosphorus	7723-14-0	No data			No data	
Sulfate	14808-79-8	No data			No data	
Total Sulfur	63705-05-5	No data			No data	
Priority Pollutants						
Carbon Dioxide	124-38-9	No data			No data	
Nitrogen Dioxide	10102-44-0	No data			No data	
Ozone	10028-15-6	No data			No data	
Particulate Matter	No CAS #	No data			No data	
Sulfur Dioxode	7446-09-5	No data			No data	
Radionuclides[e]						
Actinium-227	14952-40-0	NA			NA	
Americium-241	1596-10-2	NA			NA	
Americium-243	14993-75-0	NA			NA	
Antimony-125	14234-35-6	NA			NA	
Barium-137	13981-97-0	NA			NA	
Cadmium-113	None	NA			NA	
Carbon-14	14762-75-5	NA			NA	
Cesium-134	13967-70-9	NA			NA	
Cesium-137	10045-97-3	NA			NA	
Cobalt-60	10198-40-0	NA			NA	
Curium-242	15510-73-3	NA			NA	
Curium-243	15757-87-6	NA			NA	
Curium-244	13981-15-2	NA			NA	
Europium-152	14683-23-9	NA			NA	
Europium-154	15585-10-1	NA			NA	
Europium-155	14391-16-3	NA			NA	
Iodine-129	15046-84-1	NA			NA	
Neptunium-237	13994-20-2	NA			NA	

Table C3-1. Toxicity Reference Values (TRVs) for Plants

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997a[b] (mg/kg)	Recommended TRV[c] (mg/kg)
Nickel-59	14336-70-0	NA			NA	
Nickel-63	13981-37-8	NA			NA	
Niobium-93	7440-03-1	NA			NA	
Plutonium-238	13981-16-3	NA			NA	
Plutonium-239	15117-48-3	NA			NA	
Plutonium-240	14119-33-6	NA			NA	
Plutonium-241	14119-32-5	NA			NA	
Plutonium-242	13982-10-0	NA			NA	
Protactinium-231	14331-85-2	NA			NA	
Radium-226	13982-63-3	NA			NA	
Radium-228	15262-20-1	NA			NA	
Ruthenium-106	13967-48-1	NA			NA	
Samarium-151	15715-94-3	NA			NA	
Selenium-79	None	NA			NA	
Strontium-90	10098-97-2	NA			NA	
Technetium-99	14133-79-7	NA			NA	
Thorium-229	15594-54-4	NA			NA	
Thorium-232	7440-29-1	NA			NA	
Tin-126	15832-50-5	NA			NA	
Tritium	10028-17-8	NA			NA	
Uranium-232	14158-29-3	NA			NA	
Uranium-233	13968-55-3	NA			NA	
Uranium-234	13966-29-5	NA			NA	
Uranium-235	15117-96-1	NA			NA	
Uranium-236	13982-70-2	NA			NA	
Uranium-238	7440-61-1	NA			NA	
Yttrium-90	10098-91-6	NA			NA	
Zirconium-93	15751-77-6	NA			NA	

NA = Not applicable

[a] Published in Appendix E of EPA (1999), Table E-5

Table C3-1. Toxicity Reference Values (TRVs) for Plants

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997a[b] (mg/kg)	Recommended TRV[c] (mg/kg)

[b] Published in Table 1 of Efroymson et al. 1997a

[c] Selection criteria: EPA (1999) values are preferred over Efroymson et al. (1997a) values

[d] Benzo(a)pyrene value from Appendix E of EPA (1999), Table E-5, was used as a surrogate value

[e] TRV is not applicable to single radionuclides. Combined external and internal radiation exposure for plants from all radionuclides combined cannot exceed 1 rad/d.

Table C3-2 Toxicity Reference Values (TRVs) for Soil Invertebrates

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997b[b] (mg/kg)	Dutch Soil Cleanup Interim Act[c] (mg/kg)	Notes	Recommended TRV[d] (mg/kg)
Organic Compounds								
Aromatic Halogenated Hydrocarbons								
4-Chloro-3-methylphenol	59-50-7	No data			No data	5.00E-01	e	5.00E-01
2,3,4,6-Tetrachlorophenol	58-90-2	No data			No data	5.00E-01	e	5.00E-01
Aromatic Nonhalogenated Hydrocarbons								
2-Nitrotoluene	88-72-2	No data			No data	7.00E+00	f	7.00E+00
4-Nitrobiphenyl	92-93-3	No data			No data	No data		
Benzaldehyde	100-52-7	No data			No data	7.00E+00	f	7.00E+00
Benzene	71-43-2	No data			No data	5.00E-01		5.00E-01
Benzyl alcohol	100-51-6	No data			No data	7.00E+00	f	7.00E+00
Ethyl benzene	100-41-4	No data			No data	5.00E+00		5.00E+00
m-Xylene	108-38-3	No data			No data	5.00E+00	g	5.00E+00
o-Xylene	95-47-6	No data			No data	5.00E+00	g	5.00E+00
p-Xylene	106-42-3	No data			No data	5.00E+00	g	5.00E+00
Styrene	100-42-5	No data			No data	5.00E+00		5.00E+00
Toluene	108-88-3	No data			No data	3.00E+00		3.00E+00
Non-aromatic Nonhalogenated Hydrocarbons								
1,2-Epoxybutane	106-88-7	No data			No data	No data		
1,3-Butadiene	106-99-0	No data			No data	No data		
1,4-Dioxane	123-91-1	No data			No data	No data		
1-Methylpropyl alcohol	78-92-2	No data			No data	No data		
1-Nitropropane	108-03-2	No data			No data	No data		
2,2,4-Trimethylpentane	540-84-1	No data			No data	No data		
2-Butanone	78-93-3	No data			No data	No data		
2-Butenaldehyde (2-Butenal)	4170-30-3	No data			No data	No data		
2-Ethoxyethanol	110-80-5	No data			No data	No data		
2-Heptanone	110-43-0	No data			No data	No data		
2-Hexanone	591-78-6	No data			No data	No data		
2-Methoxyethanol	109-86-4	No data			No data	No data		
2-Methyl-2-propanol	75-65-0	No data			No data	No data		

Table C3-2 Toxicity Reference Values (TRVs) for Soil Invertebrates

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997b[b] (mg/kg)	Dutch Soil Cleanup Interim Act[c] (mg/kg)	Notes	Recommended TRV[d] (mg/kg)
2-Methyl-2-propenenitrile	126-98-7	No data			No data	No data		
2-Methylaziridine	75-55-8	No data			No data	No data		
2-Methylpropyl alcohol	78-83-1	No data			No data	No data		
2-Pentanone	107-87-9	No data			No data	No data		
2-Propanone (Acetone)	67-64-1	No data			No data	No data		
2-Propene-1-ol	107-18-6	No data			No data	No data		
2-Propyl alcohol	67-63-0	No data			No data	No data		
3-Heptanone	106-35-4	No data			No data	No data		
3-Methyl-1-butanol	123-51-3	No data			No data	No data		
3-Methyl-2-butanone	563-80-4	No data			No data	No data		
3-Pentanone	96-22-0	No data			No data	No data		
4-Heptanone	123-19-3	No data			No data	No data		
4-Methyl-2-pentanone	108-10-1	No data			No data	No data		
4-Methyl-3-penten-2-one	141-79-7	No data			No data	No data		
5-Methyl-2-hexanone	110-12-3	No data			No data	No data		
Acetaldehyde	75-07-0	No data			No data	No data		
Acetamide	60-35-5	No data			No data	No data		
Acetic acid	64-19-7	No data			No data	No data		
Acetic acid ethyl ester	141-78-6	No data			No data	No data		
Acetic acid n-butyl ester	123-86-4	No data			No data	No data		
Acetonitrile	75-05-8	No data			No data	No data		
Acrolein	107-02-8	No data			No data	No data		
Acrylonitrile	107-13-1	No data			No data	No data		
Bis(isopropyl)ether	108-20-3	No data			No data	No data		
Butane	106-97-8	No data			No data	No data		
Carbon disulfide	75-15-0	No data			No data	No data		
Cyanogen	460-19-5	No data			No data	No data		
Cyclohexane	110-82-7	No data			No data	6.00E+00		6.00E+00
Cyclohexanone	108-94-1	No data			No data	No data		
Cyclohexene	110-83-8	No data			No data	No data		
Cyclopentane	287-92-3	No data			No data	No data		

Table C3-2 Toxicity Reference Values (TRVs) for Soil Invertebrates

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997b[b] (mg/kg)	Dutch Soil Cleanup Interim Act[c] (mg/kg)	Notes	Recommended TRV[d] (mg/kg)
Ethyl alcohol	64-17-5	No data			No data	No data		
Ethyl ether	60-29-7	No data			No data	No data		
Ethyl methacrylate	97-63-2	No data			No data	No data		
Formaldehyde	50-00-0	No data			No data	No data		
Formamide	75-12-7	No data			No data	No data		
Formic acid	64-18-6	No data			No data	No data		
Formic acid, methyl ester	107-31-3	No data			No data	No data		
Glycidylaldehyde	765-34-4	No data			No data	No data		
Methyl acetate	79-20-9	No data			No data	No data		
Methyl alcohol	67-56-1	No data			No data	No data		
Methyl isocyanate	624-83-9	No data			No data	No data		
Methyl methacrylate	80-62-6	No data			No data	No data		
Methyl tert-butyl ether	1634-04-4	No data			No data	No data		
Methylacetylene	74-99-7	No data			No data	No data		
Methylcyclohexane	108-87-2	No data			No data	No data		
N,N-Dimethylacetamide	127-19-5	No data			No data	No data		
n-Butyl alcohol	71-36-3	No data			No data	No data		
n-Heptane	142-82-5	No data			No data	No data		
n-Hexane	110-54-3	No data			No data	No data		
Nitromethane	75-52-5	No data			No data	No data		
n-Nonane	111-84-2	No data			No data	No data		
n-Octane	111-65-9	No data			No data	No data		
n-Pentane	109-66-0	No data			No data	No data		
n-Propionaldehyde	123-38-6	No data			No data	No data		
n-Propyl alcohol	71-23-8	No data			No data	No data		
n-Valeraldehyde	110-62-3	No data			No data	No data		
Oxirane	75-21-8	No data			No data	No data		
p-Cymene	99-87-6	No data			No data	7.00E+00	f	7.00E+00
Phosgene	75-44-5	No data			No data	No data		
Propargyl alcohol	107-19-7	No data			No data	No data		
Propionic acid	79-09-4	No data			No data	No data		

Table C3-2 Toxicity Reference Values (TRVs) for Soil Invertebrates

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997b[b] (mg/kg)	Dutch Soil Cleanup Interim Act[c] (mg/kg)	Notes	Recommended TRV[d] (mg/kg)
Propionitrile	107-12-0	No data			No data	No data		
Propylene gylcol monomethyl ether	107-98-2	No data			No data	No data		
p-tert-Butyltoluene	98-51-1	No data			No data	7.00E+00	f	7.00E+00
Triethylamine	121-44-8	No data			No data	No data		
Trimethylamine	75-50-3	No data			No data	No data		
Vinyl acetate	108-05-4	No data			No data	No data		
Non-aromatic Halogenated Hydrocarbons								
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	No data			No data	5.00E+00	h	5.00E+00
1,1,1,2-Tetrachloroethane	630-20-6	No data			No data	5.00E+00	h	5.00E+00
1,1,1-Trichloroethane	71-55-6	No data			No data	5.00E+00	h	5.00E+00
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	No data			No data	5.00E+00	h	5.00E+00
1,1,2,2-Tetrachloroethane	79-34-5	No data			No data	5.00E+00	h	5.00E+00
1,1,2,2-Tetrachloroethene	127-18-4	No data			No data	5.00E+00	h	5.00E+00
1,1,2-Trichloroethane	79-00-5	No data			No data	5.00E+00	h	5.00E+00
1,1,2-Trichloroethylene	79-01-6	No data			No data	5.00E+00	h	5.00E+00
1,1-Dichloroethane	75-34-3	No data			No data	5.00E+00	h	5.00E+00
1,1-Dichloroethene	75-35-4	No data			No data	5.00E+00	h	5.00E+00
1,2,2-Trichloro-1,1,2-trifluoroethane	76-13-1	No data			No data	5.00E+00	h	5.00E+00
1,2,3-Trichloropropane	96-18-4	No data			No data	5.00E+00	h	5.00E+00
1,2-Dibromo-3-chloropropane	96-12-8	No data			No data	5.00E+00	h	5.00E+00
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	No data			No data	5.00E+00	h	5.00E+00
1,2-Dichloroethane	107-06-2	No data			No data	5.00E+00	h	5.00E+00
1,2-Dichloroethylene	540-59-0	No data			No data	5.00E+00	h	5.00E+00
1,2-Dichloropropane	78-87-5	No data			7.00E+02	5.00E+00	h	7.00E+02
1,3-Dichloropropene	542-75-6	No data			No data	5.00E+00	h	5.00E+00
1,4-Dichloro-2-butene	764-41-0	No data			No data	5.00E+00	h	5.00E+00
1-Chloroethene	75-01-4	No data			No data	5.00E+00	h	5.00E+00
2,2-Dichloropropionic acid	75-99-0	No data			No data	5.00E+00	h	5.00E+00
2-Chloropropane	75-29-6	No data			No data	5.00E+00	h	5.00E+00
3-Chloropropene (allyl chloride)	107-05-1	No data			No data	5.00E+00	h	5.00E+00
Bromochloromethane	74-97-5	No data			No data	5.00E+00	h	5.00E+00

Table C3-2 Toxicity Reference Values (TRVs) for Soil Invertebrates

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997b[b] (mg/kg)	Dutch Soil Cleanup Interim Act[c] (mg/kg)	Notes	Recommended TRV[d] (mg/kg)
Bromodichloromethane	75-27-4	No data			No data	5.00E+00	h	5.00E+00
Bromoethene	593-60-2	No data			No data	No		
Bromoform	75-25-2	No data			No data	No		
Bromomethane	74-83-9	No data			No data	No		
Carbon tetrachloride	56-23-5	No data			No data	5.00E+00	h	5.00E+00
Chlorodibromomethane	124-48-1	No data			No data	5.00E+00	h	5.00E+00
Chlorodifluoromethane	75-45-6	No data			No data	5.00E+00	h	5.00E+00
Chloroethane	75-00-3	No data			No data	5.00E+00	h	5.00E+00
Chloroform	67-66-3	No data			No data	5.00E+00	h	5.00E+00
Chloromethane	74-87-3	No data			No data	5.00E+00	h	5.00E+00
Chloropentafluoroethane	76-15-3	No data			No data	5.00E+00	h	5.00E+00
cis-1,2-Dichloroethene	156-59-2	No data			No data	5.00E+00	h	5.00E+00
cis-1,3-Dichloropropene	10061-01-5	No data			No data	5.00E+00	h	5.00E+00
Cyanogen bromide	506-68-3	No data			No data	No		
Cyanogen chloride	506-77-4	No data			No data	5.00E+00	h	5.00E+00
Dichlorodifluoromethane	75-71-8	No data			No data	5.00E+00	h	5.00E+00
Dichlorofluoromethane	75-43-4	No data			No data	5.00E+00	h	5.00E+00
Dichloromethane	75-09-2	No data			No data	5.00E+00	h	5.00E+00
Difluorodibromomethane	75-61-6	No data			No data	No		
Hexafluoroacetone	684-16-2	No data			No data	No		
Iodomethane	74-88-4	No data			No data	No		
Methylene bromide	74-95-3	No data			No data	No		
Pentachloroethane	76-01-7	No data			No data	5.00E+00	h	5.00E+00
trans-1,2-Dichloroethene	156-60-5	No data			No data	5.00E+00	h	5.00E+00
trans-1,3-Dichloropropene	10061-02-6	No data			No data	5.00E+00	h	5.00E+00
Trichloroacetic acid	76-03-9	No data			No data	5.00E+00	h	5.00E+00
Trichlorofluoroethane	27154-33-2	No data			No data	5.00E+00	h	5.00E+00
Trichlorofluoromethane	75-69-4	No data			No data	5.00E+00	h	5.00E+00
Trifluorobromomethane	75-63-8	No data			No data	No		
Dioxin and Furan Compounds								
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	35822-46-9	No data			No data	8.00E+00	i	8.00E+00

Table C3-2 Toxicity Reference Values (TRVs) for Soil Invertebrates

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997b[b] (mg/kg)	Dutch Soil Cleanup Interim Act[c] (mg/kg)	Notes	Recommended TRV[d] (mg/kg)
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	No data			No data	8.00E+00	i	8.00E+00
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	No data			No data	8.00E+00	i	8.00E+00
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	39227-28-6	No data			No data	8.00E+00	i	8.00E+00
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	No data			No data	8.00E+00	i	8.00E+00
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	57653-85-7	No data			No data	8.00E+00	i	8.00E+00
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	No data			No data	8.00E+00	i	8.00E+00
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	19408-74-3	No data			No data	8.00E+00	i	8.00E+00
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	No data			No data	8.00E+00	i	8.00E+00
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	40321-76-4	No data			No data	8.00E+00	i	8.00E+00
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	No data			No data	8.00E+00	i	8.00E+00
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	No data			No data	8.00E+00	i	8.00E+00
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	No data			No data	8.00E+00	i	8.00E+00
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1746-01-6	5.00E-01			No data	8.00E+00	i	5.00E-01
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	No data			No data	8.00E+00	i	8.00E+00
Dibenzofuran	132-64-9	No data			No data	8.00E+00	i	8.00E+00
Octachlorodibenzo(p)dioxin	3268-87-9	No data			No data	8.00E+00	i	8.00E+00
Octachlorodibenzofuran	39001-02-0	No data			No data	8.00E+00	i	8.00E+00
PCBs								
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	No data			No data	1.00E+00	j	1.00E+00
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	No data			No data	1.00E+00	j	1.00E+00
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	No data			No data	1.00E+00	j	1.00E+00
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	No data			No data	1.00E+00	j	1.00E+00
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	No data			No data	1.00E+00	j	1.00E+00
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	No data			No data	1.00E+00	j	1.00E+00
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	No data			No data	1.00E+00	j	1.00E+00
2',3,4,4',5-Pentachlorobiphenyl	no cas #	No data			No data	1.00E+00	j	1.00E+00
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	No data			No data	1.00E+00	j	1.00E+00
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	No data			No data	1.00E+00	j	1.00E+00
3,3',4,4',5-Pentachlorobiphenyl	no cas #	No data			No data	1.00E+00	j	1.00E+00
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	No data			No data	1.00E+00	j	1.00E+00
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	No data			No data	1.00E+00	j	1.00E+00

Table C3-2 Toxicity Reference Values (TRVs) for Soil Invertebrates

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997b[b] (mg/kg)	Dutch Soil Cleanup Interim Act[c] (mg/kg)	Notes	Recommended TRV[d] (mg/kg)
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	No data			No data	1.00E+00	j	1.00E+00
Polychlorinated biphenyls (PCBs)[k]	1336-36-3	2.51E+00			No data	1.00E+00	j	2.51E+00
Phthalates								
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	No data			No data	7.00E+00	f	7.00E+00
Butylbenzyl phthalate	85-68-7	No data			No data	7.00E+00	f	7.00E+00
Dibutyl phthalate	84-74-2	No data			No data	7.00E+00	f	7.00E+00
Diethyl phthalate	84-66-2	No data			No data	7.00E+00	f	7.00E+00
Dimethylphthalate	131-11-3	No data			2.00E+02	7.00E+00	f	2.00E+02
n-Dioctyl phthalate	117-84-0	No data			No data	7.00E+00	f	7.00E+00
Light Polycyclic Aromatic Hydrocarbons (molecular weight <200 g/mole)								
2-Chloronapthalene	91-58-7	No data			No data	1.00E+00	l	1.00E+00
2-Methyl naphthalene	91-57-6	No data			No data	2.00E+01	m	2.00E+01
5-Nitroacenaphthene	602-87-9	No data			No data	2.00E+01	m	2.00E+01
Acenaphthene	83-32-9	No data			No data	2.00E+01	m	2.00E+01
Acenaphthylene	208-96-8	No data			No data	2.00E+01	m	2.00E+01
Anthracene	120-12-7	No data			No data	1.00E+01		1.00E+01
Fluorene	86-73-7	No data			3.00E+01	2.00E+01	m	3.00E+01
Indene	95-13-6	No data			No data	2.00E+01	m	2.00E+01
Naphthalene	91-20-3	No data			No data	5.00E+00		5.00E+00
Phenanthrene	85-01-8	No data			No data	5.00E+00		5.00E+00
Pyrene	129-00-0	No data			No data	1.00E+01		1.00E+01
Heavy Polycyclic Aromatic Hydrocarbons (molecular weight >200 g/mole)								
3-Methylcholanthrene	56-49-5	2.50E+01	n		No data	2.00E+01	m	2.50E+01
5-Methylchrysene	3697-24-3	2.50E+01	n		No data	2.00E+01	m	2.50E+01
Benzo(a)anthracene	56-55-3	2.50E+01	n		No data	2.00E+01	m	2.50E+01
Benzo(a)pyrene	50-32-8	2.50E+01			No data	1.00E+00		2.50E+01
Benzo(b)fluoranthene	205-99-2	2.50E+01	n		No data	2.00E+01	m	2.50E+01
Benzo(e)pyrene	192-97-2	2.50E+01	n		No data	2.00E+01	m	2.50E+01
Benzo(g,h,i)perylene	191-24-2	2.50E+01	n		No data	2.00E+01	m	2.50E+01
Benzo(j)fluoranthene	205-82-3	2.50E+01	n		No data	2.00E+01	m	2.50E+01

Table C3-2 Toxicity Reference Values (TRVs) for Soil Invertebrates

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997b[b] (mg/kg)	Dutch Soil Cleanup Interim Act[c] (mg/kg)	Notes	Recommended TRV[d] (mg/kg)
Benzo(k)fluoranthene	207-08-9	2.50E+01	n		No data	2.00E+01	m	2.50E+01
Benzo[a,i]pyrene	191-30-0	2.50E+01	n		No data	2.00E+01	m	2.50E+01
Chrysene	218-01-9	2.50E+01	n		No data	2.00E+01	m	2.50E+01
Dibenz(a,h)anthracene	53-70-3	2.50E+01	n		No data	2.00E+01	m	2.50E+01
Dibenz[a,h]acridine	226-36-8	2.50E+01	n		No data	2.00E+01	m	2.50E+01
Dibenz[a,j]acridine	224-42-0	2.50E+01	n		No data	2.00E+01	m	2.50E+01
Dibenzo(a,e)fluoranthene	5385-75-1	2.50E+01	n		No data	2.00E+01	m	2.50E+01
Dibenzo(a,h)fluoranthene	no cas #	2.50E+01	n		No data	2.00E+01	m	2.50E+01
Dibenzo[a,e]pyrene	192-65-4	2.50E+01	n		No data	2.00E+01	m	2.50E+01
Dibenzo[a,h]pyrene	189-64-0	2.50E+01	n		No data	2.00E+01	m	2.50E+01
Dibenzo[a,i]pyrene	189-55-9	2.50E+01	n		No data	2.00E+01	m	2.50E+01
Fluoranthene	206-44-0	2.50E+01	n		No data	1.00E+01		2.50E+01
Hexachloronaphthalene	1335-87-1	2.50E+01	n		No data	1.00E+00	l	2.50E+01
Indeno(1,2,3-cd)pyrene	193-39-5	2.50E+01	n		No data	2.00E+01	m	2.50E+01
Octachloronaphthalene	2234-13-1	2.50E+01	n		No data	1.00E+00	l	2.50E+01
Pentachloronaphthalene	1321-64-8	2.50E+01	n		No data	1.00E+00	l	2.50E+01
Tetrachloronaphthalene	1335-88-2	2.50E+01	n		No data	1.00E+00	l	2.50E+01
Trichloronaphthalene	1321-65-9	2.50E+01	n		No data	1.00E+00	l	2.50E+01
Light Substituted Benzene Compounds (MW <200 g/mole)								
1,2,3-Trichlorobenzene	87-61-6	No data			2.00E+01	1.00E+00	o	2.00E+01
1,2,4-Trichlorobenzene	120-82-1	No data			2.00E+01	1.00E+00	o	2.00E+01
1,2,4-Trimethyl benzene	95-63-6	No data			No data	7.00E+00	f	7.00E+00
1,2-Dichlorobenzene	95-50-1	No data			No data	1.00E+00	o	1.00E+00
1,3,5-Trimethyl benzene	108-67-8	No data			No data	7.00E+00	f	7.00E+00
1,3-Dichlorobenzene	541-73-1	No data			No data	1.00E+00	o	1.00E+00
1,3-Dinitrobenzene	99-65-0	2.26E+00	p		No data	7.00E+00	f	2.26E+00
1,4-Dichlorobenzene	106-46-7	No data			2.00E+01	1.00E+00	o	2.00E+01
1,4-Dinitrobenzene	100-25-4	No data			No data	7.00E+00	f	7.00E+00
2,4,5-Trichlorophenol	95-95-4	No data			9.00E+00	5.00E-01	e	9.00E+00
2,4,6-Trichlorophenol	88-06-2	No data			1.00E+01	5.00E-01	e	1.00E+01
2,4-Dichlorophenol	120-83-2	No data			No data	5.00E-01	e	5.00E-01

Table C3-2 Toxicity Reference Values (TRVs) for Soil Invertebrates

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997b[b] (mg/kg)	Dutch Soil Cleanup Interim Act[c] (mg/kg)	Notes	Recommended TRV[d] (mg/kg)
2,4-Dimethylphenol	105-67-9	No data			No data	1.00E+00	q	1.00E+00
2,4-Dinitrophenol	51-28-5	No data			No data	1.00E+00	q	1.00E+00
2,4-Dinitrotoluene	121-14-2	No data			No data	7.00E+00	f	7.00E+00
2,6-Dinitrotoluene	606-20-2	No data			No data	7.00E+00	f	7.00E+00
2-Chlorophenol	95-57-8	No data			No data	5.00E-01	e	5.00E-01
2-Chlorotoluene	95-49-8	No data			No data	7.00E+00	f	7.00E+00
2-Nitrophenol	88-75-5	No data			No data	1.00E+00	q	1.00E+00
4,6-Dinitro-o-cresol	534-52-1	No data			No data	1.00E+00	q	1.00E+00
4-Chlorotoluene	106-43-4	No data			No data	7.00E+00	f	7.00E+00
4-Nitrophenol	100-02-7	No data			7.00E+00	1.00E+00	q	7.00E+00
alpha-Methylstyrene	98-83-9	No data			No data	7.00E+00	f	7.00E+00
Aniline	62-53-3	No data			No data	7.00E+00	f	7.00E+00
Benzotrichloride	98-07-7	No data			No data	1.00E+00	o	1.00E+00
Benzyl chloride	100-44-7	No data			No data	8.00E+00	i	8.00E+00
Bromobenzene	108-86-1	No data			No data	8.00E+00	i	8.00E+00
Chlorobenzene	108-90-7	No data			4.00E+01	1.00E+00		4.00E+01
Cumene	98-82-8	No data			No data	7.00E+00	f	7.00E+00
m-Cresol	108-39-4	No data			No data	1.00E+00	q	1.00E+00
n-Butyl benzene	104-51-8	No data			No data	7.00E+00	f	7.00E+00
Nitrobenzene	98-95-3	2.26E+00			4.00E+01	7.00E+00	f	2.26E+00
n-Propyl benzene	103-65-1	No data			No data	7.00E+00	f	7.00E+00
o-Cresol	95-48-7	No data			No data	1.00E+00	r	1.00E+00
o-Dinitrobenzene	528-29-0	No data			No data	7.00E+00	f	7.00E+00
o-Nitroaniline	88-74-4	No data			No data	7.00E+00	f	7.00E+00
o-Toluidine	95-53-4	No data			No data	7.00E+00	f	7.00E+00
p-Chloroaniline	106-47-8	No data			No data	8.00E+00	i	8.00E+00
p-Cresol	106-44-5	No data			No data	1.00E+00	q	1.00E+00
Phenol	108-95-2	No data			3.00E+01	1.00E+00	q	3.00E+01
p-Nitrochlorobenzene	100-00-5	No data			No data	1.00E+00	o	1.00E+00
p-Toluidine	106-49-0	No data			No data	7.00E+00	f	7.00E+00
sec-Butyl benzene	135-98-8	No data			No data	7.00E+00	f	7.00E+00

Table C3-2 Toxicity Reference Values (TRVs) for Soil Invertebrates

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997b[b] (mg/kg)	Dutch Soil Cleanup Interim Act[c] (mg/kg)	Notes	Recommended TRV[d] (mg/kg)
tert-Butyl benzene	98-06-6	No data			No data	7.00E+00	f	7.00E+00
Toluene-2,6-diamine	823-40-5	No data			No data	7.00E+00	f	7.00E+00
Trimethyl benzene	25551-13-7	No data			No data	7.00E+00	f	7.00E+00
Other Light Semivolatile Compounds (molecular weight <200 g/mole)								
1,1'-Biphenyl	92-52-4	No data			No data	No data		
1,1-Dimethylhydrazine	57-14-7	No data			No data	No data		
1,2-Dimethylhydrazine	540-73-8	No data			No data	No data		
1,2-Diphenylhydrazine	122-66-7	No data			No data	No data		
1,3-Propane sultone	1120-71-4	No data			No data	No data		
2,4-Toluene diisocyante	584-84-9	No data			No data	No data		
2-Chloroacetophenone	532-27-4	No data			No data	8.00E+00	i	8.00E+00
2-Propenoic acid	79-10-7	No data			No data	No data		
4,4-Methylenedianiline	101-77-9	No data			No data	No data		
Acetophenone	98-86-2	No data			No data	No data		
Benzoic acid	65-85-0	No data			No data	No data		
bis(2-Chloroethoxy)methane	111-91-1	No data			No data	5.00E+00	h	5.00E+00
bis(2-Chloroethyl) ether	111-44-4	No data			No data	5.00E+00	h	5.00E+00
Chlorocyclopentadiene	41851-50-7	No data			No data	5.00E+00	h	5.00E+00
Cyclohexanol	108-93-0	No data			No data	No data		
Dichloroisopropyl ether	108-60-1	No data			No data	5.00E+00	h	5.00E+00
Dichloromethyl ether	542-88-1	No data			No data	5.00E+00	h	5.00E+00
Dichloropentadiene	no cas #	No data			No data	5.00E+00	h	5.00E+00
Dimethyl sulfate	77-78-1	No data			No data	No data		
Dimethylaniline	121-69-7	No data			No data	No data		
Di-n-propylnitrosamine	621-64-7	No data			No data	No data		
Diphenyl ether	101-84-8	No data			No data	No data		
Epichlorohydrin	106-89-8	No data			No data	5.00E+00	h	5.00E+00
Ethyl carbamate (urethane)	51-79-6	No data			No data	No data		
Ethyl methanesulfonate	62-50-0	No data			No data	No data		
Ethylene dibromide	106-93-4	No data			No data	No data		
Ethylene glycol	107-21-1	No data			No data	No data		

Table C3-2 Toxicity Reference Values (TRVs) for Soil Invertebrates

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997b[b] (mg/kg)	Dutch Soil Cleanup Interim Act[c] (mg/kg)	Notes	Recommended TRV[d] (mg/kg)
Ethylene glycol monobutyl ether	111-76-2	No data			No data	No data		
Ethylene glycol monoethyl ether acetate	111-15-9	No data			No data	No data		
Ethylene thiourea	96-45-7	No data			No data	No data		
Furfural	98-01-1	No data			No data	No data		
Maleic hydrazide	123-33-1	No data			No data	No data		
Malononitrile	109-77-3	No data			No data	No data		
Methyl styrene (mixed isomers)	25013-15-4	No data			No data	7.00E+00	f	7.00E+00
Methylhydrazine	60-34-4	No data			No data	No data		
N,N-Diphenylamine	122-39-4	No data			No data	No data		
Nitric acid, propyl ester	627-13-4	No data			No data	No data		
N-Nitrosodi-n-butylamine	924-16-3	No data			No data	No data		
N-Nitrosomorpholine	59-89-2	No data			No data	No data		
N-Nitroso-N,N-dimethylamine	62-75-9	No data			No data	No data		
o-Anisidine	90-04-0	No data			No data	No data		
Oxalic acid	144-62-7	No data			No data	No data		
Phthalic anhydride	85-44-9	No data			No data	No data		
p-Phthalic acid	100-21-0	No data			No data	No data		
Pyridine	110-86-1	No data			No data	2.00E+00		2.00E+00
Quinoline	91-22-5	No data			No data	No data		
Quinone	106-51-4	No data			No data	No data		
Safrole	94-59-7	No data			No data	No data		
Tetrahydrofuran	109-99-9	No data			No data	4.00E+00		4.00E+00
Other Heavy Semivolatile Compounds (molecular weight >200 g/mole)								
1,2,4,5-Tetrachlorobenzene	95-94-3	No data			No data	No data		
1,3,5-Trinitrobenzene	99-35-4	No data			No data	No data		
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	No data			No data	1.00E+00	q	1.00E+00
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	No data			No data	1.00E+00	q	1.00E+00
2-sec-Butyl-4,6-dinitrophenol	88-85-7	No data			No data	1.00E+00	q	1.00E+00
3,3-Dichlorobenzidine	91-94-1	No data			No data	No data		
3,3'-Dimethoxybenzidine	119-90-4	No data			No data	No data		
4-Bromophenylphenyl ether	101-55-3	No data			No data	No data		

Table C3-2 Toxicity Reference Values (TRVs) for Soil Invertebrates

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997b[b] (mg/kg)	Dutch Soil Cleanup Interim Act[c] (mg/kg)	Notes	Recommended TRV[d] (mg/kg)
Ammonium perfluorooctanoate	3825-26-1	No data			No data	No data		
Azobenzene	103-33-3	No data			No data	No data		
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	No data			No data	No data		
Captan	133-06-2	No data			No data	5.00E-01	r	5.00E-01
Chlorobenzilate	510-15-6	No data			No data	No data		
Dibutylphosphate	107-66-4	No data			No data	No data		
Dimethyl aminoazobenzene	60-11-7	No data			No data	No data		
Hexachlorobenzene	118-74-1	No data			No data	No data		
Hexachlorobutadiene	87-68-3	No data			No data	No data		
Hexachlorocyclopentadiene	77-47-4	No data			No data	No data		
Hexachloroethane	67-72-1	No data			No data	No data		
Hexachlorophene	70-30-4	No data			No data	No data		
Hexamethylene-1,5-diisocyanate	822-06-0	No data			No data	No data		
Mirex	2385-85-5	No data			No data	5.00E-01	r	5.00E-01
Nitrofen	1836-75-5	No data			No data	5.00E-01	r	5.00E-01
Pentachlorobenzene	608-93-5	1.15E+00			2.00E+01	No data		1.15E+00
Pentachloronitrobenzene	82-68-8	No data			No data	No data		
Pentachlorophenol	87-86-5	1.00E+01			6.00E+00	5.00E-01	e	1.00E+01
Picric acid	88-89-1	No data			No data	No data		
Pronamide	23950-58-5	No data			No data	5.00E-01	r	5.00E-01
Strychnine	57-24-9	No data			No data	No data		
Terphenyls	26140-60-3	No data			No data	No data		
Tributyl phosphate	126-73-8	No data			No data	No data		
Trifluralin	1582-09-8	No data			No data	No data		
Triphenylamine	603-34-9	No data			No data	No data		
Herbicides and Organochlorinated Pesticides								
2,4,5-T	93-76-5	No data			No data	5.00E-01	r	5.00E-01
2,4-D and esters	94-75-7	No data			No data	5.00E-01	r	5.00E-01
4,4-DDD	72-54-8	No data			No data	5.00E-01	r	5.00E-01
4,4-DDE	72-55-9	No data			No data	5.00E-01	r	5.00E-01
4,4-DDT	50-29-3	No data			No data	5.00E-01	r	5.00E-01

Table C3-2 Toxicity Reference Values (TRVs) for Soil Invertebrates

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997b[b] (mg/kg)	Dutch Soil Cleanup Interim Act[c] (mg/kg)	Notes	Recommended TRV[d] (mg/kg)
Aldrin	309-00-2	No data			No data	5.00E-01	r	5.00E-01
alpha-BHC	319-84-6	No data			No data	5.00E-01	r	5.00E-01
beta-BHC	319-85-7	No data			No data	5.00E-01	r	5.00E-01
Chlordane	57-74-9	No data			No data	5.00E-01	r	5.00E-01
Delta-BHC	319-86-8	No data			No data	5.00E-01	r	5.00E-01
Dieldrin	60-57-1	No data			No data	5.00E-01	r	5.00E-01
Endothall	145-73-3	No data			No data	No		
Endrin	72-20-8	No data			No data	5.00E-01	r	5.00E-01
gamma-BHC (Lindane)	58-89-9	No data			No data	5.00E-01	r	5.00E-01
Heptachlor	76-44-8	No data			No data	5.00E-01	r	5.00E-01
Isodrin	465-73-6	No data			No data	5.00E-01	r	5.00E-01
Methoxychlor	72-43-5	No data			No data	5.00E-01	r	5.00E-01
Silvex (2,4,5-TP)	93-72-1	No data			No data	5.00E-01	r	5.00E-01
Toxaphene	8001-35-2	No data			No data	5.00E-01	r	5.00E-01
Inorganic Chemicals and Compounds								
Metals								
Aluminum	7429-90-5	No data			No data	No data		
Antimony	7440-36-0	No data			No data	No data		
Arsenic	7440-38-2	2.50E-01			6.00E+01	3.00E+01		2.50E-01
Barium	7440-39-3	No data			No data	4.00E+02		4.00E+02
Beryllium	7440-41-7	No data			No data	No data		
Bismuth	7440-69-9	No data			No data	No data		
Boron	7440-42-8	No data			No data	No data		
Cadmium	7440-43-9	1.00E+01			2.00E+01	5.00E+00		1.00E+01
Calcium	7440-70-2	No data			No data	No data		
Chromium (and VI)	18540-29-9	2.00E-01			4.00E-01	2.50E+02		2.00E-01
Cobalt	7440-48-4	No data			No data	5.00E+01		5.00E+01
Copper	7440-50-8	3.20E+01			5.00E+01	1.00E+02		3.20E+01
Iron	7439-89-6	No data			No data	No data		
Lead	7439-92-1	1.00E+02			5.00E+02	1.50E+02		1.00E+02

Table C3-2 Toxicity Reference Values (TRVs) for Soil Invertebrates

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997b[b] (mg/kg)	Dutch Soil Cleanup Interim Act[c] (mg/kg)	Notes	Recommended TRV[d] (mg/kg)
Lithium	7439-93-2	No data			No data	No data		
Magnesium	7439-95-4	No data			No data	No data		
Manganese	7439-96-5	No data			No data	No data		
Mercury	7439-97-6	2.50E+00			1.00E-01	2.00E+00		2.50E+00
Molybdenum	7439-98-7	No data			No data	4.00E+01		4.00E+01
Nickel	7440-02-0	1.00E+02			2.00E+02	1.00E+02		1.00E+02
Potassium	7440-09-7	No data			No data	No data		
Rhodium	7440-16-6	No data			No data	No data		
Selenium	7782-49-2	7.70E+00			7.00E+01	No data		7.70E+00
Silicon	7440-21-3	No data			No data	No data		
Silver	7440-22-4	No data			No data	No data		
Sodium	7440-23-5	No data			No data	No data		
Strontium	7440-24-6	No data			No data	No data		
Tantalum	7440-25-7	No data			No data	No data		
Thallium	7440-28-0	No data			No data	No data		
Tin	7440-31-5	No data			No data	5.00E+01		5.00E+01
Tungsten	7440-33-7	No data			No data	No data		
Uranium	7440-61-1	No data			No data	No data		
Vanadium	7440-62-2	No data			No data	No data		
Yttrium	7440-65-5	No data			No data	No data		
Zinc	7440-66-6	1.99E+02			2.00E+02	5.00E+02		1.99E+02
Zirconium	7440-67-7	No data			No data	No data		
Non-metals and Anions								
Ammonia/Ammonium	7664-41-7	No data			No data	No data		
Bromide	24959-67-9	No data			No data	5.00E+01		5.00E+01
Chloride	16887-00-6	No data			No data	No data		
Cyanide	57-12-5	No data			No data	1.00E+01		1.00E+01
Fluoride	16984-48-8	No data			No data	4.00E+02		4.00E+02
Hydroxide	14280-30-9	No data			No data	No data		
Iodine	7553-56-2	No data			No data	No data		
Nitrate	14797-55-8	No data			No data	No data		

Table C3-2 Toxicity Reference Values (TRVs) for Soil Invertebrates

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997b[b] (mg/kg)	Dutch Soil Cleanup Interim Act[c] (mg/kg)	Notes	Recommended TRV[d] (mg/kg)
Nitrite	14797-65-0	No data			No data	No data		
Phosphate	14265-44-2	No data			No data	No data		
Phosphorus	7723-14-0	No data			No data	No data		
Sulfate	14808-79-8	No data			No data	No data		
Total Sulfur	63705-05-5	No data			No data	2.00E+01		2.00E+01
Priority Pollutants								
Carbon Dioxide	124-38-9	No data			No data	No data		
Nitrogen Dioxide	10102-44-0	No data			No data	No data		
Ozone	10028-15-6	No data			No data	No data		
Particulate Matter	No CAS #	No data			No data	No data		
Sulfur Dioxode	7446-09-5	No data			No data	No data		
Radionuclides[s]								
Actinium-227	14952-40-0	NA			NA	NA		NA
Americium-241	1596-10-2	NA			NA	NA		NA
Americium-243	14993-75-0	NA			NA	NA		NA
Antimony-125	14234-35-6	NA			NA	NA		NA
Barium-137	13981-97-0	NA			NA	NA		NA
Cadmium-113	None	NA			NA	NA		NA
Carbon-14	14762-75-5	NA			NA	NA		NA
Cesium-134	13967-70-9	NA			NA	NA		NA
Cesium-137	10045-97-3	NA			NA	NA		NA
Cobalt-60	10198-40-0	NA			NA	NA		NA
Curium-242	15510-73-3	NA			NA	NA		NA
Curium-243	15757-87-6	NA			NA	NA		NA
Curium-244	13981-15-2	NA			NA	NA		NA
Europium-152	14683-23-9	NA			NA	NA		NA
Europium-154	15585-10-1	NA			NA	NA		NA
Europium-155	14391-16-3	NA			NA	NA		NA
Iodine-129	15046-84-1	NA			NA	NA		NA
Neptunium-237	13994-20-2	NA			NA	NA		NA

Table C3-2 Toxicity Reference Values (TRVs) for Soil Invertebrates

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997b[b] (mg/kg)	Dutch Soil Cleanup Interim Act[c] (mg/kg)	Notes	Recommended TRV[d] (mg/kg)
Nickel-59	14336-70-0	NA			NA	NA		NA
Nickel-63	13981-37-8	NA			NA	NA		NA
Niobium-93	7440-03-1	NA			NA	NA		NA
Plutonium-238	13981-16-3	NA			NA	NA		NA
Plutonium-239	15117-48-3	NA			NA	NA		NA
Plutonium-240	14119-33-6	NA			NA	NA		NA
Plutonium-241	14119-32-5	NA			NA	NA		NA
Plutonium-242	13982-10-0	NA			NA	NA		NA
Protactinium-231	14331-85-2	NA			NA	NA		NA
Radium-226	13982-63-3	NA			NA	NA		NA
Radium-228	15262-20-1	NA			NA	NA		NA
Ruthenium-106	13967-48-1	NA			NA	NA		NA
Samarium-151	15715-94-3	NA			NA	NA		NA
Selenium-79	None	NA			NA	NA		NA
Strontium-90	10098-97-2	NA			NA	NA		NA
Technetium-99	14133-79-7	NA			NA	NA		NA
Thorium-229	15594-54-4	NA			NA	NA		NA
Thorium-232	7440-29-1	NA			NA	NA		NA
Tin-126	15832-50-5	NA			NA	NA		NA
Tritium	10028-17-8	NA			NA	NA		NA
Uranium-232	14158-29-3	NA			NA	NA		NA
Uranium-233	13968-55-3	NA			NA	NA		NA
Uranium-234	13966-29-5	NA			NA	NA		NA
Uranium-235	15117-96-1	NA			NA	NA		NA
Uranium-236	13982-70-2	NA			NA	NA		NA
Uranium-238	7440-61-1	NA			NA	NA		NA
Yttrium-90	10098-91-6	NA			NA	NA		NA
Zirconium-93	15751-77-6	NA			NA	NA		NA

NA = Not applicable

[a] Published in Appendix E of EPA (1999), Table E-6

Table C3-2 Toxicity Reference Values (TRVs) for Soil Invertebrates

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg)	Notes	Ecology Guidance TRV (mg/kg)	Efroymson et al. 1997b[b] (mg/kg)	Dutch Soil Cleanup Interim Act[c] (mg/kg)	Notes	Recommended TRV[d] (mg/kg)

[b] Published in Table 1 of Efroymson et al. 1997b

[c] Column B values (moderate soil contamination that requires additional study) published in Table 1 of Beyer 1990. Note that this is a guidance level for soil cleanup, not a toxicity level specifically for soil invertebrates.

[d] Selection criteria: the order of preferrence is EPA (1999), Efroymson et al. (1997c), and Dutch Interim Act values

[e] Value for chlorophenols (each)

[f] Value for total monocyclic aromatic hydrocarbons

[g] Value for xylene (total)

[h] Value for aliphatic chlorinated hydrocarbons (each)

[i] Value for total chlorinated hydrocarbons

[j] Value for total PCBs

[k] Value for Aroclor 1254 in Appendix E of EPA (1999), Table E-6, was used for PCB mixtures

[l] Value for polyclic chlorinated hydrocarbons (total)

[m] Value for total polycyclic aromatic hydrocarbons

[n] Value for high-molecular-weight polynuclear aromatic hydrocarbons from Appendix E of EPA (1999), Table E-6

[o] Value for chlorobenzenes (each)

[p] Nitrobenzene value from Appendix E of EPA (1999), Table E-6, was used as a surrogate value.

[q] Value for phenols (excluding chlorophenols)

[r] Value for oganochlorinated pesticides (each)

[s] TRV is not applicable to single radionuclides. Combined external and internal radiation exposure for earthworms from all radionuclides combined cannot exceed 1 rad/d.

Table C3-3. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Bird Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
Organics											
Aromatic Halogenated Hydrocarbons											
2,3,4,6-Tetrachlorophenol	58-90-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
4-Chloro-3-methylphenol	59-50-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Aromatic Nonhalogenated Hydrocarbons											
2-Nitrotoluene	88-72-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
4-Nitrobiphenyl	92-93-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Benzaldehyde	100-52-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Benzene	71-43-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Benzyl alcohol	100-51-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Ethyl benzene	100-41-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
m-Xylene	108-38-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
o-Xylene	95-47-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
p-Xylene	106-42-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Styrene	100-42-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Toluene	108-88-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Non-aromatic Nonhalogenated Hydrocarbons											
1,2-Epoxybutane	106-88-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,3-Butadiene	106-99-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,4-Dioxane	123-91-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1-Methylpropyl alcohol	78-92-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1-Nitropropane	108-03-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2,2,4-Trimethylpentane	540-84-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2-Butanone	78-93-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2-Butenaldehyde (2-Butenal)	4170-30-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2-Ethoxyethanol	110-80-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2-Heptanone	110-43-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2-Hexanone	591-78-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2-Methoxyethanol	109-86-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV

Table C3-3. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Bird Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
2-Methyl-2-propanol	75-65-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2-Methyl-2-propenenitrile	126-98-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2-Methylaziridine	75-55-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2-Methylpropyl alcohol	78-83-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2-Pentanone	107-87-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2-Propanone (Acetone)	67-64-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2-Propene-1-ol	107-18-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2-Propyl alcohol	67-63-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
3-Heptanone	106-35-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
3-Methyl-1-butanol	123-51-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
3-Methyl-2-butanone	563-80-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
3-Pentanone	96-22-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
4-Heptanone	123-19-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
4-Methyl-2-pentanone	108-10-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
4-Methyl-3-penten-2-one	141-79-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
5-Methyl-2-hexanone	110-12-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Acetaldehyde	75-07-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Acetamide	60-35-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Acetic acid	64-19-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Acetic acid ethyl ester	141-78-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Acetic acid n-butyl ester	123-86-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Acetonitrile	75-05-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Acrolein	107-02-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Acrylonitrile	107-13-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Bis(isopropyl)ether	108-20-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Butane	106-97-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Carbon disulfide	75-15-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Cyanogen	460-19-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Cyclohexane	110-82-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV

Table C3-3. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Bird Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
Cyclohexanone	108-94-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Cyclohexene	110-83-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Cyclopentane	287-92-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Ethyl alcohol	64-17-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Ethyl ether	60-29-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Ethyl methacrylate	97-63-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Formaldehyde	50-00-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Formamide	75-12-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Formic acid	64-18-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Formic acid, methyl ester	107-31-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Glycidylaldehyde	765-34-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Methyl acetate	79-20-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Methyl alcohol	67-56-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Methyl isocyanate	624-83-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Methyl methacrylate	80-62-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Methyl tert-butyl ether	1634-04-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Methylacetylene	74-99-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Methylcyclohexane	108-87-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
N,N-Dimethylacetamide	127-19-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
n-Butyl alcohol	71-36-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
n-Heptane	142-82-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
n-Hexane	110-54-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Nitromethane	75-52-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
n-Nonane	111-84-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
n-Octane	111-65-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
n-Pentane	109-66-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
n-Propionaldehyde	123-38-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
n-Propyl alcohol	71-23-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
n-Valeraldehyde	110-62-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV

Table C3-3. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Bird Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
Oxirane	75-21-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
p-Cymene	99-87-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Phosgene	75-44-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Propargyl alcohol	107-19-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Propionic acid	79-09-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Propionitrile	107-12-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Propylene gylcol monomethyl ether	107-98-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
p-tert-Butyltoluene	98-51-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Triethylamine	121-44-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Trimethylamine	75-50-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Vinyl acetate	108-05-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Non-aromatic Halogenated Hydrocarbons											
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,1,1,2-Tetrachloroethane	630-20-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,1,1-Trichloroethane	71-55-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,1,2,2-Tetrachloroethane	79-34-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,1,2,2-Tetrachloroethene	127-18-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,1,2-Trichloroethane	79-00-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,1,2-Trichloroethylene	79-01-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,1-Dichloroethane	75-34-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,1-Dichloroethene	75-35-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,2,2-Trichloro-1,1,2-trifluoroethane	76-13-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,2,3-Trichloropropane	96-18-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,2-Dibromo-3-chloropropane	96-12-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,2-Dichloroethane	107-06-2	Chicken	1.60E+00	1.72E+01	chronic	NOAEL	Reproduction	Alumot et al. (1976b) in [1]	1.00E+00	1.00E+00	1.72E+01
1,2-Dichloroethylene	540-59-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,2-Dichloropropane	78-87-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV

Table C3-3. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Bird Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
1,3-Dichloropropene	542-75-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,4-Dichloro-2-butene	764-41-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1-Chloroethene	75-01-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2,2-Dichloropropionic acid	75-99-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2-Chloropropane	75-29-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
3-Chloropropene (Allyl chloride)	107-05-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Bromochloromethane	74-97-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Bromodichloromethane	75-27-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Bromoethene	593-60-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Bromoform	75-25-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Bromomethane	74-83-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Carbon tetrachloride	56-23-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Chlorodibromomethane	124-48-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Chlorodifluoromethane	75-45-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Chloroethane	75-00-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Chloroform	67-66-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Chloromethane	74-87-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Chloropentafluoroethane	76-15-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
cis-1,2-Dichloroethene	156-59-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
cis-1,3-Dichloropropene	10061-01-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Cyanogen bromide	506-68-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Cyanogen chloride	506-77-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Dichlorodifluoromethane	75-71-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Dichlorofluoromethane	75-43-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Dichloromethane	75-09-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Difluorodibromomethane	75-61-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Hexafluoroacetone	684-16-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Iodomethane	74-88-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Methylene bromide	74-95-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV

Table C3-3. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Bird Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
Pentachloroethane	76-01-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
trans-1,2-Dichloroethylene	156-60-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
trans-1,3-Dichloropropene	10061-02-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Trichloroacetic acid	76-03-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Trichlorofluoroethane	27154-33-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Trichlorofluoromethane	75-69-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Trifluorobromomethane	75-63-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Dioxin and Furan Compounds											
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin	35822-46-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin	39227-28-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin	57653-85-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin	19408-74-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,2,3,7,8-Pentachlorodibenzo-p-dioxin	40321-76-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2,3,7,8-Tetrachlorodibenzo-p-dioxin	1746-01-6	Ring-necked Pheasant	1.00E+00	1.40E-05	chronic	NOAEL	Reproduction	Nosek et al. (1992) in [1]	1.00E+00	1.00E+00	1.40E-05
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	Chick (1 day old)	1.21E-01	1.00E-04	subchronic	LOAEL	Mortality	McKinney et al. (1976) in [1]	1.00E-01	1.00E-01	1.00E-06
Dibenzofuran	132-64-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Octachlorodibenzo(p)dioxin	3268-87-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Octachlorodibenzofuran	39001-02-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
PCBs											
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV

Table C3-3. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Bird Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2',3,4,4',5-Pentachlorobiphenyl	no cas #	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
3,3',4,4',5-Pentachlorobiphenyl	no cas #	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Polychlorinated biphenyls (PCBs)[a]	1336-36-3	Ring-necked pheasant	1.00E+00	1.80E+00	chronic	LOAEL	Reproduction	Dahlgren et al. (1972) in [1]	1.0	0.1	1.80E-01
Phthalates											
Bis(2-ethylhexyl)phthalate	117-81-7	Ringed dove	1.55E-01	1.10E+00	chronic	NOAEL	Reproduction	Peakall (1974) in [1]	1.00E+00	1.00E+00	1.10E+00
Butylbenzylphthalate	85-68-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Di-n-butylphthalate	84-74-2	Ringed dove	1.55E-01	1.11E+00	chronic	LOAEL	Reproduction	Peakall (1974) in [1]	1.00E+00	1.00E-01	1.11E-01
Diethylphthalate	84-66-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Dimethylphthalate	131-11-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
n-Dioctyl phthalate	117-84-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Light Polycyclic Aromatic Hydrocarbons											
2-Chloronapthalene	91-58-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2-Methylnaphthalene	91-57-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
5-Nitroacenaphthene	602-87-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Acenaphthene	83-32-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Acenaphthylene	208-96-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV

Table C3-3. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Bird Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
Anthracene	120-12-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Fluorene	86-73-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Indene	95-13-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Naphthalene	91-20-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Phenanthrene	85-01-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Pyrene	129-00-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Heavy Polycyclic Aromatic Hydrocarbons											
3-Methylcholanthrene	56-49-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
5-Methylchrysene	3697-24-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Benzo(a)anthracene	56-55-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Benzo(a)pyrene	50-32-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Benzo(b)fluoranthene	205-99-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Benzo(e)pyrene	192-97-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Benzo(g,h,i)perylene	191-24-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Benzo(j)fluoranthene	205-82-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Benzo(k)fluoranthene	207-08-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Benzo[a,i]pyrene	191-30-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Chrysene	218-01-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Dibenzo(a,h)anthracene	53-70-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Dibenz[a,h]acridine	226-36-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Dibenz[a,j]acridine	224-42-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Dibenzo(a,e)fluoranthene	5385-75-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Dibenzo(a,h)fluoranthene	no cas #	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Dibenzo[a,e]pyrene	192-65-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Dibenzo[a,h]pyrene	189-64-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Dibenzo[a,i]pyrene	189-55-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Fluoranthene	206-44-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Hexachloronaphthalene	1335-87-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Indeno(1,2,3-cd)pyrene	193-39-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV

Table C3-3. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Bird Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
Octachloronaphthalene	2234-13-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Pentachloronaphthalene	1321-64-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Tetrachloronaphthalene	1335-88-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Trichloronaphthalene	1321-65-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Light Substituted Benzene Compounds											
1,2,3-Trichlorobenzene	87-61-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,2,4-Trichlorobenzene	120-82-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,2,4-Trimethyl benzene	95-63-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,2-Dichlorobenzene	95-50-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,3,5-Trimethyl benzene	108-67-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,3-Dichlorobenzene	541-73-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,3-Dinitrobenzene	99-65-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,4-Dichlorobenzene	106-46-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,4-Dinitrobenzene	100-25-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2,4,5-Trichlorophenol	95-95-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2,4,6-Trichlorophenol	88-06-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2,4-Dichlorophenol	120-83-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2,4-Dimethylphenol	105-67-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2,4-Dinitrophenol	51-28-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2,4-Dinitrotoluene	121-14-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2,6-Dinitrotoluene	606-20-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2-Chlorophenol	95-57-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2-Chlorotoluene	95-49-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2-Nitrophenol	88-75-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
4,6-Dinitro-o-cresol	534-52-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
4-Chlorotoluene	106-43-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
4-Nitrophenol	100-02-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
alpha-Methylstyrene	98-83-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Aniline	62-53-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV

Table C3-3. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Bird Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
Benzotrichloride	98-07-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Benzyl chloride	100-44-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Bromobenzene	108-86-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Chlorobenzene	108-90-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Cumene	98-82-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
m-Cresol	108-39-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
n-Butyl benzene	104-51-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Nitrobenzene	98-95-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
n-Propyl benzene	103-65-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
o-Cresol	95-48-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
o-Dinitrobenzene	528-29-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
o-Nitroaniline	88-74-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
o-Toluidine	95-53-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
p-Chloroaniline	106-47-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
p-Cresol	106-44-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Phenol	108-95-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
p-Nitrochlorobenzene	100-00-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
p-Toluidine	106-49-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
sec-Butyl benzene	135-98-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
tert-Butyl benzene	98-06-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Toluene-2,6-diamine	823-40-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Trimethyl benzene	25551-13-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Other Light Semivolatile Compounds											
1,1'-Biphenyl	92-52-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,1-Dimethylhydrazine	57-14-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,2-Dimethylhydrazine	540-73-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,2-Diphenylhydrazine	122-66-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,3-Propane sultone	1120-71-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2,4-Toluene diisocyante	584-84-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV

Table C3-3. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Bird Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
2-Chloroacetophenone	532-27-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2-Propenoic acid	79-10-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
4,4-Methylenedianiline	101-77-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Acetophenone	98-86-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Benzoic acid	65-85-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Bis(2-chloroethoxy)methane	111-91-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Bis(2-chloroethyl) ether	111-44-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Chlorocyclopentadiene	41851-50-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Cyclohexanol	108-93-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Dichloroisopropyl ether	108-60-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Dichloromethyl ether	542-88-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Dichloropentadiene	no cas #	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Dimethyl sulfate	77-78-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Dimethylaniline	121-69-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
di-n-Propylnitrosamine	621-64-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Diphenyl ether	101-84-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Epichlorohydrin	106-89-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Ethyl Carbamate (Urethane)	51-79-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Ethyl methanesulfonate	62-50-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Ethylene dibromide	106-93-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Ethylene glycol	107-21-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Ethylene glycol monobutyl ether	111-76-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Ethylene glycol monoethyl ether acetate	111-15-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Ethylene thiourea	96-45-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Furfural	98-01-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Maleic hydrazide	123-33-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Malononitrile	109-77-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Methyl styrene (mixed isomers)	25013-15-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Methylhydrazine	60-34-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV

Table C3-3. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Bird Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
N,N-Diphenylamine	122-39-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Nitric acid, propyl ester	627-13-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
N-Nitrosodi-n-butylamine	924-16-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
N-Nitrosomorpholine	59-89-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
N-Nitroso-N,N-dimethylamine	62-75-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
o-Anisidine	90-04-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Oxalic acid	144-62-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Phthalic anhydride	85-44-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
p-Phthalic acid	100-21-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Pyridine	110-86-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Quinoline	91-22-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Quinone	106-51-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Safrole	94-59-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Tetrahydrofuran	109-99-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Other Heavy Semivolatile Compounds											
1,2,4,5-Tetrachlorobenzene	95-94-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
1,3,5-Trinitrobenzene	99-35-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2-sec-Butyl-4,6-dinitrophenol	88-85-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
3,3-Dichlorobenzidine	91-94-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
3,3'-Dimethoxybenzidine	119-90-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
4-Bromophenylphenyl ether	101-55-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Ammonium perfluorooctanoate	3825-26-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Azobenzene	103-33-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Chlorobenzilate	510-15-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Dibutylphosphate	107-66-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Dimethyl aminoazobenzene	60-11-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV

Table C3-3. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Bird Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
Hexachlorobenzene	118-74-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Hexachlorobutadiene	87-68-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Hexachlorocyclopentadiene	77-47-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Hexachloroethane	67-72-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Hexachlorophene	70-30-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Hexamethylene-1,5-diisocyanate	822-06-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Mirex	2385-85-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Nitrofen	1836-75-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Pentachlorobenzene	608-93-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Pentachloronitrobenzene	82-68-8	Chicken	1.50E+00	7.07E+00	35 weeks	NOAEL	Reproduction	Dunn et al. (1979) in [1]	1.00E+00	1.00E+00	7.07E+00
Pentachlorophenol	87-86-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Picric acid	88-89-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Pronamide	23950-58-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Strychnine	57-24-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Terphenyls	26140-60-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Tributyl phosphate	126-73-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Trifluralin	1582-09-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Triphenylamine	603-34-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Herbicides and Organochlorinated Pesticides											
2,4,5-Trichlorophenoxyacetic acid (2,4,5-T)	93-76-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
2,4-D	94-75-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
4,4'-DDD	72-54-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
4,4'-DDE	72-55-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
4,4'-DDT	50-29-3	Brown pelican	3.50E+00	2.80E-02	chronic	LOAEL	Reproduction	Anderson et al. (1975) in [1]	1.00E+00	1.00E-01	2.80E-03
Aldrin	309-00-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
alpha-BHC	319-84-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
beta-BHC	319-85-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Captan	133-06-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV

Table C3-3. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Bird Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
Chlordane	57-74-9	Red-winged blackbird	6.40E-02	2.14E+00	chronic	NOAEL	Mortality	Stickel et al. (1983) in [1]	1.00E+00	1.00E+00	2.14E+00
delta-BHC	319-86-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Dieldrin	60-57-1	Barn owl	4.66E-01	7.70E-02	chronic	NOAEL	Reproduction	Mendenhall et al. (1983) in [1]	1.00E+00	1.00E+00	7.70E-02
Endothall	145-73-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Endrin	72-20-8	Mallard duck	1.15E+00	3.00E-01	chronic	NOAEL	Reproduction	Spann et al. (1986) in [1]	1.00E+00	1.00E+00	3.00E-01
gamma-BHC (Lindane)	58-89-9	Mallard Duck	1.00E+00	2.00E+00	chronic	NOAEL	Reproduction	Chakravarty et al. (1986) in [1]	1.00E+00	1.00E+00	2.00E+00
Heptachlor	76-44-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Isodrin	465-73-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Methoxychlor	72-43-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Silvex (2,4,5-TP)	93-72-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Toxaphene	8001-35-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Inorganics											
Metals											
Aluminum	7429-90-5	Ringed dove	1.55E-01	1.10E+02	chronic	NOAEL	Reproduction	Carriere et al. (1986) in [1]	1.00E+00	1.00E+00	1.10E+02
Antimony	7440-36-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Arsenic	7440-38-2	Mallard duck	1.00E+00	5.14E+00	chronic	NOAEL	Mortality	USFWS (1979) in [1]	1.00E+00	1.00E+00	5.14E+00
Barium	7440-39-3	Chick (14 day old)	1.21E-01	2.08E+02	subchronic	NOAEL	Mortality	Johnson et al. (1960) in [1]	1.00E-01	1.00E+00	2.08E+01
Beryllium	7440-41-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Bismuth	7440-69-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Boron	7440-42-8	Mallard duck	1.00E+00	2.88E+01	chronic	NOAEL	Reproduction	Smith and Anders (1989) in [1]	1.00E+00	1.00E+00	2.88E+01
Cadmium	7440-43-9	Mallard duck	1.15E+00	1.45E+00	chronic	NOAEL	Reproduction	White and Finley (1978) in [1]	1.00E+00	1.00E+00	1.45E+00
Calcium	7440-70-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV

Table C3-3. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Bird Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
Chromium (and VI)	7440-47-3	Black duck	1.25E+00	1.00E+00	chronic	NOAEL	Reproduction	Haseltine et al. (unpubl.) in [1]	1.0	1.0	1.00E+00
Cobalt	7440-48-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Copper	7440-50-8	Chick (5 week old)	5.34E-01	4.70E+01	chronic	NOAEL	Mortality	Mehring et al. (1960) in [1]	1.00E+00	1.00E+00	4.70E+01
Iron	7439-89-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Lead	7439-92-1	Quail	1.50E-01	1.13E+00	chronic	NOAEL	Reproduction	Edens et al. (1976) in [1]	1.00E+00	1.00E+00	1.13E+00
Lithium	7439-93-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Magnesium	7439-95-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Manganese	7439-96-5	Quail	7.20E-02	9.77E+02	chronic	NOAEL	Growth	Laskey and Edens (1985) in [1]	1.00E+00	1.00E+00	9.77E+02
Mercury	7439-97-6	Quail	1.50E-01	4.50E-01	chronic	NOAEL	Reproduction	Hill and Schaffner (1976) in [1]	1.00E+00	1.00E+00	4.50E-01
Molybdenum	7439-98-7	Chicken	1.50E+00	3.53E+01	chronic	LOAEL	Reproduction	Lepore and Miller (1965) in [1]	1.00E+00	1.00E-01	3.53E+00
Nickel	7440-02-0	Mallard duckling	7.82E-01	7.74E+01	chronic	NOAEL	Growth	Cain and Pafford (1981) in [1]	1.00E+00	1.00E+00	7.74E+01
Potassium	7440-09-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Rhodium	7440-16-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Selenium	7782-49-2	Mallard duck	1.00E+00	5.00E-01	chronic	NOAEL	Reproduction	Heinz et al. (1989) in [1]	1.00E+00	1.00E+00	5.00E-01
Silicon	7440-21-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Silver	7440-22-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Sodium	7440-23-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Strontium	7440-24-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Tantalum	7440-25-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Thallium	7440-28-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Tin	7440-31-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Tungsten	7440-33-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV

Table C3-3. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Bird Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
Uranium	7440-61-1	Black duck	1.25E+00	1.60E+02	subchronic	NOAEL	Metabolic	Haseltine and Sileo (1983) in [1]	1.00E-01	1.00E+00	1.60E+01
Vanadium	7440-62-2	Mallard duck	1.17E+00	1.14E+01	chronic	NOAEL	Mortality	White and Dieter (1978) in [1]	1.00E+00	1.00E+00	1.14E+01
Yttrium	7440-65-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Zinc	7440-66-6	Leghorn chicken	1.94E+00	1.45E+01	chronic	NOAEL	Reproduction	Stahl et al. (1990) in [1]	1.00E+00	1.00E+00	1.45E+01
Zirconium	7440-67-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Non-metals and Ions											
Ammonia/Ammonium	7664-41-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Bromide	24959-67-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Chloride	16887-00-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Cyanide	57-12-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Fluoride	16984-48-8	Screech owl	1.81E-01	7.80E+00	chronic	NOAEL	Reproduction	Pattee et al. (1988) in [1]	1.00E+00	1.00E+00	7.80E+00
Hydroxide	14280-30-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Iodine	7553-56-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Nitrate	14797-55-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Nitrite	14797-65-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Phosphate	14265-44-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Phosphorus	7723-14-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Sulfate	14808-79-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Total Sulfur	63705-05-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Criteria Pollutants											
Carbon dioxide	124-38-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Nitrogen dioxide	10102-44-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Ozone	10028-15-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Particulate matter	no cas #	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Sulfur dioxide	7446-09-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No TRV
Radionuclides	TRV is not applicable to single radionuclides. Combined external and internal radiation exposure for birds from all radionuclides combined cannot exceed 0.1 rad/d.										

Table C3-3. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Bird Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF

TRV = toxicity reference value

DCF = Duration conversion factor; 1 if chronic, 0.1 if subchronic (Sample et al. 1996)

ECF = Endpoint conversion factor; 1 if NOAEL, 0.1 if LOAEL (Sample et al. 1996)

NOAEL = No observed adverse effect level

NA = No data available

[a] Data for Aroclor 1254 used as representative of PCB mixtures

LOAEL = Lowest observed adverse effect level

[1] = Sample et al. (1996)

[2] = Clayton and Clayton (1981)

[3] = IRIS (1996)

[4] = IRIS (1998)

Table C3-4. Toxicity Reference Values (TRVs) for Birds

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
Organics						
Aromatic Halogenated Hydrocarbons						
2,3,4,6-Tetrachlorophenol	58-90-2	No TRV			No TRV	No TRV
4-Chloro-3-methylphenol	59-50-7	No TRV			No TRV	No TRV
Aromatic Nonhalogenated Hydrocarbons						
2-Nitrotoluene	88-72-2	No TRV			No TRV	No TRV
4-Nitrobiphenyl	92-93-3	No TRV			No TRV	No TRV
Benzaldehyde	100-52-7	No TRV			No TRV	No TRV
Benzene	71-43-2	No TRV			No TRV	No TRV
Benzyl alcohol	100-51-6	No TRV			No TRV	No TRV
Ethyl benzene	100-41-4	No TRV			No TRV	No TRV
m-Xylene	108-38-3	No TRV			No TRV	No TRV
o-Xylene	95-47-6	No TRV			No TRV	No TRV
p-Xylene	106-42-3	No TRV			No TRV	No TRV
Styrene	100-42-5	No TRV			No TRV	No TRV
Toluene	108-88-3	No TRV			No TRV	No TRV
Non-aromatic Nonhalogenated Hydrocarbons						
1,2-Epoxybutane	106-88-7	No TRV			No TRV	No TRV
1,3-Butadiene	106-99-0	No TRV			No TRV	No TRV
1,4-Dioxane	123-91-1	No TRV			No TRV	No TRV
1-Methylpropyl alcohol	78-92-2	No TRV			No TRV	No TRV
1-Nitropropane	108-03-2	No TRV			No TRV	No TRV
2,2,4-Trimethylpentane	540-84-1	No TRV			No TRV	No TRV
2-Butanone	78-93-3	No TRV			No TRV	No TRV
2-Butenaldehyde (2-Butenal)	4170-30-3	No TRV			No TRV	No TRV
2-Ethoxyethanol	110-80-5	No TRV			No TRV	No TRV
2-Heptanone	110-43-0	No TRV			No TRV	No TRV
2-Hexanone	591-78-6	No TRV			No TRV	No TRV
2-Methoxyethanol	109-86-4	No TRV			No TRV	No TRV
2-Methyl-2-propanol	75-65-0	No TRV			No TRV	No TRV

Table C3-4. Toxicity Reference Values (TRVs) for Birds

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
2-Methyl-2-propenenitrile	126-98-7	No TRV			No TRV	No TRV
2-Methylaziridine	75-55-8	No TRV			No TRV	No TRV
2-Methylpropyl alcohol	78-83-1	No TRV			No TRV	No TRV
2-Pentanone	107-87-9	No TRV			No TRV	No TRV
2-Propanone (Acetone)	67-64-1	5.20E+01			No TRV	5.20E+01
2-Propene-1-ol	107-18-6	No TRV			No TRV	No TRV
2-Propyl alcohol	67-63-0	No TRV			No TRV	No TRV
3-Heptanone	106-35-4	No TRV			No TRV	No TRV
3-Methyl-1-butanol	123-51-3	No TRV			No TRV	No TRV
3-Methyl-2-butanone	563-80-4	No TRV			No TRV	No TRV
3-Pentanone	96-22-0	No TRV			No TRV	No TRV
4-Heptanone	123-19-3	No TRV			No TRV	No TRV
4-Methyl-2-pentanone	108-10-1	No TRV			No TRV	No TRV
4-Methyl-3-penten-2-one	141-79-7	No TRV			No TRV	No TRV
5-Methyl-2-hexanone	110-12-3	No TRV			No TRV	No TRV
Acetaldehyde	75-07-0	No TRV			No TRV	No TRV
Acetamide	60-35-5	No TRV			No TRV	No TRV
Acetic acid	64-19-7	No TRV			No TRV	No TRV
Acetic acid ethyl ester	141-78-6	No TRV			No TRV	No TRV
Acetic acid n-butyl ester	123-86-4	No TRV			No TRV	No TRV
Acetonitrile	75-05-8	No TRV			No TRV	No TRV
Acrolein	107-02-8	No TRV			No TRV	No TRV
Acrylonitrile	107-13-1	No TRV			No TRV	No TRV
Bis(isopropyl)ether	108-20-3	No TRV			No TRV	No TRV
Butane	106-97-8	No TRV			No TRV	No TRV
Carbon disulfide	75-15-0	No TRV			No TRV	No TRV
Cyanogen	460-19-5	No TRV			No TRV	No TRV
Cyclohexane	110-82-7	No TRV			No TRV	No TRV
Cyclohexanone	108-94-1	No TRV			No TRV	No TRV
Cyclohexene	110-83-8	No TRV			No TRV	No TRV
Cyclopentane	287-92-3	No TRV			No TRV	No TRV

Table C3-4. Toxicity Reference Values (TRVs) for Birds

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
Ethyl alcohol	64-17-5	No TRV			No TRV	No TRV
Ethyl ether	60-29-7	No TRV			No TRV	No TRV
Ethyl methacrylate	97-63-2	No TRV			No TRV	No TRV
Formaldehyde	50-00-0	No TRV			No TRV	No TRV
Formamide	75-12-7	No TRV			No TRV	No TRV
Formic acid	64-18-6	No TRV			No TRV	No TRV
Formic acid, methyl ester	107-31-3	No TRV			No TRV	No TRV
Glycidylaldehyde	765-34-4	No TRV			No TRV	No TRV
Methyl acetate	79-20-9	No TRV			No TRV	No TRV
Methyl alcohol	67-56-1	No TRV			No TRV	No TRV
Methyl isocyanate	624-83-9	No TRV			No TRV	No TRV
Methyl methacrylate	80-62-6	No TRV			No TRV	No TRV
Methyl tert-butyl ether	1634-04-4	No TRV			No TRV	No TRV
Methylacetylene	74-99-7	No TRV			No TRV	No TRV
Methylcyclohexane	108-87-2	No TRV			No TRV	No TRV
N,N-Dimethylacetamide	127-19-5	No TRV			No TRV	No TRV
n-Butyl alcohol	71-36-3	No TRV			No TRV	No TRV
n-Heptane	142-82-5	No TRV			No TRV	No TRV
n-Hexane	110-54-3	No TRV			No TRV	No TRV
Nitromethane	75-52-5	No TRV			No TRV	No TRV
n-Nonane	111-84-2	No TRV			No TRV	No TRV
n-Octane	111-65-9	No TRV			No TRV	No TRV
n-Pentane	109-66-0	No TRV			No TRV	No TRV
n-Propionaldehyde	123-38-6	No TRV			No TRV	No TRV
n-Propyl alcohol	71-23-8	No TRV			No TRV	No TRV
n-Valeraldehyde	110-62-3	No TRV			No TRV	No TRV
Oxirane	75-21-8	No TRV			No TRV	No TRV
p-Cymene	99-87-6	No TRV			No TRV	No TRV
Phosgene	75-44-5	No TRV			No TRV	No TRV
Propargyl alcohol	107-19-7	No TRV			No TRV	No TRV
Propionic acid	79-09-4	No TRV			No TRV	No TRV

Table C3-4. Toxicity Reference Values (TRVs) for Birds

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
Propionitrile	107-12-0	No TRV			No TRV	No TRV
Propylene gylcol monomethyl ether	107-98-2	No TRV			No TRV	No TRV
p-tert-Butyltoluene	98-51-1	No TRV			No TRV	No TRV
Triethylamine	121-44-8	No TRV			No TRV	No TRV
Trimethylamine	75-50-3	No TRV			No TRV	No TRV
Vinyl acetate	108-05-4	No TRV			No TRV	No TRV
Non-aromatic Halogenated Hydrocarbons						
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	No TRV			No TRV	No TRV
1,1,1,2-Tetrachloroethane	630-20-6	No TRV			No TRV	No TRV
1,1,1-Trichloroethane	71-55-6	No TRV			No TRV	No TRV
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	No TRV			No TRV	No TRV
1,1,2,2-Tetrachloroethane	79-34-5	No TRV			No TRV	No TRV
1,1,2,2-Tetrachloroethene	127-18-4	No TRV			No TRV	No TRV
1,1,2-Trichloroethane	79-00-5	No TRV			No TRV	No TRV
1,1,2-Trichloroethylene	79-01-6	No TRV			No TRV	No TRV
1,1-Dichloroethane	75-34-3	No TRV			No TRV	No TRV
1,1-Dichloroethene	75-35-4	No TRV			No TRV	No TRV
1,2,2-Trichloro-1,1,2-trifluoroethane	76-13-1	No TRV			No TRV	No TRV
1,2,3-Trichloropropane	96-18-4	No TRV			No TRV	No TRV
1,2-Dibromo-3-chloropropane	96-12-8	No TRV			No TRV	No TRV
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	No TRV			No TRV	No TRV
1,2-Dichloroethane	107-06-2	No TRV			1.72E+01	1.72E+01
1,2-Dichloroethylene	540-59-0	No TRV			No TRV	No TRV
1,2-Dichloropropane	78-87-5	No TRV			No TRV	No TRV
1,3-Dichloropropene	542-75-6	No TRV			No TRV	No TRV
1,4-Dichloro-2-butene	764-41-0	No TRV			No TRV	No TRV
1-Chloroethene	75-01-4	No TRV			No TRV	No TRV
2,2-Dichloropropionic acid	75-99-0	No TRV			No TRV	No TRV
2-Chloropropane	75-29-6	No TRV			No TRV	No TRV
3-Chloropropene (Allyl chloride)	107-05-1	No TRV			No TRV	No TRV
Bromochloromethane	74-97-5	No TRV			No TRV	No TRV

Table C3-4. Toxicity Reference Values (TRVs) for Birds

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
Bromodichloromethane	75-27-4	No TRV			No TRV	No TRV
Bromoethene	593-60-2	No TRV			No TRV	No TRV
Bromoform	75-25-2	No TRV			No TRV	No TRV
Bromomethane	74-83-9	No TRV			No TRV	No TRV
Carbon tetrachloride	56-23-5	No TRV			No TRV	No TRV
Chlorodibromomethane	124-48-1	No TRV			No TRV	No TRV
Chlorodifluoromethane	75-45-6	No TRV			No TRV	No TRV
Chloroethane	75-00-3	No TRV			No TRV	No TRV
Chloroform	67-66-3	No TRV			No TRV	No TRV
Chloromethane	74-87-3	No TRV			No TRV	No TRV
Chloropentafluoroethane	76-15-3	No TRV			No TRV	No TRV
cis-1,2-Dichloroethene	156-59-2	No TRV			No TRV	No TRV
cis-1,3-Dichloropropene	10061-01-5	No TRV			No TRV	No TRV
Cyanogen bromide	506-68-3	No TRV			No TRV	No TRV
Cyanogen chloride	506-77-4	No TRV			No TRV	No TRV
Dichlorodifluoromethane	75-71-8	No TRV			No TRV	No TRV
Dichlorofluoromethane	75-43-4	No TRV			No TRV	No TRV
Dichloromethane	75-09-2	No TRV			No TRV	No TRV
Difluorodibromomethane	75-61-6	No TRV			No TRV	No TRV
Hexafluoroacetone	684-16-2	No TRV			No TRV	No TRV
Iodomethane	74-88-4	No TRV			No TRV	No TRV
Methylene bromide	74-95-3	No TRV			No TRV	No TRV
Pentachloroethane	76-01-7	No TRV			No TRV	No TRV
trans-1,2-Dichloroethylene	156-60-5	No TRV			No TRV	No TRV
trans-1,3-Dichloropropene	10061-02-6	No TRV			No TRV	No TRV
Trichloroacetic acid	76-03-9	No TRV			No TRV	No TRV
Trichlorofluoroethane	27154-33-2	No TRV			No TRV	No TRV
Trichlorofluoromethane	75-69-4	No TRV			No TRV	No TRV
Trifluorobromomethane	75-63-8	No TRV			No TRV	No TRV
Dioxin and Furan Compounds						
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin	35822-46-9	No TRV			No TRV	No TRV

Table C3-4. Toxicity Reference Values (TRVs) for Birds

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	No TRV			No TRV	No TRV
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	No TRV			No TRV	No TRV
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin	39227-28-6	No TRV			No TRV	No TRV
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	No TRV			No TRV	No TRV
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin	57653-85-7	No TRV			No TRV	No TRV
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	No TRV			No TRV	No TRV
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin	19408-74-3	No TRV			No TRV	No TRV
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	No TRV			No TRV	No TRV
1,2,3,7,8-Pentachlorodibenzo-p-dioxin	40321-76-4	No TRV			No TRV	No TRV
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	No TRV			No TRV	No TRV
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	No TRV			No TRV	No TRV
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	No TRV			No TRV	No TRV
2,3,7,8-Tetrachlorodibenzo-p-dioxin	1746-01-6	1.00E-05			1.40E-05	1.00E-05
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	No TRV			1.00E-06	1.00E-06
Dibenzofuran	132-64-9	No TRV			No TRV	No TRV
Octachlorodibenzo(p)dioxin	3268-87-9	No TRV			No TRV	No TRV
Octachlorodibenzofuran	39001-02-0	No TRV			No TRV	No TRV
PCBs						
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	No TRV			No TRV	No TRV
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	No TRV			No TRV	No TRV
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	No TRV			No TRV	No TRV
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	No TRV			No TRV	No TRV
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	No TRV			No TRV	No TRV
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	No TRV			No TRV	No TRV
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	No TRV			No TRV	No TRV
2',3,4,4',5-Pentachlorobiphenyl	no cas #	No TRV			No TRV	No TRV
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	No TRV			No TRV	No TRV
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	No TRV			No TRV	No TRV
3,3',4,4',5-Pentachlorobiphenyl	no cas #	No TRV			No TRV	No TRV
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	No TRV			No TRV	No TRV
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	No TRV			No TRV	No TRV

Table C3-4. Toxicity Reference Values (TRVs) for Birds

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	No TRV			No TRV	No TRV
Polychlorinated biphenyls (PCBs)[d]	1336-36-3	7.20E-02			1.80E-01	7.20E-02
Phthalates						
Bis(2-ethylhexyl)phthalate	117-81-7	1.11E-01			1.10E+00	1.11E-01
Butylbenzylphthalate	85-68-7	No TRV			No TRV	No TRV
Di-n-butylphthalate	84-74-2	No TRV			1.11E-01	1.11E-01
Diethylphthalate	84-66-2	No TRV			No TRV	No TRV
Dimethylphthalate	131-11-3	No TRV			No TRV	No TRV
n-Dioctyl phthalate	117-84-0	No TRV			No TRV	No TRV
Light Polycyclic Aromatic Hydrocarbons						
2-Chloronapthalene	91-58-7	No TRV			No TRV	No TRV
2-Methylnaphthalene	91-57-6	No TRV			No TRV	No TRV
5-Nitroacenaphthene	602-87-9	No TRV			No TRV	No TRV
Acenaphthene	83-32-9	No TRV			No TRV	No TRV
Acenaphthylene	208-96-8	No TRV			No TRV	No TRV
Anthracene	120-12-7	No TRV			No TRV	No TRV
Fluorene	86-73-7	No TRV			No TRV	No TRV
Indene	95-13-6	No TRV			No TRV	No TRV
Naphthalene	91-20-3	No TRV			No TRV	No TRV
Phenanthrene	85-01-8	No TRV			No TRV	No TRV
Pyrene	129-00-0	No TRV			No TRV	No TRV
Heavy Polycyclic Aromatic Hydrocarbons[e]						
3-Methylcholanthrene	56-49-5	No TRV			No TRV	No TRV
5-Methylchrysene	3697-24-3	No TRV			No TRV	No TRV
Benzo(a)anthracene	56-55-3	7.90E-04			No TRV	7.90E-04
Benzo(a)pyrene	50-32-8	1.00E-03			No TRV	1.00E-03
Benzo(b)fluoranthene	205-99-2	1.40E-04	f		No TRV	1.40E-04
Benzo(e)pyrene	192-97-2	No TRV			No TRV	No TRV
Benzo(g,h,i)perylene	191-24-2	No TRV			No TRV	No TRV
Benzo(j)fluoranthene	205-82-3	No TRV			No TRV	No TRV

Table C3-4. Toxicity Reference Values (TRVs) for Birds

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
Benzo(k)fluoranthene	207-08-9	1.40E-04			No TRV	1.40E-04
Benzo[a,i]pyrene	191-30-0	No TRV			No TRV	No TRV
Chrysene	218-01-9	1.00E-03			No TRV	1.00E-03
Dibenzo(a,h)anthracene	53-70-3	3.90E-04			No TRV	3.90E-04
Dibenz[a,h]acridine	226-36-8	No TRV			No TRV	No TRV
Dibenz[a,j]acridine	224-42-0	No TRV			No TRV	No TRV
Dibenzo(a,e)fluoranthene	5385-75-1	No TRV			No TRV	No TRV
Dibenzo(a,h)fluoranthene	no cas #	No TRV			No TRV	No TRV
Dibenzo[a,e]pyrene	192-65-4	No TRV			No TRV	No TRV
Dibenzo[a,h]pyrene	189-64-0	No TRV			No TRV	No TRV
Dibenzo[a,i]pyrene	189-55-9	No TRV			No TRV	No TRV
Fluoranthene	206-44-0	No TRV			No TRV	No TRV
Hexachloronaphthalene	1335-87-1	No TRV			No TRV	No TRV
Indeno(1,2,3-cd)pyrene	193-39-5	1.00E-03			No TRV	1.00E-03
Octachloronaphthalene	2234-13-1	No TRV			No TRV	No TRV
Pentachloronaphthalene	1321-64-8	No TRV			No TRV	No TRV
Tetrachloronaphthalene	1335-88-2	No TRV			No TRV	No TRV
Trichloronaphthalene	1321-65-9	No TRV			No TRV	No TRV
Light Substituted Benzene Compounds						
1,2,3-Trichlorobenzene	87-61-6	No TRV			No TRV	No TRV
1,2,4-Trichlorobenzene	120-82-1	No TRV			No TRV	No TRV
1,2,4-Trimethyl benzene	95-63-6	No TRV			No TRV	No TRV
1,2-Dichlorobenzene	95-50-1	No TRV			No TRV	No TRV
1,3,5-Trimethyl benzene	108-67-8	No TRV			No TRV	No TRV
1,3-Dichlorobenzene	541-73-1	No TRV			No TRV	No TRV
1,3-Dinitrobenzene	99-65-0	4.22E-04			No TRV	4.22E-04
1,4-Dichlorobenzene	106-46-7	No TRV			No TRV	No TRV
1,4-Dinitrobenzene	100-25-4	No TRV			No TRV	No TRV
2,4,5-Trichlorophenol	95-95-4	No TRV			No TRV	No TRV
2,4,6-Trichlorophenol	88-06-2	No TRV			No TRV	No TRV
2,4-Dichlorophenol	120-83-2	No TRV			No TRV	No TRV

Table C3-4. Toxicity Reference Values (TRVs) for Birds

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
2,4-Dimethylphenol	105-67-9	No TRV			No TRV	No TRV
2,4-Dinitrophenol	51-28-5	No TRV			No TRV	No TRV
2,4-Dinitrotoluene	121-14-2	No TRV			No TRV	No TRV
2,6-Dinitrotoluene	606-20-2	No TRV			No TRV	No TRV
2-Chlorophenol	95-57-8	No TRV			No TRV	No TRV
2-Chlorotoluene	95-49-8	No TRV			No TRV	No TRV
2-Nitrophenol	88-75-5	No TRV			No TRV	No TRV
4,6-Dinitro-o-cresol	534-52-1	No TRV			No TRV	No TRV
4-Chlorotoluene	106-43-4	No TRV			No TRV	No TRV
4-Nitrophenol	100-02-7	No TRV			No TRV	No TRV
alpha-Methylstyrene	98-83-9	No TRV			No TRV	No TRV
Aniline	62-53-3	No TRV			No TRV	No TRV
Benzotrichloride	98-07-7	No TRV			No TRV	No TRV
Benzyl chloride	100-44-7	No TRV			No TRV	No TRV
Bromobenzene	108-86-1	No TRV			No TRV	No TRV
Chlorobenzene	108-90-7	No TRV			No TRV	No TRV
Cumene	98-82-8	No TRV			No TRV	No TRV
m-Cresol	108-39-4	No TRV			No TRV	No TRV
n-Butyl benzene	104-51-8	No TRV			No TRV	No TRV
Nitrobenzene	98-95-3	No TRV			No TRV	No TRV
n-Propyl benzene	103-65-1	No TRV			No TRV	No TRV
o-Cresol	95-48-7	No TRV			No TRV	No TRV
o-Dinitrobenzene	528-29-0	No TRV			No TRV	No TRV
o-Nitroaniline	88-74-4	No TRV			No TRV	No TRV
o-Toluidine	95-53-4	No TRV			No TRV	No TRV
p-Chloroaniline	106-47-8	No TRV			No TRV	No TRV
p-Cresol	106-44-5	No TRV			No TRV	No TRV
Phenol	108-95-2	No TRV			No TRV	No TRV
p-Nitrochlorobenzene	100-00-5	No TRV			No TRV	No TRV
p-Toluidine	106-49-0	No TRV			No TRV	No TRV
sec-Butyl benzene	135-98-8	No TRV			No TRV	No TRV

Table C3-4. Toxicity Reference Values (TRVs) for Birds

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
tert-Butyl benzene	98-06-6	No TRV			No TRV	No TRV
Toluene-2,6-diamine	823-40-5	No TRV			No TRV	No TRV
Trimethyl benzene	25551-13-7	No TRV			No TRV	No TRV
Other Light Semivolatile Compounds						
1,1'-Biphenyl	92-52-4	No TRV			No TRV	No TRV
1,1-Dimethylhydrazine	57-14-7	No TRV			No TRV	No TRV
1,2-Dimethylhydrazine	540-73-8	No TRV			No TRV	No TRV
1,2-Diphenylhydrazine	122-66-7	No TRV			No TRV	No TRV
1,3-Propane sultone	1120-71-4	No TRV			No TRV	No TRV
2,4-Toluene diisocyante	584-84-9	No TRV			No TRV	No TRV
2-Chloroacetophenone	532-27-4	No TRV			No TRV	No TRV
2-Propenoic acid	79-10-7	No TRV			No TRV	No TRV
4,4-Methylenedianiline	101-77-9	No TRV			No TRV	No TRV
Acetophenone	98-86-2	No TRV			No TRV	No TRV
Benzoic acid	65-85-0	No TRV			No TRV	No TRV
Bis(2-chloroethoxy)methane	111-91-1	No TRV			No TRV	No TRV
Bis(2-chloroethyl) ether	111-44-4	No TRV			No TRV	No TRV
Chlorocyclopentadiene	41851-50-7	No TRV			No TRV	No TRV
Cyclohexanol	108-93-0	No TRV			No TRV	No TRV
Dichloroisopropyl ether	108-60-1	No TRV			No TRV	No TRV
Dichloromethyl ether	542-88-1	No TRV			No TRV	No TRV
Dichloropentadiene	no cas #	No TRV			No TRV	No TRV
Dimethyl sulfate	77-78-1	No TRV			No TRV	No TRV
Dimethylaniline	121-69-7	No TRV			No TRV	No TRV
di-n-Propylnitrosamine	621-64-7	No TRV			No TRV	No TRV
Diphenyl ether	101-84-8	No TRV			No TRV	No TRV
Epichlorohydrin	106-89-8	No TRV			No TRV	No TRV
Ethyl Carbamate (Urethane)	51-79-6	No TRV			No TRV	No TRV
Ethyl methanesulfonate	62-50-0	No TRV			No TRV	No TRV
Ethylene dibromide	106-93-4	No TRV			No TRV	No TRV
Ethylene glycol	107-21-1	No TRV			No TRV	No TRV

Table C3-4. Toxicity Reference Values (TRVs) for Birds

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
Ethylene glycol monobutyl ether	111-76-2	No TRV			No TRV	No TRV
Ethylene glycol monoethyl ether acetate	111-15-9	No TRV			No TRV	No TRV
Ethylene thiourea	96-45-7	No TRV			No TRV	No TRV
Furfural	98-01-1	No TRV			No TRV	No TRV
Maleic hydrazide	123-33-1	No TRV			No TRV	No TRV
Malononitrile	109-77-3	No TRV			No TRV	No TRV
Methyl styrene (mixed isomers)	25013-15-4	No TRV			No TRV	No TRV
Methylhydrazine	60-34-4	No TRV			No TRV	No TRV
N,N-Diphenylamine	122-39-4	No TRV			No TRV	No TRV
Nitric acid, propyl ester	627-13-4	No TRV			No TRV	No TRV
N-Nitrosodi-n-butylamine	924-16-3	No TRV			No TRV	No TRV
N-Nitrosomorpholine	59-89-2	No TRV			No TRV	No TRV
N-Nitroso-N,N-dimethylamine	62-75-9	No TRV			No TRV	No TRV
o-Anisidine	90-04-0	No TRV			No TRV	No TRV
Oxalic acid	144-62-7	No TRV			No TRV	No TRV
Phthalic anhydride	85-44-9	No TRV			No TRV	No TRV
p-Phthalic acid	100-21-0	No TRV			No TRV	No TRV
Pyridine	110-86-1	No TRV			No TRV	No TRV
Quinoline	91-22-5	No TRV			No TRV	No TRV
Quinone	106-51-4	No TRV			No TRV	No TRV
Safrole	94-59-7	No TRV			No TRV	No TRV
Tetrahydrofuran	109-99-9	No TRV			No TRV	No TRV
Other Heavy Semivolatile Compounds						
1,2,4,5-Tetrachlorobenzene	95-94-3	No TRV			No TRV	No TRV
1,3,5-Trinitrobenzene	99-35-4	No TRV			No TRV	No TRV
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	No TRV			No TRV	No TRV
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	No TRV			No TRV	No TRV
2-sec-Butyl-4,6-dinitrophenol	88-85-7	No TRV			No TRV	No TRV
3,3-Dichlorobenzidine	91-94-1	No TRV			No TRV	No TRV
3,3'-Dimethoxybenzidine	119-90-4	No TRV			No TRV	No TRV
4-Bromophenylphenyl ether	101-55-3	No TRV			No TRV	No TRV

Table C3-4. Toxicity Reference Values (TRVs) for Birds

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
Ammonium perfluorooctanoate	3825-26-1	No TRV			No TRV	No TRV
Azobenzene	103-33-3	No TRV			No TRV	No TRV
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	No TRV			No TRV	No TRV
Chlorobenzilate	510-15-6	No TRV			No TRV	No TRV
Dibutylphosphate	107-66-4	No TRV			No TRV	No TRV
Dimethyl aminoazobenzene	60-11-7	No TRV			No TRV	No TRV
Hexachlorobenzene	118-74-1	2.25E-01			No TRV	2.25E-01
Hexachlorobutadiene	87-68-3	3.19E+00			No TRV	3.19E+00
Hexachlorocyclopentadiene	77-47-4	No TRV			No TRV	No TRV
Hexachloroethane	67-72-1	No TRV			No TRV	No TRV
Hexachlorophene	70-30-4	5.75E+00			No TRV	5.75E+00
Hexamethylene-1,5-diisocyanate	822-06-0	No TRV			No TRV	No TRV
Mirex	2385-85-5	No TRV			No TRV	No TRV
Nitrofen	1836-75-5	No TRV			No TRV	No TRV
Pentachlorobenzene	608-93-5	No TRV			No TRV	No TRV
Pentachloronitrobenzene	82-68-8	6.88E+01			7.07E+00	6.88E+01
Pentachlorophenol	87-86-5	4.03E+00			No TRV	4.03E+00
Picric acid	88-89-1	No TRV			No TRV	No TRV
Pronamide	23950-58-5	No TRV			No TRV	No TRV
Strychnine	57-24-9	No TRV			No TRV	No TRV
Terphenyls	26140-60-3	No TRV			No TRV	No TRV
Tributyl phosphate	126-73-8	No TRV			No TRV	No TRV
Trifluralin	1582-09-8	No TRV			No TRV	No TRV
Triphenylamine	603-34-9	No TRV			No TRV	No TRV
Herbicides and Organochlorinated Pesticides						
2,4,5-Trichlorophenoxyacetic acid (2,4,5-T)	93-76-5	No TRV			No TRV	No TRV
2,4-D	94-75-7	No TRV			No TRV	No TRV
4,4'-DDD	72-54-8	No TRV			No TRV	No TRV
4,4'-DDE	72-55-9	8.45E-01	g		No TRV	8.45E-01
4,4'-DDT	50-29-3	No TRV			2.80E-03	2.80E-03
Aldrin	309-00-2	No TRV			No TRV	No TRV

Table C3-4. Toxicity Reference Values (TRVs) for Birds

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
alpha-BHC	319-84-6	No TRV			No TRV	No TRV
beta-BHC	319-85-7	No TRV			No TRV	No TRV
Captan	133-06-2	No TRV			No TRV	No TRV
Chlordane	57-74-9	No TRV			2.14E+00	2.14E+00
delta-BHC	319-86-8	No TRV			No TRV	No TRV
Dieldrin	60-57-1	No TRV			7.70E-02	7.70E-02
Endothall	145-73-3	No TRV			No TRV	No TRV
Endrin	72-20-8	No TRV			3.00E-01	3.00E-01
gamma-BHC (Lindane)	58-89-9	No TRV			2.00E+00	2.00E+00
Heptachlor	76-44-8	6.50E-02			No TRV	6.50E-02
Isodrin	465-73-6	No TRV			No TRV	No TRV
Methoxychlor	72-43-5	No TRV			No TRV	No TRV
Silvex (2,4,5-TP)	93-72-1	No TRV			No TRV	No TRV
Toxaphene	8001-35-2	No TRV			No TRV	No TRV
Inorganics						
Metals						
Aluminum	7429-90-5	1.00E+02			1.10E+02	1.00E+02
Antimony	7440-36-0	No TRV			No TRV	No TRV
Arsenic	7440-38-2	2.46E+00			5.14E+00	2.46E+00
Barium	7440-39-3	2.08E+01			2.08E+01	2.08E+01
Beryllium	7440-41-7	No TRV			No TRV	No TRV
Bismuth	7440-69-9	No TRV			No TRV	No TRV
Boron	7440-42-8	No TRV			2.88E+01	2.88E+01
Cadmium	7440-43-9	1.45E+00			1.45E+00	1.45E+00
Calcium	7440-70-2	No TRV			No TRV	No TRV
Chromium (and VI)	7440-47-3	1.00E+00			1.00E+00	1.00E+00
Cobalt	7440-48-4	No TRV			No TRV	No TRV
Copper	7440-50-8	4.70E+01			4.70E+01	4.70E+01
Iron	7439-89-6	No TRV			No TRV	No TRV
Lead	7439-92-1	2.50E-02			1.13E+00	2.50E-02

Table C3-4. Toxicity Reference Values (TRVs) for Birds

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
Lithium	7439-93-2	No TRV			No TRV	No TRV
Magnesium	7439-95-4	No TRV			No TRV	No TRV
Manganese	7439-96-5	No TRV			9.77E+02	9.77E+02
Mercury	7439-97-6	3.25E+00			4.50E-01	3.25E+00
Molybdenum	7439-98-7	No TRV			3.53E+00	3.53E+00
Nickel	7440-02-0	6.50E+01			7.74E+01	6.50E+01
Potassium	7440-09-7	No TRV			No TRV	No TRV
Rhodium	7440-16-6	No TRV			No TRV	No TRV
Selenium	7782-49-2	5.00E-01			5.00E-01	5.00E-01
Silicon	7440-21-3	No TRV			No TRV	No TRV
Silver	7440-22-4	1.78E+02			No TRV	1.78E+02
Sodium	7440-23-5	No TRV			No TRV	No TRV
Strontium	7440-24-6	No TRV			No TRV	No TRV
Tantalum	7440-25-7	No TRV			No TRV	No TRV
Thallium	7440-28-0	3.50E-01			No TRV	3.50E-01
Tin	7440-31-5	No TRV			No TRV	No TRV
Tungsten	7440-33-7	No TRV			No TRV	No TRV
Uranium	7440-61-1	No TRV			1.60E+01	1.60E+01
Vanadium	7440-62-2	No TRV			1.14E+01	1.14E+01
Yttrium	7440-65-5	No TRV			No TRV	No TRV
Zinc	7440-66-6	1.31E+02			1.45E+01	1.31E+02
Zirconium	7440-67-7	No TRV			No TRV	No TRV
Non-metals and Ions						
Ammonia/Ammonium	7664-41-7	No TRV			No TRV	No TRV
Bromide	24959-67-9	No TRV			No TRV	No TRV
Chloride	16887-00-6	No TRV			No TRV	No TRV
Cyanide	57-12-5	4.00E-02			No TRV	4.00E-02
Fluoride	16984-48-8	No TRV			7.80E+00	7.80E+00
Hydroxide	14280-30-9	No TRV			No TRV	No TRV
Iodine	7553-56-2	No TRV			No TRV	No TRV
Nitrate	14797-55-8	No TRV			No TRV	No TRV

Table C3-4. Toxicity Reference Values (TRVs) for Birds

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
Nitrite	14797-65-0	No TRV			No TRV	No TRV
Phosphate	14265-44-2	No TRV			No TRV	No TRV
Phosphorus	7723-14-0	No TRV			No TRV	No TRV
Sulfate	14808-79-8	No TRV			No TRV	No TRV
Total Sulfur	63705-05-5	No TRV			No TRV	No TRV
Criteria Pollutants						
Carbon dioxide	124-38-9	No TRV			No TRV	No TRV
Nitrogen dioxide	10102-44-0	No TRV			No TRV	No TRV
Ozone	10028-15-6	No TRV			No TRV	No TRV
Particulate matter	no cas #	No TRV			No TRV	No TRV
Sulfur dioxide	7446-09-5	No TRV			No TRV	No TRV
Radionuclides	TRV is not applicable to single radionuclides. Combined external and internal radiation exposure for birds from all radionuclides combined cannot exceed 0.1 rad/d.					

TRV = toxicity reference value

[a] Published in Appendix E of EPA (1999), Table E-8.

[b] Published in Table C3-3 of this work plan

[c] Order of preference is EPA (1999), then SAIC compilation

[d] Toxicity of Aroclor 1254 to ring dove used as representative of PCB mixtures

[e] Total exposure to all high molecular weight polycylic aromatic hydrocarbons combined is limited to 0.00014 mg/kg/d.

[f] Published value for benzo(k)fluoranthene used as a surrogate value.

[g] Published value for 1,1'-DDE (sic) used as a surrogate value.

Table C3-5. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
Organics											
Aromatic Halogenated Hydrocarbons											
2,3,4,6-Tetrachlorophenol	58-90-2	Rat	3.50E-01	2.50E+01	subchronic	NOAEL	Organ wt.	U.S. EPA (1986) in [3]	1.00E-01	1.00E+00	2.50E+00
4-Chloro-3-methylphenol	59-50-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Aromatic Nonhalogenated Hydrocarbons											
2-Nitrotoluene	88-72-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
4-Nitrobiphenyl	92-93-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Benzaldehyde	100-52-7	Rat	3.50E-01	1.43E+02	Subchronic	NOAEL	Forestomach lesions, kidney toxicity	Kluwe et al. (1983) in [4]	1.00E-01	1.00E+00	1.43E+01
Benzene	71-43-2	Mouse	3.00E-02	2.64E+02	chronic	LOAEL	Reproduction	Nawrot and Staples (1979) in [1]	1.00E+00	1.00E-01	2.64E+01
Benzyl alcohol	100-51-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Ethyl benzene	100-41-4	Rat	3.50E-01	9.71E+01	Subchronic	NOAEL	Liver and kidney toxicity	Wolf et al. (1956) in [3]	1.00E-01	1.00E+00	9.71E+00
m-Xylene	108-38-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
o-Xylene	95-47-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
p-Xylene	106-42-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Styrene	100-42-5	Dog	1.00E+01	2.00E+02	chronic	NOAEL	Liver toxicity	Quast et al. (1979) in [4]	1.00E+00	1.00E+00	2.00E+02
Toluene	108-88-3	Mouse	3.00E-02	2.60E+02	chronic	LOAEL	Reproduction	Nawrot and Staples (1979) in [1]	1.00E+00	1.00E-01	2.60E+01
Non-aromatic Nonhalogenated Hydrocarbons											
1,2-Epoxybutane	106-88-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,3-Butadiene	106-99-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,4-Dioxane	123-91-1	Rat	3.50E-01	5.00E-01	chronic	NOAEL	Reproduction	Giavini et al. (1985) in [1]	1.00E+00	1.00E+00	5.00E-01
1-Methylpropyl alcohol	78-92-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1-Nitropropane	108-03-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
2,2,4-Trimethylpentane	540-84-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
2-Butanone	78-93-3	Rat	3.50E-01	1.77E+03	Chronic	NOAEL	Decreased fetal birth weight	Cox et al. (1975) in [4]	1.00E+00	1.00E-01	1.77E+02
2-Butenaldehyde (2-Butenal)	4170-30-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
2-Ethoxyethanol	110-80-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV

Table C3-5. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
2-Heptanone	110-43-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
2-Hexanone	591-78-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
2-Methoxyethanol	109-86-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
2-Methyl-2-propanol	75-65-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
2-Methyl-2-propenenitrile	126-98-7	Dog	1.00E+01	3.40E-01	Subchronic	NOAEL	Increased SGOT and SGPT levels	Pozzani et al. (1968) in [4]	1.00E-01	1.00E+00	3.40E-02
2-Methylaziridine	75-55-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
2-Methylpropyl alcohol	78-83-1	Rat	3.50E-01	3.16E+02	subchronic	NOAEL	Hypoactivity and ataxia	U.S. EPA (1986) in [3]	1.00E-01	1.00E+00	3.16E+01
2-Pentanone	107-87-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
2-Propanone (Acetone)	67-64-1	Rat	3.50E-01	1.00E+02	subchronic	NOAEL	Liver and kidney toxicity	EPA (1986c) in [1]	1.00E-01	1.00E+00	1.00E+01
2-Propene-1-ol	107-18-6	Rat	3.50E-01	4.80E+00	subchronic	NOAEL	Increased liver and kidney weights	Carpanini et al. (1978) in [3]	1.00E-01	1.00E+00	4.80E-01
2-Propyl alcohol	67-63-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
3-Heptanone	106-35-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
3-Methyl-1-butanol	123-51-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
3-Methyl-2-butanone	563-80-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
3-Pentanone	96-22-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
4-Heptanone	123-19-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
4-Methyl-2-pentanone	108-10-1	Rat	3.50E-01	2.50E+02	subchronic	NOAEL	Liver/Kidney	Microbiological Associates (1986) in [1]	1.00E-01	1.00E+00	2.50E+01
4-Methyl-3-penten-2-one	141-79-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
5-Methyl-2-hexanone	110-12-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Acetaldehyde	75-07-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Acetamide	60-35-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Acetic acid	64-19-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Acetic acid ethyl ester	141-78-6	Rat	3.50E-01	9.00E+02	Subchronic	NOAEL	Mortality	U.S. EPA (1986) in [4]	1.00E-01	1.00E+00	9.00E+01
Acetic acid n-butyl ester	123-86-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Acetonitrile	75-05-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Acrolein	107-02-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Acrylonitrile	107-13-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Bis(isopropyl)ether	108-20-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV

Table C3-5. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
Butane	106-97-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Carbon disulfide	75-15-0	Rabbit	NA	1.10E+01	Chronic	NOAEL	Fetal toxicity/malformations	Hardin et al. (1981) in [3]	1.00E+00	1.00E+00	1.10E+01
Cyanogen	460-19-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Cyclohexane	110-82-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Cyclohexanone	108-94-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Cyclohexene	110-83-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Cyclopentane	287-92-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Ethyl alcohol	64-17-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Ethyl ether	60-29-7	Rat	3.50E-01	5.00E+02	subchronic	NOAEL	Depressed body weights	U.S. EPA (1986) in [3]	1.00E-01	1.00E+00	5.00E+01
Ethyl methacrylate	97-63-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Formaldehyde	50-00-0	Rat	3.50E-01	1.50E+01	Chronic	NOAEL	Histopathology	Til et al. (1989) in [4]	1.00E+00	1.00E-01	1.50E+00
Formamide	75-12-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Formic acid	64-18-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Formic acid, methyl ester	107-31-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Glycidylaldehyde	765-34-4	Rat	3.50E-01	1.09E+00	Subchronic	NOAEL	Retardation, enlarged adrenals, hydropic renal pelvis and hematopoietic effects	Hine et al. (1961) in [4]	1.00E-01	1.00E+00	1.09E-01
Methyl acetate	79-20-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Methyl alcohol	67-56-1	Rat	3.50E-01	5.00E+02	Subchronic	NOAEL	Increased SAP and SGPT, and decreased brain weight	U.S. EPA (1986) in [4]	1.00E-01	1.00E+00	5.00E+01
Methyl isocyanate	624-83-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Methyl methacrylate	80-62-6	Rat	3.50E-01	1.36E+02	chronic	NOAEL	NA	Borzelleca et al. (1964)	1.00E+00	1.00E+00	1.36E+02
Methyl tert-butyl ether	1634-04-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Methylacetylene	74-99-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Methylcyclohexane	108-87-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
N,N-Dimethylacetamide	127-19-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV

Table C3-5. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
n-Butyl alcohol	71-36-3	Rat	3.50E-01	1.25E+02	Subchronic	NOAEL	Hypoactivityand ataxia	U.S. EPA (1986) in [4]	1.00E-01	1.00E+00	1.25E+01
n-Heptane	142-82-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
n-Hexane	110-54-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Nitromethane	75-52-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
n-Nonane	111-84-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
n-Octane	111-65-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
n-Pentane	109-66-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
n-Propionaldehyde	123-38-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
n-Propyl alcohol	71-23-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
n-Valeraldehyde	110-62-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Oxirane	75-21-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
p-Cymene	99-87-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Phosgene	75-44-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Propargyl alcohol	107-19-7	Rat	3.50E-01	5.00E+00	Subchronic	NOAEL	Renal and hepatotoxicity	U.S. EPA (1987) in [4]	1.00E-01	1.00E+00	5.00E-01
Propionic acid	79-09-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Propionitrile	107-12-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Propylene gylcol monomethyl ether	107-98-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
p-tert-Butyltoluene	98-51-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Triethylamine	121-44-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Trimethylamine	75-50-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Vinyl acetate	108-05-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Non-aromatic Halogenated Hydrocarbons											
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,1,1,2-Tetrachloroethane	630-20-6	NA	NA	8.93E+01	Chronic	LOAEL	Mineralizaation of the kidneys in males, hepatic clear cell change in females	NTP (1983) in [4]	1.00E+00	1.00E-01	8.93E+00
1,1,1-Trichloroethane	71-55-6	Mouse	3.50E-02	1.00E+03	Chronic	NOAEL	Reproduction	Lane et al. (1982) in [1]	1.00E+00	1.00E+00	1.00E+03
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,1,2,2-Tetrachloroethane	79-34-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV

Table C3-5. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
1,1,2,2-Tetrachloroethene	127-18-4	Mouse	3.00E-02	1.40E+01	Chronic	NOAEL	Hepatotoxicity in mice	Buben and O'Flaherty (1985) in [4]	1.00E+00	1.00E-01	1.40E+00
1,1,2-Trichloroethane	79-00-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,1,2-Trichloroethylene	79-01-6	Mouse	3.00E-02	1.00E+02	subchronic	LOAEL	Hepatotoxicity in mice	Buben and O'Flaherty (1985) in [1]	0.07[a]	1.00E-01	7.00E-01
1,1-Dichloroethane	75-34-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,1-Dichloroethene	75-35-4	Rat	3.50E-01	3.00E+01	chronic	NOAEL	Mortality	Quast et al. (1983) in [1]	1.00E+00	1.00E+00	3.00E+01
1,2,2-Trichloro-1,1,2-trifluoroethane	76-13-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,2,3-Trichloropropane	96-18-4	Rat	3.50E-01	5.71E+00	Subchronic	NOAEL	Alterations in clinical chemistry and reduction in red cell mass	NTP (1983) in [4]	1.00E-01	1.00E+00	5.71E-01
1,2-Dibromo-3-chloropropane	96-12-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,2-Dichloroethane	107-06-2	Mouse	3.50E-02	5.00E+01	chronic	NOAEL	Reproduction	Lane et al. (1982) in [1]	1.00E+00	1.00E+00	5.00E+01
1,2-Dichloroethylene	540-59-0	Mouse	3.00E-02	4.52E+02	Subchronic	NOAEL	Body weight	Palmer et al. (1979) in [1]	1.00E-01	1.00E+00	4.52E+01
1,2-Dichloropropane	78-87-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,3-Dichloropropene	542-75-6	Rat	3.50E-01	3.00E+00	Subchronic	NOAEL	Increased organ weights	Dow Chemical (1973) in [4]	1.00E-01	1.00E+00	3.00E-01
1,4-Dichloro-2-butene	764-41-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1-Chloroethene	75-01-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
2,2-Dichloropropionic acid	75-99-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
2-Chloropropane	75-29-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
3-Chloropropene (Allyl chloride)	107-05-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Bromochloromethane	74-97-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Bromodichloromethane	75-27-4	Mouse	3.00E-02	1.79E+01	Chronic	LOAEL	Renal cytomegaly	NTP (1986) in [4]	1.00E+00	1.00E-01	1.79E+00
Bromoethene	593-60-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Bromoform	75-25-2	Rat	3.50E-01	1.79E+01	Subchronic	NOAEL	Hepatic lesions	NTP (1989) in [4]	1.00E-01	1.00E+00	1.79E+00
Bromomethane	74-83-9	Rat	3.50E-01	1.40E+00	Subchronic	NOAEL	Epithelial hyperplasia of the forestomach	Danse et al. (1984) in [4]	1.00E-01	1.00E+00	1.40E-01

Table C3-5. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
Carbon tetrachloride	56-23-5	Rat	3.50E-01	7.10E-01	Subchronic	NOAEL	Liver lesions	Bruckner et al. (1986) in [4]	1.00E-01	1.00E+00	7.10E-02
Chlorodibromomethane	124-48-1	Rat	3.50E-01	2.14E+01	Subchronic	NOAEL	Hepatic lesions	NTP (1985) in [4]	1.00E-01	1.00E+00	2.14E+00
Chlorodifluoromethane	75-45-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Chloroethane	75-00-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Chloroform	67-66-3	Rat	3.50E-01	1.50E+02	subchronic	NOAEL	Gonad atrophy	Palmer et al. (1979) in [1]	1.00E-01	1.00E+00	1.50E+01
Chloromethane	74-87-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Chloropentafluoroethane	76-15-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
cis-1,2-Dichloroethene	156-59-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
cis-1,3-Dichloropropene	10061-01-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Cyanogen bromide	506-68-3	Rat	3.50E-01	4.40E+01	Chronic	NOAEL	Thyroid effects and myelin degeneration	Philbrick et al. (1979) in [4]	1.00E+00	1.00E-01	4.40E+00
Cyanogen chloride	506-77-4	Rat	3.50E-01	2.53E+01	Chronic	NOAEL	Thyroid effects and myelin degeneration	Philbrick et al. (1979) in [4]	1.00E+00	1.00E-01	2.53E+00
Dichlorodifluoromethane	75-71-8	Rat	3.50E-01	1.50E+01	Chronic	NOAEL	Reduced body weight	Sherman (1974) in [4]	1.00E+00	1.00E-01	1.50E+00
Dichlorofluoromethane	75-43-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Dichloromethane	75-09-2	Rat	3.50E-01	6.16E+00	Chronic	NOAEL	Liver toxicity	National Coffee Association (1982) in [4]	1.00E+00	1.00E-01	6.16E-01
Difluorodibromomethane	75-61-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Hexafluoroacetone	684-16-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Iodomethane	74-88-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Methylene bromide	74-95-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Pentachloroethane	76-01-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
trans-1,2-Dichloroethylene	156-60-5	Mouse	3.00E-02	1.70E+01	Chronic	NOAEL	Increased serum alkaline phosphatase in male mice	Barnes et al. (1985) in [4]	1.00E+00	1.00E-01	1.70E+00
trans-1,3-Dichloropropene	10061-02-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Trichloroacetic acid	76-03-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Trichlorofluoroethane	27154-33-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Trichlorofluoromethane	75-69-4	Rat	3.50E-01	3.49E+02	Chronic	LOAEL	Survival and histopathology	NCI (1978) in [4]	1.00E+00	1.00E-01	3.49E+01

Table C3-5. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
Trifluorobromomethane	75-63-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Dioxin and Furan Compounds											
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin	35822-46-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin	39227-28-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin	57653-85-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	Rat	3.50E-01	1.60E-03	subchronic	NOAEL	Organ weight	Poiger et al. (1989) in [1]	1.00E-01	1.00E+00	1.60E-04
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin	19408-74-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,2,3,7,8-Pentachlorodibenzo-p-dioxin	40321-76-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	Rat	3.50E-01	1.60E-03	subchronic	NOAEL	Organ weight	Poiger et al. (1989) in [1]	1.00E-01	1.00E+00	1.60E-04
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	Rat	3.50E-01	1.60E-04	subchronic	NOAEL	Organ weight	Poiger et al. (1989) in [1]	1.00E-01	1.00E+00	1.60E-05
2,3,7,8-Tetrachlorodibenzo-p-dioxin	1746-01-6	Rat	3.50E-01	1.00E-06	chronic	NOAEL	Reproduction	Murray et al. (1979) in [1]	1.00E+00	1.00E+00	1.00E-06
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Dibenzofuran	132-64-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Octachlorodibenzo(p)dioxin	3268-87-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Octachlorodibenzofuran	39001-02-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
PCBs											
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
2',3,4,4',5-Pentachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV

Table C3-5. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
3,3',4,4',5-Pentachlorobiphenyl	no cas #	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Polychlorinated biphenyls (PCBs)[b]	1336-36-3	Oldfield mouse	1.40E-02	6.80E-01	chronic	LOAEL	Reproduction	McCoy et al. (1995) in [1]	1.00E+00	1.00E-01	6.80E-02
Phthalates											
Bis(2-ethylhexyl)phthalate	117-81-7	Mouse	3.00E-02	1.83E+01	chronic	NOAEL	Reproduction	Lamb et al. (1987) in [1]	1.00E+00	1.00E+00	1.83E+01
Butylbenzylphthalate	85-68-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Di-n-butylphthalate	84-74-2	Mouse	3.00E-02	5.50E+02	chronic	NOAEL	Reproduction	Lamb et al. (1987) in [1]	1.00E+00	1.00E+00	5.50E+02
Diethylphthalate	84-66-2	Mouse	3.00E-02	4.58E+03	chronic	NOAEL	Reproduction	Lamb et al. (1987) in [1]	1.00E+00	1.00E+00	4.58E+03
Dimethylphthalate	131-11-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
n-Dioctyl phthalate	117-84-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Light Polycyclic Aromatic Hydrocarbons											
2-Chloronapthalene	91-58-7	Mouse	3.00E-02	2.50E+02	Chronic	NOAEL	Dyspnea, abnormal appearance, liver enlargement	U.S. EPA (1989) in [4]	1.00E+00	1.00E+00	2.50E+02
2-Methylnaphthalene	91-57-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
5-Nitroacenaphthene	602-87-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Acenaphthene	83-32-9	Mouse	3.00E-02	1.75E+02	subchronic	NOAEL	Hepatotoxicity	U.S. EPA (1989) in [3]	1.00E-01	1.00E+00	1.75E+01
Acenaphthylene	208-96-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Anthracene	120-12-7	Mouse	3.00E-02	1.00E+03	Subchronic	NOAEL	NA	U.S. EPA (1989) in [3]	1.00E-01	1.00E+00	1.00E+02
Fluorene	86-73-7	NA	NA	1.25E+02	subchronic	NOAEL	Blood	U.S. EPA (1989) in [3]	1.00E-01	1.00E+00	1.25E+01
Indene	95-13-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Naphthalene	91-20-3	NA	NA	7.10E+01	Subchronic	NOAEL	Decreased body weights in males	BCL (1980a) in [4]	1.00E-01	1.00E+00	7.10E+00
Phenanthrene	85-01-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Pyrene	129-00-0	NA	NA	7.50E+01	Subchronic	NOAEL	Kidney effects	U.S. EPA (1989) in [4]	1.00E-01	1.00E+00	7.50E+00

Table C3-5. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
Heavy Polycyclic Aromatic Hydrocarbons											
3-Methylcholanthrene	56-49-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
5-Methylchrysene	3697-24-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Benzo(a)anthracene	56-55-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Benzo(a)pyrene	50-32-8	Mouse	3.00E-02	1.00E+01	chronic	LOAEL	Reproduction	Mackenzie and Angevine (1981) in [1]	1.00E+00	1.00E-01	1.00E+00
Benzo(b)fluoranthene	205-99-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Benzo(e)pyrene	192-97-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Benzo(g,h,i)perylene	191-24-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Benzo(j)fluoranthene	205-82-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Benzo(k)fluoranthene	207-08-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Benzo[a,i]pyrene	191-30-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Chrysene	218-01-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Dibenzo(a,h)anthracene	53-70-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Dibenz[a,h]acridine	226-36-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Dibenz[a,j]acridine	224-42-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Dibenzo(a,e)fluoranthene	5385-75-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Dibenzo(a,h)fluoranthene	no cas #	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Dibenzo[a,e]pyrene	192-65-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Dibenzo[a,h]pyrene	189-64-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Dibenzo[a,i]pyrene	189-55-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Fluoranthene	206-44-0	Mouse	3.00E-02	1.25E+02	Subchronic	NOAEL	Nephropathy, hematological alterations, and clinical effects	U.S. EPA (1988) in [4]	1.00E-01	1.00E+00	1.25E+01
Hexachloronaphthalene	1335-87-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Indeno(1,2,3-cd)pyrene	193-39-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Octachloronaphthalene	2234-13-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Pentachloronaphthalene	1321-64-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Tetrachloronaphthalene	1335-88-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Trichloronaphthalene	1321-65-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV

Table C3-5. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
Light Substituted Benzene Compounds											
1,2,3-Trichlorobenzene	87-61-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,2,4-Trichlorobenzene	120-82-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,2,4-Trimethyl benzene	95-63-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,2-Dichlorobenzene	95-50-1	Rat	3.50E-01	8.57E+01	Chronic	NOAEL	No adverse effects observed	NTP (1985 in [4]	1.00E+00	1.00E+00	8.57E+01
1,3,5-Trimethyl benzene	108-67-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,3-Dichlorobenzene	541-73-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,3-Dinitrobenzene	99-65-0	Rat	4.50E-01	1.13E+00	subchronic	NOAEL	Reproduction	Cody et al. (1981) in [4]	1.00E-01	NA	1.10E-01
1,4-Dichlorobenzene	106-46-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,4-Dinitrobenzene	100-25-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
2,4,5-Trichlorophenol	95-95-4	Rat	3.50E-01	1.00E+02	Subchronic	NOAEL	Liver and kidney pathology	McCollister et al. (1961) in [4]	1.00E-01	1.00E+00	1.00E+01
2,4,6-Trichlorophenol	88-06-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
2,4-Dichlorophenol	120-83-2	Rat	3.50E-01	3.00E-01	Subchronic	NOAEL	Decreased delayed hypersensitivity response	Exon and Koller (1985) in [4]	1.00E-01	1.00E+00	3.00E-02
2,4-Dimethylphenol	105-67-9	Mouse	3.00E-02	5.00E+00	Subchronic	NOAEL	Lethargy, prostration, and ataxia	U.S. EPA (1989) in [3]	1.00E-01	1.00E+00	5.00E-01
2,4-Dinitrophenol	51-28-5	Human	7.00E+01	2.00E+00	Chronic	LOAEL	Cataract formation	Horner (1942) in [4]	1.00E+00	1.00E-01	2.00E-01
2,4-Dinitrotoluene	121-14-2	Mouse	3.00E-02	1.35E+01	chronic	NOAEL	Reproduction	Ellis et al. (1979)	1.00E+00	1.00E+00	1.35E+01
2,6-Dinitrotoluene	606-20-2	Rat	3.50E-01	7.00E+00	subchronic	NOAEL	Reproduction	ATSDR (1989)	1.00E-01	1.00E+00	7.00E-01
2-Chlorophenol	95-57-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
2-Chlorotoluene	95-49-8	Rat	3.50E-01	2.00E+01	Chronic	NOAEL	Decrease in body weight gain	Gibson et al. (1974a) in [4]	1.00E+00	1.00E+00	2.00E+01
2-Nitrophenol	88-75-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
4,6-Dinitro-o-cresol	534-52-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
4-Chlorotoluene	106-43-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
4-Nitrophenol	100-02-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
alpha-Methylstyrene	98-83-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Aniline	62-53-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Benzotrichloride	98-07-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV

Table C3-5. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
Benzyl chloride	100-44-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Bromobenzene	108-86-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Chlorobenzene	108-90-7	Dog	1.00E+01	1.90E+01	chronic	NOAEL	Histopathologic changes in liver	Monsanto Co. (1967a; Knapp et al. (1971) in [3]	1.00E+00	1.00E+00	1.90E+01
Cumene	98-82-8	Rat	3.50E-01	1.10E+02	Chronic	NOAEL	Increased average kidney weight in female rats	Wolf et al. (1956) in [4]	1.00E+00	1.00E+00	1.10E+02
m-Cresol	108-39-4	Rat	3.50E-01	5.00E+01	Subchronic	NOAEL	Decreased body weights and neurotoxicity	U.S. EPA (1986 (1987) in [4]	1.00E-01	1.00E+00	5.00E+00
n-Butyl benzene	104-51-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Nitrobenzene	98-95-3	Rat/Mouse	3.50E-01	4.60E+00	Subchronic	LOAEL	Hematologic, adrenal, renal and hepatic lesions	CIIT (1984) in [4]	1.00E-01	1.00E-01	4.60E-02
n-Propyl benzene	103-65-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
o-Cresol	95-48-7	Rat	3.50E-01	5.00E+01	Subchronic	NOAEL	Decreased body weights and neurotoxicity	U.S. EPA (1986 (1987) in [4]	1.00E-01	1.00E+00	5.00E+00
o-Dinitrobenzene	528-29-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
o-Nitroaniline	88-74-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
o-Toluidine	95-53-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
p-Chloroaniline	106-47-8	Rat	3.50E-01	1.25E+01	Chronic	LOAEL	Nonneoplastic lesions of splenic capsule	NCI (1979) in [4]	1.00E+00	1.00E-01	1.25E+00
p-Cresol	106-44-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Phenol	108-95-2	Rat	3.50E-01	6.00E+01	Chronic	NOAEL	Reduced fetal body weight in rats	NTP (1983) in [4]	1.00E+00	1.00E+00	6.00E+01
p-Nitrochlorobenzene	100-00-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
p-Toluidine	106-49-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
sec-Butyl benzene	135-98-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
tert-Butyl benzene	98-06-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Toluene-2,6-diamine	823-40-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Trimethyl benzene	25551-13-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Other Light Semivolatile Compounds											
1,1'-Biphenyl	92-52-4	Rat	3.50E-01	5.00E+01	Chronic	NOAEL	Kidney damage	Ambrose et al. (1960) in [4]	1.00E+00	1.00E-01	5.00E+00

Table C3-5. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
1,1-Dimethylhydrazine	57-14-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,2-Dimethylhydrazine	540-73-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,2-Diphenylhydrazine	122-66-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
1,3-Propane sultone	1120-71-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
2,4-Toluene diisocyante	584-84-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
2-Chloroacetophenone	532-27-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
2-Propenoic acid	79-10-7	Rat	3.50E-01	5.30E+01	Chronic	NOAEL	Reduced pup weight	BASF (1993) in [4]	1.00E+00	1.00E-01	5.30E+00
4,4-Methylenedianiline	101-77-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Acetophenone	98-86-2	Rat	3.50E-01	4.23E+02	Subchronic	NOAEL	General toxicity	Hagen et al. (1967) in [4]	1.00E-01	1.00E+00	4.23E+01
Benzoic acid	65-85-0	Mouse	3.00E-02	4.00E+01	chronic	LOAEL	unknown	Shtenberg and Ignat'ev (1970) in [3]	1.00E+00	1.00E-01	4.00E+00
Bis(2-chloroethoxy)methane	111-91-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Bis(2-chloroethyl) ether	111-44-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Chlorocyclopentadiene	41851-50-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Cyclohexanol	108-93-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Dichloroisopropyl ether	108-60-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Dichloromethyl ether	542-88-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Dichloropentadiene	no cas #	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Dimethyl sulfate	77-78-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Dimethylaniline	121-69-7	Mouse	3.00E-02	2.23E+01	Subchronic	LOAEL	Splenomegalyand hematopoiesis	Abdo et al. (1984) in [3]	1.00E-01	1.00E-01	2.23E-01
di-n-Propylnitrosamine	621-64-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Diphenyl ether	101-84-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Epichlorohydrin	106-89-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Ethyl Carbamate (Urethane)	51-79-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Ethyl methanesulfonate	62-50-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Ethylene dibromide	106-93-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Ethylene glycol	107-21-1	Rat	3.50E-01	2.00E+02	Chronic	NOAEL	Kidney toxicity	DePass et al. (1986a) in [4]	1.00E+00	1.00E-01	2.00E+01
Ethylene glycol monobutyl ether	111-76-2	Rat/Mouse	3.50E-01	5.10E+00	Subchronic	NOAEL	Changes in mean corpuscular	NTP (1993) in [4]	1.00E-01	1.00E+00	5.10E-01
Ethylene glycol monoethyl ether acetate	111-15-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV

Table C3-5. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
Ethylene thiourea	96-45-7	Rat	3.50E-01	2.50E-01	Chronic	LOAEL	Increased incidence of thyroid hyperplasia	Graham et al. (1975) in [4]	1.00E+00	1.00E-01	2.50E-02
Furfural	98-01-1	Rat	3.50E-01	7.90E+00	Subchronic	LOAEL	Mild hepatocellular vacuolization	NTP (1981a) in [4]	1.00E+00	1.00E-01	7.90E-01
Maleic hydrazide	123-33-1	Rat	3.50E-01	5.00E+02	chronic	LOAEL	Renal dysfunction	Uniroyal Chemical Co. (1981) in [3]	1.00E+00	1.00E-01	5.00E+01
Malononitrile	109-77-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Methyl styrene (mixed isomers)	25013-15-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Methylhydrazine	60-34-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
N,N-Diphenylamine	122-39-4	Dog	1.00E+01	2.50E+00	Chronic	NOAEL	Increased liver and kidney weights	Thomas et. al. (1967) in [4]	1.00E+00	1.00E-01	2.50E-01
Nitric acid, propyl ester	627-13-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
N-Nitrosodi-n-butylamine	924-16-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
N-Nitrosomorpholine	59-89-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
N-Nitroso-N,N-dimethylamine	62-75-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
o-Anisidine	90-04-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Oxalic acid	144-62-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Phthalic anhydride	85-44-9	Mouse	3.00E-02	1.56E+03	Chronic	LOAEL	Lung and kidney histopathology	NCI (1979) in [4]	1.00E+00	1.00E-01	1.56E+02
p-Phthalic acid	100-21-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Pyridine	110-86-1	Rat	3.50E-01	1.00E+00	Chronic	NOAEL	Increased liver weight	U.S. EPA (1986 in [4]	1.00E+00	1.00E-01	1.00E-01
Quinoline	91-22-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Quinone	106-51-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Safrole	94-59-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Tetrahydrofuran	109-99-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Other Heavy Semivolatile Compounds											
1,2,4,5-Tetrachlorobenzene	95-94-3	Rat	3.50E-01	3.40E-01	Subchronic	NOAEL	Kidney lesions	Chu et al. (1984) in [4]	1.00E-01	1.00E+00	3.40E-02
1,3,5-Trinitrobenzene	99-35-4	Mouse	1.85E-02	6.74E+01	subchronic	NOAEL	Reproduction	Pathology Associates Inc. (1994) in [4]	1.00E-01	NA	6.70E+00
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	Human	7.00E+01	2.00E+00	Subchronic	LOAEL	Cataract formation	Horner (1942) in [4]	1.00E+00	1.00E-01	2.00E-01

Table C3-5. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
2-sec-Butyl-4,6-dinitrophenol	88-85-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
3,3-Dichlorobenzidine	91-94-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
3,3'-Dimethoxybenzidine	119-90-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
4-Bromophenylphenyl ether	101-55-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Ammonium perfluorooctanoate	3825-26-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Azobenzene	103-33-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Chlorobenzilate	510-15-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Dibutylphosphate	107-66-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Dimethyl aminoazobenzene	60-11-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Hexachlorobenzene	118-74-1	Rat	3.50E-01	8.00E-02	Chronic	NOAEL	Liver effects	Arnold et al. (1985) in [4]	1.00E+00	1.00E-01	8.00E-03
Hexachlorobutadiene	87-68-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Hexachlorocyclopentadiene	77-47-4	Rat	3.50E-01	7.00E+00	Subchronic	NOAEL	Stomach lesions	Abdo et al. (1984) in [4]	1.00E-01	1.00E+00	7.00E-01
Hexachloroethane	67-72-1	Rat	3.50E-01	1.00E+00	Subchronic	NOAEL	Atrophy and degeneration of the renal tubules	Gorzinski et al. (1985) in [4]	1.00E-01	1.00E+00	1.00E-01
Hexachlorophene	70-30-4	Dog	1.00E+01	7.50E-01	Chronic	LOAEL	Swollen salivary lands, status spongiosis in brain and optic nerve	Nationwide Chemical Corp. (1974) in [4]	1.00E+00	1.00E-01	7.50E-02
Hexamethylene-1,5-diisocyanate	822-06-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Mirex	2385-85-5	Rat	3.50E-01	7.00E-02	chronic	NOAEL	Liver cytomegaly	NTP (1990) in [3]	1.00E+00	1.00E+00	7.00E-02
Nitrofen	1836-75-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Pentachlorobenzene	608-93-5	Rat	3.50E-01	8.30E+00	Subchronic	LOAEL	Liver and kidney toxicity	Linder et al. (1980) in [4]	1.00E+00	1.00E-01	8.30E-01
Pentachloronitrobenzene	82-68-8	Dog	1.00E+01	7.50E-01	Chronic	NOAEL	Liver toxicity	Olin Mathieson Corp. (1968a) in [4]	1.00E+00	1.00E-01	7.50E-02
Pentachlorophenol	87-86-5	Rat	3.50E-01	2.40E-01	chronic	NOAEL	Reproduction	Schwetz et al. (1978) in [1]	1.00E+00	1.00E+00	2.40E-01
Picric acid	88-89-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Pronamide	23950-58-5	Dog	1.00E+01	7.50E+00	Chronic	NOAEL	No effects	Rohm & Hass, Co. (1970a) in [4]	1.00E+00	1.00E-01	7.50E-01

Table C3-5. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
Strychnine	57-24-9	Rat	3.50E-01	2.50E+00	Subchronic	LOAEL	Toxicity/histopathology	Seidl and Zbinden (1982) in [4]	1.00E+00	1.00E-01	2.50E-01
Terphenyls	26140-60-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Tributyl phosphate	126-73-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Trifluralin	1582-09-8	Dog	1.00E+01	7.50E-01	chronic	NOAEL	Increased liver weights	Hoechst Aktiengesellschaft (1984a) in [3]	1.00E-01	1.00E+00	7.50E-02
Triphenylamine	603-34-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Herbicides and Organochlorinated Pesticides											
2,4,5-Trichlorophenoxyacetic acid (2,4,5-T)	93-76-5	Rat	3.50E-01	3.00E+00	chronic	NOAEL	Urinary coproporphyrins	Smith et al. (1981) in [3]	1.00E+00	1.00E+00	3.00E+00
2,4-D	94-75-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
4,4'-DDD	72-54-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
4,4'-DDE	72-55-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
4,4'-DDT	50-29-3	Rat	3.50E-01	8.00E-01	chronic	NOAEL	Reproduction	Fitzhugh (1948) in [1]	1.00E+00	1.00E+00	8.00E-01
Aldrin	309-00-2	Rat	3.50E-01	2.00E-01	chronic	NOAEL	Reproduction	EPA (1988a) in [1]	1.00E+00	1.00E+00	2.00E-01
alpha-BHC	319-84-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
beta-BHC	319-85-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Captan	133-06-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Chlordane	57-74-9	Mouse	3.00E-02	4.58E+00	chronic	NOAEL	Reproduction	Keplinger et al. (1968) in [1]	1.00E+00	1.00E+00	4.58E+00
delta-BHC	319-86-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Dieldrin	60-57-1	Rat	3.50E-01	2.00E-01	Chronic	Loael	Reproduction	Treon and Cleveland (1955) in [1]	1.00E+00	1.00E-01	2.00E-02
Endothall	145-73-3	Dog	1.00E+01	2.00E+00	Chronic	NOAEL	Increased weight of stomach and small intestine	Pennwalt Agchem. (1965) in [4]	1.00E+00	1.00E+00	2.00E+00
Endrin	72-20-8	Mouse	3.00E-02	9.20E-01	chronic	LOAEL	Reproduction	Good and Ware (1969) in [1]	1.00E+00	1.00E-01	9.20E-02
gamma-BHC (Lindane)	58-89-9	Rat	3.50E-01	8.00E+00	chronic	NOAEL	Reproduction	Palmer et al. (1978) in [1]	1.00E+00	1.00E+00	8.00E+00
Heptachlor	76-44-8	Mink	1.00E+00	1.00E+00	chronic	LOAEL	Reproduction	Crumet al. (1993) in [1]	1.00E+00	1.00E-01	1.00E-01
Isodrin	465-73-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV

Table C3-5. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
Methoxychlor	72-43-5	NA	NA	5.01E+00	chronic	NOAEL	Excessive loss of litters	Kincaid Enterprises (1986) in [4]	1.00E+00	1.00E+00	5.01E+00
Silvex (2,4,5-TP)	93-72-1	Dog	1.00E+01	7.50E-01	chronic	NOAEL	Liver	Mullison (1966, Gehring and Betso (1978)) in [3]	1.00E+00	1.00E+00	7.50E-01
Toxaphene	8001-35-2	Rat	3.50E-01	8.00E+00	Chronic	NOAEL	Reproduction	Kennedy et al. (1973) in [1]	1.00E+00	1.00E+00	8.00E+00
Inorganics											
Metals											
Aluminum	7429-90-5	Mouse	3.00E-02	1.93E+01	chronic	LOAEL	Reproduction	Ondreicka et al. (1966) in [1]	1.00E+00	1.00E-01	1.93E+00
Antimony	7440-36-0	Mouse	3.00E-02	1.25E+00	chronic	LOAEL	Longevity	Schroeder et al. (1968b) in [1]	1.00E+00	1.00E-01	1.25E-01
Arsenic	7440-38-2	Mouse	3.00E-02	1.26E+00	chronic	LOAEL	Reproduction	Schroeder and Mitchner (1971) in [1]	1.00E+00	1.00E-01	1.26E-01
Barium	7440-39-3	Rat	4.35E-01	5.06E+00	chronic	NOAEL	Growth	Perry et al. (1983) in [1]	1.00E+00	1.00E+00	5.06E+00
Beryllium	7440-41-7	Rat	3.50E-01	6.60E-01	chronic	NOAEL	Longevity	Schroeder and Mitchner (1975) in [1]	1.00E+00	1.00E+00	6.60E-01
Bismuth	7440-69-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Boron	7440-42-8	Rat	3.50E-01	2.80E+01	chronic	NOAEL	Reproduction	Weir and Fisher (1972) in [1]	1.00E+00	1.00E+00	2.80E+01
Cadmium	7440-43-9	Rat	3.03E-01	1.00E+00	chronic	NOAEL	Reproduction	Sutou et al. (1980b) in [1]	1.00E+00	1.00E+00	1.00E+00
Calcium	7440-70-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Chromium (and VI)	7440-47-3	Rat	3.50E-01	2.74E+03	chronic	NOAEL	Reproduction	Ivankovic and Preussmann (1975) in [1]	1.00E+00	1.00E+00	2.74E+03
Cobalt	7440-48-4	Rat	3.50E-01	1.00E+00	subchronic	NOAEL	Mortality	Underhill et al. (1931) in [2]	1.00E-01	1.00E+00	1.00E-01
Copper	7440-50-8	Mink	1.00E+00	1.17E+01	chronic	NOAEL	Reproduction	Aulerich et al. (1982) in [1]	1.00E+00	1.00E+00	1.17E+01
Iron	7439-89-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Lead	7439-92-1	Rat	3.50E-01	8.00E+00	chronic	NOAEL	Reproduction	Azar et al. (1973) in [1]	1.00E+00	1.00E+00	8.00E+00
Lithium	7439-93-2	Rat	3.50E-01	9.39E+00	Chronic	NOAEL	Reproduction	Marathe and Thomas (1986) in [1]	1.00E+00	1.00E+00	9.39E+00
Magnesium	7439-95-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV

Table C3-5. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
Manganese	7439-96-5	Rat	3.50E-01	8.80E+01	chronic	NOAEL	Reproduction	Laskey et al. (1982) in [1]	1.00E+00	1.00E+00	8.80E+01
Mercury	7439-97-6	Mink	1.00E+00	1.01E+00	chronic	NOAEL	Reproduction	Aulerich et al. (1974) in [1]	1.00E+00	1.00E+00	1.01E+00
Molybdenum	7439-98-7	Mouse	3.00E-02	2.58E+00	chronic	LOAEL	Reproduction	Schroeder and Mitchner (1971) in [1]	1.00E+00	1.00E-01	2.58E-01
Nickel	7440-02-0	Rat	3.50E-01	4.00E+01	chronic	NOAEL	Reproduction	Ambrose et al. (1976) in [1]	1.00E+00	1.00E+00	4.00E+01
Potassium	7440-09-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Rhodium	7440-16-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Selenium	7782-49-2	Rat	3.50E-01	2.00E-01	chronic	NOAEL	Reproduction	Rosenfeld and Beath (1954) in [1]	1.00E+00	1.00E+00	2.00E-01
Silicon	7440-21-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Silver	7440-22-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Sodium	7440-23-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Strontium	7440-24-6	Rat	3.50E-01	2.63E+02	Chronic	NOAEL	Body weight	Skoryna (1981) in [1]	1.00E+00	1.00E+00	2.63E+02
Tantalum	7440-25-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Thallium	7440-28-0	Rat	3.65E-01	7.40E-01	subchronic	LOAEL	Reproduction	Formigli et al. (1986) in [1]	1.00E-01	1.00E-01	7.40E-03
Tin	7440-31-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Tungsten	7440-33-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Uranium	7440-61-1	Mouse	3.00E-02	3.07E+00	Chronic	NOAEL	Reproduction	Paternain et al. (1989) in [1]	1.00E+00	1.00E+00	3.07E+00
Vanadium	7440-62-2	Rat	2.60E-01	2.10E+00	chronic	LOAEL	Reproduction	Domingo et al. (1986) in [1]	1.00E+00	1.00E-01	2.10E-01
Yttrium	7440-65-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Zinc	7440-66-6	Rat	3.50E-01	1.60E+02	chronic	NOAEL	Reproduction	Schlicker and Cox (1968) in [1]	1.00E+00	1.00E+00	1.60E+02
Zirconium	7440-67-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Non-metals and Ions											
Ammonia/Ammonium	7664-41-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Bromide	24959-67-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Chloride	16887-00-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV

Table C3-5. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF
Cyanide	57-12-5	Rat	2.73E-01	6.87E+01	chronic	NOAEL	Reproduction	Tewe and Maner (1981) in [1]	1.00E+00	1.00E+00	6.87E+01
Fluoride	16984-48-8	Mink	1.00E+00	3.14E+01	Chronic	NOAEL	Reproduction	Aulerich et al. (1987) in [1]	1.00E+00	1.00E+00	3.14E+01
Hydroxide	14280-30-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Iodine	7553-56-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Nitrate	14797-55-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Nitrite	14797-65-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Phosphate	14265-44-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Phosphorus	7723-14-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Sulfate	14808-79-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Total Sulfur	63705-05-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Criteria Pollutants											
Carbon dioxide	124-38-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Nitrogen dioxide	10102-44-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Ozone	10028-15-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Particulate matter	no cas #	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Sulfur dioxide	7446-09-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	No TRV
Radionuclides	TRV is not applicable to single radionuclides. Combined external and internal radiation exposure for mammals from all radionuclides combined cannot exceed 0.1 rad/d.										

TRV = toxicity reference value

DCF = Duration conversion factor; 1 if chronic, 0.1 if subchronic (Sample et al. 1996)

LOAEL = Lowest observed adverse effect level

[1] = Sample et al. (1996)

Table C3-5. Derivation of SAIC-Compiled Toxicity Reference Values (TRVs) for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Test Species Body Weight (kg) BWt	Benchmark (mg/kgBW/d)	Test Duration	Endpoint	Effect	Source	Duration Conversion Factor DCF	Endpoint Conversion Factor ECF	TRV (mg/kgBW/d) benchmark x DCF x ECF

ECF = Endpoint conversion factor; 1 if NOAEL, 0.1 if LOAEL (Sample et al. 1996)

NOAEL = No observed adverse effect level

NA = No data available

[a] Subchronic-to-chronic conversion factor of 0.1 multiplied by 0.7 to adjust for 5 d/wk exposure.

[b] Data for Aroclor 1254 used as representative of PCB mixtures

[2] = Clayton and Clayton (1981)

[3] = IRIS (1996)

[4] = IRIS (1998)

Table C3-6. Toxicity Reference Values (TRVs) for Mammals

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
Organics						
Aromatic Halogenated Hydrocarbons						
2,3,4,6-Tetrachlorophenol	58-90-2	No TRV			2.50E+00	2.50E+00
4-Chloro-3-methylphenol	59-50-7	No TRV			No TRV	No TRV
Aromatic Nonhalogenated Hydrocarbons						
2-Nitrotoluene	88-72-2	No TRV			No TRV	No TRV
4-Nitrobiphenyl	92-93-3	No TRV			No TRV	No TRV
Benzaldehyde	100-52-7	No TRV			1.43E+01	1.43E+01
Benzene	71-43-2	No TRV			2.64E+01	2.64E+01
Benzyl alcohol	100-51-6	No TRV			No TRV	No TRV
Ethyl benzene	100-41-4	No TRV			9.71E+00	9.71E+00
m-Xylene	108-38-3	No TRV			No TRV	No TRV
o-Xylene	95-47-6	No TRV			No TRV	No TRV
p-Xylene	106-42-3	No TRV			No TRV	No TRV
Styrene	100-42-5	No TRV			2.00E+02	2.00E+02
Toluene	108-88-3	No TRV			2.60E+01	2.60E+01
Non-aromatic Nonhalogenated Hydrocarbons						
1,2-Epoxybutane	106-88-7	No TRV			No TRV	No TRV
1,3-Butadiene	106-99-0	No TRV			No TRV	No TRV
1,4-Dioxane	123-91-1	1.06E+02	d		5.00E-01	1.06E+02
1-Methylpropyl alcohol	78-92-2	No TRV			No TRV	No TRV
1-Nitropropane	108-03-2	No TRV			No TRV	No TRV
2,2,4-Trimethylpentane	540-84-1	No TRV			No TRV	No TRV
2-Butanone	78-93-3	No TRV			1.77E+02	1.77E+02
2-Butenaldehyde (2-Butenal)	4170-30-3	No TRV			No TRV	No TRV
2-Ethoxyethanol	110-80-5	No TRV			No TRV	No TRV
2-Heptanone	110-43-0	No TRV			No TRV	No TRV
2-Hexanone	591-78-6	No TRV			No TRV	No TRV
2-Methoxyethanol	109-86-4	No TRV			No TRV	No TRV
2-Methyl-2-propanol	75-65-0	No TRV			No TRV	No TRV

Table C3-6. Toxicity Reference Values (TRVs) for Mammals

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
2-Methyl-2-propenenitrile	126-98-7	No TRV			3.40E-02	3.40E-02
2-Methylaziridine	75-55-8	No TRV			No TRV	No TRV
2-Methylpropyl alcohol	78-83-1	No TRV			3.16E+01	3.16E+01
2-Pentanone	107-87-9	No TRV			No TRV	No TRV
2-Propanone (Acetone)	67-64-1	1.00E+01	d		1.00E+01	1.00E+01
2-Propene-1-ol	107-18-6	No TRV			4.80E-01	4.80E-01
2-Propyl alcohol	67-63-0	No TRV			No TRV	No TRV
3-Heptanone	106-35-4	No TRV			No TRV	No TRV
3-Methyl-1-butanol	123-51-3	No TRV			No TRV	No TRV
3-Methyl-2-butanone	563-80-4	No TRV			No TRV	No TRV
3-Pentanone	96-22-0	No TRV			No TRV	No TRV
4-Heptanone	123-19-3	No TRV			No TRV	No TRV
4-Methyl-2-pentanone	108-10-1	No TRV			2.50E+01	2.50E+01
4-Methyl-3-penten-2-one	141-79-7	No TRV			No TRV	No TRV
5-Methyl-2-hexanone	110-12-3	No TRV			No TRV	No TRV
Acetaldehyde	75-07-0	No TRV			No TRV	No TRV
Acetamide	60-35-5	No TRV			No TRV	No TRV
Acetic acid	64-19-7	No TRV			No TRV	No TRV
Acetic acid ethyl ester	141-78-6	No TRV			9.00E+01	9.00E+01
Acetic acid n-butyl ester	123-86-4	No TRV			No TRV	No TRV
Acetonitrile	75-05-8	No TRV			No TRV	No TRV
Acrolein	107-02-8	No TRV			No TRV	No TRV
Acrylonitrile	107-13-1	4.60E-01	d		No TRV	4.60E-01
Bis(isopropyl)ether	108-20-3	No TRV			No TRV	No TRV
Butane	106-97-8	No TRV			No TRV	No TRV
Carbon disulfide	75-15-0	No TRV			1.10E+01	1.10E+01
Cyanogen	460-19-5	No TRV			No TRV	No TRV
Cyclohexane	110-82-7	No TRV			No TRV	No TRV
Cyclohexanone	108-94-1	No TRV			No TRV	No TRV
Cyclohexene	110-83-8	No TRV			No TRV	No TRV

Table C3-6. Toxicity Reference Values (TRVs) for Mammals

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
Cyclopentane	287-92-3	No TRV			No TRV	No TRV
Ethyl alcohol	64-17-5	No TRV			No TRV	No TRV
Ethyl ether	60-29-7	No TRV			5.00E+01	5.00E+01
Ethyl methacrylate	97-63-2	No TRV			No TRV	No TRV
Formaldehyde	50-00-0	2.30E+00	d		1.50E+00	2.30E+00
Formamide	75-12-7	No TRV			No TRV	No TRV
Formic acid	64-18-6	No TRV			No TRV	No TRV
Formic acid, methyl ester	107-31-3	No TRV			No TRV	No TRV
Glycidylaldehyde	765-34-4	No TRV			1.09E-01	1.09E-01
Methyl acetate	79-20-9	No TRV			No TRV	No TRV
Methyl alcohol	67-56-1	No TRV			5.00E+01	5.00E+01
Methyl isocyanate	624-83-9	No TRV			No TRV	No TRV
Methyl methacrylate	80-62-6	No TRV			1.36E+02	1.36E+02
Methyl tert-butyl ether	1634-04-4	No TRV			No TRV	No TRV
Methylacetylene	74-99-7	No TRV			No TRV	No TRV
Methylcyclohexane	108-87-2	No TRV			No TRV	No TRV
N,N-Dimethylacetamide	127-19-5	No TRV			No TRV	No TRV
n-Butyl alcohol	71-36-3	No TRV			1.25E+01	1.25E+01
n-Heptane	142-82-5	No TRV			No TRV	No TRV
n-Hexane	110-54-3	No TRV			No TRV	No TRV
Nitromethane	75-52-5	No TRV			No TRV	No TRV
n-Nonane	111-84-2	No TRV			No TRV	No TRV
n-Octane	111-65-9	No TRV			No TRV	No TRV
n-Pentane	109-66-0	No TRV			No TRV	No TRV
n-Propionaldehyde	123-38-6	No TRV			No TRV	No TRV
n-Propyl alcohol	71-23-8	No TRV			No TRV	No TRV
n-Valeraldehyde	110-62-3	No TRV			No TRV	No TRV
Oxirane	75-21-8	No TRV			No TRV	No TRV
p-Cymene	99-87-6	No TRV			No TRV	No TRV
Phosgene	75-44-5	No TRV			No TRV	No TRV

Table C3-6. Toxicity Reference Values (TRVs) for Mammals

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
Propargyl alcohol	107-19-7	No TRV			5.00E-01	5.00E-01
Propionic acid	79-09-4	No TRV			No TRV	No TRV
Propionitrile	107-12-0	No TRV			No TRV	No TRV
Propylene gylcol monomethyl ether	107-98-2	No TRV			No TRV	No TRV
p-tert-Butyltoluene	98-51-1	No TRV			No TRV	No TRV
Triethylamine	121-44-8	No TRV			No TRV	No TRV
Trimethylamine	75-50-3	No TRV			No TRV	No TRV
Vinyl acetate	108-05-4	No TRV			No TRV	No TRV
Non-aromatic Halogenated Hydrocarbons						
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	No TRV			No TRV	No TRV
1,1,1,2-Tetrachloroethane	630-20-6	No TRV			8.93E+00	8.93E+00
1,1,1-Trichloroethane	71-55-6	No TRV			1.00E+03	1.00E+03
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	No TRV			No TRV	No TRV
1,1,2,2-Tetrachloroethane	79-34-5	No TRV			No TRV	No TRV
1,1,2,2-Tetrachloroethene	127-18-4	No TRV			1.40E+00	1.40E+00
1,1,2-Trichloroethane	79-00-5	No TRV			No TRV	No TRV
1,1,2-Trichloroethylene	79-01-6	No TRV			1.00E+00	1.00E+00
1,1-Dichloroethane	75-34-3	No TRV			No TRV	No TRV
1,1-Dichloroethene	75-35-4	No TRV			3.00E+01	3.00E+01
1,2,2-Trichloro-1,1,2-trifluoroethane	76-13-1	No TRV			No TRV	No TRV
1,2,3-Trichloropropane	96-18-4	No TRV			5.71E-01	5.71E-01
1,2-Dibromo-3-chloropropane	96-12-8	No TRV			No TRV	No TRV
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	No TRV			No TRV	No TRV
1,2-Dichloroethane	107-06-2	No TRV			5.00E+01	5.00E+01
1,2-Dichloroethylene	540-59-0	No TRV			4.52E+01	4.52E+01
1,2-Dichloropropane	78-87-5	No TRV			No TRV	No TRV
1,3-Dichloropropene	542-75-6	No TRV			3.00E-01	3.00E-01
1,4-Dichloro-2-butene	764-41-0	No TRV			No TRV	No TRV
1-Chloroethene	75-01-4	No TRV			No TRV	No TRV
2,2-Dichloropropionic acid	75-99-0	No TRV			No TRV	No TRV

Table C3-6. Toxicity Reference Values (TRVs) for Mammals

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
2-Chloropropane	75-29-6	No TRV			No TRV	No TRV
3-Chloropropene (Allyl chloride)	107-05-1	No TRV			No TRV	No TRV
Bromochloromethane	74-97-5	No TRV			No TRV	No TRV
Bromodichloromethane	75-27-4	No TRV			1.79E+00	1.79E+00
Bromoethene	593-60-2	No TRV			No TRV	No TRV
Bromoform	75-25-2	No TRV			1.79E+00	1.79E+00
Bromomethane	74-83-9	No TRV			1.40E-01	1.40E-01
Carbon tetrachloride	56-23-5	No TRV			7.10E-02	7.10E-02
Chlorodibromomethane	124-48-1	No TRV			2.14E+00	2.14E+00
Chlorodifluoromethane	75-45-6	No TRV			No TRV	No TRV
Chloroethane	75-00-3	No TRV			No TRV	No TRV
Chloroform	67-66-3	6.00E+01	e		1.50E+01	6.00E+01
Chloromethane	74-87-3	No TRV			No TRV	No TRV
Chloropentafluoroethane	76-15-3	No TRV			No TRV	No TRV
cis-1,2-Dichloroethene	156-59-2	No TRV			No TRV	No TRV
cis-1,3-Dichloropropene	10061-01-5	No TRV			No TRV	No TRV
Cyanogen bromide	506-68-3	No TRV			4.40E+00	4.40E+00
Cyanogen chloride	506-77-4	No TRV			2.53E+00	2.53E+00
Dichlorodifluoromethane	75-71-8	No TRV			1.50E+00	1.50E+00
Dichlorofluoromethane	75-43-4	No TRV			No TRV	No TRV
Dichloromethane	75-09-2	No TRV			6.16E-01	6.16E-01
Difluorodibromomethane	75-61-6	No TRV			No TRV	No TRV
Hexafluoroacetone	684-16-2	No TRV			No TRV	No TRV
Iodomethane	74-88-4	No TRV			No TRV	No TRV
Methylene bromide	74-95-3	No TRV			No TRV	No TRV
Pentachloroethane	76-01-7	No TRV			No TRV	No TRV
trans-1,2-Dichloroethylene	156-60-5	No TRV			1.70E+00	1.70E+00
trans-1,3-Dichloropropene	10061-02-6	No TRV			No TRV	No TRV
Trichloroacetic acid	76-03-9	No TRV			No TRV	No TRV
Trichlorofluoroethane	27154-33-2	No TRV			No TRV	No TRV

Table C3-6. Toxicity Reference Values (TRVs) for Mammals

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
Trichlorofluoromethane	75-69-4	No TRV			3.49E+01	3.49E+01
Trifluorobromomethane	75-63-8	No TRV			No TRV	No TRV
Dioxin and Furan Compounds						
1,2,3,4,6,7,8-Heptachlorodibenzo-p-dioxin	35822-46-9	No TRV			No TRV	No TRV
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	No TRV			No TRV	No TRV
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	No TRV			No TRV	No TRV
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin	39227-28-6	No TRV			No TRV	No TRV
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	No TRV			No TRV	No TRV
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin	57653-85-7	No TRV			No TRV	No TRV
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	No TRV			1.60E-04	1.60E-04
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin	19408-74-3	No TRV			No TRV	No TRV
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	No TRV			No TRV	No TRV
1,2,3,7,8-Pentachlorodibenzo-p-dioxin	40321-76-4	No TRV			No TRV	No TRV
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	No TRV			1.60E-04	1.60E-04
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	No TRV			No TRV	No TRV
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	No TRV			1.60E-05	1.60E-05
2,3,7,8-Tetrachlorodibenzo-p-dioxin	1746-01-6	1.00E-06	d		1.00E-06	1.00E-06
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	No TRV			No TRV	No TRV
Dibenzofuran	132-64-9	No TRV			No TRV	No TRV
Octachlorodibenzo(p)dioxin	3268-87-9	No TRV			No TRV	No TRV
Octachlorodibenzofuran	39001-02-0	No TRV			No TRV	No TRV
PCBs						
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	No TRV			No TRV	No TRV
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	No TRV			No TRV	No TRV
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	2.06E-03	f		No TRV	2.06E-03
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	2.06E-03	f		No TRV	2.06E-03
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	No TRV			No TRV	No TRV
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	No TRV			No TRV	No TRV
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	No TRV			No TRV	No TRV
2',3,4,4',5-Pentachlorobiphenyl	no cas #	No TRV			No TRV	No TRV

Table C3-6. Toxicity Reference Values (TRVs) for Mammals

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	No TRV			No TRV	No TRV
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	2.06E-03	f		No TRV	2.06E-03
3,3',4,4',5-Pentachlorobiphenyl	no cas #	No TRV			No TRV	No TRV
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	2.06E-03	f		No TRV	2.06E-03
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	No TRV			No TRV	No TRV
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	No TRV			No TRV	No TRV
Polychlorinated biphenyls (PCBs)	1336-36-3	2.06E-03	f		6.80E-02[g]	2.06E-03
Phthalates						
Bis(2-ethylhexyl)phthalate	117-81-7	6.00E+01	d		1.83E+01	6.00E+01
Butylbenzylphthalate	85-68-7	No TRV			No TRV	No TRV
Di-n-butylphthalate	84-74-2	No TRV			5.50E+02	5.50E+02
Diethylphthalate	84-66-2	No TRV			4.58E+03	4.58E+03
Dimethylphthalate	131-11-3	No TRV			No TRV	No TRV
n-Dioctyl phthalate	117-84-0	7.50E+03	e		No TRV	7.50E+03
Light Polycyclic Aromatic Hydrocarbons						
2-Chloronapthalene	91-58-7	No TRV			2.50E+02	2.50E+02
2-Methylnaphthalene	91-57-6	No TRV			No TRV	No TRV
5-Nitroacenaphthene	602-87-9	No TRV			No TRV	No TRV
Acenaphthene	83-32-9	No TRV			1.75E+01	1.75E+01
Acenaphthylene	208-96-8	No TRV			No TRV	No TRV
Anthracene	120-12-7	No TRV			1.00E+02	1.00E+02
Fluorene	86-73-7	No TRV			1.25E+01	1.25E+01
Indene	95-13-6	No TRV			No TRV	No TRV
Naphthalene	91-20-3	No TRV			7.10E+00	7.10E+00
Phenanthrene	85-01-8	No TRV			No TRV	No TRV
Pyrene	129-00-0	No TRV			7.50E+00	7.50E+00
Heavy Polycyclic Aromatic Hydrocarbons[h]						
3-Methylcholanthrene	56-49-5	No TRV			No TRV	No TRV
5-Methylchrysene	3697-24-3	No TRV			No TRV	No TRV

Table C3-6. Toxicity Reference Values (TRVs) for Mammals

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
Benzo(a)anthracene	56-55-3	1.67E-01	e		No TRV	1.67E-01
Benzo(a)pyrene	50-32-8	1.00E-01	e		1.00E+00	1.00E-01
Benzo(b)fluoranthene	205-99-2	No TRV			No TRV	No TRV
Benzo(e)pyrene	192-97-2	No TRV			No TRV	No TRV
Benzo(g,h,i)perylene	191-24-2	No TRV			No TRV	No TRV
Benzo(j)fluoranthene	205-82-3	No TRV			No TRV	No TRV
Benzo(k)fluoranthene	207-08-9	No TRV			No TRV	No TRV
Benzo[a,i]pyrene	191-30-0	No TRV			No TRV	No TRV
Chrysene	218-01-9	No TRV			No TRV	No TRV
Dibenzo(a,h)anthracene	53-70-3	2.00E-03	d		No TRV	2.00E-03
Dibenz[a,h]acridine	226-36-8	No TRV			No TRV	No TRV
Dibenz[a,j]acridine	224-42-0	No TRV			No TRV	No TRV
Dibenzo(a,e)fluoranthene	5385-75-1	No TRV			No TRV	No TRV
Dibenzo(a,h)fluoranthene	no cas #	No TRV			No TRV	No TRV
Dibenzo[a,e]pyrene	192-65-4	No TRV			No TRV	No TRV
Dibenzo[a,h]pyrene	189-64-0	No TRV			No TRV	No TRV
Dibenzo[a,i]pyrene	189-55-9	No TRV			No TRV	No TRV
Fluoranthene	206-44-0	No TRV			1.25E+01	1.25E+01
Hexachloronaphthalene	1335-87-1	No TRV			No TRV	No TRV
Indeno(1,2,3-cd)pyrene	193-39-5	No TRV			No TRV	No TRV
Octachloronaphthalene	2234-13-1	No TRV			No TRV	No TRV
Pentachloronaphthalene	1321-64-8	No TRV			No TRV	No TRV
Tetrachloronaphthalene	1335-88-2	No TRV			No TRV	No TRV
Trichloronaphthalene	1321-65-9	No TRV			No TRV	No TRV
Light Substituted Benzene Compounds						
1,2,3-Trichlorobenzene	87-61-6	No TRV			No TRV	No TRV
1,2,4-Trichlorobenzene	120-82-1	No TRV			No TRV	No TRV
1,2,4-Trimethyl benzene	95-63-6	No TRV			No TRV	No TRV
1,2-Dichlorobenzene	95-50-1	No TRV			8.57E+01	8.57E+01
1,3,5-Trimethyl benzene	108-67-8	No TRV			No TRV	No TRV

Table C3-6. Toxicity Reference Values (TRVs) for Mammals

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
1,3-Dichlorobenzene	541-73-1	No TRV			No TRV	No TRV
1,3-Dinitrobenzene	99-65-0	1.05E+00	d		1.10E-01	1.05E+00
1,4-Dichlorobenzene	106-46-7	No TRV			No TRV	No TRV
1,4-Dinitrobenzene	100-25-4	No TRV			No TRV	No TRV
2,4,5-Trichlorophenol	95-95-4	No TRV			1.00E+01	1.00E+01
2,4,6-Trichlorophenol	88-06-2	No TRV			No TRV	No TRV
2,4-Dichlorophenol	120-83-2	No TRV			3.00E-02	3.00E-02
2,4-Dimethylphenol	105-67-9	No TRV			5.00E-01	5.00E-01
2,4-Dinitrophenol	51-28-5	No TRV			2.00E-01	2.00E-01
2,4-Dinitrotoluene	121-14-2	7.00E-01	i		1.35E+01	7.00E-01
2,6-Dinitrotoluene	606-20-2	4.00E-01	i		7.00E-01	4.00E-01
2-Chlorophenol	95-57-8	No TRV			No TRV	No TRV
2-Chlorotoluene	95-49-8	No TRV			2.00E+01	2.00E+01
2-Nitrophenol	88-75-5	No TRV			No TRV	No TRV
4,6-Dinitro-o-cresol	534-52-1	No TRV			No TRV	No TRV
4-Chlorotoluene	106-43-4	No TRV			No TRV	No TRV
4-Nitrophenol	100-02-7	No TRV			No TRV	No TRV
alpha-Methylstyrene	98-83-9	No TRV			No TRV	No TRV
Aniline	62-53-3	No TRV			No TRV	No TRV
Benzotrichloride	98-07-7	No TRV			No TRV	No TRV
Benzyl chloride	100-44-7	No TRV			No TRV	No TRV
Bromobenzene	108-86-1	No TRV			No TRV	No TRV
Chlorobenzene	108-90-7	No TRV			1.90E+01	1.90E+01
Cumene	98-82-8	No TRV			1.10E+02	1.10E+02
m-Cresol	108-39-4	No TRV			5.00E+00	5.00E+00
n-Butyl benzene	104-51-8	No TRV			No TRV	No TRV
Nitrobenzene	98-95-3	No TRV			4.60E-02	4.60E-02
n-Propyl benzene	103-65-1	No TRV			No TRV	No TRV
o-Cresol	95-48-7	No TRV			5.00E+00	5.00E+00
o-Dinitrobenzene	528-29-0	No TRV			No TRV	No TRV

Table C3-6. Toxicity Reference Values (TRVs) for Mammals

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
o-Nitroaniline	88-74-4	No TRV			No TRV	No TRV
o-Toluidine	95-53-4	No TRV			No TRV	No TRV
p-Chloroaniline	106-47-8	No TRV			1.25E+00	1.25E+00
p-Cresol	106-44-5	No TRV			No TRV	No TRV
Phenol	108-95-2	No TRV			6.00E+01	6.00E+01
p-Nitrochlorobenzene	100-00-5	No TRV			No TRV	No TRV
p-Toluidine	106-49-0	No TRV			No TRV	No TRV
sec-Butyl benzene	135-98-8	No TRV			No TRV	No TRV
tert-Butyl benzene	98-06-6	No TRV			No TRV	No TRV
Toluene-2,6-diamine	823-40-5	No TRV			No TRV	No TRV
Trimethyl benzene	25551-13-7	No TRV			No TRV	No TRV
Other Light Semivolatile Compounds						
1,1'-Biphenyl	92-52-4	No TRV			5.00E+00	5.00E+00
1,1-Dimethylhydrazine	57-14-7	No TRV			No TRV	No TRV
1,2-Dimethylhydrazine	540-73-8	No TRV			No TRV	No TRV
1,2-Diphenylhydrazine	122-66-7	No TRV			No TRV	No TRV
1,3-Propane sultone	1120-71-4	No TRV			No TRV	No TRV
2,4-Toluene diisocyante	584-84-9	No TRV			No TRV	No TRV
2-Chloroacetophenone	532-27-4	No TRV			No TRV	No TRV
2-Propenoic acid	79-10-7	No TRV			5.30E+00	5.30E+00
4,4-Methylenedianiline	101-77-9	No TRV			No TRV	No TRV
Acetophenone	98-86-2	No TRV			4.23E+01	4.23E+01
Benzoic acid	65-85-0	No TRV			4.00E+00	4.00E+00
Bis(2-chloroethoxy)methane	111-91-1	No TRV			No TRV	No TRV
Bis(2-chloroethyl) ether	111-44-4	No TRV			No TRV	No TRV
Chlorocyclopentadiene	41851-50-7	No TRV			No TRV	No TRV
Cyclohexanol	108-93-0	No TRV			No TRV	No TRV
Dichloroisopropyl ether	108-60-1	No TRV			No TRV	No TRV
Dichloromethyl ether	542-88-1	No TRV			No TRV	No TRV
Dichloropentadiene	no cas #	No TRV			No TRV	No TRV

Table C3-6. Toxicity Reference Values (TRVs) for Mammals

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
Dimethyl sulfate	77-78-1	No TRV			No TRV	No TRV
Dimethylaniline	121-69-7	No TRV			2.23E-01	2.23E-01
di-n-Propylnitrosamine	621-64-7	No TRV			No TRV	No TRV
Diphenyl ether	101-84-8	No TRV			No TRV	No TRV
Epichlorohydrin	106-89-8	No TRV			No TRV	No TRV
Ethyl Carbamate (Urethane)	51-79-6	No TRV			No TRV	No TRV
Ethyl methanesulfonate	62-50-0	No TRV			No TRV	No TRV
Ethylene dibromide	106-93-4	No TRV			No TRV	No TRV
Ethylene glycol	107-21-1	No TRV			2.00E+01	2.00E+01
Ethylene glycol monobutyl ether	111-76-2	No TRV			5.10E-01	5.10E-01
Ethylene glycol monoethyl ether acetate	111-15-9	No TRV			No TRV	No TRV
Ethylene thiourea	96-45-7	No TRV			2.50E-02	2.50E-02
Furfural	98-01-1	No TRV			7.90E-01	7.90E-01
Maleic hydrazide	123-33-1	No TRV			5.00E+01	5.00E+01
Malononitrile	109-77-3	No TRV			No TRV	No TRV
Methyl styrene (mixed isomers)	25013-15-4	No TRV			No TRV	No TRV
Methylhydrazine	60-34-4	No TRV			No TRV	No TRV
N,N-Diphenylamine	122-39-4	No TRV			2.50E-01	2.50E-01
Nitric acid, propyl ester	627-13-4	No TRV			No TRV	No TRV
N-Nitrosodi-n-butylamine	924-16-3	No TRV			No TRV	No TRV
N-Nitrosomorpholine	59-89-2	No TRV			No TRV	No TRV
N-Nitroso-N,N-dimethylamine	62-75-9	No TRV			No TRV	No TRV
o-Anisidine	90-04-0	No TRV			No TRV	No TRV
Oxalic acid	144-62-7	No TRV			No TRV	No TRV
Phthalic anhydride	85-44-9	No TRV			1.56E+02	1.56E+02
p-Phthalic acid	100-21-0	No TRV			No TRV	No TRV
Pyridine	110-86-1	No TRV			1.00E-01	1.00E-01
Quinoline	91-22-5	No TRV			No TRV	No TRV
Quinone	106-51-4	No TRV			No TRV	No TRV
Safrole	94-59-7	No TRV			No TRV	No TRV

Table C3-6. Toxicity Reference Values (TRVs) for Mammals

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
Tetrahydrofuran	109-99-9	No TRV			No TRV	No TRV
Other Heavy Semivolatile Compounds						
1,2,4,5-Tetrachlorobenzene	95-94-3	No TRV			3.40E-02	3.40E-02
1,3,5-Trinitrobenzene	99-35-4	No TRV			6.70E+00	6.70E+00
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	No TRV			No TRV	No TRV
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	No TRV			2.00E-01	2.00E-01
2-sec-Butyl-4,6-dinitrophenol	88-85-7	No TRV			No TRV	No TRV
3,3-Dichlorobenzidine	91-94-1	No TRV			No TRV	No TRV
3,3'-Dimethoxybenzidine	119-90-4	No TRV			No TRV	No TRV
4-Bromophenylphenyl ether	101-55-3	No TRV			No TRV	No TRV
Ammonium perfluorooctanoate	3825-26-1	No TRV			No TRV	No TRV
Azobenzene	103-33-3	No TRV			No TRV	No TRV
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	No TRV			No TRV	No TRV
Chlorobenzilate	510-15-6	No TRV			No TRV	No TRV
Dibutylphosphate	107-66-4	No TRV			No TRV	No TRV
Dimethyl aminoazobenzene	60-11-7	No TRV			No TRV	No TRV
Hexachlorobenzene	118-74-1	1.60E+00	d		8.00E-03	1.60E+00
Hexachlorobutadiene	87-68-3	2.00E-01	d		No TRV	2.00E-01
Hexachlorocyclopentadiene	77-47-4	3.80E+00	d		7.00E-01	3.80E+00
Hexachloroethane	67-72-1	No TRV			1.00E-01	1.00E-01
Hexachlorophene	70-30-4	5.60E+00			7.50E-02	5.60E+00
Hexamethylene-1,5-diisocyanate	822-06-0	No TRV			No TRV	No TRV
Mirex	2385-85-5	No TRV			7.00E-02	7.00E-02
Nitrofen	1836-75-5	No TRV			No TRV	No TRV
Pentachlorobenzene	608-93-5	7.25E+00	d		8.30E-01	7.25E+00
Pentachloronitrobenzene	82-68-8	4.58E+02	e		7.50E-02	4.58E+02
Pentachlorophenol	87-86-5	3.00E-01	d		2.40E-01	3.00E-01
Picric acid	88-89-1	No TRV			No TRV	No TRV
Pronamide	23950-58-5	No TRV			7.50E-01	7.50E-01
Strychnine	57-24-9	No TRV			2.50E-01	2.50E-01

Table C3-6. Toxicity Reference Values (TRVs) for Mammals

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
Terphenyls	26140-60-3	No TRV			No TRV	No TRV
Tributyl phosphate	126-73-8	No TRV			No TRV	No TRV
Trifluralin	1582-09-8	No TRV			7.50E-02	7.50E-02
Triphenylamine	603-34-9	No TRV			No TRV	No TRV
Herbicides and Organochlorinated Pesticides						
2,4,5-Trichlorophenoxyacetic acid (2,4,5-T)	93-76-5	No TRV			3.00E+00	3.00E+00
2,4-D	94-75-7	No TRV			No TRV	No TRV
4,4'-DDD	72-54-8	No TRV			No TRV	No TRV
4,4'-DDE	72-55-9	1.00E+00	d		No TRV	1.00E+00
4,4'-DDT	50-29-3	No TRV			8.00E-01	8.00E-01
Aldrin	309-00-2	No TRV			2.00E-01	2.00E-01
alpha-BHC	319-84-6	No TRV			No TRV	No TRV
beta-BHC	319-85-7	No TRV			No TRV	No TRV
Captan	133-06-2	No TRV			No TRV	No TRV
Chlordane	57-74-9	No TRV			4.58E+00	4.58E+00
delta-BHC	319-86-8	No TRV			No TRV	No TRV
Dieldrin	60-57-1	No TRV			2.00E-02	2.00E-02
Endothall	145-73-3	No TRV			2.00E+00	2.00E+00
Endrin	72-20-8	No TRV			9.20E-02	9.20E-02
gamma-BHC (Lindane)	58-89-9	No TRV			8.00E+00	8.00E+00
Heptachlor	76-44-8	2.50E-03	d		1.00E-01	2.50E-03
Isodrin	465-73-6	No TRV			No TRV	No TRV
Methoxychlor	72-43-5	No TRV			5.01E+00	5.01E+00
Silvex (2,4,5-TP)	93-72-1	No TRV			7.50E-01	7.50E-01
Toxaphene	8001-35-2	No TRV			8.00E+00	8.00E+00
Inorganics						
Metals						
Aluminum	7429-90-5	1.93E+00	d		1.93E+00	1.93E+00
Antimony	7440-36-0	6.60E-02	d		1.25E-01	6.60E-02
Arsenic	7440-38-2	1.25E+00	i		1.26E-01	1.25E+00

Table C3-6. Toxicity Reference Values (TRVs) for Mammals

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
Barium	7440-39-3	5.10E-01	d		5.06E+00	5.10E-01
Beryllium	7440-41-7	6.60E-01	d		6.60E-01	6.60E-01
Bismuth	7440-69-9	No TRV			No TRV	No TRV
Boron	7440-42-8	No TRV			2.80E+01	2.80E+01
Cadmium	7440-43-9	2.52E-02	e		1.00E+00	2.52E-02
Calcium	7440-70-2	No TRV			No TRV	No TRV
Chromium (and VI)	7440-47-3	3.50E+00	d		2.74E+03	3.50E+00
Cobalt	7440-48-4	No TRV			1.00E-01	1.00E-01
Copper	7440-50-8	1.20E+01	j		1.17E+01	1.20E+01
Iron	7439-89-6	No TRV			No TRV	No TRV
Lead	7439-92-1	3.75E-02	e		8.00E+00	3.75E-02
Lithium	7439-93-2	No TRV			9.39E+00	9.39E+00
Magnesium	7439-95-4	No TRV			No TRV	No TRV
Manganese	7439-96-5	No TRV			8.80E+01	8.80E+01
Mercury	7439-97-6	1.01E+00	j		1.01E+00	1.01E+00
Molybdenum	7439-98-7	No TRV			2.58E-01	2.58E-01
Nickel	7440-02-0	5.00E+01	d		4.00E+01	5.00E+01
Potassium	7440-09-7	No TRV			No TRV	No TRV
Rhodium	7440-16-6	No TRV			No TRV	No TRV
Selenium	7782-49-2	7.60E-02	e		2.00E-01	7.60E-02
Silicon	7440-21-3	No TRV			No TRV	No TRV
Silver	7440-22-4	3.75E-01	e		No TRV	3.75E-01
Sodium	7440-23-5	No TRV			No TRV	No TRV
Strontium	7440-24-6	No TRV			2.63E+02	2.63E+02
Tantalum	7440-25-7	No TRV			No TRV	No TRV
Thallium	7440-28-0	1.31E-02	d		7.40E-03	1.31E-02
Tin	7440-31-5	No TRV			No TRV	No TRV
Tungsten	7440-33-7	No TRV			No TRV	No TRV
Uranium	7440-61-1	No TRV			3.07E+00	3.07E+00
Vanadium	7440-62-2	No TRV			2.10E-01	2.10E-01

Table C3-6. Toxicity Reference Values (TRVs) for Mammals

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)
Yttrium	7440-65-5	No TRV			No TRV	No TRV
Zinc	7440-66-6	1.04E+01	e		1.60E+02	1.04E+01
Zirconium	7440-67-7	No TRV			No TRV	No TRV
Non-metals and Ions						
Ammonia/Ammonium	7664-41-7	No TRV			No TRV	No TRV
Bromide	24959-67-9	No TRV			No TRV	No TRV
Chloride	16887-00-6	No TRV			No TRV	No TRV
Cyanide	57-12-5	2.40E+01	d		6.87E+01	2.40E+01
Fluoride	16984-48-8	No TRV			3.14E+01	3.14E+01
Hydroxide	14280-30-9	No TRV			No TRV	No TRV
Iodine	7553-56-2	No TRV			No TRV	No TRV
Nitrate	14797-55-8	No TRV			No TRV	No TRV
Nitrite	14797-65-0	No TRV			No TRV	No TRV
Phosphate	14265-44-2	No TRV			No TRV	No TRV
Phosphorus	7723-14-0	No TRV			No TRV	No TRV
Sulfate	14808-79-8	No TRV			No TRV	No TRV
Total Sulfur	63705-05-5	No TRV			No TRV	No TRV
Criteria Pollutants						
Carbon dioxide	124-38-9	No TRV			No TRV	No TRV
Nitrogen dioxide	10102-44-0	No TRV			No TRV	No TRV
Ozone	10028-15-6	No TRV			No TRV	No TRV
Particulate matter	no cas #	No TRV			No TRV	No TRV
Sulfur dioxide	7446-09-5	No TRV			No TRV	No TRV
Radionuclides	TRV is not applicable to single radionuclides. Combined external and internal radiation exposure for mammals from all radionuclides combined cannot exceed 0.1 rad/d.					

TRV = toxicity reference value

[a] Published in Appendix E of EPA (1999), Table E-7.

[b] Published in Table C3-5 of this work plan

Table C3-6. Toxicity Reference Values (TRVs) for Mammals

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a] (mg/kg/d)	Notes	Ecology Guidance TRV (mg/kg/d)	SAIC Compiled TRV[b] (mg/kg/d)	Recommended TRV[c] (mg/kg/d)

[c] Order of preference is EPA (1999), then SAIC compilation

[d] Toxicity to rat

[e] Toxicity to mouse

[f] Based on toxicity of 3,4,5-hexachlorobiphenyl to mink

[g] Toxicity of Aroclor 1254 to oldfield mouse used as representative of PCB mixtures

[h] Total exposure to all high molecular weight polycylic aromatic hydrocarbons combined is limited to 0.1 mg/kg/d.

[i] Toxicity to dog

[j] Toxicity to mink

Table C3-7. Derivation of Inhalation TRVs for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Assumed Test Species Body Weight (kg)[a]	Published Test Benchmark/Unit/Duration		Molecular Weight	Converted Test Benchmark (mg/m^3)[b]	Test Duration[c]	Test Endpoint	Effect	Source	Proposed Duration Conversion Factor DCF[d]	Proposed Endpoint Conversion Factor ECF[e]	TRV (mg/m^3) = Benchmark x DCF x ECF
Organic Compounds														
Aromatic Halogenated Hydrocarbons														
4-Chloro-3-methylphenol	59-50-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2,3,4,6-Tetrachlorophenol	58-90-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Aromatic Nonhalogenated Hydrocarbons														
2-Nitrotoluene	88-72-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
4-Nitrobiphenyl	92-93-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Benzaldehyde	100-52-7	Rat	3.50E-01	2.60E+01	mg/m3/5H/17W-I	1.06E+02	2.60E+01	Subchronic	TCLo	Blood	RTECS	1.00E-01	1.00E-01	2.60E-01
Benzene	71-43-2	Mouse	3.00E-02	5.00E+00	ppm/6-15D	7.81E+01	1.60E+01	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	1.60E-01
Benzyl alcohol	100-51-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Ethyl benzene	100-41-4	Rabbit	3.80E+00	1.00E+00	g/m3/24H	1.06E+02	1.00E+00	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	1.00E-02
m-Xylene	108-38-3	Mouse	3.00E-02	5.00E+02	mg/m3/12H/6-15D	1.06E+02	5.00E+02	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	5.00E+00
o-Xylene	95-47-6	Rat	3.50E-01	1.50E+02	mg/m3/24H/7-14D	1.06E+02	1.50E+02	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	1.50E+00
p-Xylene	106-42-3	Rat	3.50E-01	1.50E+02	mg/m3/24H/7-14D	1.06E+02	1.50E+02	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	1.50E+00
Styrene	100-42-5	Mouse	3.00E-02	9.50E+03	mg/m3/4H	1.04E+02	9.50E+03	Acute	LC50	Mortality	RTECS	1.00E-01	1.00E-02	9.50E+00
Toluene	108-88-3	Rabbit	3.80E+00	1.00E+02	ppm/6H/6-18D	9.21E+01	3.77E+02	Subchronic	TDLo	Reproduction	RTECS	1.00E-01	1.00E-02	3.77E-01
Non-aromatic Nonhalogenated Hydrocarbons														
1,2-Epoxybutane	106-88-7	Mouse	3.00E-02	2.60E+01	mg/m3/6H/5D/2yr	7.21E+01	2.60E+01	Chronic	TCLo	Nasal tissue	IRIS	1.00E+00	1.00E-01	2.60E+00
1,3-Butadiene	106-99-0	Mouse	3.00E-02	1.00E+03	ppm/6H/6-15D	5.41E+01	2.21E+03	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	2.21E+01
1,4-Dioxane	123-91-1	Rat	3.50E-01	2.05E+04	ug/m3/13W-I	8.81E+01	2.05E+01	Chronic	TCLo	Organ dysfunction	RTECS	1.00E+00	1.00E-01	2.05E+00
1-Methylpropyl alcohol	78-92-2	Rat	3.50E-01	5.00E+03	ppm/7H/1-19D	7.41E+01	1.52E+04	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	1.52E+02
1-Nitropropane	108-03-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2,2,4-Trimethylpentane	540-84-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2-Butanone	78-93-3	Rat	3.50E-01	1.00E+03	ppm/7H/6-15D	7.21E+01	2.95E+03	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	2.95E+01

Table C3-7. Derivation of Inhalation TRVs for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Assumed Test Species Body Weight (kg)[a]	Published Test Benchmark/Unit/Duration		Molecular Weight	Converted Test Benchmark (mg/m^3)[b]	Test Duration[c]	Test Endpoint	Effect	Source	Proposed Duration Conversion Factor DCF[d]	Proposed Endpoint Conversion Factor ECF[e]	TRV (mg/m^3) = Benchmark x DCF x ECF
2-Butenaldehyde (2-Butenal)	4170-30-3	Rat	3.50E-01	2.00E+01	mg/m3/7H/8W-I	7.01E+01	2.00E+01	Chronic	TCLo	Organ dysfunction	RTECS	1.00E+00	1.00E-01	2.00E+00
2-Ethoxyethanol	110-80-5	Rabbit	3.80E+00	1.03E+02	ppm	9.01E+01	3.80E+02	Chronic	–	Organ dysfunction	IRIS	1.00E+00	1.00E-01	3.80E+01
2-Heptanone	110-43-0	Rat	3.50E-01	4.00E+03	ppm/4H	1.14E+02	1.87E+04	Acute	TCLo	Mortality	RTECS	1.00E-01	1.00E-01	1.87E+02
2-Hexanone	591-78-6	Rat	3.50E-01	8.00E+03	ppm/4H	1.00E+02	3.28E+04	Acute	LC50	Mortality	RTECS	1.00E-01	1.00E-02	3.28E+01
2-Methoxyethanol	109-86-4	Rat	3.50E-01	3.00E+02	ppm/6H/13W-I	7.60E+01	9.33E+02	Chronic	TCLo	Metabolism	RTECS	1.00E+00	1.00E-01	9.33E+01
2-Methyl-2-propanol	75-65-0	Rat	3.50E-01	2.00E+03	ppm/7H/1-19D	7.41E+01	6.06E+03	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	6.06E+01
2-Methyl-2-propenenitrile	126-98-7	Rat	3.50E-01	1.00E+02	ppm/6H/6-20D	6.71E+01	2.74E+02	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	2.74E+00
2-Methylaziridine	75-55-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2-Methylpropyl alcohol	78-83-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2-Pentanone	107-87-9	Rat	3.50E-01	2.00E+03	ppm/4H	8.61E+01	7.05E+03	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	7.05E+01
2-Propanone (Acetone)	67-64-1	Rat	3.50E-01	1.99E+02	mg/m3/8H/45D-I	5.81E+01	3.15E+01	Chronic	TCLo	Behavioral	RTECS	1.00E+00	1.00E+00	3.15E+01
2-Propene-1-ol	107-18-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2-Propyl alcohol	67-63-0	Rat	3.50E-01	3.50E+03	ppm/7H/1-19D	6.01E+01	8.60E+03	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	8.60E+01
3-Heptanone	106-35-4	Rat	3.50E-01	2.00E+03	ppm/4H	1.14E+02	9.34E+03	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	9.34E+01
3-Methyl-1-butanol	123-51-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
3-Methyl-2-butanone	563-80-4	Rat	3.50E-01	5.70E+03	ppm/4H	8.61E+01	2.01E+04	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	2.01E+02
3-Pentanone	96-22-0	Rat	3.50E-01	8.00E+03	ppm/4H	8.61E+01	2.82E+04	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	2.82E+02
4-Heptanone	123-19-3	Rat	3.50E-01	2.69E+03	ppm/6H	1.14E+02	1.26E+04	Acute	LC50	Mortality	RTECS	1.00E-01	1.00E-02	1.26E+01
4-Methyl-2-pentanone	108-10-1	Mouse	3.00E-02	3.00E+03	ppm/6H/6-15D	1.00E+02	1.23E+04	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	1.23E+02
4-Methyl-3-penten-2-one	141-79-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
5-Methyl-2-hexanone	110-12-3	Rat	3.50E-01	1.00E+03	ppm/6H/16D-I	1.14E+02	4.67E+03	Subchronic	TCLo	Organ dysfunction	RTECS	1.00E-01	1.00E-01	4.67E+01
Acetaldehyde	75-07-0	Rat	3.50E-01	1.50E+02	ppm	4.45E+01	2.73E+02	Chronic	–	Nasal tissue	IRIS	1.00E+00	1.00E-01	2.73E+01
Acetamide	60-35-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Acetic acid	64-19-7	Rat	3.50E-01	1.60E+04	ppm/4H	6.01E+01	3.93E+04	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	3.93E+02
Acetic acid ethyl ester	141-78-6	Cat	4.50E+00	6.10E+01	gm/m3	8.81E+01	6.10E+04	Acute	LCLo	Mortality	RTECS	1.00E+00	1.00E-01	6.10E+03
Acetic acid n-butyl ester	123-86-4	Rat	3.50E-01	1.50E+03	ppm/7H/7-16D	1.16E+02	7.13E+03	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	7.13E+01
Acetonitrile	75-05-8	Mouse	3.00E-02	8.00E+02	ppm/6H/13W-I	4.11E+01	1.34E+03	Chronic	TCLo	Mortality	RTECS	1.00E+00	1.00E-01	1.34E+02

Table C3-7. Derivation of Inhalation TRVs for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Assumed Test Species Body Weight (kg)[a]	Published Test Benchmark/Unit/Duration		Molecular Weight	Converted Test Benchmark (mg/m^3)[b]	Test Duration[c]	Test Endpoint	Effect	Source	Proposed Duration Conversion Factor DCF[d]	Proposed Endpoint Conversion Factor ECF[e]	TRV (mg/m^3) = Benchmark x DCF x ECF
Acrolein	107-02-8	Rat	3.50E-01	4.90E+00	ppb/6H/13W-I	5.61E+01	1.12E+01	Chronic	TCLo	Mortality	RTECS	1.00E+00	1.00E-01	1.12E+00
Acrylonitrile	107-13-1	Rat	3.50E-01	2.50E+01	ppm/6H/6-20D	5.31E+01	5.43E+01	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	5.43E-01
Bis(isopropyl)ether	108-20-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Butane	106-97-8	Rat	3.50E-01	6.58E+02	gm/m3/4H	5.81E+01	6.58E+05	Acute	LC50	Mortality	RTECS	1.00E-01	1.00E-02	6.58E+02
Carbon disulfide	75-15-0	Rat	3.50E-01	3.00E+01	ug/m3/8H/1-22D	7.61E+01	3.00E-02	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	3.00E-04
Cyanogen	460-19-5	Rat	3.50E-01	2.50E+01	ppm/6H/26W-I	5.20E+01	5.32E+01	Chronic	TCLo	Metabolism	RTECS	1.00E+00	1.00E-01	5.32E+00
Cyclohexane	110-82-7	Mouse	3.00E-02	7.00E+01	gm/m3/2H	8.42E+01	7.00E+04	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	7.00E+02
Cyclohexanone	108-94-1	Rat	3.50E-01	1.05E+02	mg/m3/4H/1-20D	9.82E+01	1.05E+02	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	1.05E+00
Cyclohexene	110-83-8	Rat	3.50E-01	6.37E+03	ppm/4H	8.21E+01	2.14E+04	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	2.14E+02
Cyclopentane	287-92-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Ethyl alcohol	64-17-5	Rat	3.50E-01	2.00E+04	ppm/7H/1-22D	4.61E+01	3.77E+04	Chronic	TCLo	Reproduction	RTECS	1.00E+00	1.00E-01	3.77E+03
Ethyl ether	60-29-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Ethyl methacrylate	97-63-2	Rat	3.50E-01	8.30E+03	ppm/4H	1.14E+02	3.87E+04	Acute	LC50	Mortality	RTECS	1.00E-01	1.00E-02	3.87E+01
Formaldehyde	50-00-0	Rat	3.50E-01	1.20E+01	ug/m3/24H/15D	3.00E+01	1.20E-02	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	1.20E-04
Formamide	75-12-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Formic acid	64-18-6	Rat	3.50E-01	6.40E+01	ppm/6H/13W-I	4.60E+01	1.20E+02	Chronic	TCLo	Organ dysfunction	RTECS	1.00E+00	1.00E-01	1.20E+01
Formic acid, methyl ester	107-31-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Glycidylaldehyde	765-34-4	Rat	3.50E-01	4.00E+01	ppm/4H/12W-I	7.21E+01	1.18E+02	Chronic	TCLo	Metabolism	RTECS	1.00E+00	1.00E-01	1.18E+01
Methyl acetate	79-20-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Methyl alcohol	67-56-1	Monkey	5.00E+00	1.00E+03	ppm	3.20E+01	1.31E+03	Chronic	LCLo	Mortality	RTECS	1.00E+00	1.00E-01	1.31E+02
Methyl isocyanate	624-83-9	Mouse	3.00E-02	1.00E+00	ppm/6H/14 17D	5.71E+01	2.33E+00	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	2.33E-02
Methyl methacrylate	80-62-6	Rat	3.50E-01	3.50E+01	ppm/6H/5d/wk/2yr	1.00E+02	1.43E+02	Chronic	TCLo	Nasal tissue	IRIS	1.00E+00	1.00E-01	1.43E+01
Methyl tert-butyl ether	1634-04-4	Rat	3.50E-01	8.00E+03	ppm/6H	8.82E+01	2.88E+04	Acute	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	2.88E+02
Methylacetylene	74-99-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data

Table C3-7. Derivation of Inhalation TRVs for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Assumed Test Species Body Weight (kg)[a]	Published Test Benchmark/Unit/Duration		Molecular Weight	Converted Test Benchmark (mg/m³)[b]	Test Duration[c]	Test Endpoint	Effect	Source	Proposed Duration Conversion Factor DCF[d]	Proposed Endpoint Conversion Factor ECF[e]	TRV (mg/m³) = Benchmark x DCF x ECF
Methylcyclohexane	108-87-2	Rabbit	3.80E+00	1.01E+04	ppm/6H/2W-I	9.82E+01	4.04E+04	Subchronic	TCLo	Mortality	RTECS	1.00E-01	1.00E-01	4.04E+02
N,N-Dimethylacetamide	127-19-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
n-Butyl alcohol	71-36-3	Rat	3.50E-01	6.00E+03	ppm/7H/1-19D	7.41E+01	1.82E+04	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	1.82E+02
n-Heptane	142-82-5	Rat	3.50E-01	4.00E+03	ppm/6H/28D-I	1.00E+02	1.64E+04	Chronic	TCLo	Metabolism	RTECS	1.00E+00	1.00E-01	1.64E+03
n-Hexane	110-54-3	Rat	3.50E-01	1.00E+03	ppm/6H/8-16D	8.62E+01	3.52E+03	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	3.52E+01
Nitromethane	75-52-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
n-Nonane	111-84-2	Rat	3.50E-01	1.60E+03	ppm/6H/13W-I	1.28E+02	8.39E+03	Chronic	TCLo	Metabolism	RTECS	1.00E+00	1.00E-01	8.39E+02
n-Octane	111-65-9	Rat	3.50E-01	1.18E+02	gm/m3/4H	1.14E+02	1.18E+05	Acute	LC50	Mortality	RTECS	1.00E-01	1.00E-02	1.18E+02
n-Pentane	109-66-0	Mouse	3.00E-02	3.25E+02	gm/m3/2H	7.22E+01	3.25E+05	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	3.25E+03
n-Propionaldehyde	123-38-6	Rat	3.50E-01	8.00E+03	ppm/4H	5.81E+01	1.90E+04	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	1.90E+02
n-Propyl alcohol	71-23-8	Rat	3.50E-01	4.00E+03	ppm/4H	6.01E+01	9.83E+03	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	9.83E+01
n-Valeraldehyde	110-62-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Oxirane	75-21-8	Rat	3.50E-01	3.60E+03	ug/m3/24H/60D	4.41E+01	3.60E+00	Chronic	TCLo	Reproduction	RTECS	1.00E+00	1.00E-01	3.60E-01
p-Cymene	99-87-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Phosgene	75-44-5	Rat	3.50E-01	5.00E+02	ppb/6H/12W-I	9.89E+01	2.02E+00	Chronic	TCLo	Organ dysfunction	RTECS	1.00E+00	1.00E-01	2.02E-01
Propargyl alcohol	107-19-7	Rat	3.50E-01	7.00E+00	mg/m3/4H/12W-I	5.61E+01	7.00E+00	Chronic	TCLo	Organ dysfunction	RTECS	1.00E+00	1.00E-01	7.00E-01
Propionic acid	79-09-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Propionitrile	107-12-0	Rat	3.50E-01	1.00E+02	ppm/6H/6-20D	5.51E+01	2.25E+02	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	2.25E+00
Propylene gylcol monomethyl ether	107-98-2	Rat	3.50E-01	1.00E+03	ppm	9.01E+01	3.69E+03	Chronic	--	Sedation	IRIS	1.00E+00	1.00E-01	3.69E+02
p-tert-Butyltoluene	98-51-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Triethylamine	121-44-8	Rat	3.50E-01	1.00E+03	ppm/4H	1.01E+02	4.14E+03	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	4.14E+01
Trimethylamine	75-50-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Vinyl acetate	108-05-4	Rat	3.50E-01	2.50E+03	ppm/4H/1Y-I	8.61E+01	8.80E+03	Chronic	TCLo	Mortality	RTECS	1.00E+00	1.00E-01	8.80E+02
Non-aromatic Halogenated Hydrocarbons														
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
1,1,1,2-Tetrachloroethane	630-20-6	Rat	3.50E-01	2.10E+03	ppm/4H	1.68E+02	1.44E+04	Acute	LC50	Mortality	RTECS	1.00E-01	1.00E-02	1.44E+01

Table C3-7. Derivation of Inhalation TRVs for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Assumed Test Species Body Weight (kg)[a]	Published Test Benchmark/Unit/Duration		Molecular Weight	Converted Test Benchmark (mg/m³)[b]	Test Duration[c]	Test Endpoint	Effect	Source	Proposed Duration Conversion Factor DCF[d]	Proposed Endpoint Conversion Factor ECF[e]	TRV (mg/m³) = Benchmark x DCF x ECF
1,1,1-Trichloroethane	71-55-6	Guinea pig	3.50E-01	6.50E+02	ppm/7H/58D-I	1.33E+02	3.55E+03	Chronic	TCLo	Metabolism	RTECS	1.00E+00	1.00E-01	3.55E+02
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
1,1,2,2-Tetrachloroethane	79-34-5	Rat	3.50E-01	1.00E+03	ppm/4H	1.68E+02	6.87E+03	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	6.87E+01
1,1,2,2-Tetrachloroethene	127-18-4	Rat	3.50E-01	3.00E+02	ppm/7H/6-15D	1.66E+02	2.03E+03	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	2.03E+01
1,1,2-Trichloroethane	79-00-5	Rat	3.50E-01	5.00E+02	ppm/4H	1.33E+02	2.73E+03	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	2.73E+01
1,1,2-Trichloroethylene	79-01-6	Rat	3.50E-01	1.00E+02	ppm/4H/6-22D	1.31E+02	5.37E+02	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	5.37E+00
1,1-Dichloroethane	75-34-3	Rat	3.50E-01	6.00E+03	ppm/7H/6-15D	9.90E+01	2.43E+04	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	2.43E+02
1,1-Dichloroethene	75-35-4	Rat	3.50E-01	5.50E+01	ppm/6H/55D	9.70E+01	2.18E+02	Chronic	TCLo	Reproduction	RTECS	1.00E+00	1.00E-01	2.18E+01
1,2,2-Trichloro-1,1,2-trifluoroethane	76-13-1	Rat	3.50E-01	2.00E+03	ppm/6H/2W-I	1.87E+02	1.53E+04	Subchronic	TCLo	Organ dysfunction	RTECS	1.00E-01	1.00E-01	1.53E+02
1,2,3-Trichloropropane	96-18-4	Rat	3.50E-01	2.00E+00	mg/m3/24H/94D-C	1.47E+02	2.00E+00	Chronic	TCLo	Organ dysfunction	RTECS	1.00E+00	1.00E-01	2.00E-01
1,2-Dibromo-3-chloropropane	96-12-8	Rabbit	3.80E+00	1.00E-01	ppm	2.36E+02	9.60E-01	Chronic	--	Reproduction	IRIS	1.00E+00	1.00E-01	9.60E-02
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	Guinea pig	3.50E-01	2.00E+05	ppm/8H	1.71E+02	1.40E+06	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	1.40E+04
1,2-Dichloroethane	107-06-2	Rat	3.50E-01	2.00E+02	ppm/7H/6W-I	9.90E+01	8.09E+02	Chronic	TCLo	Mortality	RTECS	1.00E+00	1.00E-01	8.09E+01
1,2-Dichloroethylene	540-59-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
1,2-Dichloropropane	78-87-5	Rat	3.50E-01	1.00E+03	ppm/7H/20W-I	1.13E+02	4.62E+03	Chronic	TCLo	Mortality	RTECS	1.00E+00	1.00E-01	4.62E+02
1,3-Dichloropropene	542-75-6	Mouse	3.00E-02	5.00E+00	ppm	1.11E+02	2.27E+01	Chronic	--	Nasal tissue	IRIS	1.00E+00	1.00E-01	2.27E+00
1,4-Dichloro-2-butene	764-41-0	Rat	3.50E-01	8.70E+03	ug/m3/17W-I	1.25E+02	8.70E+00	Chronic	TCLo	Neurological	RTECS	1.00E-01	1.00E-01	8.70E-02
1-Chloroethene	75-01-4	Rat	3.50E-01	1.00E+02	ppm/6H/26W	6.25E+01	2.56E+02	Chronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	2.56E+00
2,2-Dichloropropionic acid	75-99-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2-Chloropropane	75-29-6	Rat	3.50E-01	2.36E+02	gm/m3/30M/1W-I	7.86E+01	2.36E+05	Subchronic	TCLo	Organ dysfunction	RTECS	1.00E-01	1.00E-01	2.36E+03
3-Chloropropene (allyl chloride)	107-05-1	Rat	3.50E-01	3.00E+02	ppm/7H/6-15D	7.65E+01	9.39E+02	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	9.39E+00
Bromochloromethane	74-97-5	Rat	3.50E-01	5.00E+02	ppm/6H/26W-I	1.29E+02	2.64E+03	Chronic	TCLo	Metabolism	RTECS	1.00E+00	1.00E-01	2.64E+02
Bromodichloromethane	75-27-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Bromoethene	593-60-2	Rat	3.50E-01	9.70E+00	ppm	1.07E+02	4.24E+01	Chronic	--	Organ dysfunction	IRIS	1.00E+00	1.00E-01	4.24E+00

Table C3-7. Derivation of Inhalation TRVs for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Assumed Test Species Body Weight (kg)[a]	Published Test Benchmark/Unit/Duration		Molecular Weight	Converted Test Benchmark (mg/m^3)[b]	Test Duration[c]	Test Endpoint	Effect	Source	Proposed Duration Conversion Factor DCF[d]	Proposed Endpoint Conversion Factor ECF[e]	TRV (mg/m^3) = Benchmark x DCF x ECF
Bromoform	75-25-2	Mammal	Unspecified	1.21E+02	mg/m3	2.53E+02	1.21E+02	Acute	LC50	Mortality	RTECS	1.00E-01	1.00E-02	1.21E-01
Bromomethane	74-83-9	Mouse	3.00E-02	2.00E+02	ppm/6H/14D-I	9.50E+01	7.77E+02	Subchronic	TCLo	Mortality	RTECS	1.00E-01	1.00E-01	7.77E+00
Carbon tetrachloride	56-23-5	Rat	3.50E-01	2.50E+02	ppm/8H/10 15D	1.54E+02	1.57E+03	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	1.57E+01
Chlorodibromomethane	124-48-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Chlorodifluoromethane	75-45-6	Rat	3.50E-01	5.00E+00	ppm/6-15D	8.65E+01	1.77E+01	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	1.77E-01
Chloroethane	75-00-3	Rat	3.50E-01	6.00E+01	mg/m3/4H	6.45E+01	6.00E+01	Acute	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	6.00E-01
Chloroform	67-66-3	Rat	3.50E-01	2.01E+04	ug/m3/1H/7-14D	1.19E+02	2.01E+01	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	2.01E-01
Chloromethane	74-87-3	Mouse	3.00E-02	5.00E+02	ppm/6H/6-17D	5.05E+01	1.03E+03	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	1.03E+01
Chloropentafluoroethane	76-15-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
cis-1,2-Dichloroethene	156-59-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
cis-1,3-Dichloropropene	10061-01-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Cyanogen bromide	506-68-3	Mouse	3.00E-02	5.00E+02	mg/m3/10M	1.06E+02	5.00E+02	Chronic	LCLo	Mortality	RTECS	1.00E+00	1.00E-01	5.00E+01
Cyanogen chloride	506-77-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Dichlorodifluoromethane	75-71-8	Rat	3.50E-01	4.00E+03	mg/m3/90D-C	1.21E+02	4.00E+03	Chronic	TCLo	Mortality	RTECS	1.00E+00	1.00E-01	4.00E+02
Dichlorofluoromethane	75-43-4	Rat	3.50E-01	1.00E+04	ppm/6H	1.03E+02	4.21E+04	Acute	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	4.21E+02
Dichloromethane	75-09-2	Rat	3.50E-01	1.25E+03	ppm/7H/6-15D	8.49E+01	4.34E+03	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	4.34E+01
Difluorodibromomethane	75-61-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Hexafluoroacetone	684-16-2	Rat	3.50E-01	1.20E+01	ppm/6H/30D	1.66E+02	8.15E+01	Chronic	TCLo	Reproduction	RTECS	1.00E+00	1.00E-01	8.15E+00
Iodomethane	74-88-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Methylene bromide	74-95-3	Rat	3.50E-01	4.00E+01	gm/m3/2H	1.74E+02	4.00E-02	Acute	LC50	Mortality	RTECS	1.00E-01	1.00E-02	4.00E-05
Pentachloroethane	76-01-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
trans-1,2-Dichloroethene	156-60-5	Rat	3.50E-01	6.00E+03	ppm/6H	9.70E+01	2.38E+04	Chronic	TCLo	Reproduction	RTECS	1.00E+00	1.00E-01	2.38E+03
trans-1,3-Dichloropropene	10061-02-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Trichloroacetic acid	76-03-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Trichlorofluoroethane	27154-33-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Trichlorofluoromethane	75-69-4	Rat	3.50E-01	1.20E+04	ppm/4H/10D-I	1.37E+02	6.74E+04	Subchronic	TCLo	Organ dysfunction	RTECS	1.00E-01	1.00E-01	6.74E+02

Table C3-7. Derivation of Inhalation TRVs for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Assumed Test Species Body Weight (kg)[a]	Published Test Benchmark/Unit/Duration		Molecular Weight	Converted Test Benchmark (mg/m³)[b]	Test Duration[c]	Test Endpoint	Effect	Source	Proposed Duration Conversion Factor DCF[d]	Proposed Endpoint Conversion Factor ECF[e]	TRV (mg/m³) = Benchmark x DCF x ECF
Trifluorobromomethane	75-63-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Dioxin and Furan Compounds														
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	35822-46-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	39227-28-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	57653-85-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	19408-74-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	40321-76-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1746-01-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Dibenzofuran	132-64-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Octachlorodibenzo(p)dioxin	3268-87-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Octachlorodibenzofuran	39001-02-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
PCBs														
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data

Table C3-7. Derivation of Inhalation TRVs for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Assumed Test Species Body Weight (kg)[a]	Published Test Benchmark/Unit/Duration		Molecular Weight	Converted Test Benchmark (mg/m^3)[b]	Test Duration[c]	Test Endpoint	Effect	Source	Proposed Duration Conversion Factor DCF[d]	Proposed Endpoint Conversion Factor ECF[e]	TRV (mg/m^3) = Benchmark x DCF x ECF
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2',3,4,4',5-Pentachlorobiphenyl	no cas #	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
3,3',4,4',5-Pentachlorobiphenyl	no cas #	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Polychlorinated biphenyls (PCBs)[k]	1336-36-3	Rat	3.50E-01	9.30E+02	ug/m3/8H/20W-I	--	9.30E-01	Chronic	TCLo	Organ dysfunction	RTECS	1.00E+00	1.00E-01	9.30E-02
		Phthalates												
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	Rat	3.50E-01	9.40E+02	mg/m3/6H/4W-I	3.91E+02	9.40E+02	Chronic	TCLo	Organ dysfunction	RTECS	1.00E+00	1.00E-01	9.40E+01
Butylbenzyl phthalate	85-68-7	Rat	3.50E-01	2.10E+03	mg/m3/6H/4W-I	3.12E+02	2.10E+03	Chronic	TCLo	Metabolism	RTECS	1.00E+00	1.00E-01	2.10E+02
Dibutyl phthalate	84-74-2	Rat	3.50E-01	9.00E+02	mg/m3/6H/35D-I	2.78E+02	9.00E+02	Chronic	TCLo	Metabolism	RTECS	1.00E+00	1.00E-01	9.00E+01
Diethyl phthalate	84-66-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Dimethylphthalate	131-11-3	Cat	4.50E+00	9.30E+03	mg/m3/6.5H	1.94E+02	9.30E+03	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	9.30E+01
n-Dioctyl phthalate	117-84-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Light Polycyclic Aromatic Hydrocarbons (molecular weight <200 g/mole)														
2-Chloronapthalene	91-58-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2-Methyl naphthalene	91-57-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
5-Nitroacenaphthene	602-87-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Acenaphthene	83-32-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Acenaphthylene	208-96-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Anthracene	120-12-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Fluorene	86-73-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Indene	95-13-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data

Table C3-7. Derivation of Inhalation TRVs for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Assumed Test Species Body Weight (kg)[a]	Published Test Benchmark/Unit/Duration		Molecular Weight	Converted Test Benchmark (mg/m^3)[b]	Test Duration[c]	Test Endpoint	Effect	Source	Proposed Duration Conversion Factor DCF[d]	Proposed Endpoint Conversion Factor ECF[e]	TRV (mg/m^3) = Benchmark x DCF x ECF
Naphthalene	91-20-3	Mouse	3.00E-02	3.00E+01	ppm/6H/2Y-I	1.28E+02	1.57E+02	Chronic	TCLo	Tumorigenic	RTECS	1.00E+00	1.00E-01	1.57E+01
Phenanthrene	85-01-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Pyrene	129-00-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Heavy Polycyclic Aromatic Hydrocarbons (molecular weight >200 g/mole)														
3-Methylcholanthrene	56-49-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
5-Methylchrysene	3697-24-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Benzo(a)anthracene	56-55-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Benzo(a)pyrene	50-32-8	Mouse	3.00E-02	2.00E+02	ng/m3/6H/13W-I	2.52E+02	2.00E-04	Chronic	TCLo	Tumorigenic	RTECS	1.00E+00	1.00E-01	2.00E-05
Benzo(b)fluoranthene	205-99-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Benzo(e)pyrene	192-97-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Benzo(g,h,i)perylene	191-24-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Benzo(j)fluoranthene	205-82-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Benzo(k)fluoranthene	207-08-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Benzo[a,i]pyrene	191-30-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Chrysene	218-01-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Dibenz(a,h)anthracene	53-70-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Dibenz[a,h]acridine	226-36-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Dibenz[a,j]acridine	224-42-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Dibenzo(a,e)fluoranthene	5385-75-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Dibenzo(a,h)fluoranthene	no cas #	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Dibenzo[a,e]pyrene	192-65-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Dibenzo[a,h]pyrene	189-64-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Dibenzo[a,i]pyrene	189-55-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Fluoranthene	206-44-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Hexachloronaphthalene	1335-87-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Indeno(1,2,3-cd)pyrene	193-39-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Octachloronaphthalene	2234-13-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Pentachloronaphthalene	1321-64-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Tetrachloronaphthalene	1335-88-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Trichloronaphthalene	1321-65-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data

Table C3-7. Derivation of Inhalation TRVs for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Assumed Test Species Body Weight (kg)[a]	Published Test Benchmark/Unit/Duration		Molecular Weight	Converted Test Benchmark (mg/m³)[b]	Test Duration[c]	Test Endpoint	Effect	Source	Proposed Duration Conversion Factor DCF[d]	Proposed Endpoint Conversion Factor ECF[e]	TRV (mg/m³) = Benchmark x DCF x ECF
Light Substituted Benzene Compounds (MW <200 g/mole)														
1,2,3-Trichlorobenzene	87-61-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
1,2,4-Trichlorobenzene	120-82-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
1,2,4-Trimethyl benzene	95-63-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
1,2-Dichlorobenzene	95-50-1	Rat	3.50E-01	2.00E+02	ppm/6H/6-15 D	1.47E+02	1.20E+03	Subchronic	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	1.20E+01
1,3,5-Trimethyl benzene	108-67-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
1,3-Dichlorobenzene	541-73-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
1,3-Dinitrobenzene	99-65-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
1,4-Dichlorobenzene	106-46-7	Rabbit	3.80E+00	8.00E+02	ppm/6H/6-18D	1.47E+02	4.81E+03	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	4.81E+01
1,4-Dinitrobenzene	100-25-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2,4,5-Trichlorophenol	95-95-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2,4,6-Trichlorophenol	88-06-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2,4-Dichlorophenol	120-83-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2,4-Dimethylphenol	105-67-9	Rat	3.50E-01	3.00E+01	mg/m3	1.22E+02	3.00E+01	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	3.00E-01
2,4-Dinitrophenol	51-28-5	Dog	1.27E+01	3.00E+02	mg/m3/30 M	1.84E+02	3.00E+02	Chronic	LCLo	Mortality	RTECS	1.00E+00	1.00E-01	3.00E+01
2,4-Dinitrotoluene	121-14-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2,6-Dinitrotoluene	606-20-2	Rat	3.50E-01	2.40E+02	mg/m3/6H	1.82E+02	2.40E+02	Acute	LC50	Mortality	RTECS	1.00E-01	1.00E-02	2.40E-01
2-Chlorophenol	95-57-8	Rat	3.50E-01	3.10E+03	mg/m3/6H	1.29E+02	3.10E+03	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	3.10E+01
2-Chlorotoluene	95-49-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2-Nitrophenol	88-75-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
4,6-Dinitro-o-cresol	534-52-1	Cat	4.50E+00	2.00E+00	mg/m3/4H/4W	1.98E+02	2.00E+00	Chronic	TCLo	Organ dysfunction	RTECS	1.00E+00	1.00E-01	2.00E-01
4-Chlorotoluene	106-43-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
4-Nitrophenol	100-02-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
alpha-Methylstyrene	98-83-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Aniline	62-53-3	Rat	3.50E-01	5.00E+00	ppm	9.31E+01	1.90E+01	Chronic	--	Organ dysfunction	IRIS	1.00E+00	1.00E-01	1.90E+00
Benzotrichloride	98-07-7	Mouse	3.00E-02	1.62E+00	ppm/20M/22W	1.95E+02	1.30E+01	Chronic	TCLo	Tumorigenic	RTECS	1.00E+00	1.00E-01	1.30E+00
Benzyl chloride	100-44-7	Mouse	3.00E-02	4.60E+01	ppm/6H/14 D	1.27E+02	2.38E+02	Subchronic	TCLo	Sense organs	RTECS	1.00E-01	1.00E-01	2.38E+00
Bromobenzene	108-86-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data

Table C3-7. Derivation of Inhalation TRVs for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Assumed Test Species Body Weight (kg)[a]	Published Test Benchmark/Unit/Duration		Molecular Weight	Converted Test Benchmark (mg/m^3)[b]	Test Duration[c]	Test Endpoint	Effect	Source	Proposed Duration Conversion Factor DCF[d]	Proposed Endpoint Conversion Factor ECF[e]	TRV (mg/m^3) = Benchmark x DCF x ECF
Chlorobenzene	108-90-7	Rabbit	3.80E+00	1.00E+01	ppm/6H/6-18D	1.13E+02	4.60E+01	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	4.60E-01
Cumene	98-82-8	Mouse	3.00E-02	1.00E+01	gm/m3/7H	1.20E+02	1.00E-02	Acute	LC50	Mortality	RTECS	1.00E-01	1.00E-02	1.00E-05
m-Cresol	108-39-4	Rabbit	3.80E+00	2.00E+00	mg/m3/24H/6W	1.08E+02	2.00E+00	Chronic	TCLo	Organ dysfunction	RTECS	1.00E+00	1.00E-01	2.00E-01
n-Butyl benzene	104-51-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Nitrobenzene	98-95-3	Rat	3.50E-01	1.26E+03	ug/m3/4H/1-21D	1.23E+02	1.26E+00	Chronic	TCLo	Reproduction	RTECS	1.00E+00	1.00E-01	1.26E-01
n-Propyl benzene	103-65-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
o-Cresol	95-48-7	Mouse	3.00E-02	1.79E+02	mg/m3/2H	1.08E+02	1.79E+02	Acute	LC50	Mortality	RTECS	1.00E-01	1.00E-02	1.79E-01
o-Dinitrobenzene	528-29-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
o-Nitroaniline	88-74-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
o-Toluidine	95-53-4	Rat	3.50E-01	8.62E+02	ppm/4H	1.07E+02	3.77E+03	Acute	LC50'	Mortality	RTECS	1.00E-01	1.00E-02	3.77E+00
p-Chloroaniline	106-47-8	Rat	3.50E-01	2.34E+03	mg/m3/4H	1.28E+02	2.34E+03	Acute	LC50	Mortality	RTECS	1.00E-01	1.00E-02	2.34E+00
p-Cresol	106-44-5	Rat	3.50E-01	7.10E+02	mg/m3/1H	1.08E+02	7.10E+02	Acute	LC50	Mortality	RTECS	1.00E-01	1.00E-02	7.10E-01
Phenol	108-95-2	Rat	3.50E-01	5.00E+00	mg/m3/4H//17W-I	9.41E+01	5.00E+00	Chronic	TCLo	Organ dysfunction	RTECS	1.00E+00	1.00E-01	5.00E-01
p-Nitrochlorobenzene	100-00-5	Rat	3.50E-01	2.40E+01	ppm/6H/13W	1.58E+02	1.55E+02	Chronic	TCLo	Reproduction	RTECS	1.00E+00	1.00E-01	1.55E+01
p-Toluidine	106-49-0	Rat	3.50E-01	6.40E+02	mg/m3/1H	1.07E+02	6.40E+02	Acute	LC50	Mortality	RTECS	1.00E-01	1.00E-02	6.40E-01
sec-Butyl benzene	135-98-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
tert-Butyl benzene	98-06-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Toluene-2,6-diamine	823-40-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Trimethyl benzene	25551-13-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Other Light Semivolatile Compounds (MW+A275 <200 g/mole)														
1,1'-Biphenyl	92-52-4	Rat	3.50E-01	2.00E+02	mg/m3	1.54E+02	2.00E+02	Chronic	LC	Mortality	RTECS	1.00E+00	1.00E-01	2.00E+01
1,1-Dimethylhydrazine	57-14-7	Rat	3.50E-01	3.40E+01	gm/m3/13M/40D	6.01E+01	3.40E+04	Chronic	TCLo	Organ dysfunction	RTECS	1.00E+00	1.00E-01	3.40E+03
1,2-Dimethylhydrazine	540-73-8	Rat	3.50E-01	2.80E+02	ppm/4H	6.01E+01	6.88E+02	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	6.88E+00
1,2-Diphenylhydrazine	122-66-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
1,3-Propane sultone	1120-71-4	Rat	3.50E-01	2.14E+03	mg/m3/6H	1.22E+02	2.14E+03	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	2.14E+01

Table C3-7. Derivation of Inhalation TRVs for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Assumed Test Species Body Weight (kg)[a]	Published Test Benchmark/Unit/Duration		Molecular Weight	Converted Test Benchmark (mg/m^3)[b]	Test Duration[c]	Test Endpoint	Effect	Source	Proposed Duration Conversion Factor DCF[d]	Proposed Endpoint Conversion Factor ECF[e]	TRV (mg/m^3) = Benchmark x DCF x ECF
2,4-Toluene diisocyante	584-84-9	Rat	3.50E-01	2.04E+02	ug/m3/24H/84D	1.74E+02	2.04E-01	Chronic	TCLo	Muscle tissue	RTECS	1.00E+00	1.00E-01	2.04E-02
2-Chloroacetophenone	532-27-4	Rat	3.50E-01	1.00E+00	mg/m3	1.55E+02	1.00E+00	Chronic	--	Nasal tissue	IRIS	1.00E+00	1.00E-01	1.00E-01
2-Propenoic acid	79-10-7	Monkey	5.00E+00	7.50E+01	ppm/6H	7.21E+01	2.21E+02	Acute	LC	Sense organs	RTECS	1.00E-01	1.00E-01	2.21E+00
4,4-Methylenedianiline	101-77-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Acetophenone	98-86-2	Rat	3.50E-01	2.10E+02	ppm/8H	1.21E+02	1.03E+03	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	1.03E+01
Benzoic acid	65-85-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
bis(2-Chloroethoxy)methane	111-91-1	Rat	3.50E-01	6.20E+01	ppm/4H	1.73E+02	4.39E+02	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	4.39E+00
bis(2-Chloroethyl) ether	111-44-4	Rat	3.50E-01	3.30E+02	mg/m3/4H	1.43E+02	3.30E+02	Acute	LC50	Mortality	RTECS	1.00E-01	1.00E-02	3.30E-01
Chlorocyclopentadiene	41851-50-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Cyclohexanol	108-93-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Dichloroisopropyl ether	108-60-1	Rat	3.50E-01	7.00E+02	ppm/5H	1.71E+02	4.90E+03	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	4.90E+01
Dichloromethyl ether	542-88-1	Rat	3.50E-01	1.00E+02	ppb/6H/4W I	1.15E+02	4.70E-01	Chronic	TCLo	Tumorigenic	RTECS	1.00E+00	1.00E-01	4.70E-02
Dichloropentadiene	no cas #	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Dimethyl sulfate	77-78-1	Rat	3.50E-01	1.70E+01	mg/m3/19W	1.26E+02	1.70E+01	Chronic	TCLo	Tumorigenic	RTECS	1.00E+00	1.00E-01	1.70E+00
Dimethylaniline	121-69-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Di-n-propylnitrosamine	621-64-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Diphenyl ether	101-84-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Epichlorohydrin	106-89-8	Rat/Mouse	3.00E-02	5.00E+00	ppm	9.25E+01	1.89E+01	Chronic	--	Nasal tissue	IRIS	1.00E+00	1.00E-01	1.89E+00
Ethyl carbamate (urethane)	51-79-6	Mouse	3.00E-02	1.38E+02	ppm/130d-1	8.91E+01	5.03E+02	Chronic	TCLo	Tumorigenic	RTECS	1.00E+00	1.00E-01	5.03E+01
Ethyl methanesulfonate	62-50-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Ethylene dibromide	106-93-4	Mouse	3.00E-02	3.00E+00	ppm/6H/13W-1	1.88E+02	2.31E+01	Chronic	TCLo	Mortality	RTECS	1.00E+00	1.00E-01	2.31E+00
Ethylene glycol	107-21-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Ethylene glycol monobutyl ether	111-76-2	Rat	3.50E-01	3.80E+02	mg/m3	1.18E+02	3.80E+02	Chronic	--	Blood effects	IRIS	1.00E+00	1.00E-01	3.80E+01
Ethylene glycol monoethyl ether acetate	111-15-9	Rat	3.50E-01	1.21E+04	mg/m3/8H	1.32E+02	1.21E+04	Acute	LC50	Mortality	RTECS	1.00E-01	1.00E-02	1.21E+01
Ethylene thiourea	96-45-7	Rat	3.50E-01	2.72E+04	ug/m3/3H	1.02E+02	2.72E+01	Acute	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	2.72E-01
Furfural	98-01-1	Rat	3.50E-01	1.75E+02	ppm/6H	9.61E+01	6.88E+02	Acute	LC50	Mortality	RTECS	1.00E-01	1.00E-02	6.88E-01
Maleic hydrazide	123-33-1	Rat	3.50E-01	2.00E+01	gm/m3	1.12E+02	2.00E-02	Acute	LC50	Mortality	RTECS	1.00E-01	1.00E-02	2.00E-05
Malononitrile	109-77-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data

Table C3-7. Derivation of Inhalation TRVs for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Assumed Test Species Body Weight (kg)[a]	Published Test Benchmark/Unit/Duration		Molecular Weight	Converted Test Benchmark (mg/m^3)[b]	Test Duration[c]	Test Endpoint	Effect	Source	Proposed Duration Conversion Factor DCF[d]	Proposed Endpoint Conversion Factor ECF[e]	TRV (mg/m^3) = Benchmark x DCF x ECF
Methyl styrene (mixed isomers)	25013-15-4	Mouse	3.00E-02	2.00E+02	ppm/6H/15D-I	1.18E+02	9.67E+02	Subchronic	TCLo	Organ dysfunction	RTECS	1.00E-01	1.00E-01	9.67E+00
Methylhydrazine	60-34-4	Dog	1.27E+01	2.00E-01	ppm/24H/30W-C	4.61E+01	3.77E-01	Chronic	TCLo	Metabolism	RTECS	1.00E+00	1.00E-01	3.77E-02
N,N-Diphenylamine	122-39-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Nitric acid, propyl ester	627-13-4	Rat	3.50E-01	5.82E+03	ppm/4H	1.05E+02	2.50E+04	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	2.50E+02
N-Nitrosodi-n-butylamine	924-16-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
N-Nitrosomorpholine	59-89-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
N-Nitroso-N,N-dimethylamine	62-75-9	Mouse	3.00E-02	2.00E+02	ug/m3/45W-C	7.41E+01	2.00E-01	Chronic	TCLo	Tumorigenic	RTECS	1.00E+00	1.00E-01	2.00E-02
o-Anisidine	90-04-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Oxalic acid	144-62-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Phthalic anhydride	85-44-9	Rat	3.50E-01	1.00E+00	mg/m3	1.48E+02	1.00E+00	Acute	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	1.00E-02
p-Phthalic acid	100-21-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Pyridine	110-86-1	Rat	3.50E-01	1.00E+00	mg/m3/24H/60D-C	7.91E+01	1.00E+00	Chronic	TCLo	Blood effects	RTECS	1.00E+00	1.00E-01	1.00E-01
Quinoline	91-22-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Quinone	106-51-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Safrole	94-59-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Tetrahydrofuran	109-99-9	Mouse	3.00E-02	1.80E+03	ppm/6H/6-17D	7.21E+01	5.31E+03	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	5.31E+01
Other Heavy Semivolatile Compounds (MW >200 g/mole)														
1,2,4,5-Tetrachlorobenzene	95-94-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
1,3,5-Trinitrobenzene	99-35-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2-sec-Butyl-4,6-dinitrophenol	88-85-7	Cat	4.50E+00	4.50E+01	mg/m3/3H	2.40E+02	4.50E+01	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	4.50E-01
3,3-Dichlorobenzidine	91-94-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
3,3'-Dimethoxybenzidine	119-90-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
4-Bromophenylphenyl ether	101-55-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Ammonium perfluorooctanoate	3825-26-1	Rat	3.50E-01	2.50E+01	mg/m3/6H/6-15D	4.31E+02	2.50E+01	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	2.50E-01

Table C3-7. Derivation of Inhalation TRVs for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Assumed Test Species Body Weight (kg)[a]	Published Test Benchmark/Unit/Duration		Molecular Weight	Converted Test Benchmark (mg/m^3)[b]	Test Duration[c]	Test Endpoint	Effect	Source	Proposed Duration Conversion Factor DCF[d]	Proposed Endpoint Conversion Factor ECF[e]	TRV (mg/m^3) = Benchmark x DCF x ECF
Azobenzene	103-33-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Captan	133-06-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Chlorobenzilate	510-15-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Dibutylphosphate	107-66-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Dimethyl aminoazobenzene	60-11-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Hexachlorobenzene	118-74-1	Rabbit	3.80E+00	1.80E+03	mg/m3	2.85E+02	1.80E+03	Acute	LC50	Mortality	RTECS	1.00E-01	1.00E-02	1.80E+00
Hexachlorobutadiene	87-68-3	Rat	3.50E-01	1.50E+01	ppm/6H/6-20D	2.61E+02	1.60E+02	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	1.60E+00
Hexachlorocyclopentadiene	77-47-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Hexachloroethane	67-72-1	Guinea pig	3.50E-01	2.60E+02	ppm/6H/6 W-I	2.37E+02	2.52E+03	Chronic	TCLo	Organ dysfunction	RTECS	1.00E+00	1.00E-01	2.52E+02
Hexachlorophene	70-30-4	Rat	3.50E-01	3.35E+04	ug/m3/24H	4.07E+02	3.35E+01	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	3.35E-01
Hexamethylene-1,5-diisocyanate	822-06-0	Rat	3.50E-01	5.00E-03	ppm	1.68E+02	3.40E-02	Chronic	--	Nasal tissue	IRIS	1.00E+00	1.00E-01	3.40E-03
Mirex	2385-85-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Nitrofen	1836-75-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Pentachlorobenzene	608-93-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Pentachloronitrobenzene	82-68-8	Rat	3.50E-01	1.40E+03	mg/m3	2.95E+02	1.40E+03	Acute	LC50	Mortality	RTECS	1.00E-01	1.00E-02	1.40E+00
Pentachlorophenol	87-86-5	Mouse	3.00E-02	2.25E+02	mg/m3	2.66E+02	2.25E+02	Acute	LC50	Mortality	RTECS	1.00E-01	1.00E-02	2.25E-01
Picric acid	88-89-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Pronamide	23950-58-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Strychnine	57-24-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Terphenyls	26140-60-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Tributyl phosphate	126-73-8	Rat	3.50E-01	5.00E+02	mg/m3/6H/30D-I	2.66E+02	5.00E+02	Chronic	TCLo	Organ dysfunction	RTECS	1.00E+00	1.00E-01	5.00E+01
Trifluralin	1582-09-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Triphenylamine	603-34-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Herbicides and Organochlorinated Pesticides														
2,4,5-T	93-76-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
2,4-D and esters	94-75-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
4,4-DDD	72-54-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data

Table C3-7. Derivation of Inhalation TRVs for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Assumed Test Species Body Weight (kg)[a]	Published Test Benchmark/Unit/Duration		Molecular Weight	Converted Test Benchmark (mg/m³)[b]	Test Duration[c]	Test Endpoint	Effect	Source	Proposed Duration Conversion Factor DCF[d]	Proposed Endpoint Conversion Factor ECF[e]	TRV (mg/m³) = Benchmark x DCF x ECF
4,4-DDE	72-55-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
4,4-DDT	50-29-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Aldrin	309-00-2	Rat	3.50E-01	5.80E+03	ug/m3/4H	3.65+02	5.80E+00	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	5.80E-02
alpha-BHC	319-84-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
beta-BHC	319-85-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Chlordane	57-74-9	Rat	3.50E-01	2.82E+04	ug/m3/8H/28D-I	4.10E+02	2.82E+01	Chronic	TCLo	Organ dysfunction	RTECS	1.00E+00	1.00E-01	2.82E+00
Delta-BHC	319-86-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Dieldrin	60-57-1	Rat	3.50E-01	1.30E+01	mg/m3/4H	3.81E+02	1.30E+01	Acute	LC50	Mortality	RTECS	1.00E-01	1.00E-02	1.30E-02
Endothall	145-73-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Endrin	72-20-8	Mouse	3.00E-02	3.60E-01	ppm/7H/22W-I	3.81E+02	5.61E+00	Chronic	TCLo	Mortality	RTECS	1.00E+00	1.00E-01	5.61E-01
gamma-BHC (Lindane)	58-89-9	Cat	4.50E+00	2.00E+00	mg/m3/4H/9W-I	2.91E+02	2.00E+00	Chronic	TCLo	Immune system	RTECS	1.00E+00	1.00E-01	2.00E-01
Heptachlor	76-44-8	Cat	4.50E+00	1.50E+02	mg/m3/4H	3.73E+02	1.50E+02	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	1.50E+00
Isodrin	465-73-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Methoxychlor	72-43-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Silvex (2,4,5-TP)	93-72-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Toxaphene	8001-35-2	Mouse	3.00E-02	2.00E+03	mg/m3/2H	4.14E+02	2.00E+03	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	2.00E+01
Inorganic Chemicals and Compounds														
		Metals												
Aluminum	7429-90-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Antimony	7440-36-0	Rat	3.50E-01	5.00E+01	mg/m3/7H/52W-I	1.22E+02	5.00E+01	Chronic	TCLo	Tumorigenic	RTECS	1.00E+00	1.00E-01	5.00E+00
Arsenic	7440-38-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Barium	7440-39-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Beryllium	7440-41-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Bismuth	7440-69-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Boron	7440-42-8	Rat	3.50E-01	1.06E+04	ug/m3/4H/17W-I	1.08E+01	1.06E+01	Chronic	TCLo	Metabolism	RTECS	1.00E+00	1.00E-01	1.06E+00
Cadmium	7440-43-9	Rat	3.50E-01	2.50E+01	mg/m3/30M/60D-I	1.12E+02	2.50E+01	Chronic	LC50	Mortality	RTECS	1.00E+00	1.00E-02	2.50E-01

Table C3-7. Derivation of Inhalation TRVs for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Assumed Test Species Body Weight (kg)[a]	Published Test Benchmark/Unit/Duration		Molecular Weight	Converted Test Benchmark (mg/m^3)[b]	Test Duration[c]	Test Endpoint	Effect	Source	Proposed Duration Conversion Factor DCF[d]	Proposed Endpoint Conversion Factor ECF[e]	TRV (mg/m^3) = Benchmark x DCF x ECF
Calcium	7440-70-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Chromium (and VI)	18540-29-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Cobalt	7440-48-4	Rat	3.50E-01	2.00E+02	mg/m3/17W-I	5.89E+01	2.00E+02	Chronic	TCLo	Organ dysfunction	RTECS	1.00E+00	1.00E-01	2.00E+01
Copper	7440-50-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Iron	7439-89-6	Rat	3.50E-01	2.50E+02	mg/m3/6H/4W-I	5.59E+01	2.50E+02	Chronic	TCLo	Organ dysfunction	RTECS	1.00E+00	1.00E-01	2.50E+01
Lead	7439-92-1	Rat	3.50E-01	3.00E+00	mg/m3/24H	2.07E+02	3.00E+00	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	3.00E-02
Lithium	7439-93-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Magnesium	7439-95-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Manganese	7439-96-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Mercury	7439-97-6	Rat	3.50E-01	8.90E+02	ng/m3/24H	2.01E+02	8.90E-04	Subchronic	TCLo	Reproduction	RTECS	1.00E-01	1.00E-01	8.90E-06
Molybdenum	7439-98-7	Rat	3.50E-01	1.95E+04	ug/m3	9.59E+01	1.95E+01	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	1.95E-01
Nickel	7440-02-0	Rat	3.50E-01	1.00E+02	ug/m3/24H/17W-C	5.87E+01	1.00E-01	Chronic	TCLo	Organ dysfunction	RTECS	1.00E+00	1.00E-01	1.00E-02
Potassium	7440-09-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Rhodium	7440-16-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Selenium	7782-49-2	Rat	3.50E-01	1.47E+02	mg/m3	7.90E+01	1.47E+02	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	1.47E+00
Silicon	7440-21-3	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Silver	7440-22-4	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Sodium	7440-23-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Strontium	7440-24-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Tantalum	7440-25-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Thallium	7440-28-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Tin	7440-31-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Tungsten	7440-33-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Uranium	7440-61-1	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Vanadium	7440-62-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Yttrium	7440-65-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Zinc	7440-66-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Zirconium	7440-67-7	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Non-metals and Anions														

Table C3-7. Derivation of Inhalation TRVs for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Assumed Test Species Body Weight (kg)[a]	Published Test Benchmark/Unit/Duration		Molecular Weight	Converted Test Benchmark (mg/m^3)[b]	Test Duration[c]	Test Endpoint	Effect	Source	Proposed Duration Conversion Factor DCF[d]	Proposed Endpoint Conversion Factor ECF[e]	TRV (mg/m^3) = Benchmark x DCF x ECF
Ammonia/Ammonium	7664-41-7	Rat	3.50E-01	3.00E+02	ppm/6H/5D-I	1.70E+01	2.09E+02	Subchronic	TCLo	Organ dysfunction	RTECS	1.00E-01	1.00E-01	2.09E+00
Bromide	24959-67-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Chloride	16887-00-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Cyanide	57-12-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Fluoride	16984-48-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Hydroxide	14280-30-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Iodine	7553-56-2	Rat	3.50E-01	1.37E+02	ppm/1H	2.54E+02	1.42E+03	Acute	LCLo	Mortality	RTECS	1.00E-01	1.00E-01	1.42E+01
Nitrate	14797-55-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Nitrite	14797-65-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Phosphate	14265-44-2	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Phosphorus	7723-14-0	Rabbit	3.80E+00	1.60E+02	mg/m3/30M/60D-I	1.24E+02	1.60E+02	Chronic	TCLo	Blood	RTECS	1.00E+00	1.00E-01	1.60E+01
Sulfate	14808-79-8	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Total Sulfur	63705-05-5	Mammal	unspecified	1.66E+03	mg/m3	3.21E+01	1.66E+03	Acute	LC50	Mortality	RTECS	1.00E-01	1.00E-02	1.66E+00
Priority Pollutants														
Carbon Dioxide	124-38-9	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Nitrogen Dioxide	10102-44-0	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Ozone	10028-15-6	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Particulate Matter	No CAS #	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Sulfur Dioxode	7446-09-5	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
Radionuclides														
Actinium-227	14952-40-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Americium-241	1596-10-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Americium-243	14993-75-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Antimony-125	14234-35-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Barium-137	13981-97-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Cadmium-113	None	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Carbon-14	14762-75-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Cesium-134	13967-70-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Cesium-137	10045-97-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Cobalt-60	10198-40-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

Table C3-7. Derivation of Inhalation TRVs for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Assumed Test Species Body Weight (kg)[a]	Published Test Benchmark/Unit/Duration		Molecular Weight	Converted Test Benchmark (mg/m^3)[b]	Test Duration[c]	Test Endpoint	Effect	Source	Proposed Duration Conversion Factor DCF[d]	Proposed Endpoint Conversion Factor ECF[e]	TRV (mg/m^3) = Benchmark x DCF x ECF
Curium-242	15510-73-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Curium-243	15757-87-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Curium-244	13981-15-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Europium-152	14683-23-9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Europium-154	15585-10-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Europium-155	14391-16-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Iodine-129	15046-84-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Neptunium-237	13994-20-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Nickel-59	14336-70-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Nickel-63	13981-37-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Niobium-93	7440-03-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Plutonium-238	13981-16-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Plutonium-239	15117-48-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Plutonium-240	14119-33-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Plutonium-241	14119-32-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Plutonium-242	13982-10-0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Protactinium-231	14331-85-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Radium-226	13982-63-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Radium-228	15262-20-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Ruthenium-106	13967-48-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Samarium-151	15715-94-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Selenium-79	None	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Strontium-90	10098-97-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Technetium-99	14133-79-7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Thorium-229	15594-54-4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Thorium-232	7440-29-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Tin-126	15832-50-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Tritium	10028-17-8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Uranium-232	14158-29-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Uranium-233	13968-55-3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Uranium-234	13966-29-5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

Table C3-7. Derivation of Inhalation TRVs for Mammal Test Species

Constituent of Potential Concern	CAS Registry Number	Test Species	Assumed Test Species Body Weight (kg)[a]	Published Test Benchmark/Unit/Duration		Molecular Weight	Converted Test Benchmark (mg/m^3)[b]	Test Duration[c]	Test Endpoint	Effect	Source	Proposed Duration Conversion Factor DCF[d]	Proposed Endpoint Conversion Factor ECF[e]	TRV (mg/m^3) = Benchmark x DCF x ECF
Uranium-235	15117-96-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Uranium-236	13982-70-2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Uranium-238	7440-61-1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Yttrium-90	10098-91-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Zirconium-93	15751-77-6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

[a] Body weights not reported; mouse, rat, rabbit and dog assumed to be EPA standard reference body weights as reported in Table 1 in Sample et al. (1996); other assumptions include guinea pig = rat; cat = red fox (Storm et al. 1976); monkey = rhesus monkey (Barsotti et al. 1976). Body weights and inhalation rates will be used to calculate TRV doses (mg/kg/d).

[b] mg/m^3 = ppm x (molecular weight/24450) x 1000 L/m^3 (Clayton and Clayton 1981); mg/m^3 = (ug/m^3)/1000 (ug/mg)

[c] if Published Test Duration < 24 hours = acute; if < 20 days = subchronic; if > 20 days = chronic; assumed to be chronic if no duration information except when Test Endpoint is LC50 assumed to be acute.

[d] DCF = 1 if chronic, 0.1 if subchronic or acute

[e] ECF = 1 if NOAEL, 0.1 if TCLo or LCLo, 0.01 if LC_{50}

TCLo = Toxic concentration, low

LC50 = Lethal concentration, 50 percent kill

LCLo = Lethal concentration, low

-- = no data; for test endpoint assumed to be TCLo

RTECS = Registry of Toxic Effects of Chemical Substances (NIOSH), downloaded in January 1999

IRIS = USEPA Integrated Risk Information System, downloaded in January 2000

No data = No inhalation data available from the available Iris or Rtecs source

Table C3-8. Sediment TRVs for Sediment-Dwelling Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Jones et al. 1997[b]		Recommended TRV[c]
				Ontario MOE (LOW)	NOAA (ER-L)	
Organic Compounds (All units are in ug/kg)						
Aromatic Halogenated Hydrocarbons						
4-Chloro-3-methylphenol	59-50-7	No data		No data	No data	
2,3,4,6-Tetrachlorophenol	58-90-2	No data		No data	No data	
Aromatic Nonhalogenated Hydrocarbons						
2-Nitrotoluene	88-72-2	No data		No data	No data	
4-Nitrobiphenyl	92-93-3	No data		No data	No data	
Benzaldehyde	100-52-7	No data		No data	No data	
Benzene	71-43-2	No data		No data	No data	
Benzyl alcohol	100-51-6	No data		No data	No data	
Ethyl benzene	100-41-4	No data		No data	No data	
m-Xylene	108-38-3	No data		No data	No data	
o-Xylene	95-47-6	No data		No data	No data	
p-Xylene	106-42-3	No data		No data	No data	
Styrene	100-42-5	No data		No data	No data	
Toluene	108-88-3	No data		No data	No data	
Non-aromatic Nonhalogenated Hydrocarbons						
1,2-Epoxybutane	106-88-7	No data		No data	No data	
1,3-Butadiene	106-99-0	No data		No data	No data	
1,4-Dioxane	123-91-1	2.17E+03[d]		No data	No data	2.17E+03
1-Methylpropyl alcohol	78-92-2	No data		No data	No data	
1-Nitropropane	108-03-2	No data		No data	No data	
2,2,4-Trimethylpentane	540-84-1	No data		No data	No data	
2-Butanone	78-93-3	No data		No data	No data	
2-Butenaldehyde (2-Butenal)	4170-30-3	No data		No data	No data	
2-Ethoxyethanol	110-80-5	No data		No data	No data	
2-Heptanone	110-43-0	No data		No data	No data	
2-Hexanone	591-78-6	No data		No data	No data	
2-Methoxyethanol	109-86-4	No data		No data	No data	
2-Methyl-2-propanol	75-65-0	No data		No data	No data	

Table C3-8. Sediment TRVs for Sediment-Dwelling Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Jones et al. 1997[b]		Recommended TRV[c]
				Ontario MOE (LOW)	NOAA (ER-L)	
2-Methyl-2-propenenitrile	126-98-7	No data		No data	No data	
2-Methylaziridine	75-55-8	No data		No data	No data	
2-Methylpropyl alcohol	78-83-1	No data		No data	No data	
2-Pentanone	107-87-9	No data		No data	No data	
2-Propanone (Acetone)	67-64-1	5.71E+01[d]		No data	No data	5.71E+01
2-Propene-1-ol	107-18-6	No data		No data	No data	
2-Propyl alcohol	67-63-0	No data		No data	No data	
3-Heptanone	106-35-4	No data		No data	No data	
3-Methyl-1-butanol	123-51-3	No data		No data	No data	
3-Methyl-2-butanone	563-80-4	No data		No data	No data	
3-Pentanone	96-22-0	No data		No data	No data	
4-Heptanone	123-19-3	No data		No data	No data	
4-Methyl-2-pentanone	108-10-1	No data		No data	No data	
4-Methyl-3-penten-2-one	141-79-7	No data		No data	No data	
5-Methyl-2-hexanone	110-12-3	No data		No data	No data	
Acetaldehyde	75-07-0	No data		No data	No data	
Acetamide	60-35-5	No data		No data	No data	
Acetic acid	64-19-7	No data		No data	No data	
Acetic acid ethyl ester	141-78-6	No data		No data	No data	
Acetic acid n-butyl ester	123-86-4	No data		No data	No data	
Acetonitrile	75-05-8	No data		No data	No data	
Acrolein	107-02-8	No data		No data	No data	
Acrylonitrile	107-13-1	2.31E+01[d]		No data	No data	2.31E+01
Bis(isopropyl)ether	108-20-3	No data		No data	No data	
Butane	106-97-8	No data		No data	No data	
Carbon disulfide	75-15-0	No data		No data	No data	
Cyanogen	460-19-5	No data		No data	No data	
Cyclohexane	110-82-7	No data		No data	No data	
Cyclohexanone	108-94-1	No data		No data	No data	
Cyclohexene	110-83-8	No data		No data	No data	

Table C3-8. Sediment TRVs for Sediment-Dwelling Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Jones et al. 1997[b]		Recommended TRV[c]
				Ontario MOE (LOW)	NOAA (ER-L)	
Cyclopentane	287-92-3	No data		No data	No data	
Ethyl alcohol	64-17-5	No data		No data	No data	
Ethyl ether	60-29-7	No data		No data	No data	
Ethyl methacrylate	97-63-2	No data		No data	No data	
Formaldehyde	50-00-0	5.20E+00[d]		No data	No data	5.20E+00
Formamide	75-12-7	No data		No data	No data	
Formic acid	64-18-6	No data		No data	No data	
Formic acid, methyl ester	107-31-3	No data		No data	No data	
Glycidylaldehyde	765-34-4	No data		No data	No data	
Methyl acetate	79-20-9	No data		No data	No data	
Methyl alcohol	67-56-1	No data		No data	No data	
Methyl isocyanate	624-83-9	No data		No data	No data	
Methyl methacrylate	80-62-6	No data		No data	No data	
Methyl tert-butyl ether	1634-04-4	No data		No data	No data	
Methylacetylene	74-99-7	No data		No data	No data	
Methylcyclohexane	108-87-2	No data		No data	No data	
N,N-Dimethylacetamide	127-19-5	No data		No data	No data	
n-Butyl alcohol	71-36-3	No data		No data	No data	
n-Heptane	142-82-5	No data		No data	No data	
n-Hexane	110-54-3	No data		No data	No data	
Nitromethane	75-52-5	No data		No data	No data	
n-Nonane	111-84-2	No data		No data	No data	
n-Octane	111-65-9	No data		No data	No data	
n-Pentane	109-66-0	No data		No data	No data	
n-Propionaldehyde	123-38-6	No data		No data	No data	
n-Propyl alcohol	71-23-8	No data		No data	No data	
n-Valeraldehyde	110-62-3	No data		No data	No data	
Oxirane	75-21-8	No data		No data	No data	
p-Cymene	99-87-6	No data		No data	No data	
Phosgene	75-44-5	No data		No data	No data	

Table C3-8. Sediment TRVs for Sediment-Dwelling Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Jones et al. 1997[b]		Recommended TRV[c]
				Ontario MOE (LOW)	NOAA (ER-L)	
Propargyl alcohol	107-19-7	No data		No data	No data	
Propionic acid	79-09-4	No data		No data	No data	
Propionitrile	107-12-0	No data		No data	No data	
Propylene gylcol monomethyl ether	107-98-2	No data		No data	No data	
p-tert-Butyltoluene	98-51-1	No data		No data	No data	
Triethylamine	121-44-8	No data		No data	No data	
Trimethylamine	75-50-3	No data		No data	No data	
Vinyl acetate	108-05-4	No data		No data	No data	
Non-aromatic Halogenated Hydrocarbons						
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	No data		No data	No data	
1,1,1,2-Tetrachloroethane	630-20-6	No data		No data	No data	
1,1,1-Trichloroethane	71-55-6	No data		No data	No data	
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	No data		No data	No data	
1,1,2,2-Tetrachloroethane	79-34-5	No data		No data	No data	
1,1,2,2-Tetrachloroethene	127-18-4	No data		No data	No data	
1,1,2-Trichloroethane	79-00-5	No data		No data	No data	
1,1,2-Trichloroethylene	79-01-6	No data		No data	No data	
1,1-Dichloroethane	75-34-3	No data		No data	No data	
1,1-Dichloroethene	75-35-4	No data		No data	No data	
1,2,2-Trichloro-1,1,2-trifluoroethane	76-13-1	No data		No data	No data	
1,2,3-Trichloropropane	96-18-4	No data		No data	No data	
1,2-Dibromo-3-chloropropane	96-12-8	No data		No data	No data	
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	No data		No data	No data	
1,2-Dichloroethane	107-06-2	No data		No data	No data	
1,2-Dichloroethylene	540-59-0	No data		No data	No data	
1,2-Dichloropropane	78-87-5	No data		No data	No data	
1,3-Dichloropropene	542-75-6	No data		No data	No data	
1,4-Dichloro-2-butene	764-41-0	No data		No data	No data	
1-Chloroethene	75-01-4	No data		No data	No data	
2,2-Dichloropropionic acid	75-99-0	No data		No data	No data	

Table C3-8. Sediment TRVs for Sediment-Dwelling Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Jones et al. 1997[b]		Recommended TRV[c]
				Ontario MOE (LOW)	NOAA (ER-L)	
2-Chloropropane	75-29-6	No data		No data	No data	
3-Chloropropene (allyl chloride)	107-05-1	No data		No data	No data	
Bromochloromethane	74-97-5	No data		No data	No data	
Bromodichloromethane	75-27-4	No data		No data	No data	
Bromoethene	593-60-2	No data		No data	No data	
Bromoform	75-25-2	No data		No data	No data	
Bromomethane	74-83-9	No data		No data	No data	
Carbon tetrachloride	56-23-5	No data		No data	No data	
Chlorodibromomethane	124-48-1	No data		No data	No data	
Chlorodifluoromethane	75-45-6	No data		No data	No data	
Chloroethane	75-00-3	No data		No data	No data	
Chloroform	67-66-3	5.94E+01[d]		No data	No data	5.94E+01
Chloromethane	74-87-3	No data		No data	No data	
Chloropentafluoroethane	76-15-3	No data		No data	No data	
cis-1,2-Dichloroethene	156-59-2	No data		No data	No data	
cis-1,3-Dichloropropene	10061-01-5	No data		No data	No data	
Cyanogen bromide	506-68-3	No data		No data	No data	
Cyanogen chloride	506-77-4	No data		No data	No data	
Dichlorodifluoromethane	75-71-8	No data		No data	No data	
Dichlorofluoromethane	75-43-4	No data		No data	No data	
Dichloromethane	75-09-2	No data		No data	No data	
Difluorodibromomethane	75-61-6	No data		No data	No data	
Hexafluoroacetone	684-16-2	No data		No data	No data	
Iodomethane	74-88-4	No data		No data	No data	
Methylene bromide	74-95-3	No data		No data	No data	
Pentachloroethane	76-01-7	No data		No data	No data	
trans-1,2-Dichloroethene	156-60-5	No data		No data	No data	
trans-1,3-Dichloropropene	10061-02-6	No data		No data	No data	
Trichloroacetic acid	76-03-9	No data		No data	No data	
Trichlorofluoroethane	27154-33-2	No data		No data	No data	

Table C3-8. Sediment TRVs for Sediment-Dwelling Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Jones et al. 1997[b]		Recommended TRV[c]
				Ontario MOE (LOW)	NOAA (ER-L)	
Trichlorofluoromethane	75-69-4	No data		No data	No data	
Trifluorobromomethane	75-63-8	No data		No data	No data	
Dioxin and Furan Compounds						
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	35822-46-9	No data		No data	No data	
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	No data		No data	No data	
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	No data		No data	No data	
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	39227-28-6	No data		No data	No data	
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	No data		No data	No data	
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	57653-85-7	No data		No data	No data	
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	No data		No data	No data	
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	19408-74-3	No data		No data	No data	
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	No data		No data	No data	
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	40321-76-4	No data		No data	No data	
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	No data		No data	No data	
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	No data		No data	No data	
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	No data		No data	No data	
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1746-01-6	4.10E-01[d]		No data	No data	4.10E-01
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	No data		No data	No data	
Dibenzofuran	132-64-9	No data		No data	No data	
Octachlorodibenzo(p)dioxin	3268-87-9	No data		No data	No data	
Octachlorodibenzofuran	39001-02-0	No data		No data	No data	
PCBs						
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	No data		No data	No data	
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	No data		No data	No data	
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	No data		No data	No data	
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	No data		No data	No data	
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	No data		No data	No data	
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	No data		No data	No data	
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	No data		No data	No data	
2',3,4,4',5-Pentachlorobiphenyl	no cas #	No data		No data	No data	

Table C3-8. Sediment TRVs for Sediment-Dwelling Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Jones et al. 1997[b]		Recommended TRV[c]
				Ontario MOE (LOW)	NOAA (ER-L)	
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	No data		No data	No data	
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	No data		No data	No data	
3,3',4,4',5-Pentachlorobiphenyl	no cas #	No data		No data	No data	
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	No data		No data	No data	
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	No data		No data	No data	
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	No data		No data	No data	
Polychlorinated biphenyls (PCBs)[k]	1336-36-3	No data		7.00E+01[e]	No data	7.00E+01
Phthalates						
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	1.33E+04[d]		No data	No data	1.33E+04
Butylbenzyl phthalate	85-68-7	No data		No data	No data	
Dibutyl phthalate	84-74-2	No data		No data	No data	
Diethyl phthalate	84-66-2	No data		No data	No data	
Dimethylphthalate	131-11-3	No data		No data	No data	
n-Dioctyl phthalate	117-84-0	1.16E+10[d]		No data	No data	1.16E+10
Light Polycyclic Aromatic Hydrocarbons (molecular weight <200 g/mole)						
2-Chloronapthalene	91-58-7	No data		No data	No data	
2-Methyl naphthalene	91-57-6	No data		No data	7.00E+01	7.00E+01
5-Nitroacenaphthene	602-87-9	No data		No data	No data	
Acenaphthene	83-32-9	No data		No data	4.60E+01	4.60E+01
Acenaphthylene	208-96-8	No data		No data	4.40E+01	4.40E+01
Anthracene	120-12-7	No data		2.20E+02	8.53E+01	2.20E+02
Fluorene	86-73-7	No data		1.90E+02	1.90E+01	1.90E+02
Indene	95-13-6	No data		No data	No data	
Naphthalene	91-20-3	No data		No data	1.60E+02	1.60E+02
Phenanthrene	85-01-8	No data		5.60E+02	2.40E+02	5.60E+02
Pyrene	129-00-0	No data		4.90E+02	6.65E+02	4.90E+02
Heavy Polycyclic Aromatic Hydrocarbons (molecular weight >200 g/mole)						
3-Methylcholanthrene	56-49-5	No data		No data	No data	
5-Methylchrysene	3697-24-3	No data		No data	No data	
Benzo(a)anthracene	56-55-3	1.90E+01		3.20E+02	2.61E+02	1.90E+01

Table C3-8. Sediment TRVs for Sediment-Dwelling Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Jones et al. 1997[b]		Recommended TRV[c]
				Ontario MOE (LOW)	NOAA (ER-L)	
Benzo(a)pyrene	50-32-8	8.40E+01		3.70E+02	4.30E+02	8.40E+01
Benzo(b)fluoranthene	205-99-2	3.70E+01		No data	No data	3.70E+01
Benzo(e)pyrene	192-97-2	No data		No data	No data	
Benzo(g,h,i)perylene	191-24-2	No data		1.70E+02	No data	1.70E+02
Benzo(j)fluoranthene	205-82-3	No data		No data	No data	
Benzo(k)fluoranthene	207-08-9	3.70E+01		2.40E+02	No data	3.70E+01
Benzo[a,i]pyrene	191-30-0	No data		No data	No data	
Chrysene	218-01-9	3.00E+01		3.40E+02	3.84E+02	3.00E+01
Dibenz(a,h)anthracene	53-70-3	1.00E+01		6.00E+01	6.34E+01	1.00E+01
Dibenz[a,h]acridine	226-36-8	No data		No data	No data	
Dibenz[a,j]acridine	224-42-0	No data		No data	No data	
Dibenzo(a,e)fluoranthene	5385-75-1	No data		No data	No data	
Dibenzo(a,h)fluoranthene	no cas #	No data		No data	No data	
Dibenzo[a,e]pyrene	192-65-4	No data		No data	No data	
Dibenzo[a,h]pyrene	189-64-0	No data		No data	No data	
Dibenzo[a,i]pyrene	189-55-9	No data		No data	No data	
Fluoranthene	206-44-0	No data		7.50E+02	6.00E+02	7.50E+02
Hexachloronaphthalene	1335-87-1	No data		No data	No data	
Indeno(1,2,3-cd)pyrene	193-39-5	3.00E+01		2.00E+02	No data	3.00E+01
Octachloronaphthalene	2234-13-1	No data		No data	No data	
Pentachloronaphthalene	1321-64-8	No data		No data	No data	
Tetrachloronaphthalene	1335-88-2	No data		No data	No data	
Trichloronaphthalene	1321-65-9	No data		No data	No data	
Light Substituted Benzene Compounds (MW <200 g/mole)						
1,2,3-Trichlorobenzene	87-61-6	No data		No data	No data	
1,2,4-Trichlorobenzene	120-82-1	No data		No data	No data	
1,2,4-Trimethyl benzene	95-63-6	No data		No data	No data	
1,2-Dichlorobenzene	95-50-1	No data		No data	No data	
1,3,5-Trimethyl benzene	108-67-8	No data		No data	No data	
1,3-Dichlorobenzene	541-73-1	No data		No data	No data	

Table C3-8. Sediment TRVs for Sediment-Dwelling Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Jones et al. 1997[b]		Recommended TRV[c]
				Ontario MOE (LOW)	NOAA (ER-L)	
1,3-Dinitrobenzene	99-65-0	2.14E+01[d]		No data	No data	2.14E+01
1,4-Dichlorobenzene	106-46-7	No data		No data	No data	
1,4-Dinitrobenzene	100-25-4	No data		No data	No data	
2,4,5-Trichlorophenol	95-95-4	No data		No data	No data	
2,4,6-Trichlorophenol	88-06-2	No data		No data	No data	
2,4-Dichlorophenol	120-83-2	No data		No data	No data	
2,4-Dimethylphenol	105-67-9	No data		No data	No data	
2,4-Dinitrophenol	51-28-5	No data		No data	No data	
2,4-Dinitrotoluene	121-14-2	4.69E+01[d]		No data	No data	4.69E+01
2,6-Dinitrotoluene	606-20-2	1.01E+02[d]		No data	No data	1.01E+02
2-Chlorophenol	95-57-8	No data		No data	No data	
2-Chlorotoluene	95-49-8	No data		No data	No data	
2-Nitrophenol	88-75-5	No data		No data	No data	
4,6-Dinitro-o-cresol	534-52-1	No data		No data	No data	
4-Chlorotoluene	106-43-4	No data		No data	No data	
4-Nitrophenol	100-02-7	No data		No data	No data	
alpha-Methylstyrene	98-83-9	No data		No data	No data	
Aniline	62-53-3	No data		No data	No data	
Benzotrichloride	98-07-7	No data		No data	No data	
Benzyl chloride	100-44-7	No data		No data	No data	
Bromobenzene	108-86-1	No data		No data	No data	
Chlorobenzene	108-90-7	No data		No data	No data	
Cumene	98-82-8	No data		No data	No data	
m-Cresol	108-39-4	No data		No data	No data	
n-Butyl benzene	104-51-8	No data		No data	No data	
Nitrobenzene	98-95-3	1.28E+03[d]		No data	No data	1.28E+03
n-Propyl benzene	103-65-1	No data		No data	No data	
o-Cresol	95-48-7	No data		No data	No data	
o-Dinitrobenzene	528-29-0	No data		No data	No data	
o-Nitroaniline	88-74-4	No data		No data	No data	

Table C3-8. Sediment TRVs for Sediment-Dwelling Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Jones et al. 1997[b]		Recommended TRV[c]
				Ontario MOE (LOW)	NOAA (ER-L)	
o-Toluidine	95-53-4	No data		No data	No data	
p-Chloroaniline	106-47-8	No data		No data	No data	
p-Cresol	106-44-5	No data		No data	No data	
Phenol	108-95-2	No data		No data	No data	
p-Nitrochlorobenzene	100-00-5	No data		No data	No data	
p-Toluidine	106-49-0	No data		No data	No data	
sec-Butyl benzene	135-98-8	No data		No data	No data	
tert-Butyl benzene	98-06-6	No data		No data	No data	
Toluene-2,6-diamine	823-40-5	No data		No data	No data	
Trimethyl benzene	25551-13-7	No data		No data	No data	
Other Light Semivolatile Compounds (molecular weight <200 g/mole)						
1,1'-Biphenyl	92-52-4	No data		No data	No data	
1,1-Dimethylhydrazine	57-14-7	No data		No data	No data	
1,2-Dimethylhydrazine	540-73-8	No data		No data	No data	
1,2-Diphenylhydrazine	122-66-7	No data		No data	No data	
1,3-Propane sultone	1120-71-4	No data		No data	No data	
2,4-Toluene diisocyante	584-84-9	No data		No data	No data	
2-Chloroacetophenone	532-27-4	No data		No data	No data	
2-Propenoic acid	79-10-7	No data		No data	No data	
4,4-Methylenedianiline	101-77-9	No data		No data	No data	
Acetophenone	98-86-2	No data		No data	No data	
Benzoic acid	65-85-0	No data		No data	No data	
bis(2-Chloroethoxy)methane	111-91-1	No data		No data	No data	
bis(2-Chloroethyl) ether	111-44-4	No data		No data	No data	
Chlorocyclopentadiene	41851-50-7	No data		No data	No data	
Cyclohexanol	108-93-0	No data		No data	No data	
Dichloroisopropyl ether	108-60-1	No data		No data	No data	
Dichloromethyl ether	542-88-1	No data		No data	No data	
Dichloropentadiene	no cas #	No data		No data	No data	
Dimethyl sulfate	77-78-1	No data		No data	No data	

Table C3-8. Sediment TRVs for Sediment-Dwelling Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Jones et al. 1997[b]		Recommended TRV[c]
				Ontario MOE (LOW)	NOAA (ER-L)	
Dimethylaniline	121-69-7	No data		No data	No data	
Di-n-propylnitrosamine	621-64-7	No data		No data	No data	
Diphenyl ether	101-84-8	No data		No data	No data	
Epichlorohydrin	106-89-8	No data		No data	No data	
Ethyl carbamate (urethane)	51-79-6	No data		No data	No data	
Ethyl methanesulfonate	62-50-0	No data		No data	No data	
Ethylene dibromide	106-93-4	No data		No data	No data	
Ethylene glycol	107-21-1	No data		No data	No data	
Ethylene glycol monobutyl ether	111-76-2	No data		No data	No data	
Ethylene glycol monoethyl ether acetate	111-15-9	No data		No data	No data	
Ethylene thiourea	96-45-7	No data		No data	No data	
Furfural	98-01-1	No data		No data	No data	
Maleic hydrazide	123-33-1	No data		No data	No data	
Malononitrile	109-77-3	No data		No data	No data	
Methyl styrene (mixed isomers)	25013-15-4	No data		No data	No data	
Methylhydrazine	60-34-4	No data		No data	No data	
N,N-Diphenylamine	122-39-4	No data		No data	No data	
Nitric acid, propyl ester	627-13-4	No data		No data	No data	
N-Nitrosodi-n-butylamine	924-16-3	No data		No data	No data	
N-Nitrosomorpholine	59-89-2	No data		No data	No data	
N-Nitroso-N,N-dimethylamine	62-75-9	No data		No data	No data	
o-Anisidine	90-04-0	No data		No data	No data	
Oxalic acid	144-62-7	No data		No data	No data	
Phthalic anhydride	85-44-9	No data		No data	No data	
p-Phthalic acid	100-21-0	No data		No data	No data	
Pyridine	110-86-1	No data		No data	No data	
Quinoline	91-22-5	No data		No data	No data	
Quinone	106-51-4	No data		No data	No data	
Safrole	94-59-7	No data		No data	No data	
Tetrahydrofuran	109-99-9	No data		No data	No data	

Table C3-8. Sediment TRVs for Sediment-Dwelling Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Jones et al. 1997[b]		Recommended TRV[c]
				Ontario MOE (LOW)	NOAA (ER-L)	
Other Heavy Semivolatile Compounds (molecular weight >200 g/mole)						
1,2,4,5-Tetrachlorobenzene	95-94-3	No data		No data	No data	
1,3,5-Trinitrobenzene	99-35-4	No data		No data	No data	
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	No data		No data	No data	
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	No data		No data	No data	
2-sec-Butyl-4,6-dinitrophenol	88-85-7	No data		No data	No data	
3,3-Dichlorobenzidine	91-94-1	No data		No data	No data	
3,3'-Dimethoxybenzidine	119-90-4	No data		No data	No data	
4-Bromophenylphenyl ether	101-55-3	No data		No data	No data	
Ammonium perfluorooctanoate	3825-26-1	No data		No data	No data	
Azobenzene	103-33-3	No data		No data	No data	
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	No data		No data	No data	
Captan	133-06-2	No data		No data	No data	
Chlorobenzilate	510-15-6	No data		No data	No data	
Dibutylphosphate	107-66-4	No data		No data	No data	
Dimethyl aminoazobenzene	60-11-7	No data		No data	No data	
Hexachlorobenzene	118-74-1	2.00E+01		No data	No data	2.00E+01
Hexachlorobutadiene	87-68-3	2.58E+02[d]		No data	No data	2.58E+02
Hexachlorocyclopentadiene	77-47-4	1.98E+02[d]		No data	No data	1.98E+02
Hexachloroethane	67-72-1	No data		No data	No data	
Hexachlorophene	70-30-4	6.34E+04[d]		No data	No data	6.34E+04
Hexamethylene-1,5-diisocyanate	822-06-0	No data		No data	No data	
Mirex	2385-85-5	No data		7.00E+00	No data	7.00E+00
Nitrofen	1836-75-5	No data		No data	No data	
Pentachlorobenzene	608-93-5	6.04E+02[d]		No data	No data	6.04E+02
Pentachloronitrobenzene	82-68-8	2.36E+03[d]		No data	No data	2.36E+03
Pentachlorophenol	87-86-5	7.00E+03		No data	No data	7.00E+03
Picric acid	88-89-1	No data		No data	No data	
Pronamide	23950-58-5	No data		No data	No data	
Strychnine	57-24-9	No data		No data	No data	

Table C3-8. Sediment TRVs for Sediment-Dwelling Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Jones et al. 1997[b]		Recommended TRV[c]
				Ontario MOE (LOW)	NOAA (ER-L)	
Terphenyls	26140-60-3	No data		No data	No data	
Tributyl phosphate	126-73-8	No data		No data	No data	
Trifluralin	1582-09-8	No data		No data	No data	
Triphenylamine	603-34-9	No data		No data	No data	
Herbicides and Organochlorinated Pesticides						
2,4,5-T	93-76-5	No data		No data	No data	
2,4-D and esters	94-75-7	No data		No data	No data	
4,4-DDD	72-54-8	No data		8.00E+00[f]	2.00E+00	8.00E+00
4,4-DDE	72-55-9	5.00E+00		5.00E+00	2.20E+00	5.00E+00
4,4-DDT	50-29-3	No data		8.00E+00[f]	1.00E+00	8.00E+00
Aldrin	309-00-2	No data		2.00E+00	No data	2.00E+00
alpha-BHC	319-84-6	No data		6.00E+00	No data	6.00E+00
beta-BHC	319-85-7	No data		5.00E+00	No data	5.00E+00
Chlordane	57-74-9	No data		7.00E+00	5.00E-01[f]	7.00E+00
Delta-BHC	319-86-8	No data		No data	No data	
Dieldrin	60-57-1	No data		2.00E+00	2.00E-02[f]	2.00E+00
Endothall	145-73-3	No data		No data	No data	
Endrin	72-20-8	No data		3.00E+00	2.00E-02[f]	3.00E+00
gamma-BHC (Lindane)	58-89-9	No data		3.00E+00[e,g]	No data	3.00E+00
Heptachlor	76-44-8	3.08E-01		No data	No data	3.08E-01
Isodrin	465-73-6	No data		No data	No data	
Methoxychlor	72-43-5	No data		No data	No data	
Silvex (2,4,5-TP)	93-72-1	No data		No data	No data	
Toxaphene	8001-35-2	No data		No data	No data	
Inorganic Chemicals and Compounds (All units are in mg/kg)						
Metals						
Aluminum	7429-90-5	1.40E+04		No data	No data	1.40E+04
Antimony	7440-36-0	6.40E+01		No data	2.00E+00[f]	6.40E+01
Arsenic	7440-38-2	6.00E+00		6.00E+00	8.20E+00	6.00E+00
Barium	7440-39-3	2.00E+01[h]		No data	No data	2.00E+01

Table C3-8. Sediment TRVs for Sediment-Dwelling Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Jones et al. 1997[b]		Recommended TRV[c]
				Ontario MOE (LOW)	NOAA (ER-L)	
Beryllium	7440-41-7	No data		No data	No data	
Bismuth	7440-69-9	No data		No data	No data	
Boron	7440-42-8	No data		No data	No data	
Cadmium	7440-43-9	6.00E-01		6.00E-01	1.20E+00	6.00E-01
Calcium	7440-70-2	No data		No data	No data	
Chromium (and VI)	18540-29-9	2.60E+01		2.60E+01	8.10E+01	2.60E+01
Cobalt	7440-48-4	No data		No data	No data	
Copper	7440-50-8	1.60E+01		1.60E+01	3.40E+01	1.60E+01
Iron	7439-89-6	No data		2.00E+04	No data	2.00E+04
Lead	7439-92-1	3.10E+01		3.10E+01	4.67E+01	3.10E+01
Lithium	7439-93-2	No data		No data	No data	
Magnesium	7439-95-4	No data		No data	No data	
Manganese	7439-96-5	No data		4.60E+02	No data	4.60E+02
Mercury	7439-97-6	2.00E-01		2.00E-01	1.50E-01	2.00E-01
Molybdenum	7439-98-7	No data		No data	No data	
Nickel	7440-02-0	1.60E+01		1.60E+01	2.09E+01	1.60E+01
Potassium	7440-09-7	No data		No data	No data	
Rhodium	7440-16-6	No data		No data	No data	
Selenium	7782-49-2	1.00E-01		No data	No data	1.00E-01
Silicon	7440-21-3	No data		No data	No data	
Silver	7440-22-4	4.50E+00		No data	1.00E+00	4.50E+00
Sodium	7440-23-5	No data		No data	No data	
Strontium	7440-24-6	No data		No data	No data	
Tantalum	7440-25-7	No data		No data	No data	
Thallium	7440-28-0	No data		No data	No data	
Tin	7440-31-5	No data		No data	No data	
Tungsten	7440-33-7	No data		No data	No data	
Uranium	7440-61-1	No data		No data	No data	
Vanadium	7440-62-2	No data		No data	No data	
Yttrium	7440-65-5	No data		No data	No data	

Table C3-8. Sediment TRVs for Sediment-Dwelling Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Jones et al. 1997[b]		Recommended TRV[c]
				Ontario MOE (LOW)	NOAA (ER-L)	
Zinc	7440-66-6	1.10E+02		1.20E+02	1.50E+02	1.10E+02
Zirconium	7440-67-7	No data		No data	No data	
Non-metals and Anions						
Ammonia/Ammonium	7664-41-7	No data		No data	No data	
Bromide	24959-67-9	No data		No data	No data	
Chloride	16887-00-6	No data		No data	No data	
Cyanide	57-12-5	1.00E-01[h]		No data	No data	1.00E-01
Fluoride	16984-48-8	No data		No data	No data	
Hydroxide	14280-30-9	No data		No data	No data	
Iodine	7553-56-2	No data		No data	No data	
Nitrate	14797-55-8	No data		No data	No data	
Nitrite	14797-65-0	No data		No data	No data	
Phosphate	14265-44-2	No data		No data	No data	
Phosphorus	7723-14-0	No data		No data	No data	
Sulfate	14808-79-8	No data		No data	No data	
Total Sulfur	63705-05-5	No data		No data	No data	
Priority Pollutants						
Carbon Dioxide	124-38-9	No data		No data	No data	
Nitrogen Dioxide	10102-44-0	No data		No data	No data	
Ozone	10028-15-6	No data		No data	No data	
Particulate Matter	No CAS #	No data		No data	No data	
Sulfur Dioxode	7446-09-5	No data		No data	No data	
Radionuclides[i]						
Actinium-227	14952-40-0	NA		NA	NA	
Americium-241	1596-10-2	NA		NA	NA	
Americium-243	14993-75-0	NA		NA	NA	
Antimony-125	14234-35-6	NA		NA	NA	
Barium-137	13981-97-0	NA		NA	NA	
Cadmium-113	None	NA		NA	NA	
Carbon-14	14762-75-5	NA		NA	NA	

Table C3-8. Sediment TRVs for Sediment-Dwelling Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Jones et al. 1997[b]		Recommended TRV[c]
				Ontario MOE (LOW)	NOAA (ER-L)	
Cesium-134	13967-70-9	NA		NA	NA	
Cesium-137	10045-97-3	NA		NA	NA	
Cobalt-60	10198-40-0	NA		NA	NA	
Curium-242	15510-73-3	NA		NA	NA	
Curium-243	15757-87-6	NA		NA	NA	
Curium-244	13981-15-2	NA		NA	NA	
Europium-152	14683-23-9	NA		NA	NA	
Europium-154	15585-10-1	NA		NA	NA	
Europium-155	14391-16-3	NA		NA	NA	
Iodine-129	15046-84-1	NA		NA	NA	
Neptunium-237	13994-20-2	NA		NA	NA	
Nickel-59	14336-70-0	NA		NA	NA	
Nickel-63	13981-37-8	NA		NA	NA	
Niobium-93	7440-03-1	NA		NA	NA	
Plutonium-238	13981-16-3	NA		NA	NA	
Plutonium-239	15117-48-3	NA		NA	NA	
Plutonium-240	14119-33-6	NA		NA	NA	
Plutonium-241	14119-32-5	NA		NA	NA	
Plutonium-242	13982-10-0	NA		NA	NA	
Protactinium-231	14331-85-2	NA		NA	NA	
Radium-226	13982-63-3	NA		NA	NA	
Radium-228	15262-20-1	NA		NA	NA	
Ruthenium-106	13967-48-1	NA		NA	NA	
Samarium-151	15715-94-3	NA		NA	NA	
Selenium-79	None	NA		NA	NA	
Strontium-90	10098-97-2	NA		NA	NA	
Technetium-99	14133-79-7	NA		NA	NA	
Thorium-229	15594-54-4	NA		NA	NA	
Thorium-232	7440-29-1	NA		NA	NA	
Tin-126	15832-50-5	NA		NA	NA	

Table C3-8. Sediment TRVs for Sediment-Dwelling Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Jones et al. 1997[b]		Recommended TRV[c]
				Ontario MOE (LOW)	NOAA (ER-L)	
Tritium	10028-17-8	NA		NA	NA	
Uranium-232	14158-29-3	NA		NA	NA	
Uranium-233	13968-55-3	NA		NA	NA	
Uranium-234	13966-29-5	NA		NA	NA	
Uranium-235	15117-96-1	NA		NA	NA	
Uranium-236	13982-70-2	NA		NA	NA	
Uranium-238	7440-61-1	NA		NA	NA	
Yttrium-90	10098-91-6	NA		NA	NA	
Zirconium-93	15751-77-6	NA		NA	NA	

NA = Not applicable

[a] Published in Appendix E of EPA 1999, Table E-3.

[b] Published in Tables 1 and 4 of Jones et al. (1997)

[c] Order of preference is EPA 1999, then MOE (Low), then NOAA (ER-L)

[d] TRV was calculated using EqP approach (EPA 1993) assuming a fractional organic content of 0.04

[e] Denotes tentative guideline

[f] Source document is Long and Morgan (1991)

[g] 10th percentile of screening level concentration

[h] TRV is a U.S. EPA Region 5 guideline value for classification of sediments for determining the suitability of dredged sediments for open water disposal, as cited in Hall and Suter II (1994)

[i] TRV is not applicable to single radionuclides. Combined external and internal radiation exposure for sediment-dwelling biota from all radionuclides combined cannot exceed 1 rad/d.

Table C3-9. Surface Water TRVs for Aquatic Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Suter & Tsao 1996[b]		Recommended TRV[c]
				NAWQC Chronic	Tier II Chronic	
Organic Compounds (All units are in ug/L)						
Aromatic Halogenated Hydrocarbons						
4-Chloro-3-methylphenol	59-50-7	No data		No data	No data	
2,3,4,6-Tetrachlorophenol	58-90-2	No data		No data	No data	
Aromatic Nonhalogenated Hydrocarbons						
2-Nitrotoluene	88-72-2	No data		No data	No data	
4-Nitrobiphenyl	92-93-3	No data		No data	No data	
Benzaldehyde	100-52-7	No data		No data	No data	
Benzene	71-43-2	No data		No data	1.30E+02	1.30E+02
Benzyl alcohol	100-51-6	No data		No data	8.60E+00	8.60E+00
Ethyl benzene	100-41-4	No data		No data	7.30E+00	7.30E+00
m-Xylene	108-38-3	No data		No data	1.80E+00[d]	1.80E+00
o-Xylene	95-47-6	No data		No data	No data	
p-Xylene	106-42-3	No data		No data	No data	
Styrene	100-42-5	No data		No data	No data	
Toluene	108-88-3	No data		No data	9.80E+00	9.80E+00
Non-aromatic Nonhalogenated Hydrocarbons						
1,2-Epoxybutane	106-88-7	No data		No data	No data	
1,3-Butadiene	106-99-0	No data		No data	No data	
1,4-Dioxane	123-91-1	6.21E+04[e]		No data	No data	6.21E+04
1-Methylpropyl alcohol	78-92-2	No data		No data	No data	
1-Nitropropane	108-03-2	No data		No data	No data	
2,2,4-Trimethylpentane	540-84-1	No data		No data	No data	
2-Butanone	78-93-3	No data		No data	1.40E+00	1.40E+00
2-Butenaldehyde (2-Butenal)	4170-30-3	No data		No data	No data	
2-Ethoxyethanol	110-80-5	No data		No data	No data	
2-Heptanone	110-43-0	No data		No data	No data	
2-Hexanone	591-78-6	No data		No data	9.90E+01	9.90E+01

Table C3-9. Surface Water TRVs for Aquatic Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Suter & Tsao 1996[b]		Recommended TRV[c]
				NAWQC Chronic	Tier II Chronic	
2-Methoxyethanol	109-86-4	No data		No data	No data	
2-Methyl-2-propanol	75-65-0	No data		No data	No data	
2-Methyl-2-propenenitrile	126-98-7	No data		No data	No data	
2-Methylaziridine	75-55-8	No data		No data	No data	
2-Methylpropyl alcohol	78-83-1	No data		No data	No data	
2-Pentanone	107-87-9	No data		No data	No data	
2-Propanone (Acetone)	67-64-1	1.50E+03[e]		No data	1.50E+03[e]	1.50E+03
2-Propene-1-ol	107-18-6	No data		No data	No data	
2-Propyl alcohol	67-63-0	No data		No data	No data	
3-Heptanone	106-35-4	No data		No data	No data	
3-Methyl-1-butanol	123-51-3	No data		No data	No data	
3-Methyl-2-butanone	563-80-4	No data		No data	No data	
3-Pentanone	96-22-0	No data		No data	No data	
4-Heptanone	123-19-3	No data		No data	No data	
4-Methyl-2-pentanone	108-10-1	No data		No data	1.70E+02	1.70E+02
4-Methyl-3-penten-2-one	141-79-7	No data		No data	No data	
5-Methyl-2-hexanone	110-12-3	No data		No data	No data	
Acetaldehyde	75-07-0	No data		No data	No data	
Acetamide	60-35-5	No data		No data	No data	
Acetic acid	64-19-7	No data		No data	No data	
Acetic acid ethyl ester	141-78-6	No data		No data	No data	
Acetic acid n-butyl ester	123-86-4	No data		No data	No data	
Acetonitrile	75-05-8	No data		No data	No data	
Acrolein	107-02-8	No data		No data	No data	
Acrylonitrile	107-13-1	2.60E+02[e]		No data	No data	2.60E+02
Bis(isopropyl)ether	108-20-3	No data		No data	No data	
Butane	106-97-8	No data		No data	No data	
Carbon disulfide	75-15-0	No data		No data	9.20E-01	9.20E-01
Cyanogen	460-19-5	No data		No data	No data	

Table C3-9. Surface Water TRVs for Aquatic Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Suter & Tsao 1996[b]		Recommended TRV[c]
				NAWQC Chronic	Tier II Chronic	
Cyclohexane	110-82-7	No data		No data	No data	
Cyclohexanone	108-94-1	No data		No data	No data	
Cyclohexene	110-83-8	No data		No data	No data	
Cyclopentane	287-92-3	No data		No data	No data	
Ethyl alcohol	64-17-5	No data		No data	No data	
Ethyl ether	60-29-7	No data		No data	No data	
Ethyl methacrylate	97-63-2	No data		No data	No data	
Formaldehyde	50-00-0	4.96E+01[e]		No data	No data	4.96E+01
Formamide	75-12-7	No data		No data	No data	
Formic acid	64-18-6	No data		No data	No data	
Formic acid, methyl ester	107-31-3	No data		No data	No data	
Glycidylaldehyde	765-34-4	No data		No data	No data	
Methyl acetate	79-20-9	No data		No data	No data	
Methyl alcohol	67-56-1	No data		No data	No data	
Methyl isocyanate	624-83-9	No data		No data	No data	
Methyl methacrylate	80-62-6	No data		No data	No data	
Methyl tert-butyl ether	1634-04-4	No data		No data	No data	
Methylacetylene	74-99-7	No data		No data	No data	
Methylcyclohexane	108-87-2	No data		No data	No data	
N,N-Dimethylacetamide	127-19-5	No data		No data	No data	
n-Butyl alcohol	71-36-3	No data		No data	No data	
n-Heptane	142-82-5	No data		No data	No data	
n-Hexane	110-54-3	No data		No data	5.80E-01	5.80E-01
Nitromethane	75-52-5	No data		No data	No data	
n-Nonane	111-84-2	No data		No data	No data	
n-Octane	111-65-9	No data		No data	No data	
n-Pentane	109-66-0	No data		No data	No data	
n-Propionaldehyde	123-38-6	No data		No data	No data	
n-Propyl alcohol	71-23-8	No data		No data	No data	

Table C3-9. Surface Water TRVs for Aquatic Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Suter & Tsao 1996[b]		Recommended TRV[c]
				NAWQC Chronic	Tier II Chronic	
n-Valeraldehyde	110-62-3	No data		No data	No data	
Oxirane	75-21-8	No data		No data	No data	
p-Cymene	99-87-6	No data		No data	No data	
Phosgene	75-44-5	No data		No data	No data	
Propargyl alcohol	107-19-7	No data		No data	No data	
Propionic acid	79-09-4	No data		No data	No data	
Propionitrile	107-12-0	No data		No data	No data	
Propylene gylcol monomethyl ether	107-98-2	No data		No data	No data	
p-tert-Butyltoluene	98-51-1	No data		No data	No data	
Triethylamine	121-44-8	No data		No data	No data	
Trimethylamine	75-50-3	No data		No data	No data	
Vinyl acetate	108-05-4	No data		No data	1.60E+01	1.60E+01
Non-aromatic Halogenated Hydrocarbons						
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	No data		No data	No data	
1,1,1,2-Tetrachloroethane	630-20-6	No data		No data	No data	
1,1,1-Trichloroethane	71-55-6	No data		No data	1.10E+01	1.10E+01
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	No data		No data	No data	
1,1,2,2-Tetrachloroethane	79-34-5	No data		No data	6.10E+02	6.10E+02
1,1,2,2-Tetrachloroethene	127-18-4	No data		No data	No data	
1,1,2-Trichloroethane	79-00-5	No data		No data	1.20E+03	1.20E+03
1,1,2-Trichloroethylene	79-01-6	No data		No data	No data	
1,1-Dichloroethane	75-34-3	No data		No data	4.70E+01	4.70E+01
1,1-Dichloroethene	75-35-4	No data		No data	2.50E+01	2.50E+01
1,2,2-Trichloro-1,1,2-trifluoroethane	76-13-1	No data		No data	No data	
1,2,3-Trichloropropane	96-18-4	No data		No data	No data	
1,2-Dibromo-3-chloropropane	96-12-8	No data		No data	No data	
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	No data		No data	No data	
1,2-Dichloroethane	107-06-2	No data		No data	9.10E+02	9.10E+02
1,2-Dichloroethylene	540-59-0	No data		No data	5.90E+02	5.90E+02

Table C3-9. Surface Water TRVs for Aquatic Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Suter & Tsao 1996[b]		Recommended TRV[c]
				NAWQC Chronic	Tier II Chronic	
1,2-Dichloropropane	78-87-5	No data		No data	No data	
1,3-Dichloropropene	542-75-6	No data		No data	5.50E-02	5.50E-02
1,4-Dichloro-2-butene	764-41-0	No data		No data	No data	
1-Chloroethene	75-01-4	No data		No data	No data	
2,2-Dichloropropionic acid	75-99-0	No data		No data	No data	
2-Chloropropane	75-29-6	No data		No data	No data	
3-Chloropropene (allyl chloride)	107-05-1	No data		No data	No data	
Bromochloromethane	74-97-5	No data		No data	No data	
Bromodichloromethane	75-27-4	No data		No data	No data	
Bromoethene	593-60-2	No data		No data	No data	
Bromoform	75-25-2	No data		No data	No data	
Bromomethane	74-83-9	No data		No data	No data	
Carbon tetrachloride	56-23-5	No data		No data	9.80E+00	9.80E+00
Chlorodibromomethane	124-48-1	No data		No data	No data	
Chlorodifluoromethane	75-45-6	No data		No data	No data	
Chloroethane	75-00-3	No data		No data	No data	
Chloroform	67-66-3	2.88E+01[e]		No data	2.80E+01	2.88E+01
Chloromethane	74-87-3	No data		No data	No data	
Chloropentafluoroethane	76-15-3	No data		No data	No data	
cis-1,2-Dichloroethene	156-59-2	No data		No data	No data	
cis-1,3-Dichloropropene	10061-01-5	No data		No data	No data	
Cyanogen bromide	506-68-3	No data		No data	No data	
Cyanogen chloride	506-77-4	No data		No data	No data	
Dichlorodifluoromethane	75-71-8	No data		No data	No data	
Dichlorofluoromethane	75-43-4	No data		No data	No data	
Dichloromethane	75-09-2	No data		No data	No data	
Difluorodibromomethane	75-61-6	No data		No data	No data	
Hexafluoroacetone	684-16-2	No data		No data	No data	
Iodomethane	74-88-4	No data		No data	No data	

Table C3-9. Surface Water TRVs for Aquatic Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Suter & Tsao 1996[b]		Recommended TRV[c]
				NAWQC Chronic	Tier II Chronic	
Methylene bromide	74-95-3	No data		No data	No data	
Pentachloroethane	76-01-7	No data		No data	No data	
trans-1,2-Dichloroethene	156-60-5	No data		No data	No data	
trans-1,3-Dichloropropene	10061-02-6	No data		No data	No data	
Trichloroacetic acid	76-03-9	No data		No data	No data	
Trichlorofluoroethane	27154-33-2	No data		No data	No data	
Trichlorofluoromethane	75-69-4	No data		No data	No data	
Trifluorobromomethane	75-63-8	No data		No data	No data	
Dioxin and Furan Compounds						
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	35822-46-9	No data		No data	No data	
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	No data		No data	No data	
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	No data		No data	No data	
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	39227-28-6	No data		No data	No data	
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	No data		No data	No data	
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	57653-85-7	No data		No data	No data	
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	No data		No data	No data	
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	19408-74-3	No data		No data	No data	
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	No data		No data	No data	
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	40321-76-4	No data		No data	No data	
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	No data		No data	No data	
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	No data		No data	No data	
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	No data		No data	No data	
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1746-01-6	3.80E-06[e]		No data	No data	3.80E-06
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	No data		No data	No data	
Dibenzofuran	132-64-9	No data		No data	3.70E+00	3.70E+00
Octachlorodibenzo(p)dioxin	3268-87-9	No data		No data	No data	
Octachlorodibenzofuran	39001-02-0	No data		No data	No data	
PCBs						
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	No data		No data	No data	

Table C3-9. Surface Water TRVs for Aquatic Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Suter & Tsao 1996[b]		Recommended TRV[c]
				NAWQC Chronic	Tier II Chronic	
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	No data		No data	No data	
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	No data		No data	No data	
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	No data		No data	No data	
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	No data		No data	No data	
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	No data		No data	No data	
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	No data		No data	No data	
2',3,4,4',5-Pentachlorobiphenyl	no cas #	No data		No data	No data	
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	No data		No data	No data	
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	No data		No data	No data	
3,3',4,4',5-Pentachlorobiphenyl	no cas #	No data		No data	No data	
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	No data		No data	No data	
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	No data		No data	No data	
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	No data		No data	No data	
Polychlorinated biphenyls (PCBs)[k]	1336-36-3	No data		No data	1.40E+01[f]	1.40E+01
Phthalates						
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	3.00E+00[e]		No data	3.00E+00	3.00E+00
Butylbenzyl phthalate	85-68-7	No data		No data	1.90E+01[g]	1.90E+01
Dibutyl phthalate	84-74-2	No data		No data	No data	
Diethyl phthalate	84-66-2	No data		No data	2.10E+02	2.10E+02
Dimethylphthalate	131-11-3	No data		No data	No data	
n-Dioctyl phthalate	117-84-0	3.20E+02[e]		No data	No data	3.20E+02
Light Polycyclic Aromatic Hydrocarbons (MW <200 g/mole)						
2-Chloronapthalene	91-58-7	No data		No data	No data	
2-Methyl naphthalene	91-57-6	No data		No data	No data	
5-Nitroacenaphthene	602-87-9	No data		No data	No data	
Acenaphthene	83-32-9	No data		2.30E+01[h]	No data	2.30E+01
Acenaphthylene	208-96-8	No data		No data	No data	

Table C3-9. Surface Water TRVs for Aquatic Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Suter & Tsao 1996[b]		Recommended TRV[c]
				NAWQC Chronic	Tier II Chronic	
Anthracene	120-12-7	No data		No data	7.30E-01	7.30E-01
Fluorene	86-73-7	No data		No data	3.90E+00[g]	3.90E+00
Indene	95-13-6	No data		No data	No data	
Naphthalene	91-20-3	No data		No data	1.20E+01	1.20E+01
Phenanthrene	85-01-8	No data		6.30E+00[f]	No data	6.30E+00
Pyrene	129-00-0	No data		No data	No data	
Heavy Polycyclic Aromatic Hydrocarbons (MW >200 g/mole)						
3-Methylcholanthrene	56-49-5	No data		No data	No data	
5-Methylchrysene	3697-24-3	No data		No data	No data	
Benzo(a)anthracene	56-55-3	2.70E-02		No data	2.70E-02	2.70E-02
Benzo(a)pyrene	50-32-8	1.40E-02		No data	1.40E-02	1.40E-02
Benzo(b)fluoranthene	205-99-2	2.70E-02		No data	No data	2.70E-02
Benzo(e)pyrene	192-97-2	No data		No data	No data	
Benzo(g,h,i)perylene	191-24-2	No data		No data	No data	
Benzo(j)fluoranthene	205-82-3	No data		No data	No data	
Benzo(k)fluoranthene	207-08-9	2.70E-02		No data	No data	2.70E-02
Benzo[a,i]pyrene	191-30-0	No data		No data	No data	
Chrysene	218-01-9	2.70E-02		No data	No data	2.70E-02
Dibenz(a,h)anthracene	53-70-3	2.70E-02		No data	No data	2.70E-02
Dibenz[a,h]acridine	226-36-8	No data		No data	No data	
Dibenz[a,j]acridine	224-42-0	No data		No data	No data	
Dibenzo(a,e)fluoranthene	5385-75-1	No data		No data	No data	
Dibenzo(a,h)fluoranthene	no cas #	No data		No data	No data	
Dibenzo[a,e]pyrene	192-65-4	No data		No data	No data	
Dibenzo[a,h]pyrene	189-64-0	No data		No data	No data	
Dibenzo[a,i]pyrene	189-55-9	No data		No data	No data	
Fluoranthene	206-44-0	No data		6.16E+00[h]	No data	6.16E+00

Table C3-9. Surface Water TRVs for Aquatic Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Suter & Tsao 1996[b]		Recommended TRV[c]
				NAWQC Chronic	Tier II Chronic	
Hexachloronaphthalene	1335-87-1	No data		No data	No data	
Indeno(1,2,3-cd)pyrene	193-39-5	2.70E-02		No data	No data	2.70E-02
Octachloronaphthalene	2234-13-1	No data		No data	No data	
Pentachloronaphthalene	1321-64-8	No data		No data	No data	
Tetrachloronaphthalene	1335-88-2	No data		No data	No data	
Trichloronaphthalene	1321-65-9	No data		No data	No data	
Light Substituted Benzene Compounds (MW <200 g/mole)						
1,2,3-Trichlorobenzene	87-61-6	No data		No data	No data	
1,2,4-Trichlorobenzene	120-82-1	No data		No data	1.10E+02[d]	1.10E+02
1,2,4-Trimethyl benzene	95-63-6	No data		No data	No data	
1,2-Dichlorobenzene	95-50-1	No data		No data	1.40E+01[g]	1.40E+01
1,3,5-Trimethyl benzene	108-67-8	No data		No data	No data	
1,3-Dichlorobenzene	541-73-1	No data		No data	7.10E+01[g]	7.10E+01
1,3-Dinitrobenzene	99-65-0	2.60E+01[e]		No data	No data	2.60E+01
1,4-Dichlorobenzene	106-46-7	No data		No data	1.50E+01[g]	1.50E+01
1,4-Dinitrobenzene	100-25-4	No data		No data	No data	
2,4,5-Trichlorophenol	95-95-4	No data		No data	No data	
2,4,6-Trichlorophenol	88-06-2	No data		No data	No data	
2,4-Dichlorophenol	120-83-2	No data		No data	No data	
2,4-Dimethylphenol	105-67-9	No data		No data	No data	
2,4-Dinitrophenol	51-28-5	No data		No data	No data	
2,4-Dinitrotoluene	121-14-2	2.30E+01[e]		No data	No data	2.30E+01
2,6-Dinitrotoluene	606-20-2	6.00E+01[e]		No data	No data	6.00E+01
2-Chlorophenol	95-57-8	No data		No data	No data	
2-Chlorotoluene	95-49-8	No data		No data	No data	
2-Nitrophenol	88-75-5	No data		No data	No data	

Table C3-9. Surface Water TRVs for Aquatic Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Suter & Tsao 1996[b]		Recommended TRV[c]
				NAWQC Chronic	Tier II Chronic	
4,6-Dinitro-o-cresol	534-52-1	No data		No data	No data	
4-Chlorotoluene	106-43-4	No data		No data	No data	
4-Nitrophenol	100-02-7	No data		No data	3.00E+02	3.00E+02
alpha-Methylstyrene	98-83-9	No data		No data	No data	
Aniline	62-53-3	No data		No data	No data	
Benzotrichloride	98-07-7	No data		No data	No data	
Benzyl chloride	100-44-7	No data		No data	No data	
Bromobenzene	108-86-1	No data		No data	No data	
Chlorobenzene	108-90-7	No data		No data	6.40E+01	6.40E+01
Cumene	98-82-8	No data		No data	No data	
m-Cresol	108-39-4	No data		No data	No data	
n-Butyl benzene	104-51-8	No data		No data	No data	
Nitrobenzene	98-95-3	2.70E+02[e]		No data	No data	2.70E+02
n-Propyl benzene	103-65-1	No data		No data	No data	
o-Cresol	95-48-7	No data		No data	No data	
o-Dinitrobenzene	528-29-0	No data		No data	No data	
o-Nitroaniline	88-74-4	No data		No data	No data	
o-Toluidine	95-53-4	No data		No data	No data	
p-Chloroaniline	106-47-8	No data		No data	No data	
p-Cresol	106-44-5	No data		No data	No data	
Phenol	108-95-2	No data		1.10E+02[d]	No data	1.10E+02
p-Nitrochlorobenzene	100-00-5	No data		No data	No data	
p-Toluidine	106-49-0	No data		No data	No data	
sec-Butyl benzene	135-98-8	No data		No data	No data	
tert-Butyl benzene	98-06-6	No data		No data	No data	
Toluene-2,6-diamine	823-40-5	No data		No data	No data	
Trimethyl benzene	25551-13-7	No data		No data	No data	
Other Light Semivolatile Compounds (MW <200 g/mole)						

Table C3-9. Surface Water TRVs for Aquatic Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Suter & Tsao 1996[b]		Recommended TRV[c]
				NAWQC Chronic	Tier II Chronic	
1,1'-Biphenyl	92-52-4	No data		No data	No data	
1,1-Dimethylhydrazine	57-14-7	No data		No data	No data	
1,2-Dimethylhydrazine	540-73-8	No data		No data	No data	
1,2-Diphenylhydrazine	122-66-7	No data		No data	No data	
1,3-Propane sultone	1120-71-4	No data		No data	No data	
2,4-Toluene diisocyante	584-84-9	No data		No data	No data	
2-Chloroacetophenone	532-27-4	No data		No data	No data	
2-Propenoic acid	79-10-7	No data		No data	No data	
4,4-Methylenedianiline	101-77-9	No data		No data	No data	
Acetophenone	98-86-2	No data		No data	No data	
Benzoic acid	65-85-0	No data		No data	4.20E+01	4.20E+01
bis(2-Chloroethoxy)methane	111-91-1	No data		No data	No data	
bis(2-Chloroethyl) ether	111-44-4	No data		No data	No data	
Chlorocyclopentadiene	41851-50-7	No data		No data	No data	
Cyclohexanol	108-93-0	No data		No data	No data	
Dichloroisopropyl ether	108-60-1	No data		No data	No data	
Dichloromethyl ether	542-88-1	No data		No data	No data	
Dichloropentadiene	no cas #	No data		No data	No data	
Dimethyl sulfate	77-78-1	No data		No data	No data	
Dimethylaniline	121-69-7	No data		No data	No data	
Di-n-propylnitrosamine	621-64-7	No data		No data	No data	
Diphenyl ether	101-84-8	No data		No data	No data	
Epichlorohydrin	106-89-8	No data		No data	No data	
Ethyl carbamate (urethane)	51-79-6	No data		No data	No data	
Ethyl methanesulfonate	62-50-0	No data		No data	No data	
Ethylene dibromide	106-93-4	No data		No data	No data	
Ethylene glycol	107-21-1	No data		No data	No data	
Ethylene glycol monobutyl ether	111-76-2	No data		No data	No data	
Ethylene glycol monoethyl ether acetate	111-15-9	No data		No data	No data	

Table C3-9. Surface Water TRVs for Aquatic Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Suter & Tsao 1996[b]		Recommended TRV[c]
				NAWQC Chronic	Tier II Chronic	
Ethylene thiourea	96-45-7	No data		No data	No data	
Furfural	98-01-1	No data		No data	No data	
Maleic hydrazide	123-33-1	No data		No data	No data	
Malononitrile	109-77-3	No data		No data	No data	
Methyl styrene (mixed isomers)	25013-15-4	No data		No data	No data	
Methylhydrazine	60-34-4	No data		No data	No data	
N,N-Diphenylamine	122-39-4	No data		No data	No data	
Nitric acid, propyl ester	627-13-4	No data		No data	No data	
N-Nitrosodi-n-butylamine	924-16-3	No data		No data	No data	
N-Nitrosomorpholine	59-89-2	No data		No data	No data	
N-Nitroso-N,N-dimethylamine	62-75-9	No data		No data	No data	
o-Anisidine	90-04-0	No data		No data	No data	
Oxalic acid	144-62-7	No data		No data	No data	
Phthalic anhydride	85-44-9	No data		No data	No data	
p-Phthalic acid	100-21-0	No data		No data	No data	
Pyridine	110-86-1	No data		No data	No data	
Quinoline	91-22-5	No data		No data	No data	
Quinone	106-51-4	No data		No data	No data	
Safrole	94-59-7	No data		No data	No data	
Tetrahydrofuran	109-99-9	No data		No data	No data	
Other Heavy Semivolatile Compounds (MW >200 g/mole)						
1,2,4,5-Tetrachlorobenzene	95-94-3	No data		No data	No data	
1,3,5-Trinitrobenzene	99-35-4	No data		No data	No data	
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	No data		No data	No data	
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	No data		No data	No data	
2-sec-Butyl-4,6-dinitrophenol	88-85-7	No data		No data	No data	
3,3-Dichlorobenzidine	91-94-1	No data		No data	No data	
3,3'-Dimethoxybenzidine	119-90-4	No data		No data	No data	

Table C3-9. Surface Water TRVs for Aquatic Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Suter & Tsao 1996[b]		Recommended TRV[c]
				NAWQC Chronic	Tier II Chronic	
4-Bromophenylphenyl ether	101-55-3	No data		No data	1.50E+00[g]	1.50E+00
Ammonium perfluorooctanoate	3825-26-1	No data		No data	No data	
Azobenzene	103-33-3	No data		No data	No data	
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	No data		No data	No data	
Captan	133-06-2	No data		No data	No data	
Chlorobenzilate	510-15-6	No data		No data	No data	
Dibutylphosphate	107-66-4	No data		No data	No data	
Dimethyl aminoazobenzene	60-11-7	No data		No data	No data	
Hexachlorobenzene	118-74-1	3.68E+00		No data	No data	3.68E+00
Hexachlorobutadiene	87-68-3	9.30E-01[e]		No data	No data	9.30E-01
Hexachlorocyclopentadiene	77-47-4	5.20E-01[e]		No data	No data	5.20E-01
Hexachloroethane	67-72-1	No data		No data	1.20E+00[g]	1.20E+00
Hexachlorophene	70-30-4	8.80E-01[e]		No data	No data	8.80E-01
Hexamethylene-1,5-diisocyanate	822-06-0	No data		No data	No data	
Mirex	2385-85-5	No data		No data	No data	
Nitrofen	1836-75-5	No data		No data	No data	
Pentachlorobenzene	608-93-5	4.70E-01[e]		No data	4.70E-01[g]	4.70E-01
Pentachloronitrobenzene	82-68-8	1.00E+01[e]		No data	No data	1.00E+01
Pentachlorophenol	87-86-5	1.50E+01		No data	No data	1.50E+01
Picric acid	88-89-1	No data		No data	No data	
Pronamide	23950-58-5	No data		No data	No data	
Strychnine	57-24-9	No data		No data	No data	
Terphenyls	26140-60-3	No data		No data	No data	
Tributyl phosphate	126-73-8	No data		No data	No data	
Trifluralin	1582-09-8	No data		No data	No data	
Triphenylamine	603-34-9	No data		No data	No data	
Herbicides and Organochlorinated Pesticides						

Table C3-9. Surface Water TRVs for Aquatic Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Suter & Tsao 1996[b]		Recommended TRV[c]
				NAWQC Chronic	Tier II Chronic	
2,4,5-T	93-76-5	No data		No data	No data	
2,4-D and esters	94-75-7	No data		No data	No data	
4,4-DDD	72-54-8	No data		No data	1.10E-02	1.10E-02
4,4-DDE	72-55-9	1.05E+01		No data	1.10E-02	1.05E+01
4,4-DDT	50-29-3	No data		No data	1.30E-02	1.30E-02
Aldrin	309-00-2	No data		No data	No data	
alpha-BHC	319-84-6	No data		No data	2.20E+00	2.20E+00
beta-BHC	319-85-7	No data		No data	2.20E+00	2.20E+00
Chlordane	57-74-9	No data		1.70E-01[h]	No data	1.70E-01
Delta-BHC	319-86-8	No data		No data	2.20E+00	2.20E+00
Dieldrin	60-57-1	No data		6.20E-02[h]	No data	6.20E-02
Endothall	145-73-3	No data		No data	No data	
Endrin	72-20-8	No data		6.10E-02[h]	No data	6.10E-02
gamma-BHC (Lindane)	58-89-9	No data		8.00E-02	No data	8.00E-02
Heptachlor	76-44-8	3.80E-03		No data	6.90E-03[d,f]	3.80E-03
Isodrin	465-73-6	No data		No data	No data	
Methoxychlor	72-43-5	No data		No data	1.90E-02[d]	1.90E-02
Silvex (2,4,5-TP)	93-72-1	No data		No data	No data	
Toxaphene	8001-35-2	No data		No data	No data	
Inorganic Chemicals and Compounds (All units are in mg/L)						
Metals						
Aluminum	7429-90-5	8.70E-02		8.70E+01	No data	8.70E+01
Antimony	7440-36-0	3.00E-02		No data	3.00E+01[j]	3.00E-02
Arsenic	7440-38-2	1.50E-01		1.90E+02	No data	1.90E+02
Barium	7440-39-3	1.40E-03[i]		No data	4.00E+00	1.40E-03
Beryllium	7440-41-7	6.60E-04		No data	6.60E-01	6.60E-04

Table C3-9. Surface Water TRVs for Aquatic Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Suter & Tsao 1996[b]		Recommended TRV[c]
				NAWQC Chronic	Tier II Chronic	
Bismuth	7440-69-9	No data		No data	No data	
Boron	7440-42-8	No data		No data	1.60E+00	1.60E+00
Cadmium	7440-43-9	2.20E-03		1.10E+00	No data	1.10E+00
Calcium	7440-70-2	No data		No data	No data	
Chromium (and VI)	18540-29-9	1.10E-02		1.10E+01	No data	1.10E+01
Cobalt	7440-48-4	No data		No data	2.30E+01	2.30E+01
Copper	7440-50-8	9.00E-03		1.20E+01	No data	1.20E+01
Iron	7439-89-6	No data		1.00E+03	No data	1.00E+03
Lead	7439-92-1	2.50E-03		3.20E+00	No data	3.20E+00
Lithium	7439-93-2	No data		No data	1.40E+01	1.40E+01
Magnesium	7439-95-4	No data		No data	No data	
Manganese	7439-96-5	No data		No data	1.20E+02	1.20E+02
Mercury	7439-97-6	2.80E-06		No data	1.30E+00[f]	2.80E-06
Molybdenum	7439-98-7	No data		No data	3.70E+02	3.70E+02
Nickel	7440-02-0	5.20E-02		1.60E+02	No data	1.60E+02
Potassium	7440-09-7	No data		No data	No data	
Rhodium	7440-16-6	No data		No data	No data	
Selenium	7782-49-2	5.00E-03		5.00E+00	No data	5.00E+00
Silicon	7440-21-3	No data		No data	No data	
Silver	7440-22-4	1.20E-04		No data	3.60E-01	1.20E-04
Sodium	7440-23-5	No data		No data	No data	
Strontium	7440-24-6	No data		No data	1.50E+03	1.50E+03
Tantalum	7440-25-7	No data		No data	No data	
Thallium	7440-28-0	4.00E-03		No data	1.20E+01	4.00E-03
Tin	7440-31-5	No data		No data	7.30E+01	7.30E+01
Tungsten	7440-33-7	No data		No data	No data	
Uranium	7440-61-1	No data		No data	2.60E+00	2.60E+00
Vanadium	7440-62-2	No data		No data	2.00E+01	2.00E+01
Yttrium	7440-65-5	No data		No data	No data	

Table C3-9. Surface Water TRVs for Aquatic Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Suter & Tsao 1996[b]		Recommended TRV[c]
				NAWQC Chronic	Tier II Chronic	
Zinc	7440-66-6	1.18E-01		1.10E+02	No data	1.10E+02
Zirconium	7440-67-7	No data		No data	1.70E+01	1.70E+01
Non-metals and Anions						
Ammonia/Ammonium	7664-41-7	No data		No data	No data	
Bromide	24959-67-9	No data		No data	No data	
Chloride	16887-00-6	No data		No data	No data	
Cyanide	57-12-5	5.20E-03[i]		5.20E+00	No data	5.20E+00
Fluoride	16984-48-8	No data		No data	No data	
Hydroxide	14280-30-9	No data		No data	No data	
Iodine	7553-56-2	No data		No data	No data	
Nitrate	14797-55-8	No data		No data	No data	
Nitrite	14797-65-0	No data		No data	No data	
Phosphate	14265-44-2	No data		No data	No data	
Phosphorus	7723-14-0	No data		No data	No data	
Sulfate	14808-79-8	No data		No data	No data	
Total Sulfur	63705-05-5	No data		No data	No data	
Priority Pollutants						
Carbon Dioxide	124-38-9	No data		No data	No data	
Nitrogen Dioxide	10102-44-0	No data		No data	No data	
Ozone	10028-15-6	No data		No data	No data	
Particulate Matter	No CAS #	No data		No data	No data	
Sulfur Dioxode	7446-09-5	No data		No data	No data	
Radionuclides[k]						
Actinium-227	14952-40-0	NA		NA	NA	
Americium-241	1596-10-2	NA		NA	NA	
Americium-243	14993-75-0	NA		NA	NA	
Antimony-125	14234-35-6	NA		NA	NA	
Barium-137	13981-97-0	NA		NA	NA	
Cadmium-113	None	NA		NA	NA	

Table C3-9. Surface Water TRVs for Aquatic Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Suter & Tsao 1996[b]		Recommended TRV[c]
				NAWQC Chronic	Tier II Chronic	
Carbon-14	14762-75-5	NA		NA	NA	
Cesium-134	13967-70-9	NA		NA	NA	
Cesium-137	10045-97-3	NA		NA	NA	
Cobalt-60	10198-40-0	NA		NA	NA	
Curium-242	15510-73-3	NA		NA	NA	
Curium-243	15757-87-6	NA		NA	NA	
Curium-244	13981-15-2	NA		NA	NA	
Europium-152	14683-23-9	NA		NA	NA	
Europium-154	15585-10-1	NA		NA	NA	
Europium-155	14391-16-3	NA		NA	NA	
Iodine-129	15046-84-1	NA		NA	NA	
Neptunium-237	13994-20-2	NA		NA	NA	
Nickel-59	14336-70-0	NA		NA	NA	
Nickel-63	13981-37-8	NA		NA	NA	
Niobium-93	7440-03-1	NA		NA	NA	
Plutonium-238	13981-16-3	NA		NA	NA	
Plutonium-239	15117-48-3	NA		NA	NA	
Plutonium-240	14119-33-6	NA		NA	NA	
Plutonium-241	14119-32-5	NA		NA	NA	
Plutonium-242	13982-10-0	NA		NA	NA	
Protactinium-231	14331-85-2	NA		NA	NA	
Radium-226	13982-63-3	NA		NA	NA	
Radium-228	15262-20-1	NA		NA	NA	
Ruthenium-106	13967-48-1	NA		NA	NA	
Samarium-151	15715-94-3	NA		NA	NA	
Selenium-79	None	NA		NA	NA	
Strontium-90	10098-97-2	NA		NA	NA	
Technetium-99	14133-79-7	NA		NA	NA	
Thorium-229	15594-54-4	NA		NA	NA	

Table C3-9. Surface Water TRVs for Aquatic Biota

Constituent of Potential Concern	CAS Registry Number	EPA 1999[a]	Ecology Guidance TRV	Compiled by Suter & Tsao 1996[b]		Recommended TRV[c]
				NAWQC Chronic	Tier II Chronic	
Thorium-232	7440-29-1	NA		NA	NA	
Tin-126	15832-50-5	NA		NA	NA	
Tritium	10028-17-8	NA		NA	NA	
Uranium-232	14158-29-3	NA		NA	NA	
Uranium-233	13968-55-3	NA		NA	NA	
Uranium-234	13966-29-5	NA		NA	NA	
Uranium-235	15117-96-1	NA		NA	NA	
Uranium-236	13982-70-2	NA		NA	NA	
Uranium-238	7440-61-1	NA		NA	NA	
Yttrium-90	10098-91-6	NA		NA	NA	
Zirconium-93	15751-77-6	NA		NA	NA	

NA = Not applicable

[a] Published in Appendix E of EPA (1999), Table E-1

[b] Published in Table 1 of Suter and Tsao (1996)

[c] Order of preference is: NAWQC, then EPA 1999, then Tier II chronic value.

[d] Values calculated by the Great Lakes Water Quality Initiative (EPA 1993d)

[e] TRV was calculated using EqP approach (EPA 1993), assuming a fractional organic content of 0.04

[f] This value is based on final residue values

[g] Values calculated for OSWER (1996)

[h] These numbers are FAVs and FCVs calculated by EPA for use in the derivation of sediment quality criteria (EPA 1993b)

[i] TRV is a U.S. EPA Region 5 guideline value for classification of sediments for determining the suitability of dredged sediments for open water disposal, as cited in Hall and Suter II (1994).

[j] These values are draft Final Acute Values (FAV) and Final Chronic Values (FCV) values (EPA 1988b)

[k] TRV is not applicable to single radionuclides. Combined external and internal radiation exposure for aquatic biota from all radionuclides combined cannot exceed 1 rad/d.

References for Appendix C-3

ATSDR. 1989. *Toxicological Profile for 2-4 Dinitrotoluene and 2,6-Dinitrotoluene*, Agency for Toxic Substances and Disease Registry. US Public Health Service, Atlanta, Georgia, USA.

Clayton GD and Clayton FE. 1981. *Patty's Industrial Hygiene and Toxicology: Volume 2A: Toxicology*, Third Edition. John Wiley and Sons, New York, New York, USA.

Dilley JV, Tyson CA, Spanggord RJ, Sasmore DP, Newell GW, Dacre JC. 1982. *Short-term Oral Toxicity of 2,4,6-Trinitrotoluene in Mice, Rats, and Dogs*, J. Toxicol. Environ. Health, volume 9, p 565-585.

Ellis HV III, Hagensen JH, Hodgson JR, Minor JL, Hong CB, Ellis ER, Girvin JD, Helton DO, Herndon BL, and Lee CC. 1979. *Mammalian Toxicity of Munitions Compounds, Phase III: Effects of Lifetime Exposure, Part 1: 2,4-Dinitrotoluene*, Final Report Number 7, Contract Number DAMO 17-74-C-4073, ADA077692. . Midwest Research Institute, Kansas City, Missouri, USA.

EPA. 1996. *Ecotox Thresholds, ECO Update 3(2), EPA 540/F-95/038, Office of Solid Waste and Emergency Response*. US Environmental Protection Agency, Washington, D.C., USA.

Jones DS, Suter II GW, and Hull RN. 1997. *Toxicological Benchmarks for Screening Contaminants of Potential Concern for Effects on Sediment-Associated Biota: 1997 Revision*, ES/ER/TM-95/R4, November 1997. Lockheed Martin Energy Systems, Inc., Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA.

IRIS. 1996. *Integrated Risk Information System (IRIS), On-line Database of Toxicity Measures*. Office of Research and Development, Environmental Criteria and Assessment Office, Cincinnati, Ohio, USA.

Long ER and Morgan LG. 1991. *The Potential for Biological Effects of Sediment-Sorbed Contaminants Tested in the National Status and Trends Program*. National Oceanic and Atmospheric Administration Technical Memorandum NOS OMA 52, Seattle, Washington, USA.

Long ER, Macdonald DD, Smith SL, and Calder FD.1995. *Incidence of Adverse Biological Effects within Ranges of Chemical Concentrations in Marine and Estuarine Sediments*, Environmental Management, volume 19, number 1, p 81-97.

Persuad, D, Jaagumagi R, and Hayton A. 1993. *Guideline for the Protection and Management of Aquatic Sediment Quality in Ontario*, Log 92-2309-067, PIBS 1962, August 1993. Ontario Ministry of the Environment, Water Resources Branch, Toronto, Ontario, Canada.

Quast JF, Humiston CG, Kalnins RY, and others. 1979. *Results of a toxicity study of monomeric styrene administered to beagle dogs by oral intubation for 19 months*, Final Report. Toxicology Research Laboratory, Health and Environmental Sciences. DOW Chemical Co., Midland, Michigan, USA.

Sample BE, Opresko DM, and Suter II GW. 1996. *Toxicological Benchmarks for National Laboratory*. Oak Ridge, Tennessee, USA.

Suter, GW II and Tsao CL. 1996. *Toxicological Benchmarks for Screening Potential Contaminants of Concern for Effects on Aquatic Biota: 1996 Revision*, ES/ER/TM-96/R2, 1996. Lockheed Martin Energy Systems, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA.

Talmage, SS and others. (in press), "Nitroaromatic Munition Compounds: Environmental Effects and Screening Values", *Reviews of Environmental Contamination and Toxicology*.

Will ME and Suter II GW. 1995a. *Toxicological Benchmarks for Screening Potential Contaminants of Concern for Effects on Terrestrial Plants: 1995 Revision*, ES/ER/TM-85/R1, 1995. Lockheed Martin Energy Systems, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA.

Will ME and Suter II GW. 1995b. *Toxicological Benchmarks for Screening Potential Contaminants of Concern for Effects on Soil and Litter Invertebrates and Heterotrophic Process*, ES/ER/TM-126, 1995. Martin Marietta Energy Systems, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA.

Appendix C-4

Site-Specific Parameters for the RPP-WTP Risk Assessment

Table C4-1 Site-Specific Parameters for the RPP-WTP Risk Assessment

Parameter (independent variables)	Description	Value	Units	Note
	Terrestrial			
A	Area or Stream length	NA	ha, km	Assume larger than all home ranges and foraging distances
BD	Soil Bulk Density	1.5	g/cm^3	EPA (1998a)
ER	Soil enrichment ratio	3	none	ATG (1998)
Ev	Average annual evapotranspiration	NA	cm/yr	Not needed if ks = ksg + kdecay, i.e., no erosion, runoff, etc.
Irr	Average annual irrigation	NA	cm/yr	Appears as I (Appendix B-2); not needed if no erosion, runoff, etc.
ks	Soil-loss constant due to all processes (ksg, kdecay, kse, ksr, ksl, k	NA	yr^{-1}	ks = ksg + kdecay, if no runoff, erosion, etc.
kse	Soil-loss constant due to erosion (Xe, SD, ER, Kd, BD, Zs, θsw)	0	yr^{-1}	Sect. 6.2
ksl	Soil-loss constant due to leaching (P, Irr, RO, Ev, Kd, BD, Zs, θsv	0	yr^{-1}	Sect. 6.2
ksr	Soil-loss constant due to surface runoff (RO, Kd, BD, Zs, θsw)	0	yr^{-1}	Sect. 6.2
ksv	Soil-loss constant due to volatilization	0	yr^{-1}	EPA (1998a)
P	Average annual precipitation	NA	cm/yr	Not needed if ks = ksg + kdecay, i.e., no erosion, runoff, etc.
RO	Average annual surface runoff from pervious areas	NA	cm/yr	Not needed if ks = ksg + kdecay, i.e., no erosion, runoff, etc.
C	USLE cover management factor	0.1	unitless	ATG (1998)
LS	USLE length-slope factor	1.5	none	ATG (1998)
K	USLE erodibility factor	0.36	ton/acre	ATG (1998)
PF	USLE supporting practice factor	1	unitless	ATG (1998)
RF	USLE rainfall (erosivity) factor	NA	yr^{-1}	Not needed if no erosion contribution to TSS
Xe	Unit soil loss (RF, K, LS, C, PF)	0	kg/m^2yr	Not needed if no erosion
θsw	Soil volumetric water content	0.2	ml/cm^3	EPA (1998a)

Table C4-1 Site-Specific Parameters for the RPP-WTP Risk Assessment

Parameter (independent variables)	Description	Value	Units	Note
Z_s	Soil mixing zone depth	1	cm	EPA (1998a), untilled
	Aquatic			
A_I	Impervious watershed surface area	0	m^2	Assuming stationary model
A_L	Total watershed surface area	NA	m^2	Not needed if no erosion
A_W	Water body surface area	NA	m^2	May not be needed if deposition flux is per unit meter square area
a	Sediment Delivery Equation intercept, empirical	NA	none	Not needed if no erosion
b	Sediment Delivery Equation slope, empirical	0.125	none	ATG (1998)
d_{bs}	Depth of upper sediment layer	0.03	m	ATG (1998)
d_{wc}	Depth of water column	tbd	m	$d_{wc} = d_z - d_{bs}$
d_z	Total water body depth	tbd	m	
f_{bs}	Benthic sediment fract. of total water body COPC (f_{wc})	nd	none	$f_{bs} = (1-f_{wc})$
f_{wc}	Water col. fract. of total water body COPC (Kd_{sw}, TSS, d_{wc}, d_z, θ_b)	tbd	none	
FOC	Fraction of organic carbon in soil or sediment	0.04	none	ATG (1998); also appears as OCsed
kb	Benthic burial rate constant	nd	yr^{-1}	No equation given
kv	Water body loss constant due to volatilization (Kv, d_z, Kd_{sw}, TSS)	nd	yr^{-1}	
kwt	Total water body COPC dissipation rate constant (f_{wc}, kv, f_{bs}, kb)	nd	yr^{-1}	
K_G	Gas phase transfer coefficent (C_d, W, k, λ_z, μ_a, ρ_a, D_a)	nd	m/yr	
K_L	Liguid phase transfer coefficent (D_w, u, d_z) or (C_d, W, ρ_a, ρ_w, k, λ_z	nd	m/yr	
Kv	Overall COPC transfer rate coefficient (K_L, K_G, H, R)	nd	m/yr	
L_{dep}	Load from wet and dry deposition (Q, Fv, Dywwv, Dytwp, A_w)	nd	g/yr	Equation on in Sect. 6.4
L_{dif}	Load to water body by diffusion (Kv, Q, Fv, Cywv, A_w, H, R, T_{wk})	nd	g/yr	Equation on in Sect. 6.4
L_{ri}	Load from runoff from impervious surfaces (Q, Fv, Dywwv, Dytw	0	g/yr	Sect. 6.4
L_{rp}	Load from runoff from pervious surfaces (RO, A_L, A_I, Cs, BD, Kd	0	g/yr	Sect. 6.4
L_e	Load from soil erosion (Xe, A_L, A_I, SD, ER, Cs, BD, Kd, θsw)	0	g/yr	Sect. 6.4
L_{trans}	Load from within-body transformations	0	g/yr	(EPA 1998a)

Table C4-1 Site-Specific Parameters for the RPP-WTP Risk Assessment

Parameter (independent variables)	Description	Value	Units	Note
L_T	Total COPC load to water body (L_{dep}, L_{dif}, L_{ri}, L_{rp}, L_e, L_{trans})	nd	g/yr	$L_T = L_{dep} + L_{dif}$, if no runoff and erosion
SD	Sediment delivery ratio for watershed (a, A_L, b)	0	none	Assuming no erosion; ATG (1998) has values otherwise
θ_{bs}	Bed sediment porosity	0.6	L/L	ATG (1998)
TSS	Total suspended sediment (A_L, A_I, Vf_x, Dss, A_W, SD, Xe)	10	mg/L	ATG (1998)
T_{wk}	Water body temperature	tbd	°K	
u	Current velocity	tbd	m/s	
Vf_x	Average volumetric flow rate through water body	tbd	m^3/yr	
	Air			
μ_a	Viscosity of air at given temp.	tbd	g/cm s	
ρ_a	Density of air	tbd	g/cm^3	
W	Average annual wind speed	tbd	m/s	

ATG (1998) *Mixed Waste Facility RCRA/TSCA Permit Application, Attachment 4 - Risk Assessment Work Plan*, ATG, Inc., Richland, WA, July 1998.
EPA (1998a) *Human Health Risk Assessment Protocol for Hazardous Waste Combustion Facilities*. US EPA, Washington, DC
NA = not applicable
nd = no data
tbd = to be determined
USLE = Universal Soil Loss Equation

Appendix D

Selection of COPCs and ROPCs for Human Health and Ecological Risk Assessment

PDC 6-00

Table D-1. Selection of Human Health COPCs for Inclusion in the Quantitative PRA

		Are human health toxicity data available?		Can an exposure concentration be quantified?		Can human health risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Acute Toxicity? [a]	Chronic Toxicity? [b]	Emission Rate Available to Quantify Direct Exposure? [c]	Methods and Parameters Available to Quantify Indirect Exposure? [d]	Acute risk? [e]	Chronic Risk for Direct Exposure? [f]	Chronic Risk for Indirect Exposure? [g]	Is the constituent a quantitative COPC? [h]
Organics									
Aromatic Halogenated Hydrocarbons									
2,3,4,6-Tetrachlorophenol	58-90-2	no	yes	yes	yes	no	yes	yes	yes
4-Chloro-3-methylphenol	59-50-7	yes	no	yes	yes	yes	no	no	yes
Aromatic Nonhalogenated Hydrocarbons									
2-Nitrotoluene	88-72-2	no	yes	yes	yes	no	yes	yes	yes
4-Nitrobiphenyl	92-93-3	yes	no	yes	yes	yes	no	no	yes
Benzaldehyde	100-52-7	yes	yes	no	no	no	no	no	no
Benzene	71-43-2	yes	yes	yes	yes	yes	yes	yes	yes
Benzyl alcohol	100-51-6	yes	yes	no	no	no	no	no	no
Ethyl benzene	100-41-4	yes	yes	yes	yes	yes	yes	yes	yes
m-Xylene	108-38-3	yes	yes	yes	yes	yes	yes	yes	yes
o-Xylene	95-47-6	yes	yes	yes	yes	yes	yes	yes	yes
p-Xylene	106-42-3	yes	no	yes	yes	yes	no	no	yes
Styrene	100-42-5	yes	yes	yes	yes	yes	yes	yes	yes
Toluene	108-88-3	yes	yes	yes	yes	yes	yes	yes	yes
Non-aromatic Nonhalogenated Hydrocarbons									
1,2-Epoxybutane	106-88-7	yes	yes	yes	yes	yes	yes	yes	yes
1,3-Butadiene	106-99-0	yes	yes	yes	yes	yes	yes	yes	yes
1,4-Dioxane	123-91-1	yes	yes	yes	yes	yes	yes	yes	yes
1-Methylpropyl alcohol	78-92-2	yes	no	yes	yes	yes	no	no	yes
1-Nitropropane	108-03-2	yes	no	yes	yes	yes	no	no	yes
2,2,4-Trimethylpentane	540-84-1	yes	no	yes	yes	yes	no	no	yes
2-Butanone	78-93-3	yes	yes	yes	yes	yes	yes	yes	yes
2-Butenaldehyde (2-Butenal)	4170-30-3	yes	no	yes	yes	yes	no	no	yes

Table D-1. Selection of Human Health COPCs for Inclusion in the Quantitative PRA

		Are human health toxicity data available?		Can an exposure concentration be quantified?		Can human health risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Acute Toxicity? [a]	Chronic Toxicity? [b]	Emission Rate Available to Quantify Direct Exposure? [c]	Methods and Parameters Available to Quantify Indirect Exposure? [d]	Acute risk? [e]	Chronic Risk for Direct Exposure? [f]	Chronic Risk for Indirect Exposure? [g]	Is the constituent a quantitative COPC? [h]
2-Ethoxyethanol	110-80-5	yes	yes	no	no	no	no	no	no
2-Heptanone	110-43-0	yes	no	yes	yes	yes	no	no	yes
2-Hexanone	591-78-6	yes	no	yes	yes	yes	no	no	yes
2-Methoxyethanol	109-86-4	yes	yes	no	no	no	no	no	no
2-Methyl-2-propanol	75-65-0	yes	no	yes	yes	yes	no	no	yes
2-Methyl-2-propenenitrile	126-98-7	yes	yes	yes	yes	yes	yes	yes	yes
2-Methylaziridine	75-55-8	yes	no	yes	yes	yes	no	no	yes
2-Methylpropyl alcohol	78-83-1	yes	yes	yes	yes	yes	yes	yes	yes
2-Pentanone	107-87-9	yes	no	yes	yes	yes	no	no	yes
2-Propanone (Acetone)	67-64-1	yes	yes	yes	yes	yes	yes	yes	yes
2-Propene-1-ol	107-18-6	yes	yes	yes	yes	yes	yes	yes	yes
2-Propyl alcohol	67-63-0	yes	no	yes	yes	yes	no	no	yes
3-Heptanone	106-35-4	no	no	yes	yes	no	no	no	no
3-Methyl-1-butanol	123-51-3	yes	no	yes	yes	yes	no	no	yes
3-Methyl-2-butanone	563-80-4	no	no	yes	yes	no	no	no	no
3-Pentanone	96-22-0	no	no	yes	yes	no	no	no	no
4-Heptanone	123-19-3	no	no	yes	yes	no	no	no	no
4-Methyl-2-pentanone	108-10-1	yes	yes	yes	yes	yes	yes	yes	yes
4-Methyl-3-penten-2-one	141-79-7	yes	no	yes	yes	yes	no	no	yes
5-Methyl-2-hexanone	110-12-3	yes	no	yes	yes	yes	no	no	yes
Acetaldehyde	75-07-0	yes	yes	yes	yes	yes	yes	yes	yes
Acetamide	60-35-5	yes	no	yes	yes	yes	no	no	yes
Acetic acid	64-19-7	yes	no	yes	yes	yes	no	no	yes
Acetic acid ethyl ester	141-78-6	yes	yes	yes	yes	yes	yes	yes	yes
Acetic acid n-butyl ester	123-86-4	no	no	yes	yes	no	no	no	no
Acetonitrile	75-05-8	yes	yes	yes	yes	yes	yes	yes	yes
Acrolein	107-02-8	yes	yes	yes	yes	yes	yes	yes	yes

Table D-1. Selection of Human Health COPCs for Inclusion in the Quantitative PRA

		Are human health toxicity data available?		Can an exposure concentration be quantified?		Can human health risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Acute Toxicity? [a]	Chronic Toxicity? [b]	Emission Rate Available to Quantify Direct Exposure? [c]	Methods and Parameters Available to Quantify Indirect Exposure? [d]	Acute risk? [e]	Chronic Risk for Direct Exposure? [f]	Chronic Risk for Indirect Exposure? [g]	Is the constituent a quantitative COPC? [h]
Acrylonitrile	107-13-1	yes	yes	yes	yes	yes	yes	yes	yes
Bis(isopropyl)ether	108-20-3	no	no	yes	yes	no	no	no	no
Butane	106-97-8	yes	no	yes	yes	yes	no	no	yes
Carbon disulfide	75-15-0	yes	yes	yes	yes	yes	yes	yes	yes
Cyanogen	460-19-5	yes	yes	no	no	no	no	no	no
Cyclohexane	110-82-7	yes	no	yes	yes	yes	no	no	yes
Cyclohexanone	108-94-1	yes	yes	yes	yes	yes	yes	yes	yes
Cyclohexene	110-83-8	yes	no	yes	yes	yes	no	no	yes
Cyclopentane	287-92-3	no	no	yes	yes	no	no	no	no
Ethyl alcohol	64-17-5	yes	no	yes	yes	yes	no	no	yes
Ethyl ether	60-29-7	yes	yes	yes	yes	yes	yes	yes	yes
Ethyl methacrylate	97-63-2	no	yes	no	no	no	no	no	no
Formaldehyde	50-00-0	yes	yes	yes	yes	yes	yes	yes	yes
Formamide	75-12-7	yes	no	yes	yes	yes	no	no	yes
Formic acid	64-18-6	yes	yes	yes	yes	yes	yes	yes	yes
Formic acid, methyl ester	107-31-3	no	no	yes	yes	no	no	no	no
Glycidylaldehyde	765-34-4	yes	yes	no	no	no	no	no	no
Methyl acetate	79-20-9	no	yes	yes	yes	no	yes	yes	yes
Methyl alcohol	67-56-1	yes	yes	yes	yes	yes	yes	yes	yes
Methyl isocyanate	624-83-9	yes	no	yes	yes	yes	no	no	yes
Methyl methacrylate	80-62-6	yes	yes	no	no	no	no	no	no
Methyl tert-butyl ether	1634-04-4	yes	yes	yes	yes	yes	yes	yes	yes
Methylacetylene	74-99-7	yes	no	yes	yes	yes	no	no	yes
Methylcyclohexane	108-87-2	yes	yes	yes	yes	yes	yes	yes	yes
N,N-Dimethylacetamide	127-19-5	yes	no	yes	yes	yes	no	no	yes
n-Butyl alcohol	71-36-3	no	yes	yes	yes	no	yes	yes	yes
n-Heptane	142-82-5	yes	no	yes	yes	yes	no	no	yes

Table D-1. Selection of Human Health COPCs for Inclusion in the Quantitative PRA

		Are human health toxicity data available?		Can an exposure concentration be quantified?		Can human health risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Acute Toxicity? [a]	Chronic Toxicity? [b]	Emission Rate Available to Quantify Direct Exposure? [c]	Methods and Parameters Available to Quantify Indirect Exposure? [d]	Acute risk? [e]	Chronic Risk for Direct Exposure? [f]	Chronic Risk for Indirect Exposure? [g]	Is the constituent a quantitative COPC? [h]
n-Hexane	110-54-3	yes	yes	yes	yes	yes	yes	yes	yes
Nitromethane	75-52-5	yes	no	yes	yes	yes	no	no	yes
n-Nonane	111-84-2	yes	no	yes	yes	yes	no	no	yes
n-Octane	111-65-9	no	no	yes	yes	no	no	no	no
n-Pentane	109-66-0	yes	no	yes	yes	yes	no	no	yes
n-Propionaldehyde	123-38-6	yes	no	yes	yes	yes	no	no	yes
n-Propyl alcohol	71-23-8	yes	no	yes	yes	yes	no	no	yes
n-Valeraldehyde	110-62-3	no	no	yes	yes	no	no	no	no
Oxirane	75-21-8	yes	yes	yes	yes	yes	yes	yes	yes
p-Cymene	99-87-6	no	no	no	no	no	no	no	no
Phosgene	75-44-5	yes	no	no	no	no	no	no	no
Propargyl alcohol	107-19-7	yes	yes	no	no	no	no	no	no
Propionic acid	79-09-4	yes	no	yes	yes	yes	no	no	yes
Propionitrile	107-12-0	yes	no	yes	yes	yes	no	no	yes
Propylene gylcol monomethyl ether	107-98-2	yes	yes	no	no	no	no	no	no
p-tert-Butyltoluene	98-51-1	no	no	yes	yes	no	no	no	no
Triethylamine	121-44-8	yes	yes	yes	yes	yes	yes	yes	yes
Trimethylamine	75-50-3	yes	no	yes	yes	yes	no	no	yes
Vinyl acetate	108-05-4	yes	yes	yes	yes	yes	yes	yes	yes
Non-aromatic Halogenated Hydrocarbons									
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	no	no	yes	yes	no	no	no	no
1,1,1,2-Tetrachloroethane	630-20-6	yes	yes	yes	yes	yes	yes	yes	yes
1,1,1-Trichloroethane	71-55-6	yes	yes	yes	yes	yes	yes	yes	yes
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	no	no	yes	yes	no	no	no	no
1,1,2,2-Tetrachloroethane	79-34-5	yes	yes	yes	yes	yes	yes	yes	yes
1,1,2,2-Tetrachloroethene	127-18-4	yes	yes	yes	yes	yes	yes	yes	yes

Table D-1. Selection of Human Health COPCs for Inclusion in the Quantitative PRA

		Are human health toxicity data available?		Can an exposure concentration be quantified?		Can human health risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Acute Toxicity? [a]	Chronic Toxicity? [b]	Emission Rate Available to Quantify Direct Exposure? [c]	Methods and Parameters Available to Quantify Indirect Exposure? [d]	Acute risk? [e]	Chronic Risk for Direct Exposure? [f]	Chronic Risk for Indirect Exposure? [g]	Is the constituent a quantitative COPC? [h]
1,1,2-Trichloroethane	79-00-5	yes	yes	yes	yes	yes	yes	yes	yes
1,1,2-Trichloroethylene	79-01-6	yes	yes	yes	yes	yes	yes	yes	yes
1,1-Dichloroethane	75-34-3	yes	yes	yes	yes	yes	yes	yes	yes
1,1-Dichloroethene	75-35-4	yes	yes	yes	yes	yes	yes	yes	yes
1,2,2-Trichloro-1,1,2-trifluoroethane	76-13-1	yes	yes	yes	yes	yes	yes	yes	yes
1,2,3-Trichloropropane	96-18-4	yes	yes	yes	yes	yes	yes	yes	yes
1,2-Dibromo-3-chloropropane	96-12-8	yes	yes	no	no	no	no	no	no
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	yes	no	yes	yes	yes	no	no	yes
1,2-Dichloroethane	107-06-2	yes	yes	yes	yes	yes	yes	yes	yes
1,2-Dichloroethylene	540-59-0	yes	yes	yes	yes	yes	yes	yes	yes
1,2-Dichloropropane	78-87-5	yes	yes	yes	yes	yes	yes	yes	yes
1,3-Dichloropropene	542-75-6	yes	yes	no	no	no	no	no	no
1,4-Dichloro-2-butene	764-41-0	yes	yes	no	no	no	no	no	no
1-Chloroethene	75-01-4	yes	yes	yes	yes	yes	yes	yes	yes
2,2-Dichloropropionic acid	75-99-0	no	yes	yes	yes	no	yes	yes	yes
2-Chloropropane	75-29-6	no	yes	no	no	no	no	no	no
3-Chloropropene (Allyl chloride)	107-05-1	yes	yes	yes	yes	yes	yes	yes	yes
Bromochloromethane	74-97-5	yes	no	yes	yes	yes	no	no	yes
Bromodichloromethane	75-27-4	yes	yes	yes	yes	yes	yes	yes	yes
Bromoethene	593-60-2	yes	yes	no	no	no	no	no	no
Bromoform	75-25-2	yes	yes	no	no	no	no	no	no
Bromomethane	74-83-9	yes	yes	yes	yes	yes	yes	yes	yes
Carbon tetrachloride	56-23-5	yes	yes	yes	yes	yes	yes	yes	yes
Chlorodibromomethane	124-48-1	yes	yes	no	no	no	no	no	no
Chlorodifluoromethane	75-45-6	yes	yes	yes	yes	yes	yes	yes	yes
Chloroethane	75-00-3	yes	yes	yes	yes	yes	yes	yes	yes
Chloroform	67-66-3	yes	yes	yes	yes	yes	yes	yes	yes

Table D-1. Selection of Human Health COPCs for Inclusion in the Quantitative PRA

		Are human health toxicity data available?		Can an exposure concentration be quantified?		Can human health risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Acute Toxicity? [a]	Chronic Toxicity? [b]	Emission Rate Available to Quantify Direct Exposure? [c]	Methods and Parameters Available to Quantify Indirect Exposure? [d]	Acute risk? [e]	Chronic Risk for Direct Exposure? [f]	Chronic Risk for Indirect Exposure? [g]	Is the constituent a quantitative COPC? [h]
Chloromethane	74-87-3	yes	yes	yes	yes	yes	yes	yes	yes
Chloropentafluoroethane	76-15-3	no	no	yes	yes	no	no	no	no
cis-1,2-Dichloroethene	156-59-2	yes	yes	no	no	no	no	no	no
cis-1,3-Dichloropropene	10061-01-5	yes	no	yes	yes	yes	no	no	yes
Cyanogen bromide	506-68-3	no	yes	no	no	no	no	no	no
Cyanogen chloride	506-77-4	yes	yes	no	no	no	no	no	no
Dichlorodifluoromethane	75-71-8	yes	yes	yes	yes	yes	yes	yes	yes
Dichlorofluoromethane	75-43-4	yes	no	yes	yes	yes	no	no	yes
Dichloromethane	75-09-2	yes	yes	yes	yes	yes	yes	yes	yes
Difluorodibromomethane	75-61-6	no	no	yes	yes	no	no	no	no
Hexafluoroacetone	684-16-2	yes	no	yes	yes	yes	no	no	yes
Iodomethane	74-88-4	yes	no	no	no	no	no	no	no
Methylene bromide	74-95-3	yes	yes	no	no	no	no	no	no
Pentachloroethane	76-01-7	yes	no	no	no	no	no	no	no
trans-1,2-Dichloroethylene	156-60-5	yes	yes	yes	yes	yes	yes	yes	yes
trans-1,3-Dichloropropene	10061-02-6	no	no	yes	yes	no	no	no	no
Trichloroacetic acid	76-03-9	yes	no	yes	yes	yes	no	no	yes
Trichlorofluoroethane	27154-33-2	no	no	yes	yes	no	no	no	no
Trichlorofluoromethane	75-69-4	yes	yes	yes	yes	yes	yes	yes	yes
Trifluorobromomethane	75-63-8	yes	no	yes	yes	yes	no	no	yes
Dioxin and Furan Compounds									
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	35822-46-9	no	no*	yes	yes	no	yes	yes	yes
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	no	no*	yes	yes	no	yes	yes	yes
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	no	no*	yes	yes	no	yes	yes	yes
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	39227-28-6	no	no*	yes	yes	no	yes	yes	yes
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	yes	no*	yes	yes	yes	yes	yes	yes

Table D-1. Selection of Human Health COPCs for Inclusion in the Quantitative PRA

		Are human health toxicity data available?		Can an exposure concentration be quantified?		Can human health risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Acute Toxicity? [a]	Chronic Toxicity? [b]	Emission Rate Available to Quantify Direct Exposure? [c]	Methods and Parameters Available to Quantify Indirect Exposure? [d]	Acute risk? [e]	Chronic Risk for Direct Exposure? [f]	Chronic Risk for Indirect Exposure? [g]	Is the constituent a quantitative COPC? [h]
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	57653-85-7	yes	no*	yes	yes	yes	yes	yes	yes
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	yes	no*	yes	yes	yes	yes	yes	yes
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	19408-74-3	no	no*	yes	yes	no	yes	yes	yes
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	no	no*	yes	yes	no	yes	yes	yes
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	40321-76-4	yes	no*	yes	yes	yes	yes	yes	yes
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	no	no*	yes	yes	no	yes	yes	yes
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	yes	no*	yes	yes	yes	yes	yes	yes
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	yes	no*	yes	yes	yes	yes	yes	yes
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1746-01-6	yes	yes	yes	yes	yes	yes	yes	yes
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	yes	no*	yes	yes	yes	yes	yes	yes
Dibenzofuran	132-64-9	no	no	yes	yes	no	no	no	no
Octachlorodibenzo(p)dioxin	3268-87-9	no	no*	yes	yes	no	yes	yes	yes
Octachlorodibenzofuran	39001-02-0	no	no*	yes	yes	no	yes	yes	yes
PCBs									
2,2',3,3',4,4',5- Heptachlorobiphenyl	35065-30-6	no	no	yes	yes	no	no	no	no
2,2',3,4,4',5,5'- Heptachlorobiphenyl	35065-29-3	no	no	yes	yes	no	no	no	no
2,3,3',4,4',5'- Hexachlorobiphenyl	69782-90-7	no	no*	yes	yes	no	yes	yes	yes
2,3,3',4,4',5- Hexachlorobiphenyl	38380-08-4	no	no*	yes	yes	no	yes	yes	yes
2,3,3',4,4',5,5'- Heptachlorobiphenyl	no cas #	no	no*	yes	yes	no	yes	yes	yes
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	no	no*	yes	yes	no	yes	yes	yes
2,3,4,4',5- Pentachlorobiphenyl	74472-37-0	no	no*	yes	yes	no	yes	yes	yes
2',3,4,4',5- Pentachlorobiphenyl	no cas #	no	no*	yes	yes	no	yes	yes	yes
2,3',4,4',5- Pentachlorobiphenyl	31508-00-6	no	no*	yes	yes	no	yes	yes	yes
2,3',4,4',5,5'- Hexachlorobiphenyl	no cas #	no	no*	yes	yes	no	yes	yes	yes
3,3',4,4',5- Pentachlorobiphenyl	no cas #	no	no*	yes	yes	no	yes	yes	yes
3,3',4,4',5,5'- Hexachlorobiphenyl	32774-16-6	no	no*	yes	yes	no	yes	yes	yes

Table D-1. Selection of Human Health COPCs for Inclusion in the Quantitative PRA

		Are human health toxicity data available?		Can an exposure concentration be quantified?		Can human health risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Acute Toxicity? [a]	Chronic Toxicity? [b]	Emission Rate Available to Quantify Direct Exposure? [c]	Methods and Parameters Available to Quantify Indirect Exposure? [d]	Acute risk? [e]	Chronic Risk for Direct Exposure? [f]	Chronic Risk for Indirect Exposure? [g]	Is the constituent a quantitative COPC? [h]
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	no	no*	yes	yes	no	yes	yes	yes
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	no	no*	yes	yes	no	yes	yes	yes
Polychlorinated biphenyls (PCBs)	1336-36-3	yes	yes	yes	yes	yes	yes	yes	yes
Phthalates									
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	yes	yes	yes	yes	yes	yes	yes	yes
Butylbenzyl phthalate	85-68-7	yes	yes	yes	yes	yes	yes	yes	yes
Dibutyl phthalate	84-74-2	yes	yes	yes	yes	yes	yes	yes	yes
Diethyl phthalate	84-66-2	yes	yes	yes	yes	yes	yes	yes	yes
Dimethylphthalate	131-11-3	yes	no	no	no	no	no	no	no
n-Dioctyl phthalate	117-84-0	yes	yes	yes	yes	yes	yes	yes	yes
Light Polycyclic Aromatic Hydrocarbons									
2-Chloronapthalene	91-58-7	yes	yes	yes	yes	yes	yes	yes	yes
2-Methyl naphthalene	91-57-6	yes	no	no	no	no	no	no	no
5-Nitroacenaphthene	602-87-9	no	no	yes	yes	no	no	no	no
Acenaphthene	83-32-9	yes	yes	yes	yes	yes	yes	yes	yes
Acenaphthylene	208-96-8	yes	no	yes	yes	yes	no	no	yes
Anthracene	120-12-7	yes	yes	yes	yes	yes	yes	yes	yes
Fluorene	86-73-7	yes	yes	yes	yes	yes	yes	yes	yes
Indene	95-13-6	yes	no	yes	yes	yes	no	no	yes
Naphthalene	91-20-3	yes	yes	yes	yes	yes	yes	yes	yes
Phenanthrene	85-01-8	yes	no	yes	yes	yes	no	no	yes
Pyrene	129-00-0	yes	yes	yes	yes	yes	yes	yes	yes
Heavy Polycyclic Aromatic Hydrocarbons									
3-Methylcholanthrene	56-49-5	yes	no	yes	yes	yes	no	no	yes

Table D-1. Selection of Human Health COPCs for Inclusion in the Quantitative PRA

		Are human health toxicity data available?		Can an exposure concentration be quantified?		Can human health risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Acute Toxicity? [a]	Chronic Toxicity? [b]	Emission Rate Available to Quantify Direct Exposure? [c]	Methods and Parameters Available to Quantify Indirect Exposure? [d]	Acute risk? [e]	Chronic Risk for Direct Exposure? [f]	Chronic Risk for Indirect Exposure? [g]	Is the constituent a quantitative COPC? [h]
5-Methylchrysene	3697-24-3	no	no	yes	yes	no	no	no	no
Benzo(a)anthracene	56-55-3	yes	no*	yes	yes	yes	yes	yes	yes
Benzo(a)pyrene	50-32-8	yes	yes	yes	yes	yes	yes	yes	yes
Benzo(b)fluoranthene	205-99-2	no	no*	yes	yes	no	yes	yes	yes
Benzo(e)pyrene	192-97-2	no	no	no	no	no	no	no	no
Benzo(g,h,i)perylene	191-24-2	yes	no	yes	yes	yes	no	no	yes
Benzo(j)fluoranthene	205-82-3	no	no	yes	yes	no	no	no	no
Benzo(k)fluoranthene	207-08-9	no	no*	yes	yes	no	yes	yes	yes
Benzo[a,i]pyrene	191-30-0	no	no	yes	yes	no	no	no	no
Chrysene	218-01-9	yes	no*	yes	yes	yes	yes	yes	yes
Dibenz(a,h)anthracene	53-70-3	yes	no*	yes	yes	yes	yes	yes	yes
Dibenz[a,h]acridine	226-36-8	no	no	yes	yes	no	no	no	no
Dibenz[a,j]acridine	224-42-0	no	no	yes	yes	no	no	no	no
Dibenzo(a,e)fluoranthene	5385-75-1	no	no	no	no	no	no	no	no
Dibenzo(a,h)fluoranthene	no cas #	no	no	no	no	no	no	no	no
Dibenzo[a,e]pyrene	192-65-4	no	no	yes	yes	no	no	no	no
Dibenzo[a,h]pyrene	189-64-0	no	no	yes	yes	no	no	no	no
Dibenzo[a,i]pyrene	189-55-9	no	no	yes	yes	no	no	no	no
Fluoranthene	206-44-0	yes	yes	yes	yes	yes	yes	yes	yes
Hexachloronaphthalene	1335-87-1	yes	no	yes	yes	yes	no	no	yes
Indeno(1,2,3-cd)pyrene	193-39-5	no	no*	yes	yes	no	yes	yes	yes
Octachloronaphthalene	2234-13-1	yes	no	yes	yes	yes	no	no	yes
Pentachloronaphthalene	1321-64-8	no	no	yes	yes	no	no	no	no
Tetrachloronaphthalene	1335-88-2	no	no	yes	yes	no	no	no	no
Trichloronaphthalene	1321-65-9	no	no	yes	yes	no	no	no	no
Light Substituted Benzene Compounds									

Table D-1. Selection of Human Health COPCs for Inclusion in the Quantitative PRA

		Are human health toxicity data available?		Can an exposure concentration be quantified?		Can human health risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Acute Toxicity? [a]	Chronic Toxicity? [b]	Emission Rate Available to Quantify Direct Exposure? [c]	Methods and Parameters Available to Quantify Indirect Exposure? [d]	Acute risk? [e]	Chronic Risk for Direct Exposure? [f]	Chronic Risk for Indirect Exposure? [g]	Is the constituent a quantitative COPC? [h]
1,2,3-Trichlorobenzene	87-61-6	yes	no	no	no	no	no	no	no
1,2,4-Trichlorobenzene	120-82-1	yes	yes	yes	yes	yes	yes	yes	yes
1,2,4-Trimethyl benzene	95-63-6	yes	no	no	no	no	no	no	no
1,2-Dichlorobenzene	95-50-1	yes	yes	yes	yes	yes	yes	yes	yes
1,3,5-Trimethyl benzene	108-67-8	yes	no	no	no	no	no	no	no
1,3-Dichlorobenzene	541-73-1	yes	no	yes	yes	yes	no	no	yes
1,3-Dinitrobenzene	99-65-0	yes	yes	no	no	no	no	no	no
1,4-Dichlorobenzene	106-46-7	yes	yes	yes	yes	yes	yes	yes	yes
1,4-Dinitrobenzene	100-25-4	no	yes	yes	yes	no	yes	yes	yes
2,4,5-Trichlorophenol	95-95-4	yes	yes	yes	yes	yes	yes	yes	yes
2,4,6-Trichlorophenol	88-06-2	yes	yes	yes	yes	yes	yes	yes	yes
2,4-Dichlorophenol	120-83-2	yes	yes	yes	yes	yes	yes	yes	yes
2,4-Dimethylphenol	105-67-9	no	yes	no	no	no	no	no	no
2,4-Dinitrophenol	51-28-5	yes	yes	no	no	no	no	no	no
2,4-Dinitrotoluene	121-14-2	yes	yes	no	no	no	no	no	no
2,6-Dinitrotoluene	606-20-2	yes	yes	no	no	no	no	no	no
2-Chlorophenol	95-57-8	yes	yes	yes	yes	yes	yes	yes	yes
2-Chlorotoluene	95-49-8	yes	yes	yes	yes	yes	yes	yes	yes
2-Nitrophenol	88-75-5	no	no	yes	yes	no	no	no	no
4,6-Dinitro-o-cresol	534-52-1	yes	no	no	no	no	no	no	no
4-Chlorotoluene	106-43-4	yes	no	no	no	no	no	no	no
4-Nitrophenol	100-02-7	yes	no	no	no	no	no	no	no
alpha-Methylstyrene	98-83-9	no	yes	yes	yes	no	yes	yes	yes
Aniline	62-53-3	yes	yes	no	no	no	no	no	no
Benzotrichloride	98-07-7	yes	yes	no	no	no	no	no	no
Benzyl chloride	100-44-7	yes	yes	no	no	no	no	no	no
Bromobenzene	108-86-1	yes	no	no	no	no	no	no	no

Table D-1. Selection of Human Health COPCs for Inclusion in the Quantitative PRA

		Are human health toxicity data available?		Can an exposure concentration be quantified?		Can human health risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Acute Toxicity? [a]	Chronic Toxicity? [b]	Emission Rate Available to Quantify Direct Exposure? [c]	Methods and Parameters Available to Quantify Indirect Exposure? [d]	Acute risk? [e]	Chronic Risk for Direct Exposure? [f]	Chronic Risk for Indirect Exposure? [g]	Is the constituent a quantitative COPC? [h]
Chlorobenzene	108-90-7	yes	yes	yes	yes	yes	yes	yes	yes
Cumene	98-82-8	yes	yes	yes	yes	yes	yes	yes	yes
m-Cresol	108-39-4	no	yes	yes	yes	no	yes	yes	yes
n-Butyl benzene	104-51-8	yes	no	no	no	no	no	no	no
Nitrobenzene	98-95-3	yes	yes	yes	yes	yes	yes	yes	yes
n-Propyl benzene	103-65-1	yes	no	no	no	no	no	no	no
o-Cresol	95-48-7	no	yes	yes	yes	no	yes	yes	yes
o-Dinitrobenzene	528-29-0	no	yes	no	no	no	no	no	no
o-Nitroaniline	88-74-4	no	yes	no	no	no	no	no	no
o-Toluidine	95-53-4	yes	yes	no	no	no	no	no	no
p-Chloroaniline	106-47-8	yes	yes	no	no	no	no	no	no
p-Cresol	106-44-5	no	yes	no	no	no	no	no	no
Phenol	108-95-2	yes	yes	yes	yes	yes	yes	yes	yes
p-Nitrochlorobenzene	100-00-5	yes	no	yes	yes	yes	no	no	yes
p-Toluidine	106-49-0	no	yes	no	no	no	no	no	no
sec-Butyl benzene	135-98-8	yes	no	no	no	no	no	no	no
tert-Butyl benzene	98-06-6	yes	no	no	no	no	no	no	no
Toluene-2,6-diamine	823-40-5	no	yes	no	no	no	no	no	no
Trimethyl benzene	25551-13-7	no	no	yes	yes	no	no	no	no
Other Light Semivolatile Compounds									
1,1`-Biphenyl	92-52-4	yes	yes	yes	yes	yes	yes	yes	yes
1,1-Dimethylhydrazine	57-14-7	yes	no	yes	yes	yes	no	no	yes
1,2-Dimethylhydrazine	540-73-8	yes	no	no	no	no	no	no	no
1,2-Diphenylhydrazine	122-66-7	yes	yes	no	no	no	no	no	no
1,3-Propane sultone	1120-71-4	yes	no	no	no	no	no	no	no
2,4-Toluene diisocyante	584-84-9	yes	no	no	no	no	no	no	no

Table D-1. Selection of Human Health COPCs for Inclusion in the Quantitative PRA

		Are human health toxicity data available?		Can an exposure concentration be quantified?		Can human health risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Acute Toxicity? [a]	Chronic Toxicity? [b]	Emission Rate Available to Quantify Direct Exposure? [c]	Methods and Parameters Available to Quantify Indirect Exposure? [d]	Acute risk? [e]	Chronic Risk for Direct Exposure? [f]	Chronic Risk for Indirect Exposure? [g]	Is the constituent a quantitative COPC? [h]
2-Chloroacetophenone	532-27-4	no	yes	no	no	no	no	no	no
2-Propenoic acid	79-10-7	yes	yes	yes	yes	yes	yes	yes	yes
4,4-Methylenedianiline	101-77-9	yes	no	no	no	no	no	no	no
Acetophenone	98-86-2	yes	yes	yes	yes	yes	yes	yes	yes
Benzoic acid	65-85-0	yes	yes	no	no	no	no	no	no
Bis(2-chloroethoxy)methane	111-91-1	no	no	no	no	no	no	no	no
Bis(2-chloroethyl) ether	111-44-4	yes	yes	no	no	no	no	no	no
Chlorocyclopentadiene	41851-50-7	no	no	no	no	no	no	no	no
Cyclohexanol	108-93-0	no	no	yes	yes	no	no	no	no
Dichloroisopropyl ether	108-60-1	yes	yes	no	no	no	no	no	no
Dichloromethyl ether	542-88-1	yes	yes	yes	yes	yes	yes	yes	yes
Dichloropentadiene	no cas #	no	no	no	no	no	no	no	no
Dimethyl sulfate	77-78-1	yes	no	no	no	no	no	no	no
Dimethylaniline	121-69-7	no	yes	yes	yes	no	yes	yes	yes
Di-n-Propylnitrosamine	621-64-7	yes	yes	yes	yes	yes	yes	yes	yes
Diphenyl ether	101-84-8	no	no	yes	yes	no	no	no	no
Epichlorohydrin	106-89-8	yes	yes	no	no	no	no	no	no
Ethyl Carbamate (urethane)	51-79-6	no	no	no	no	no	no	no	no
Ethyl methanesulfonate	62-50-0	no	no	no	no	no	no	no	no
Ethylene dibromide	106-93-4	yes	yes	yes	yes	yes	yes	yes	yes
Ethylene glycol	107-21-1	yes	yes	no	no	no	no	no	no
Ethylene glycol monobutyl ether	111-76-2	yes	yes	yes	yes	yes	yes	yes	yes
Ethylene glycol monoethyl ether acetate	111-15-9	yes	yes	no	no	no	no	no	no
Ethylene thiourea	96-45-7	yes	yes	no	no	no	no	no	no
Furfural	98-01-1	yes	yes	no	no	no	no	no	no
Maleic hydrazide	123-33-1	no	yes	no	no	no	no	no	no
Malononitrile	109-77-3	no	yes	no	no	no	no	no	no

Table D-1. Selection of Human Health COPCs for Inclusion in the Quantitative PRA

		Are human health toxicity data available?		Can an exposure concentration be quantified?		Can human health risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Acute Toxicity? [a]	Chronic Toxicity? [b]	Emission Rate Available to Quantify Direct Exposure? [c]	Methods and Parameters Available to Quantify Indirect Exposure? [d]	Acute risk? [e]	Chronic Risk for Direct Exposure? [f]	Chronic Risk for Indirect Exposure? [g]	Is the constituent a quantitative COPC? [h]
Methyl styrene (mixed isomers)	25013-15-4	no	yes	no	no	no	no	no	no
Methylhydrazine	60-34-4	yes	no	yes	yes	yes	no	no	yes
N,N-Diphenylamine	122-39-4	yes	yes	yes	yes	yes	yes	yes	yes
Nitric acid, propyl ester	627-13-4	no	no	yes	yes	no	no	no	no
N-Nitrosodi-n-butylamine	924-16-3	no	yes	no	no	no	no	no	no
N-Nitrosomorpholine	59-89-2	yes	no	yes	yes	yes	no	no	yes
N-Nitroso-N,N-dimethylamine	62-75-9	yes	yes	yes	yes	yes	yes	yes	yes
o-Anisidine	90-04-0	yes	no	no	no	no	no	no	no
Oxalic acid	144-62-7	yes	no	yes	yes	yes	no	no	yes
Phthalic anhydride	85-44-9	yes	yes	no	no	no	no	no	no
p-Phthalic acid	100-21-0	no	yes	yes	yes	no	yes	yes	yes
Pyridine	110-86-1	yes	yes	yes	yes	yes	yes	yes	yes
Quinoline	91-22-5	yes	yes	yes	yes	yes	yes	yes	yes
Quinone	106-51-4	yes	no	no	no	no	no	no	no
Safrole	94-59-7	no	no	no	no	no	no	no	no
Tetrahydrofuran	109-99-9	yes	no	yes	yes	yes	no	no	yes
Other Heavy Semivolatile Compounds									
1,2,4,5-Tetrachlorobenzene	95-94-3	yes	yes	no	no	no	no	no	no
1,3,5-Trinitrobenzene	99-35-4	yes	yes	no	no	no	no	no	no
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	no	no	yes	yes	no	no	no	no
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	no	yes	no	no	no	no	no	no
2-sec-Butyl-4,6-dinitrophenol	88-85-7	yes	yes	yes	yes	yes	yes	yes	yes
3,3-Dichlorobenzidine	91-94-1	yes	yes	no	no	no	no	no	no
3,3'-Dimethoxybenzidine	119-90-4	yes	yes	no	no	no	no	no	no
4-Bromophenylphenyl ether	101-55-3	no	no	yes	yes	no	no	no	no
Ammonium perfluorooctanoate	3825-26-1	no	no	yes	yes	no	no	no	no

Table D-1. Selection of Human Health COPCs for Inclusion in the Quantitative PRA

		Are human health toxicity data available?		Can an exposure concentration be quantified?		Can human health risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Acute Toxicity? [a]	Chronic Toxicity? [b]	Emission Rate Available to Quantify Direct Exposure? [c]	Methods and Parameters Available to Quantify Indirect Exposure? [d]	Acute risk? [e]	Chronic Risk for Direct Exposure? [f]	Chronic Risk for Indirect Exposure? [g]	Is the constituent a quantitative COPC? [h]
Azobenzene	103-33-3	no	yes	no	no	no	no	no	no
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulf	96-69-5	no	no	yes	yes	no	no	no	no
Captan	133-06-2	yes	yes	no	no	no	no	no	no
Chlorobenzilate	510-15-6	yes	yes	no	no	no	no	no	no
Dibutylphosphate	107-66-4	yes	no	yes	yes	yes	no	no	yes
Dimethyl aminoazobenzene	60-11-7	yes	no	no	no	no	no	no	no
Hexachlorobenzene	118-74-1	yes	yes	yes	yes	yes	yes	yes	yes
Hexachlorobutadiene	87-68-3	yes	yes	yes	yes	yes	yes	yes	yes
Hexachlorocyclopentadiene	77-47-4	yes	yes	no	no	no	no	no	no
Hexachloroethane	67-72-1	yes	yes	yes	yes	yes	yes	yes	yes
Hexachlorophene	70-30-4	yes	yes	no	no	no	no	no	no
Hexamethylene-1,5-diisocyanate	822-06-0	yes	yes	no	no	no	no	no	no
Mirex	2385-85-5	yes	yes	yes	yes	yes	yes	yes	yes
Nitrofen	1836-75-5	no	no	yes	yes	no	no	no	no
Pentachlorobenzene	608-93-5	yes	yes	no	no	no	no	no	no
Pentachloronitrobenzene	82-68-8	yes	yes	yes	yes	yes	yes	yes	yes
Pentachlorophenol	87-86-5	yes	yes	yes	yes	yes	yes	yes	yes
Picric acid	88-89-1	yes	no	yes	yes	yes	no	no	yes
Pronamide	23950-58-5	no	yes	no	no	no	no	no	no
Strychnine	57-24-9	yes	yes	no	no	no	no	no	no
Terphenyls	26140-60-3	yes	no	yes	yes	yes	no	no	yes
Tributyl phosphate	126-73-8	yes	no	yes	yes	yes	no	no	yes
Trifluralin	1582-09-8	yes	yes	yes	yes	yes	yes	yes	yes
Triphenylamine	603-34-9	no	no	yes	yes	no	no	no	no
Herbicides and Organochlorinated Pesticides									
2,4,5-T	93-76-5	yes	yes	yes	yes	yes	yes	yes	yes

Table D-1. Selection of Human Health COPCs for Inclusion in the Quantitative PRA

		Are human health toxicity data available?		Can an exposure concentration be quantified?		Can human health risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Acute Toxicity? [a]	Chronic Toxicity? [b]	Emission Rate Available to Quantify Direct Exposure? [c]	Methods and Parameters Available to Quantify Indirect Exposure? [d]	Acute risk? [e]	Chronic Risk for Direct Exposure? [f]	Chronic Risk for Indirect Exposure? [g]	Is the constituent a quantitative COPC? [h]
2,4-D and esters	94-75-7	yes	yes	yes	yes	yes	yes	yes	yes
4,4-DDD	72-54-8	yes	yes	yes	yes	yes	yes	yes	yes
4,4-DDE	72-55-9	yes	yes	yes	yes	yes	yes	yes	yes
4,4-DDT	50-29-3	yes	yes	yes	yes	yes	yes	yes	yes
Aldrin	309-00-2	yes	yes	yes	yes	yes	yes	yes	yes
alpha-BHC	319-84-6	yes	yes	yes	yes	yes	yes	yes	yes
beta-BHC	319-85-7	yes	yes	yes	yes	yes	yes	yes	yes
Chlordane	57-74-9	yes	yes	no	no	no	no	no	no
Delta-BHC	319-86-8	no	no	yes	yes	no	no	no	no
Dieldrin	60-57-1	yes	yes	yes	yes	yes	yes	yes	yes
Endothall	145-73-3	no	yes	no	no	no	no	no	no
Endrin	72-20-8	yes	yes	yes	yes	yes	yes	yes	yes
gamma-BHC (Lindane)	58-89-9	yes	yes	yes	yes	yes	yes	yes	yes
Heptachlor	76-44-8	yes	yes	yes	yes	yes	yes	yes	yes
Isodrin	465-73-6	no	no	yes	yes	no	no	no	no
Methoxychlor	72-43-5	yes	yes	yes	yes	yes	yes	yes	yes
Silvex (2,4,5-TP)	93-72-1	yes	yes	yes	yes	yes	yes	yes	yes
Toxaphene	8001-35-2	yes	yes	yes	yes	yes	yes	yes	yes
Inorganics									
Metals									
Aluminum	7429-90-5	yes	no	yes	yes	yes	no	no	yes
Antimony	7440-36-0	yes	yes	yes	yes	yes	yes	yes	yes
Arsenic	7440-38-2	yes	yes	yes	yes	yes	yes	yes	yes
Barium	7440-39-3	yes	yes	yes	yes	yes	yes	yes	yes
Beryllium	7440-41-7	yes	yes	yes	yes	yes	yes	yes	yes
Bismuth	7440-69-9	yes	no	yes	yes	yes	no	no	yes

Table D-1. Selection of Human Health COPCs for Inclusion in the Quantitative PRA

		Are human health toxicity data available?		Can an exposure concentration be quantified?		Can human health risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Acute Toxicity? [a]	Chronic Toxicity? [b]	Emission Rate Available to Quantify Direct Exposure? [c]	Methods and Parameters Available to Quantify Indirect Exposure? [d]	Acute risk? [e]	Chronic Risk for Direct Exposure? [f]	Chronic Risk for Indirect Exposure? [g]	Is the constituent a quantitative COPC? [h]
Boron	7440-42-8	yes	yes	yes	yes	yes	yes	yes	yes
Cadmium	7440-43-9	yes	yes	yes	yes	yes	yes	yes	yes
Calcium	7440-70-2	yes	no	yes	yes	yes	no	no	yes
Chromium (and VI)	18540-29-9	yes	yes	yes	yes	yes	yes	yes	yes
Cobalt	7440-48-4	yes	no	yes	yes	yes	no	no	yes
Copper	7440-50-8	yes	no	yes	yes	yes	no	no	yes
Iron	7439-89-6	yes	no	yes	yes	yes	no	no	yes
Lead	7439-92-1	yes	no**	yes	yes	yes	yes	yes	yes
Lithium	7439-93-2	yes	no	yes	yes	yes	no	no	yes
Magnesium	7439-95-4	yes	no	yes	yes	yes	no	no	yes
Manganese	7439-96-5	yes	yes	yes	yes	yes	yes	yes	yes
Mercury	7439-97-6	yes	no**	yes	yes	yes	yes	yes	yes
Molybdenum	7439-98-7	yes	yes	yes	yes	yes	yes	yes	yes
Nickel	7440-02-0	yes	yes	yes	yes	yes	yes	yes	yes
Potassium	7440-09-7	yes	no	yes	yes	yes	no	no	yes
Rhodium	7440-16-6	yes	no	yes	yes	yes	no	no	yes
Selenium	7782-49-2	yes	yes	yes	yes	yes	yes	yes	yes
Silicon	7440-21-3	yes	no	yes	yes	yes	no	no	yes
Silver	7440-22-4	yes	yes	yes	yes	yes	yes	yes	yes
Sodium	7440-23-5	yes	no	yes	yes	yes	no	no	yes
Strontium	7440-24-6	yes	yes	no	no	no	no	no	no
Tantalum	7440-25-7	no	no	yes	yes	no	no	no	no
Thallium	7440-28-0	yes	no	yes	yes	yes	no	no	yes
Tin	7440-31-5	yes	yes	yes	yes	yes	yes	yes	yes
Tungsten	7440-33-7	yes	no	yes	yes	yes	no	no	yes
Uranium	7440-61-1	yes	yes	yes	yes	yes	yes	yes	yes
Vanadium	7440-62-2	yes	yes	yes	yes	yes	yes	yes	yes

Table D-1. Selection of Human Health COPCs for Inclusion in the Quantitative PRA

		Are human health toxicity data available?		Can an exposure concentration be quantified?		Can human health risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Acute Toxicity? [a]	Chronic Toxicity? [b]	Emission Rate Available to Quantify Direct Exposure? [c]	Methods and Parameters Available to Quantify Indirect Exposure? [d]	Acute risk? [e]	Chronic Risk for Direct Exposure? [f]	Chronic Risk for Indirect Exposure? [g]	Is the constituent a quantitative COPC? [h]
Yttrium	7440-65-5	yes	no	yes	yes	yes	no	no	yes
Zinc	7440-66-6	yes	yes	yes	yes	yes	yes	yes	yes
Zirconium	7440-67-7	yes	no	yes	yes	yes	no	no	yes
Non-metals and Anions									
Ammonia/Ammonium	7664-41-7	yes	yes	yes	yes	yes	yes	yes	yes
Bromide	24959-67-9	no	no	yes	yes	no	no	no	no
Chloride	16887-00-6	no	no	yes	yes	no	no	no	no
Cyanide	57-12-5	yes	yes	yes	yes	yes	yes	yes	yes
Fluoride	16984-48-8	yes	yes	yes	yes	yes	yes	yes	yes
Hydroxide	14280-30-9	no	no	yes	yes	no	no	no	no
Iodine	7553-56-2	yes	no	yes	yes	yes	no	no	yes
Nitrate	14797-55-8	yes	yes	yes	yes	yes	yes	yes	yes
Nitrite	14797-65-0	no	yes	yes	yes	no	yes	yes	yes
Phosphate	14265-44-2	no	no	yes	yes	no	no	no	no
Phosphorus	7723-14-0	yes	no	yes	yes	yes	no	no	yes
Sulfate	14808-79-8	yes	no	yes	yes	yes	no	no	yes
Total Sulfur	63705-05-5	yes	no	yes	yes	yes	no	no	yes
Criteria Pollutants									
Carbon dioxide	124-38-9	yes	no	yes	yes	yes	no	no	yes
Nitrogen dioxide	10102-44-0	yes	no	yes	yes	yes	no	no	yes
Ozone	10028-15-6	yes	no	yes	yes	yes	no	no	yes
Particulate matter	no cas #	no	no***	yes	yes	no	yes	yes	yes
Sulfur dioxide	7446-09-5	yes	no	yes	yes	yes	no	no	yes

Table D-1. Selection of Human Health COPCs for Inclusion in the Quantitative PRA

		Are human health toxicity data available?		Can an exposure concentration be quantified?		Can human health risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Acute Toxicity? [a]	Chronic Toxicity? [b]	Emission Rate Available to Quantify Direct Exposure? [c]	Methods and Parameters Available to Quantify Indirect Exposure? [d]	Acute risk? [e]	Chronic Risk for Direct Exposure? [f]	Chronic Risk for Indirect Exposure? [g]	Is the constituent a quantitative COPC? [h]

[a] Acute inhalation exposure criteria appear in Table 4-4. If an acute inhalation exposure criteria value is available, a value of "yes" appears in this column.

[b] Chronic toxicity data (reference doses and slope factors) appear in Table 4-1 for organics and in Table 4-2 for inorganics. If a chronic toxicity value is available, a value of "yes" appears in this column.

[c] Is an emission rate estimate available to quantify direct exposure (i.e., air concentrations)? If so, then a value of "yes" appears in this column.

d Is an emission rate estimate available and are all chemical-specific parameter values available so that indirect exposures (e.g., soil, plant, animal, and water concentrations) can be modeled? If so, then a value of "yes" appears in this column.

(e.g., soil, plant, animal, and water concentrations) can be modeled? If so, then a value of "yes" appears in this column.

[e] Can acute inhalation risk be quantified? If acute toxicity values are available ("yes" appears in column 3) and direct exposures can be quantified ("yes" appears in column 5) then acute risk can be quantified and a value of "yes" appears in this column.

[f] Can chronic risk for direct exposure (i.e., from the inhalation of air pathway) be quantified? If chronic toxicity values are available ("yes" appears in column 4) and direct exposures can be quantified ("yes" appears in column 5) then chronic risk for direct exposure can be quantified and a value of "yes" appears in this column.

[g] Can chronic risk for indirect exposure (e.g., from exposures to soil, plants, animals, or water) be quantified? If chronic toxicity values are available ("yes" appears in column 4) and indirect exposures can be quantified ("yes" appears in column 6) then chronic risk for indirect exposure can be quantified and a value of "yes" appears in this column.

[h] The constituent is a quantitative COPC if acute inhalation can be quantified ("yes" appears in column 7) or chronic risk for direct exposure can be quantified ("yes" appears in column 8) or chronic risk for indirect exposure can be quantified ("yes" appears in column 9); in this case a value of "yes" appears in this column.

* Even though the COPC has no toxicity data, the COPC can be evaluated by using toxicity equivalency factors (TEFs).

** Even though the COPC has no toxicity data, the COPC can be evaluated by comparing its concentrations against USEPA Region 6 Health-Based Target Levels.

*** Even though the COPC has no toxicity data, the COPC can be evaluated by comparing its modeled annual average concentration to the average National Ambient Air Quality Standard (NAAQS) for PM_{10} of 50 ug/m^3.

Table D-2. Selection of Human Health ROPCs for Inclusion in the Quantitative PRA

		Are human health toxicity data available?	Can an exposure concentration be quantified?		Can human health risk be quantified?		
Constituent of Potential Concern	CAS Registry Number	Chronic Toxicity? [a]	Emission Rate Available to Quantify Direct Exposure? [b]	Methods and Parameters Available to Quantify Indirect Exposure? [c]	Chronic Risk for Direct Exposure? [d]	Chronic Risk for Indirect Exposure? [e]	Is the constituent a quantitative ROPC? [f]
Actinium-227	14952-40-0	yes	no	no	no	no	no
Americium-241	14596-10-2	yes	no	no	no	no	no
Americium-243	14993-75-0	yes	no	no	no	no	no
Antimony-125	14234-35-6	yes	no	no	no	no	no
Barium-137	13981-97-0	yes	yes	yes	yes	yes	yes
Cadmium-113	no cas #	yes	no	no	no	no	no
Carbon-14	14762-75-5	yes	yes	yes	yes	yes	yes
Cesium-134	13967-70-9	yes	yes	yes	yes	yes	yes
Cesium-137	10045-97-3	yes	no	no	no	no	no
Cobalt-60	10198-40-0	yes	yes	yes	yes	yes	yes
Curium-242	15510-73-3	yes	no	no	no	no	no
Curium-243	15757-87-6	yes	no	no	no	no	no
Curium-244	13981-15-2	yes	no	no	no	no	no
Europium-152	14683-23-9	yes	no	no	no	no	no
Europium-154	15585-10-1	yes	yes	yes	yes	yes	yes
Europium-155	14391-16-3	yes	yes	yes	yes	yes	yes
Iodine-129	15046-84-1	yes	yes	yes	yes	yes	yes
Neptunium-237	13994-20-2	yes	yes	yes	yes	yes	yes
Nickel-59	14336-70-0	yes	no	no	no	no	no
Nickel-63	13981-37-8	yes	no	no	no	no	no
Niobium-93	7440-03-1	yes	no	no	no	no	no
Plutonium-238	13981-16-3	yes	yes	yes	yes	yes	yes
Plutonium-239	15117-48-3	yes	yes	yes	yes	yes	yes
Plutonium-240	14119-33-6	yes	yes	yes	yes	yes	yes
Plutonium-241	14119-32-5	yes	yes	yes	yes	yes	yes
Plutonium-242	13982-10-0	yes	no	no	no	no	no

Table D-2. Selection of Human Health ROPCs for Inclusion in the Quantitative PRA

		Are human health toxicity data available?	Can an exposure concentration be quantified?		Can human health risk be quantified?		
Constituent of Potential Concern	CAS Registry Number	Chronic Toxicity? [a]	Emission Rate Available to Quantify Direct Exposure? [b]	Methods and Parameters Available to Quantify Indirect Exposure? [c]	Chronic Risk for Direct Exposure? [d]	Chronic Risk for Indirect Exposure? [e]	Is the constituent a quantitative ROPC? [f]
Protactinium-231	14331-85-2	yes	no	no	no	no	no
Radium-226	13982-63-3	yes	no	no	no	no	no
Radium-228	15262-20-1	yes	no	no	no	no	no
Ruthenium-106	13967-48-1	yes	no	no	no	no	no
Samarium-151	15715-94-3	yes	no	no	no	no	no
Selenium-79	no cas #	yes	no	no	no	no	no
Strontium-90	10098-97-2	yes	yes	yes	yes	yes	yes
Technetium-99	14133-76-7	yes	yes	yes	yes	yes	yes
Thorium-229	15594-54-4	yes	no	no	no	no	no
Thorium-232	7440-29-1	yes	no	no	no	no	no
Tin-126	15832-50-5	yes	no	no	no	no	no
Tritium	10028-17-8	yes	yes	yes	yes	yes	yes
Uranium-232	14158-29-3	yes	no	no	no	no	no
Uranium-233	13968-55-3	yes	no	no	no	no	no
Uranium-234	13966-29-5	yes	no	no	no	no	no
Uranium-235	15117-96-1	yes	yes	yes	yes	yes	yes
Uranium-236	13982-70-2	yes	yes	yes	yes	yes	yes
Uranium-238	7440-61-1	yes	no	no	no	no	no
Yttrium-90	10098-91-6	yes	yes	yes	yes	yes	yes
Zirconium-93	15751-77-6	yes	no	no	no	no	no

nd = not determined at this time

[a] Chronic toxicity data (slope factors) appear in Table 4-3 for radionuclides. If a chronic toxicity value is available, a value of "yes" appears in this column.

[b] Is an emission rate estimate available to quantify direct exposure (i.e., air concentrations)? If so, then a value of "yes"

Table D-2. Selection of Human Health ROPCs for Inclusion in the Quantitative PRA

		Are human health toxicity data available?	Can an exposure concentration be quantified?		Can human health risk be quantified?		
Constituent of Potential Concern	CAS Registry Number	Chronic Toxicity? [a]	Emission Rate Available to Quantify Direct Exposure? [b]	Methods and Parameters Available to Quantify Indirect Exposure? [c]	Chronic Risk for Direct Exposure? [d]	Chronic Risk for Indirect Exposure? [e]	Is the constituent a quantitative ROPC? [f]

appears in this column.

[c] Is an emission rate estimate available t and are all chemical-specific parameter values available so that indirect exposures (e.g., soil, plant, animal, and water concentrations) can be modeled? If so, then a value of "yes" appears in this column.

[d] Can chronic risk for direct exposure (i.e., from the inhalation of air and external exposure to air pathways) be quantified? If chronic toxicity values are available ("yes" appears in column 3) and direct exposures can be quantified ("yes" appears in column 4) then chronic risk for direct exposure can be quantified and a value of "yes" appears in this column.

[e] Can chronic risk for indirect exposure (e.g., from exposures to soil, plants, animals, or water) be quantified? If chronic toxicity values are available ("yes" appears in column 3) and indirect exposures can be quantified ("yes" appears in column 5) then chronic risk for indirect exposure can be quantified and a value of "yes" appears in this column.

[f] The constituent is a quantitative ROPC chronic risk for direct exposure can be quantified ("yes" appears in column 6) or chronic risk for indirect exposure can be quantified ("yes" appears in column 7); in this case a value of "yes" appears in this column.

Table D-3. Summary of Selection of COPCs and ROPCs for Inclusion in the Quantitative Human Health PRA

	Number that can be evaluated quantitatively	Number that cannot be evaluated quantitatively	Number that cannot yet be determined	Total
Organic COPCs	224	146	0	370
Inorganic COPCs	48	6	0	54
ROPCs	18	28	0	46
Total	290	180	0	470

Table D-4. Selection of Ecological COPCs for Inclusion in the Quantitative PRA

		Are there ecological toxicity data available?			concentration be quantified?		Can ecological risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Direct Contact?[a]	Ingestion?[b]	Inhalation?[c]	Emission Rate Available to Quantify Direct Exposure?[d]	Emission Rate Available to Quantify Indirect Exposure?[e]	Direct Contact?[f]	Ingestion?[g]	Inhalation?[h]	Is the constituent a quantitative COPC?[i]
Organics										
Aromatic Halogenated Hydrocarbons										
2,3,4,6-Tetrachlorophenol	58-90-2	yes	yes	no	yes	yes	yes	yes	no	yes
4-Chloro-3-methylphenol	59-50-7	yes	no	no	yes	yes	yes	no	no	yes
Aromatic Nonhalogenated Hydrocarbons										
2-Nitrotoluene	88-72-2	yes	no	no	yes	yes	yes	no	no	yes
4-Nitrobiphenyl	92-93-3	no	no	no	yes	yes	no	no	no	no
Benzaldehyde	100-52-7	yes	yes	yes	no	no	no	no	no	no
Benzene	71-43-2	yes	yes	yes	yes	yes	yes	yes	yes	yes
Benzyl alcohol	100-51-6	yes	no	no	no	no	no	no	no	no
Ethyl benzene	100-41-4	yes	yes	yes	yes	yes	yes	yes	yes	yes
m-Xylene	108-38-3	yes	no	yes	yes	yes	yes	no	yes	yes
o-Xylene	95-47-6	yes	no	yes	yes	yes	yes	no	yes	yes
p-Xylene	106-42-3	yes	no	yes	yes	yes	yes	no	yes	yes
Styrene	100-42-5	yes	yes	yes	yes	yes	yes	yes	yes	yes
Toluene	108-88-3	yes	yes	yes	yes	yes	yes	yes	yes	yes
Non-aromatic Nonhalogenated Hydrocarbons										
1,2-Epoxybutane	106-88-7	no	no	yes	yes	yes	no	no	yes	yes
1,3-Butadiene	106-99-0	no	no	yes	yes	yes	no	no	yes	yes
1,4-Dioxane	123-91-1	yes	yes	yes	yes	yes	yes	yes	yes	yes
1-Methylpropyl alcohol	78-92-2	no	no	yes	yes	yes	no	no	yes	yes
1-Nitropropane	108-03-2	no	no	no	yes	yes	no	no	no	no
2,2,4-Trimethylpentane	540-84-1	no	no	no	yes	yes	no	no	no	no
2-Butanone	78-93-3	yes	yes	yes	yes	yes	yes	yes	yes	yes
2-Butenaldehyde (2-Butenal)	4170-30-3	no	no	yes	yes	yes	no	no	yes	yes
2-Ethoxyethanol	110-80-5	no	no	yes	no	no	no	no	no	no

Table D-4. Selection of Ecological COPCs for Inclusion in the Quantitative PRA

		Are there ecological toxicity data available?			concentration be quantified?		Can ecological risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Direct Contact?[a]	Ingestion?[b]	Inhalation?[c]	Emission Rate Available to Quantify Direct Exposure?[d]	Emission Rate Available to Quantify Indirect Exposure?[e]	Direct Contact?[f]	Ingestion?[g]	Inhalation?[h]	Is the constituent a quantitative COPC?[i]
2-Heptanone	110-43-0	no	no	yes	yes	yes	no	no	yes	yes
2-Hexanone	591-78-6	yes	no	yes	yes	yes	yes	no	yes	yes
2-Methoxyethanol	109-86-4	no	no	yes	no	no	no	no	no	no
2-Methyl-2-propanol	75-65-0	no	no	yes	yes	yes	no	no	yes	yes
2-Methyl-2-propenenitrile	126-98-7	no	yes	yes	yes	yes	no	yes	yes	yes
2-Methylaziridine	75-55-8	no	no	no	yes	yes	no	no	no	no
2-Methylpropyl alcohol	78-83-1	no	yes	no	yes	yes	no	yes	no	yes
2-Pentanone	107-87-9	no	no	yes	yes	yes	no	no	yes	yes
2-Propanone (Acetone)	67-64-1	yes	yes	yes	yes	yes	yes	yes	yes	yes
2-Propene-1-ol	107-18-6	no	yes	no	yes	yes	no	yes	no	yes
2-Propyl alcohol	67-63-0	no	no	yes	yes	yes	no	no	yes	yes
3-Heptanone	106-35-4	no	no	yes	yes	yes	no	no	yes	yes
3-Methyl-1-butanol	123-51-3	no	no	no	yes	yes	no	no	no	no
3-Methyl-2-butanone	563-80-4	no	no	yes	yes	yes	no	no	yes	yes
3-Pentanone	96-22-0	no	no	yes	yes	yes	no	no	yes	yes
4-Heptanone	123-19-3	no	no	yes	yes	yes	no	no	yes	yes
4-Methyl-2-pentanone	108-10-1	yes	yes	yes	yes	yes	yes	yes	yes	yes
4-Methyl-3-penten-2-one	141-79-7	no	no	no	yes	yes	no	no	no	no
5-Methyl-2-hexanone	110-12-3	no	no	yes	yes	yes	no	no	yes	yes
Acetaldehyde	75-07-0	no	no	yes	yes	yes	no	no	yes	yes
Acetamide	60-35-5	no	no	no	yes	yes	no	no	no	no
Acetic acid	64-19-7	no	no	yes	yes	yes	no	no	yes	yes
Acetic acid ethyl ester	141-78-6	no	yes	yes	yes	yes	no	yes	yes	yes
Acetic acid n-butyl ester	123-86-4	no	no	yes	yes	yes	no	no	yes	yes
Acetonitrile	75-05-8	no	no	yes	yes	yes	no	no	yes	yes
Acrolein	107-02-8	no	no	yes	yes	yes	no	no	yes	yes
Acrylonitrile	107-13-1	yes	yes	yes	yes	yes	yes	yes	yes	yes
Bis(isopropyl)ether	108-20-3	no	no	no	yes	yes	no	no	no	no

Table D-4. Selection of Ecological COPCs for Inclusion in the Quantitative PRA

		Are there ecological toxicity data available?			concentration be quantified?		Can ecological risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Direct Contact?[a]	Ingestion?[b]	Inhalation?[c]	Emission Rate Available to Quantify Direct Exposure?[d]	Emission Rate Available to Quantify Indirect Exposure?[e]	Direct Contact?[f]	Ingestion?[g]	Inhalation?[h]	Is the constituent a quantitative COPC?[i]
Butane	106-97-8	no	no	yes	yes	yes	no	no	yes	yes
Carbon disulfide	75-15-0	yes	yes	yes	yes	yes	yes	yes	yes	yes
Cyanogen	460-19-5	no	no	yes	no	no	no	no	no	no
Cyclohexane	110-82-7	yes	no	yes	yes	yes	yes	no	yes	yes
Cyclohexanone	108-94-1	no	no	yes	yes	yes	no	no	yes	yes
Cyclohexene	110-83-8	no	no	yes	yes	yes	no	no	yes	yes
Cyclopentane	287-92-3	no	no	no	yes	yes	no	no	no	no
Ethyl alcohol	64-17-5	no	no	yes	yes	yes	no	no	yes	yes
Ethyl ether	60-29-7	no	yes	no	yes	yes	no	yes	no	yes
Ethyl methacrylate	97-63-2	no	no	no	no	no	no	no	no	no
Formaldehyde	50-00-0	yes	yes	yes	yes	yes	yes	yes	yes	yes
Formamide	75-12-7	no	no	no	yes	yes	no	no	no	no
Formic acid	64-18-6	no	no	yes	yes	yes	no	no	yes	yes
Formic acid, methyl ester	107-31-3	no	no	no	yes	yes	no	no	no	no
Glycidylaldehyde	765-34-4	no	yes	yes	no	no	no	no	no	no
Methyl acetate	79-20-9	no	no	no	yes	yes	no	no	no	no
Methyl alcohol	67-56-1	no	yes	yes	yes	yes	no	yes	yes	yes
Methyl isocyanate	624-83-9	no	no	yes	yes	yes	no	no	yes	yes
Methyl methacrylate	80-62-6	no	yes	yes	no	no	no	no	no	no
Methyl tert-butyl ether	1634-04-4	no	no	yes	yes	yes	no	no	yes	yes
Methylacetylene	74-99-7	no	no	no	yes	yes	no	no	no	no
Methylcyclohexane	108-87-2	no	no	yes	yes	yes	no	no	yes	yes
N,N-Dimethylacetamide	127-19-5	no	no	no	yes	yes	no	no	no	no
n-Butyl alcohol	71-36-3	no	yes	yes	yes	yes	no	yes	yes	yes
n-Heptane	142-82-5	no	no	yes	yes	yes	no	no	yes	yes
n-Hexane	110-54-3	yes	no	yes	yes	yes	yes	no	yes	yes
Nitromethane	75-52-5	no	no	no	yes	yes	no	no	no	no
n-nonane	111-84-2	no	no	yes	yes	yes	no	no	yes	yes

Table D-4. Selection of Ecological COPCs for Inclusion in the Quantitative PRA

		Are there ecological toxicity data available?			concentration be quantified?		Can ecological risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Direct Contact?[a]	Ingestion?[b]	Inhalation?[c]	Emission Rate Available to Quantify Direct Exposure?[d]	Emission Rate Available to Quantify Indirect Exposure?[e]	Direct Contact?[f]	Ingestion?[g]	Inhalation?[h]	Is the constituent a quantitative COPC?[i]
n-Octane	111-65-9	no	no	yes	yes	yes	no	no	yes	yes
n-Pentane	109-66-0	no	no	yes	yes	yes	no	no	yes	yes
n-Propionaldehyde	123-38-6	no	no	yes	yes	yes	no	no	yes	yes
n-Propyl alcohol	71-23-8	no	no	yes	yes	yes	no	no	yes	yes
n-Valeraldehyde	110-62-3	no	no	no	yes	yes	no	no	no	no
Oxirane	75-21-8	no	no	yes	yes	yes	no	no	yes	yes
p-Cymene	99-87-6	yes	no	no	no	no	no	no	no	no
Phosgene	75-44-5	no	no	yes	no	no	no	no	no	no
Propargyl alcohol	107-19-7	no	yes	yes	no	no	no	no	no	no
Propionic acid	79-09-4	no	no	no	yes	yes	no	no	no	no
Propionitrile	107-12-0	no	no	yes	yes	yes	no	no	yes	yes
Propylene gylcol monomethyl ether	107-98-2	no	no	yes	no	no	no	no	no	no
p-tert-Butyltoluene	98-51-1	yes	no	no	yes	yes	yes	no	no	yes
Triethylamine	121-44-8	no	no	yes	yes	yes	no	no	yes	yes
Trimethylamine	75-50-3	no	no	no	yes	yes	no	no	no	no
Vinyl acetate	108-05-4	yes	no	yes	yes	yes	yes	no	yes	yes
Non-aromatic Halogenated Hydrocarbons										
1,1,1,2-Tetrachloro-2,2-difluoroethane	76-11-9	yes	no	no	yes	yes	yes	no	no	yes
1,1,1,2-Tetrachloroethane	630-20-6	yes	yes	yes	yes	yes	yes	yes	yes	yes
1,1,1-Trichloroethane	71-55-6	yes	yes	yes	yes	yes	yes	yes	yes	yes
1,1,2,2-Tetrachloro-1,2-difluoroethane	76-12-0	yes	no	no	yes	yes	yes	no	no	yes
1,1,2,2-Tetrachloroethane	79-34-5	yes	no	yes	yes	yes	yes	no	yes	yes
1,1,2,2-Tetrachloroethene	127-18-4	yes	yes	yes	yes	yes	yes	yes	yes	yes
1,1,2-Trichloroethane	79-00-5	yes	no	yes	yes	yes	yes	no	yes	yes
1,1,2-Trichloroethylene	79-01-6	yes	yes	yes	yes	yes	yes	yes	yes	yes
1,1-Dichloroethane	75-34-3	yes	no	yes	yes	yes	yes	no	yes	yes
1,1-Dichloroethene	75-35-4	yes	yes	yes	yes	yes	yes	yes	yes	yes

Table D-4. Selection of Ecological COPCs for Inclusion in the Quantitative PRA

		Are there ecological toxicity data available?			concentration be quantified?		Can ecological risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Direct Contact?[a]	Ingestion?[b]	Inhalation?[c]	Emission Rate Available to Quantify Direct Exposure?[d]	Emission Rate Available to Quantify Indirect Exposure?[e]	Direct Contact?[f]	Ingestion?[g]	Inhalation?[h]	Is the constituent a quantitative COPC?[i]
1,2,2-Trichloro-1,1,2-trifluoroethane	76-13-1	yes	no	yes	yes	yes	yes	no	yes	yes
1,2,3-Trichloropropane	96-18-4	yes	yes	yes	yes	yes	yes	yes	yes	yes
1,2-Dibromo-3-chloropropane	96-12-8	yes	no	yes	no	no	no	no	no	no
1,2-Dichloro-1,1,2,2-tetrafluoroethane	76-14-2	yes	no	yes	yes	yes	yes	no	yes	yes
1,2-Dichloroethane	107-06-2	yes	yes	yes	yes	yes	yes	yes	yes	yes
1,2-Dichloroethylene	540-59-0	yes	yes	no	yes	yes	yes	yes	no	yes
1,2-Dichloropropane	78-87-5	yes	no	yes	yes	yes	yes	no	yes	yes
1,3-Dichloropropene	542-75-6	yes	yes	yes	no	no	no	no	no	no
1,4-Dichloro-2-butene	764-41-0	yes	no	yes	no	no	no	no	no	no
1-Chloroethene	75-01-4	yes	no	yes	yes	yes	yes	no	yes	yes
2,2-Dichloropropionic acid	75-99-0	yes	no	no	yes	yes	yes	no	no	yes
2-Chloropropane	75-29-6	yes	no	yes	no	no	no	no	no	no
3-Chloropropene (Allyl chloride)	107-05-1	yes	no	yes	yes	yes	yes	no	yes	yes
Bromochloromethane	74-97-5	yes	no	yes	yes	yes	yes	no	yes	yes
Bromodichloromethane	75-27-4	yes	yes	no	yes	yes	yes	yes	no	yes
Bromoethene	593-60-2	no	no	yes	no	no	no	no	no	no
Bromoform	75-25-2	no	yes	yes	no	no	no	no	no	no
Bromomethane	74-83-9	no	yes	yes	yes	yes	no	yes	yes	yes
Carbon tetrachloride	56-23-5	yes	yes	yes	yes	yes	yes	yes	yes	yes
Chlorodibromomethane	124-48-1	yes	yes	no	no	no	no	no	no	no
Chlorodifluoromethane	75-45-6	yes	no	yes	yes	yes	yes	no	yes	yes
Chloroethane	75-00-3	yes	no	yes	yes	yes	yes	no	yes	yes
Chloroform	67-66-3	yes	yes	yes	yes	yes	yes	yes	yes	yes
Chloromethane	74-87-3	yes	no	yes	yes	yes	yes	no	yes	yes
Chloropentafluoroethane	76-15-3	yes	no	no	yes	yes	yes	no	no	yes
cis-1,2-Dichloroethene	156-59-2	yes	no	no	no	no	no	no	no	no
cis-1,3-Dichloropropene	10061-01-5	yes	no	no	yes	yes	yes	no	no	yes
Cyanogen bromide	506-68-3	no	yes	yes	no	no	no	no	no	no

Table D-4. Selection of Ecological COPCs for Inclusion in the Quantitative PRA

		Are there ecological toxicity data available?			concentration be quantified?		Can ecological risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Direct Contact?[a]	Ingestion?[b]	Inhalation?[c]	Emission Rate Available to Quantify Direct Exposure?[d]	Emission Rate Available to Quantify Indirect Exposure?[e]	Direct Contact?[f]	Ingestion?[g]	Inhalation?[h]	Is the constituent a quantitative COPC?[i]
Cyanogen chloride	506-77-4	yes	yes	no	no	no	no	no	no	no
Dichlorodifluoromethane	75-71-8	yes	yes	yes	yes	yes	yes	yes	yes	yes
Dichlorofluoromethane	75-43-4	yes	no	yes	yes	yes	yes	no	yes	yes
Dichloromethane	75-09-2	yes	yes	yes	yes	yes	yes	yes	yes	yes
Difluorodibromomethane	75-61-6	no	no	no	yes	yes	no	no	no	no
Hexafluoroacetone	684-16-2	no	no	yes	yes	yes	no	no	yes	yes
Iodomethane	74-88-4	no	no	no	no	no	no	no	no	no
Methylene bromide	74-95-3	no	no	yes	no	no	no	no	no	no
Pentachloroethane	76-01-7	yes	no	no	no	no	no	no	no	no
trans-1,2-Dichloroethylene	156-60-5	yes	yes	yes	yes	yes	yes	yes	yes	yes
trans-1,3-Dichloropropene	10061-02-6	yes	no	no	yes	yes	yes	no	no	yes
Trichloroacetic acid	76-03-9	yes	no	no	yes	yes	yes	no	no	yes
Trichlorofluoroethane	27154-33-2	yes	no	no	yes	yes	yes	no	no	yes
Trichlorofluoromethane	75-69-4	yes	yes	yes	yes	yes	yes	yes	yes	yes
Trifluorobromomethane	75-63-8	no	no	no	yes	yes	no	no	no	no
Dioxin and Furan Compounds										
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	35822-46-9	yes	no	no	yes	yes	yes	no	no	yes
1,2,3,4,6,7,8-Heptachlorodibenzofuran	67562-39-4	yes	no	no	yes	yes	yes	no	no	yes
1,2,3,4,7,8,9-Heptachlorodibenzofuran	55673-89-7	yes	no	no	yes	yes	yes	no	no	yes
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	39227-28-6	yes	no	no	yes	yes	yes	no	no	yes
1,2,3,4,7,8-Hexachlorodibenzofuran	70648-26-9	yes	no	no	yes	yes	yes	no	no	yes
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	57653-85-7	yes	no	no	yes	yes	yes	no	no	yes
1,2,3,6,7,8-Hexachlorodibenzofuran	57117-44-9	yes	yes	no	yes	yes	yes	yes	no	yes
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	19408-74-3	yes	no	no	yes	yes	yes	no	no	yes
1,2,3,7,8,9-Hexachlorodibenzofuran	72918-21-9	yes	no	no	yes	yes	yes	no	no	yes
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	40321-76-4	yes	no	no	yes	yes	yes	no	no	yes
1,2,3,7,8-Pentachlorodibenzofuran	57117-41-6	yes	yes	no	yes	yes	yes	yes	no	yes

Table D-4. Selection of Ecological COPCs for Inclusion in the Quantitative PRA

		Are there ecological toxicity data available?			concentration be quantified?		Can ecological risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Direct Contact?[a]	Ingestion?[b]	Inhalation?[c]	Emission Rate Available to Quantify Direct Exposure?[d]	Emission Rate Available to Quantify Indirect Exposure?[e]	Direct Contact?[f]	Ingestion?[g]	Inhalation?[h]	Is the constituent a quantitative COPC?[i]
2,3,4,6,7,8-Hexachlorodibenzofuran	60851-34-5	yes	no	no	yes	yes	yes	no	no	yes
2,3,4,7,8-Pentachlorodibenzofuran	57117-31-4	yes	yes	no	yes	yes	yes	yes	no	yes
2,3,7,8-Tetrachlorodibenzo(p)dioxin	1746-01-6	yes	yes	no	yes	yes	yes	yes	no	yes
2,3,7,8-Tetrachlorodibenzofuran	51207-31-9	yes	yes	no	yes	yes	yes	yes	no	yes
Dibenzofuran	132-64-9	yes	no	no	yes	yes	yes	no	no	yes
Octachlorodibenzo(p)dioxin	3268-87-9	yes	no	no	yes	yes	yes	no	no	yes
Octachlorodibenzofuran	39001-02-0	yes	no	no	yes	yes	yes	no	no	yes
PCBs										
2,2',3,3',4,4',5-Heptachlorobiphenyl	35065-30-6	yes	no	no	yes	yes	yes	no	no	yes
2,2',3,4,4',5,5'-Heptachlorobiphenyl	35065-29-3	yes	no	no	yes	yes	yes	no	no	yes
2,3,3',4,4',5'-Hexachlorobiphenyl	69782-90-7	yes	yes	no	yes	yes	yes	yes	no	yes
2,3,3',4,4',5-Hexachlorobiphenyl	38380-08-4	yes	yes	no	yes	yes	yes	yes	no	yes
2,3,3',4,4',5,5'-Heptachlorobiphenyl	no cas #	yes	no	no	yes	yes	yes	no	no	yes
2,3,3',4,4'-Pentachlorobiphenyl	32598-14-4	yes	no	no	yes	yes	yes	no	no	yes
2,3,4,4',5-Pentachlorobiphenyl	74472-37-0	yes	no	no	yes	yes	yes	no	no	yes
2',3,4,4',5-Pentachlorobiphenyl	no cas #	yes	no	no	yes	yes	yes	no	no	yes
2,3',4,4',5-Pentachlorobiphenyl	31508-00-6	yes	no	no	yes	yes	yes	no	no	yes
2,3',4,4',5,5'-Hexachlorobiphenyl	no cas #	yes	yes	no	yes	yes	yes	yes	no	yes
3,3',4,4',5-Pentachlorobiphenyl	no cas #	yes	no	no	yes	yes	yes	no	no	yes
3,3',4,4',5,5'-Hexachlorobiphenyl	32774-16-6	yes	yes	no	yes	yes	yes	yes	no	yes
3,3',4,4'-Tetrachlorobiphenyl	32598-13-3	yes	no	no	yes	yes	yes	no	no	yes
3,4,4',5-Tetrachlorobiphenyl	70362-50-4	yes	no	no	yes	yes	yes	no	no	yes
Polychlorinated biphenyls (PCBs)	1336-36-3	yes	yes	yes	yes	yes	yes	yes	yes	yes
Phthalates										
Bis(2-ethylhexyl)phthalate (DEHP)	117-81-7	yes	yes	yes	yes	yes	yes	yes	yes	yes
Butylbenzyl phthalate	85-68-7	yes	no	yes	yes	yes	yes	no	yes	yes

Table D-4. Selection of Ecological COPCs for Inclusion in the Quantitative PRA

		Are there ecological toxicity data available?			concentration be quantified?		Can ecological risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Direct Contact?[a]	Ingestion?[b]	Inhalation?[c]	Emission Rate Available to Quantify Direct Exposure?[d]	Emission Rate Available to Quantify Indirect Exposure?[e]	Direct Contact?[f]	Ingestion?[g]	Inhalation?[h]	Is the constituent a quantitative COPC?[i]
Di-n-butylphthalate	84-74-2	yes	yes	yes	yes	yes	yes	yes	yes	yes
Diethyl phthalate	84-66-2	yes	yes	no	yes	yes	yes	yes	no	yes
Dimethylphthalate	131-11-3	yes	no	yes	no	no	no	no	no	no
n-Dioctyl phthalate	117-84-0	yes	yes	no	yes	yes	yes	yes	no	yes
Light Polycyclic Aromatic Hydrocarbons										
2-Chloronapthalene	91-58-7	yes	yes	no	yes	yes	yes	yes	no	yes
2-Methyl naphthalene	91-57-6	yes	no	no	no	no	no	no	no	no
5-Nitroacenaphthene	602-87-9	yes	no	no	yes	yes	yes	no	no	yes
Acenaphthene	83-32-9	yes	yes	no	yes	yes	yes	yes	no	yes
Acenaphthylene	208-96-8	yes	no	no	yes	yes	yes	no	no	yes
Anthracene	120-12-7	yes	yes	no	yes	yes	yes	yes	no	yes
Fluorene	86-73-7	yes	yes	no	yes	yes	yes	yes	no	yes
Indene	95-13-6	yes	no	no	yes	yes	yes	no	no	yes
Naphthalene	91-20-3	yes	yes	yes	yes	yes	yes	yes	yes	yes
Phenanthrene	85-01-8	yes	no	no	yes	yes	yes	no	no	yes
Pyrene	129-00-0	yes	yes	no	yes	yes	yes	yes	no	yes
Heavy Polycyclic Aromatic Hydrocarbons										
3-Methylcholanthrene	56-49-5	yes	no	no	yes	yes	yes	no	no	yes
5-Methylchrysene	3697-24-3	yes	no	no	yes	yes	yes	no	no	yes
Benzo(a)anthracene	56-55-3	yes	yes	no	yes	yes	yes	yes	no	yes
Benzo(a)pyrene	50-32-8	yes	yes	yes	yes	yes	yes	yes	yes	yes
Benzo(b)fluoranthene	205-99-2	yes	yes	no	yes	yes	yes	yes	no	yes
Benzo(e)pyrene	192-97-2	yes	no	no	no	no	no	no	no	no
Benzo(g,h,i)perylene	191-24-2	yes	no	no	yes	yes	yes	no	no	yes
Benzo(j)fluoranthene	205-82-3	yes	no	no	yes	yes	yes	no	no	yes
Benzo(k)fluoranthene	207-08-9	yes	yes	no	yes	yes	yes	yes	no	yes

Table D-4. Selection of Ecological COPCs for Inclusion in the Quantitative PRA

		Are there ecological toxicity data available?			concentration be quantified?		Can ecological risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Direct Contact?[a]	Ingestion?[b]	Inhalation?[c]	Emission Rate Available to Quantify Direct Exposure?[d]	Emission Rate Available to Quantify Indirect Exposure?[e]	Direct Contact?[f]	Ingestion?[g]	Inhalation?[h]	Is the constituent a quantitative COPC?[i]
Benzo[a,i]pyrene	191-30-0	yes	no	no	yes	yes	yes	no	no	yes
Chrysene	218-01-9	yes	yes	no	yes	yes	yes	yes	no	yes
Dibenz(a,h)anthracene	53-70-3	yes	yes	no	yes	yes	yes	yes	no	yes
Dibenz[a,h]acridine	226-36-8	yes	no	no	yes	yes	yes	no	no	yes
Dibenz[a,j]acridine	224-42-0	yes	no	no	yes	yes	yes	no	no	yes
Dibenzo(a,e)fluoranthene	5385-75-1	yes	no	no	no	no	no	no	no	no
Dibenzo(a,h)fluoranthene	no cas #	yes	no	no	no	no	no	no	no	no
Dibenzo[a,e]pyrene	192-65-4	yes	no	no	yes	yes	yes	no	no	yes
Dibenzo[a,h]pyrene	189-64-0	yes	no	no	yes	yes	yes	no	no	yes
Dibenzo[a,i]pyrene	189-55-9	yes	no	no	yes	yes	yes	no	no	yes
Fluoranthene	206-44-0	yes	yes	no	yes	yes	yes	yes	no	yes
Hexachloronaphthalene	1335-87-1	yes	no	no	yes	yes	yes	no	no	yes
Indeno(1,2,3-cd)pyrene	193-39-5	yes	yes	no	yes	yes	yes	yes	no	yes
Octachloronaphthalene	2234-13-1	yes	no	no	yes	yes	yes	no	no	yes
Pentachloronaphthalene	1321-64-8	yes	no	no	yes	yes	yes	no	no	yes
Tetrachloronaphthalene	1335-88-2	yes	no	no	yes	yes	yes	no	no	yes
Trichloronaphthalene	1321-65-9	yes	no	no	yes	yes	yes	no	no	yes
Light Substituted Benzene Compounds										
1,2,3-Trichlorobenzene	87-61-6	yes	no	no	no	no	no	no	no	no
1,2,4-Trichlorobenzene	120-82-1	yes	no	no	yes	yes	yes	no	no	yes
1,2,4-Trimethyl benzene	95-63-6	yes	no	no	no	no	no	no	no	no
1,2-Dichlorobenzene	95-50-1	yes	yes	yes	yes	yes	yes	yes	yes	yes
1,3,5-Trimethyl benzene	108-67-8	yes	no	no	no	no	no	no	no	no
1,3-Dichlorobenzene	541-73-1	yes	no	no	yes	yes	yes	no	no	yes
1,3-Dinitrobenzene	99-65-0	yes	yes	no	no	no	no	no	no	no
1,4-Dichlorobenzene	106-46-7	yes	no	yes	yes	yes	yes	no	yes	yes
1,4-Dinitrobenzene	100-25-4	yes	no	no	yes	yes	yes	no	no	yes

Table D-4. Selection of Ecological COPCs for Inclusion in the Quantitative PRA

		Are there ecological toxicity data available?			concentration be quantified?		Can ecological risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Direct Contact?[a]	Ingestion?[b]	Inhalation?[c]	Emission Rate Available to Quantify Direct Exposure?[d]	Emission Rate Available to Quantify Indirect Exposure?[e]	Direct Contact?[f]	Ingestion?[g]	Inhalation?[h]	Is the constituent a quantitative COPC?[i]
2,4,5-Trichlorophenol	95-95-4	yes	yes	no	yes	yes	yes	yes	no	yes
2,4,6-Trichlorophenol	88-06-2	yes	no	no	yes	yes	yes	no	no	yes
2,4-Dichlorophenol	120-83-2	yes	yes	no	yes	yes	yes	yes	no	yes
2,4-Dimethylphenol	105-67-9	yes	yes	yes	no	no	no	no	no	no
2,4-Dinitrophenol	51-28-5	yes	yes	yes	no	no	no	no	no	no
2,4-Dinitrotoluene	121-14-2	yes	yes	no	no	no	no	no	no	no
2,6-Dinitrotoluene	606-20-2	yes	yes	yes	no	no	no	no	no	no
2-Chlorophenol	95-57-8	yes	no	yes	yes	yes	yes	no	yes	yes
2-Chlorotoluene	95-49-8	yes	yes	no	yes	yes	yes	yes	no	yes
2-Nitrophenol	88-75-5	yes	no	no	yes	yes	yes	no	no	yes
4,6-Dinitro-o-cresol	534-52-1	yes	no	yes	no	no	no	no	no	no
4-Chlorotoluene	106-43-4	yes	no	no	no	no	no	no	no	no
4-Nitrophenol	100-02-7	yes	no	no	no	no	no	no	no	no
alpha-Methylstyrene	98-83-9	yes	no	no	yes	yes	yes	no	no	yes
Aniline	62-53-3	yes	no	yes	no	no	no	no	no	no
Benzotrichloride	98-07-7	yes	no	yes	no	no	no	no	no	no
Benzyl chloride	100-44-7	yes	no	yes	no	no	no	no	no	no
Bromobenzene	108-86-1	yes	no	no	no	no	no	no	no	no
Chlorobenzene	108-90-7	yes	yes	yes	yes	yes	yes	yes	yes	yes
Cumene	98-82-8	yes	yes	yes	yes	yes	yes	yes	yes	yes
m-Cresol	108-39-4	yes	yes	yes	yes	yes	yes	yes	yes	yes
n-Butyl benzene	104-51-8	yes	no	no	no	no	no	no	no	no
Nitrobenzene	98-95-3	yes	yes	yes	yes	yes	yes	yes	yes	yes
n-Propyl benzene	103-65-1	yes	no	no	no	no	no	no	no	no
o-Cresol	95-48-7	yes	yes	yes	yes	yes	yes	yes	yes	yes
o-Dinitrobenzene	528-29-0	yes	no	no	no	no	no	no	no	no
o-Nitroaniline	88-74-4	yes	no	no	no	no	no	no	no	no
o-Toluidine	95-53-4	yes	no	yes	no	no	no	no	no	no

Table D-4. Selection of Ecological COPCs for Inclusion in the Quantitative PRA

		Are there ecological toxicity data available?			concentration be quantified?		Can ecological risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Direct Contact?[a]	Ingestion?[b]	Inhalation?[c]	Emission Rate Available to Quantify Direct Exposure?[d]	Emission Rate Available to Quantify Indirect Exposure?[e]	Direct Contact?[f]	Ingestion?[g]	Inhalation?[h]	Is the constituent a quantitative COPC?[i]
p-Chloroaniline	106-47-8	yes	yes	yes	no	no	no	no	no	no
p-Cresol	106-44-5	yes	no	yes	no	no	no	no	no	no
Phenol	108-95-2	yes	yes	yes	yes	yes	yes	yes	yes	yes
p-Nitrochlorobenzene	100-00-5	yes	no	yes	yes	yes	yes	no	yes	yes
p-Toluidine	106-49-0	yes	no	yes	no	no	no	no	no	no
sec-Butyl benzene	135-98-8	yes	no	no	no	no	no	no	no	no
tert-Butyl benzene	98-06-6	yes	no	no	no	no	no	no	no	no
Toluene-2,6-diamine	823-40-5	yes	no	no	no	no	no	no	no	no
Trimethyl benzene	25551-13-7	yes	no	no	yes	yes	yes	no	no	yes
Other Light Semivolatile Compounds										
1,1`-Biphenyl	92-52-4	yes	yes	yes	yes	yes	yes	yes	yes	yes
1,1-Dimethylhydrazine	57-14-7	no	no	yes	yes	yes	no	no	yes	yes
1,2-Dimethylhydrazine	540-73-8	no	no	no	no	no	no	no	no	no
1,2-Diphenylhydrazine	122-66-7	no	no	no	no	no	no	no	no	no
1,3-Propane sultone	1120-71-4	no	no	yes	no	no	no	no	no	no
2,4-Toluene diisocyante	584-84-9	no	no	yes	no	no	no	no	no	no
2-Chloroacetophenone	532-27-4	yes	no	yes	no	no	no	no	no	no
2-Propenoic acid	79-10-7	no	yes	yes	yes	yes	no	yes	yes	yes
4,4-Methylenedianiline	101-77-9	no	no	no	no	no	no	no	no	no
Acetophenone	98-86-2	no	yes	yes	yes	yes	no	yes	yes	yes
Benzoic acid	65-85-0	yes	yes	no	no	no	no	no	no	no
Bis(2-chloroethoxy)methane	111-91-1	yes	no	yes	no	no	no	no	no	no
Bis(2-chloroethyl) ether	111-44-4	yes	no	yes	no	no	no	no	no	no
Chlorocyclopentadiene	41851-50-7	yes	no	no	no	no	no	no	no	no
Cyclohexanol	108-93-0	no	no	no	yes	yes	no	no	no	no
Dichloroisopropyl ether	108-60-1	yes	no	yes	no	no	no	no	no	no
Dichloromethyl ether	542-88-1	yes	no	yes	yes	yes	yes	no	yes	yes

Table D-4. Selection of Ecological COPCs for Inclusion in the Quantitative PRA

		Are there ecological toxicity data available?			concentration be quantified?		Can ecological risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Direct Contact?[a]	Ingestion?[b]	Inhalation?[c]	Emission Rate Available to Quantify Direct Exposure?[d]	Emission Rate Available to Quantify Indirect Exposure?[e]	Direct Contact?[f]	Ingestion?[g]	Inhalation?[h]	Is the constituent a quantitative COPC?[i]
Dichloropentadiene	no cas #	yes	no	no	no	no	no	no	no	no
Dimethyl sulfate	77-78-1	no	no	yes	no	no	no	no	no	no
Dimethylaniline	121-69-7	no	yes	no	yes	yes	no	yes	no	yes
di-n-Propylnitrosamine	621-64-7	no	no	no	yes	yes	no	no	no	no
Diphenyl ether	101-84-8	no	no	no	yes	yes	no	no	no	no
Epichlorohydrin	106-89-8	yes	no	yes	no	no	no	no	no	no
Ethyl Carbamate (Urethane)	51-79-6	no	no	yes	no	no	no	no	no	no
Ethyl methanesulfonate	62-50-0	no	no	no	no	no	no	no	no	no
Ethylene dibromide	106-93-4	no	no	yes	yes	yes	no	no	yes	yes
Ethylene glycol	107-21-1	no	yes	no	no	no	no	no	no	no
Ethylene glycol monobutyl ether	111-76-2	no	yes	yes	yes	yes	no	yes	yes	yes
Ethylene glycol monoethyl ether acetate	111-15-9	no	no	yes	no	no	no	no	no	no
Ethylene thiourea	96-45-7	no	yes	yes	no	no	no	no	no	no
Furfural	98-01-1	no	yes	yes	no	no	no	no	no	no
Maleic hydrazide	123-33-1	no	yes	yes	no	no	no	no	no	no
Malononitrile	109-77-3	no	no	no	no	no	no	no	no	no
Methyl styrene (mixed isomers)	25013-15-4	yes	no	yes	no	no	no	no	no	no
Methylhydrazine	60-34-4	no	no	yes	yes	yes	no	no	yes	yes
N,N-Diphenylamine	122-39-4	no	yes	no	yes	yes	no	yes	no	yes
Nitric acid, propyl ester	627-13-4	no	no	yes	yes	yes	no	no	yes	yes
N-Nitrosodi-n-butylamine	924-16-3	no	no	no	no	no	no	no	no	no
N-Nitrosomorpholine	59-89-2	no	no	no	yes	yes	no	no	no	no
N-Nitroso-N,N-dimethylamine	62-75-9	no	no	yes	yes	yes	no	no	yes	yes
o-Anisidine	90-04-0	no	no	no	no	no	no	no	no	no
Oxalic acid	144-62-7	no	no	no	yes	yes	no	no	no	no
Phthalic anhydride	85-44-9	no	yes	yes	no	no	no	no	no	no
p-Phthalic acid	100-21-0	no	no	no	yes	yes	no	no	no	no
Pyridine	110-86-1	yes	yes	yes	yes	yes	yes	yes	yes	yes

Table D-4. Selection of Ecological COPCs for Inclusion in the Quantitative PRA

		Are there ecological toxicity data available?			concentration be quantified?		Can ecological risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Direct Contact?[a]	Ingestion?[b]	Inhalation?[c]	Emission Rate Available to Quantify Direct Exposure?[d]	Emission Rate Available to Quantify Indirect Exposure?[e]	Direct Contact?[f]	Ingestion?[g]	Inhalation?[h]	Is the constituent a quantitative COPC?[i]
Quinoline	91-22-5	no	no	no	yes	yes	no	no	no	no
Quinone	106-51-4	no	no	no	no	no	no	no	no	no
Safrole	94-59-7	no	no	no	no	no	no	no	no	no
Tetrahydrofuran	109-99-9	yes	no	yes	yes	yes	yes	no	yes	yes
Other Heavy Semivolatile Compounds										
1,2,4,5-Tetrachlorobenzene	95-94-3	no	yes	no	no	no	no	no	no	no
1,3,5-Trinitrobenzene	99-35-4	no	yes	no	no	no	no	no	no	no
2,6-Bis(tert-butyl)-4-methylphenol	128-37-0	yes	no	no	yes	yes	yes	no	no	yes
2-Cyclohexyl-4,6-dinitrophenol	131-89-5	yes	yes	no	no	no	no	no	no	no
2-sec-Butyl-4,6-dinitrophenol	88-85-7	yes	no	yes	yes	yes	yes	no	yes	yes
3,3-Dichlorobenzidine	91-94-1	no	no	no	no	no	no	no	no	no
3,3'-Dimethoxybenzidine	119-90-4	no	no	no	no	no	no	no	no	no
4-Bromophenylphenyl ether	101-55-3	yes	no	no	yes	yes	yes	no	no	yes
Ammonium perfluorooctanoate	3825-26-1	no	no	yes	yes	yes	no	no	yes	yes
Azobenzene	103-33-3	no	no	no	no	no	no	no	no	no
Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulfide	96-69-5	no	no	no	yes	yes	no	no	no	no
Captan	133-06-2	yes	no	no	no	no	no	no	no	no
Chlorobenzilate	510-15-6	no	no	no	no	no	no	no	no	no
Dibutylphosphate	107-66-4	no	no	no	yes	yes	no	no	no	no
Dimethyl aminoazobenzene	60-11-7	no	no	no	no	no	no	no	no	no
Hexachlorobenzene	118-74-1	yes	yes	yes	yes	yes	yes	yes	yes	yes
Hexachlorobutadiene	87-68-3	yes	yes	yes	yes	yes	yes	yes	yes	yes
Hexachlorocyclopentadiene	77-47-4	yes	yes	no	no	no	no	no	no	no
Hexachloroethane	67-72-1	yes	yes	yes	yes	yes	yes	yes	yes	yes
Hexachlorophene	70-30-4	yes	yes	yes	no	no	no	no	no	no
Hexamethylene-1,5-diisocyanate	822-06-0	no	no	yes	no	no	no	no	no	no
Mirex	2385-85-5	yes	yes	no	yes	yes	yes	yes	no	yes

Table D-4. Selection of Ecological COPCs for Inclusion in the Quantitative PRA

		Are there ecological toxicity data available?			concentration be quantified?		Can ecological risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Direct Contact?[a]	Ingestion?[b]	Inhalation?[c]	Emission Rate Available to Quantify Direct Exposure?[d]	Emission Rate Available to Quantify Indirect Exposure?[e]	Direct Contact?[f]	Ingestion?[g]	Inhalation?[h]	Is the constituent a quantitative COPC?[i]
Nitrofen	1836-75-5	yes	no	no	yes	yes	yes	no	no	yes
Pentachlorobenzene	608-93-5	yes	yes	no	no	no	no	no	no	no
Pentachloronitrobenzene	82-68-8	yes	yes	yes	yes	yes	yes	yes	yes	yes
Pentachlorophenol	87-86-5	yes	yes	yes	yes	yes	yes	yes	yes	yes
Picric acid	88-89-1	no	no	no	yes	yes	no	no	no	no
Pronamide	23950-58-5	yes	yes	no	no	no	no	no	no	no
Strychnine	57-24-9	no	yes	no	no	no	no	no	no	no
Terphenyls	26140-60-3	no	no	no	yes	yes	no	no	no	no
Tributyl phosphate	126-73-8	no	no	yes	yes	yes	no	no	yes	yes
Trifluralin	1582-09-8	no	yes	no	yes	yes	no	yes	no	yes
Triphenylamine	603-34-9	no	no	no	yes	yes	no	no	no	no
Herbicides and Organochlorinated Pesticides										
2,4,5-T	93-76-5	yes	yes	no	yes	yes	yes	yes	no	yes
2,4-D and esters	94-75-7	yes	no	no	yes	yes	yes	no	no	yes
4,4'-DDD	72-54-8	yes	no	no	yes	yes	yes	no	no	yes
4,4'-DDE	72-55-9	yes	yes	no	yes	yes	yes	yes	no	yes
4,4'-DDT	50-29-3	yes	yes	no	yes	yes	yes	yes	no	yes
Aldrin	309-00-2	yes	yes	yes	yes	yes	yes	yes	yes	yes
alpha-BHC	319-84-6	yes	no	no	yes	yes	yes	no	no	yes
beta-BHC	319-85-7	yes	no	no	yes	yes	yes	no	no	yes
Chlordane	57-74-9	yes	yes	yes	no	no	no	no	no	no
Delta-BHC	319-86-8	yes	no	no	yes	yes	yes	no	no	yes
Dieldrin	60-57-1	yes	yes	yes	yes	yes	yes	yes	yes	yes
Endothall	145-73-3	no	yes	no	no	no	no	no	no	no
Endrin	72-20-8	yes	yes	yes	yes	yes	yes	yes	yes	yes
gamma-BHC (Lindane)	58-89-9	yes	yes	yes	yes	yes	yes	yes	yes	yes
Heptachlor	76-44-8	yes	yes	yes	yes	yes	yes	yes	yes	yes

Table D-4. Selection of Ecological COPCs for Inclusion in the Quantitative PRA

		Are there ecological toxicity data available?			concentration be quantified?		Can ecological risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Direct Contact?[a]	Ingestion?[b]	Inhalation?[c]	Emission Rate Available to Quantify Direct Exposure?[d]	Emission Rate Available to Quantify Indirect Exposure?[e]	Direct Contact?[f]	Ingestion?[g]	Inhalation?[h]	Is the constituent a quantitative COPC?[i]
Isodrin	465-73-6	yes	no	no	yes	yes	yes	no	no	yes
Methoxychlor	72-43-5	yes	yes	no	yes	yes	yes	yes	no	yes
Silvex (2,4,5-TP)	93-72-1	yes	yes	no	yes	yes	yes	yes	no	yes
Toxaphene	8001-35-2	yes	yes	yes	yes	yes	yes	yes	yes	yes
Inorganics										
Metals										
Aluminum	7429-90-5	yes	yes	no	yes	yes	yes	yes	no	yes
Antimony	7440-36-0	yes	yes	yes	yes	yes	yes	yes	yes	yes
Arsenic	7440-38-2	yes	yes	no	yes	yes	yes	yes	no	yes
Barium	7440-39-3	yes	yes	no	yes	yes	yes	yes	no	yes
Beryllium	7440-41-7	yes	yes	no	yes	yes	yes	yes	no	yes
Bismuth	7440-69-9	no	no	no	yes	yes	no	no	no	no
Boron	7440-42-8	yes	yes	yes	yes	yes	yes	yes	yes	yes
Cadmium	7440-43-9	yes	yes	yes	yes	yes	yes	yes	yes	yes
Calcium	7440-70-2	no	no	no	yes	yes	no	no	no	no
Chromium (and VI)	18540-29-9	yes	yes	no	yes	yes	yes	yes	no	yes
Cobalt	7440-48-4	yes	yes	yes	yes	yes	yes	yes	yes	yes
Copper	7440-50-8	yes	yes	no	yes	yes	yes	yes	no	yes
Iron	7439-89-6	yes	no	yes	yes	yes	yes	no	yes	yes
Lead	7439-92-1	yes	yes	yes	yes	yes	yes	yes	yes	yes
Lithium	7439-93-2	yes	yes	no	yes	yes	yes	yes	no	yes
Magnesium	7439-95-4	no	no	no	yes	yes	no	no	no	no
Manganese	7439-96-5	yes	yes	no	yes	yes	yes	yes	no	yes
Mercury	7439-97-6	yes	yes	yes	yes	yes	yes	yes	yes	yes
Molybdenum	7439-98-7	yes	yes	yes	yes	yes	yes	yes	yes	yes
Nickel	7440-02-0	yes	yes	yes	yes	yes	yes	yes	yes	yes
Potassium	7440-09-7	no	no	no	yes	yes	no	no	no	no
Rhodium	7440-16-6	yes	no	no	yes	yes	yes	no	no	yes

Table D-4. Selection of Ecological COPCs for Inclusion in the Quantitative PRA

		Are there ecological toxicity data available?			concentration be quantified?		Can ecological risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Direct Contact?[a]	Ingestion?[b]	Inhalation?[c]	Emission Rate Available to Quantify Direct Exposure?[d]	Emission Rate Available to Quantify Indirect Exposure?[e]	Direct Contact?[f]	Ingestion?[g]	Inhalation?[h]	Is the constituent a quantitative COPC?[i]
Selenium	7782-49-2	yes	yes	yes	yes	yes	yes	yes	yes	yes
Silicon	7440-21-3	no	no	no	yes	yes	no	no	no	no
Silver	7440-22-4	yes	yes	no	yes	yes	yes	yes	no	yes
Sodium	7440-23-5	no	no	no	yes	yes	no	no	no	no
Strontium	7440-24-6	yes	yes	no	nd	no	no	no	no	no
Tantalum	7440-25-7	no	no	no	yes	yes	no	no	no	no
Thallium	7440-28-0	yes	yes	no	yes	yes	yes	yes	no	yes
Tin	7440-31-5	yes	no	no	yes	yes	yes	no	no	yes
Tungsten	7440-33-7	no	no	no	yes	yes	no	no	no	no
Uranium	7440-61-1	yes	yes	no	yes	yes	yes	yes	no	yes
Vanadium	7440-62-2	yes	yes	no	yes	yes	yes	yes	no	yes
Yttrium	7440-65-5	no	no	no	yes	yes	no	no	no	no
Zinc	7440-66-6	yes	yes	no	yes	yes	yes	yes	no	yes
Zirconium	7440-67-7	yes	no	no	yes	yes	yes	no	no	yes
Non-metals and Ions										
Ammonia/Ammonium	7664-41-7	no	no	yes	yes	yes	no	no	yes	yes
Bromide	24959-67-9	yes	no	no	yes	yes	yes	no	no	yes
Chloride	16887-00-6	no	no	no	yes	yes	no	no	no	no
Cyanide	57-12-5	yes	yes	no	yes	yes	yes	yes	no	yes
Fluoride	16984-48-8	yes	yes	no	yes	yes	yes	yes	no	yes
Hydroxide	14280-30-9	no	no	no	yes	yes	no	no	no	no
Iodine	7553-56-2	yes	no	yes	yes	yes	yes	no	yes	yes
Nitrate	14797-55-8	no	no	no	yes	yes	no	no	no	no
Nitrite	14797-65-0	no	no	no	yes	yes	no	no	no	no
Phosphate	14265-44-2	no	no	no	yes	yes	no	no	no	no
Phosphorus	7723-14-0	no	no	yes	yes	yes	no	no	yes	yes
Sulfate	14808-79-8	no	no	no	yes	yes	no	no	no	no

Table D-4. Selection of Ecological COPCs for Inclusion in the Quantitative PRA

		Are there ecological toxicity data available?			concentration be quantified?		Can ecological risk be quantified?			
Constituent of Potential Concern	CAS Registry Number	Direct Contact?[a]	Ingestion?[b]	Inhalation?[c]	Emission Rate Available to Quantify Direct Exposure?[d]	Emission Rate Available to Quantify Indirect Exposure?[e]	Direct Contact?[f]	Ingestion?[g]	Inhalation?[h]	Is the constituent a quantitative COPC?[i]
Total Sulfur	63705-05-5	yes	no	yes	nd	yes	no	no	yes	yes
Criteria Pollutants										
Carbon dioxide	124-38-9	no	no	no	nd	yes	no	no	no	no
Nitrogen dioxide	10102-44-0	no	no	no	nd	yes	no	no	no	no
Ozone	10028-15-6	no	no	no	nd	yes	no	no	no	no
Particulate matter	no cas #	no	no	no	nd	yes	no	no	no	no
Sulfur dioxide	7446-09-5	no	no	no	nd	yes	no	no	no	no

nd = not determined at this time

[a] Toxicity reference values for plants, earthworms, sediment-dwelling biota and aquatic biota appear in Tables C-3-1, C-3-2, C-3-8, and C-3-9, respectively. If a TRV for any of these receptors is available, a value of "yes" appears in this column.

[b] Oral toxicity reference values for birds and mammals appear in Tables C-3-4 and C-3-6, respectively. If a TRV for any of these receptors is available, a value of "yes" appears in this column.

[c] Inhalation toxicity reference values for mammals appear in Table C-3-7. If a TRV is available, a value of "yes" appears in this column.

[d] Is an emission rate estimate available to quantify direct exposure (i.e., air concentrations)? If so, then a value of "yes" appears in this column.

[e] Is an emission rate estimate available to quantify indirect exposure (i.e., air concentrations)? If so, then a value of "yes" appears in this column.

[f] If a TRV for plants, earthworms, sediment-dwelling biota or aquatic biota is available and a laboratory analysis method is available, then a value of "yes" appears in this column.

[g] If an oral TRV for birds or mammals is available and a laboratory analysis method and all chemical-specific parameters required to model indirect exposure are available, a value of "yes" appears in this column.

[h] If an inhalation TRV for mammals is available and a laboratory analysis method and all chemical-specific parameters required to model indirect exposure are available, a value of "yes" appears in this column.

[i] If risk can be quantified for either direct exposure to plants, earthworms, sediment-dwelling biota, or aquatic biota, indirect exposure by ingestion to birds or mammals, or indirect exposure by inhalation to mammals, then a value of "yes" appears in this column.

Table D-5. Selection of Ecological ROPCs for Inclusion in the Quantitative PRA

		Are there ecological toxicity data available?			Can exposure concentration be quantified?		Can ecological risk be quantified?			
Consituent of Potential Concern	CAS Registry Number	Direct Contact?[a]	Ingestion?[b]	Inhalation?[c]	Emission Rate Available to Quantify Direct Exposure? [D]	Emission Rate Available to Quantify Indirect Exposure? [E]	Direct Contact?[f]	Ingestion?[g]	Inhalation?[h]	Is constituent a quantitative ROPC?[i]
Actinium-227	14952-40-0	yes	yes	yes	no	no	no	no	no	no
Americium-241	14596-10-2	yes	yes	yes	no	no	no	no	no	no
Americium-243	14993-75-0	yes	yes	yes	no	no	no	no	no	no
Antimony-125	14234-35-6	yes	yes	yes	no	no	no	no	no	no
Barium-137	13981-97-0	yes	yes	yes	yes	yes	yes	yes	yes	yes
Cadmium-113	no cas #	yes	yes	yes	no	no	no	no	no	no
Carbon-14	14762-75-5	yes	yes	yes	yes	yes	yes	yes	yes	yes
Cesium-134	13967-70-9	yes	yes	yes	yes	yes	yes	yes	yes	yes
Cesium-137	10045-97-3	yes	yes	yes	no	no	no	no	no	no
Cobalt-60	10198-40-0	yes	yes	yes	yes	yes	yes	yes	yes	yes
Curium-242	15510-73-3	yes	yes	yes	no	no	no	no	no	no
Curium-243	15757-87-6	yes	yes	yes	no	no	no	no	no	no
Curium-244	13981-15-2	yes	yes	yes	no	no	no	no	no	no
Europium-152	14683-23-9	yes	yes	yes	no	no	no	no	no	no
Europium-154	15585-10-1	yes	yes	yes	yes	yes	yes	yes	yes	yes
Europium-155	14391-16-3	yes	yes	yes	yes	yes	yes	yes	yes	yes
Iodine-129	15046-84-1	yes	yes	yes	yes	yes	yes	yes	yes	yes
Neptunium-237	13994-20-2	yes	yes	yes	yes	yes	yes	yes	yes	yes
Nickel-59	14336-70-0	yes	yes	yes	no	no	no	no	no	no
Nickel-63	13981-37-8	yes	yes	yes	no	no	no	no	no	no
Niobium-93	7440-03-1	yes	yes	yes	no	no	no	no	no	no
Plutonium-238	13981-16-3	yes	yes	yes	yes	yes	yes	yes	yes	yes
Plutonium-239	15117-48-3	yes	yes	yes	yes	yes	yes	yes	yes	yes
Plutonium-240	14119-33-6	yes	yes	yes	yes	yes	yes	yes	yes	yes
Plutonium-241	14119-32-5	yes	yes	yes	yes	yes	yes	yes	yes	yes
Plutonium-242	13982-10-0	yes	yes	yes	no	no	no	no	no	no
Protactinium-231	14331-85-2	yes	yes	yes	no	no	no	no	no	no
Radium-226	13982-63-3	yes	yes	yes	no	no	no	no	no	no
Radium-228	15262-20-1	yes	yes	yes	no	no	no	no	no	no
Ruthenium-106	13967-48-1	yes	yes	yes	no	no	no	no	no	no

Table D-5. Selection of Ecological ROPCs for Inclusion in the Quantitative PRA

		Are there ecological toxicity data available?			Can exposure concentration be quantified?		Can ecological risk be quantified?			
Consituent of Potential Concern	CAS Registry Number	Direct Contact?[a]	Ingestion?[b]	Inhalation?[c]	Emission Rate Available to Quantify Direct Exposure? [D]	Emission Rate Available to Quantify Indirect Exposure? [E]	Direct Contact?[f]	Ingestion?[g]	Inhalation?[h]	Is constituent a quantitative ROPC?[i]
Samarium-151	15715-94-3	yes	yes	yes	no	no	no	no	no	no
Selenium-79	no cas #	yes	yes	yes	no	no	no	no	no	no
Strontium-90	10098-97-2	yes	yes	yes	yes	yes	yes	yes	yes	yes
Technetium-99	14133-76-7	yes	yes	yes	yes	yes	yes	yes	yes	yes
Thorium-229	15594-54-4	yes	yes	yes	no	no	no	no	no	no
Thorium-232	7440-29-1	yes	yes	yes	no	no	no	no	no	no
Tin-126	15832-50-5	yes	yes	yes	no	no	no	no	no	no
Tritium	10028-17-8	yes	yes	yes	yes	yes	yes	yes	yes	yes
Uranium-232	14158-29-3	yes	yes	yes	no	no	no	no	no	no
Uranium-233	13968-55-3	yes	yes	yes	no	no	no	no	no	no
Uranium-234	13966-29-5	yes	yes	yes	no	no	no	no	no	no
Uranium-235	15117-96-1	yes	yes	yes	yes	yes	yes	yes	yes	yes
Uranium-236	13982-70-2	yes	yes	yes	yes	yes	yes	yes	yes	yes
Uranium-238	7440-61-1	yes	yes	yes	no	no	no	no	no	no
Yttrium-90	10098-91-6	yes	yes	yes	yes	yes	yes	yes	yes	yes
Zirconium-93	15751-77-6	yes	yes	yes	no	no	no	no	no	no

nd = not determined at this time

[a] If a radiological TRV for plants, soil-dwelling invertebrates, sediment-dwelling biota, or aquatic biota is available, a value of "yes" appears in this column.
[b] If an radiological ingestion TRV for birds or mammals is available, a value of "yes" appears in this column.
[c] If an radiological inhalation TRV for birds or mammals is available, a value of "yes" appears in this column.
[d] Is an emission rate estimate available to quantify direct exposure (i.e., air concentrations)? If so, then a value of "yes" appears in this column.
[e] Is an emission rate estimate available to quantify indirect exposure (i.e., air concentrations)? If so, then a value of "yes" appears in this column.

[f] If a radiological TRV for plants, soil-dwelling invertebrates, sediment-dwelling biota or aquatic biota is available and a laboratory analysis method is available, then a value of "yes" appears in this column.

[g] If an radiological ingestion TRV for birds or mammals is available and a laboratory analysis method and all chemical-specific parameters required to model indirect exposure are available, a value of "yes" appears in this column.
[h] If a radiological inhalation TRV for mammals is available and a laboratory analysis method and all chemical-specific parameters required to model indirect exposure are available, a value of "yes" appears in this column.
[i] If risk can be quantified for either direct exposure to plants, earthworms, sediment-dwelling biota, or aquatic biota, indirect exposure by ingestion to birds or mammals, or indirect exposure by inhalation to mammals, then a value of "yes" appears in this column.

Table D-6. Summary of Selection of Ecological COPCs and ROPCs for Inclusion in the Quantitative PRA

	Number that can be evaluated quantitatively	Number that cannot be evaluated quantitatively	Number that cannot yet be determined	Total
Organic COPCs	237	133	0	370
Inorganic COPCs	33	21	0	54
ROPCs	18	28	0	46
Total	288	182	0	470

Appendix E

COPCs and ROPCs for Which There are Emission Rates

Table E-1. List of Organic Constituents for Which There are Estimated Emission Rates and No Emission Rates

CAS Registry Number	Constituent of Potential Concern With Emission Rates	CAS Registry Number	Constituent of Potential Concern With Emission Rates
Aromatic Halogenated Hydrocarbons		**Dioxin and Furan Compounds**	
58-90-2	2,3,4,6-Tetrachlorophenol	132-64-9	Dibenzofuran
59-50-7	4-Chloro-3-methylphenol	1746-01-6	2,3,7,8-Tetrachlorodibenzo(p)dioxin [a]
Aromatic Nonhalogenated Hydrocarbons		**PCBs**	
100-41-4	Ethyl benzene	1336-36-3	Polychlorinated biphenyls (PCBs)
100-42-5	Styrene	1746-01-6	2,3,7,8-TCDD (coplanar PCBs) [b]
106-42-3	p-Xylene	**Phthalates**	
108-38-3	m-Xylene	117-81-7	Bis(2-ethylhexyl)phthalate (DEHP)
108-88-3	Toluene	117-84-0	n-Dioctyl phthalate
71-43-2	Benzene	84-66-2	Diethyl phthalate
88-72-2	2-Nitrotoluene	84-74-2	Dibutyl phthalate
92-93-3	4-Nitrobiphenyl	85-68-7	Butylbenzyl phthalate
95-47-6	o-Xylene	**Light Polycyclic Aromatic Hydrocarbons**	
Non-aromatic Nonhalogenated Hydrocarbons		120-12-7	Anthracene
106-35-4	3-Heptanone	129-00-0	Pyrene
106-88-7	1,2-Epoxybutane	208-96-8	Acenaphthylene
106-97-8	Butane	602-87-9	5-Nitroacenaphthene
106-99-0	1,3-Butadiene	83-32-9	Acenaphthene
107-02-8	Acrolein	85-01-8	Phenanthrene
107-12-0	Propionitrile	86-73-7	Fluorene
107-13-1	Acrylonitrile	91-20-3	Naphthalene
107-18-6	2-Propene-1-ol	91-58-7	2-Chloronapthalene
107-31-3	Formic acid, methyl ester	95-13-6	Indene
107-87-9	2-Pentanone	**Heavy Polycyclic Aromatic Hydrocarbons**	
108-03-2	1-Nitropropane	1321-64-8	Pentachloronaphthalene
108-05-4	Vinyl acetate	1321-65-9	Trichloronaphthalene
108-10-1	4-Methyl-2-pentanone	1335-87-1	Hexachloronaphthalene
108-20-3	Bis(isopropyl)ether	1335-88-2	Tetrachloronaphthalene
108-87-2	Methylcyclohexane	189-55-9	Dibenzo[a,i]pyrene
108-94-1	Cyclohexanone	189-64-0	Dibenzo[a,h]pyrene
109-66-0	n-Pentane	191-24-2	Benzo(g,h,i)perylene
110-12-3	5-Methyl-2-hexanone	191-30-0	Benzo[a,i]pyrene
110-43-0	2-Heptanone	192-65-4	Dibenzo[a,e]pyrene
110-54-3	n-Hexane	193-39-5	Indeno(1,2,3-cd)pyrene
110-62-3	n-Valeraldehyde	205-82-3	Benzo(j)fluoranthene
110-82-7	Cyclohexane	205-99-2	Benzo(b)fluoranthene
110-83-8	Cyclohexene	206-44-0	Fluoranthene
111-65-9	n-Octane	207-08-9	Benzo(k)fluoranthene
111-84-2	n-Nonane	218-01-9	Chrysene
121-44-8	Triethylamine	2234-13-1	Octachloronaphthalene
123-19-3	4-Heptanone	224-42-0	Dibenz[a,j]acridine
123-38-6	n-Propionaldehyde	226-36-8	Dibenz[a,h]acridine
123-51-3	3-Methyl-1-butanol	3697-24-3	5-Methylchrysene
123-86-4	Acetic acid n-butyl ester	50-32-8	Benzo(a)pyrene
123-91-1	1,4-Dioxane	53-70-3	Dibenz(a,h)anthracene

Table E-1. List of Organic Constituents for Which There are Estimated Emission Rates and No Emission Rates

CAS Registry Number	Constituent of Potential Concern With Emission Rates	CAS Registry Number	Constituent of Potential Concern With Emission Rates
126-98-7	2-Methyl-2-propenenitrile	56-49-5	3-Methylcholanthrene
127-19-5	N,N-Dimethylacetamide	56-55-3	Benzo(a)anthracene
141-78-6	Acetic acid ethyl ester	**Light Substituted Benzene Compounds**	
141-79-7	4-Methyl-3-penten-2-one	100-00-5	p-Nitrochlorobenzene
142-82-5	n-Heptane	100-25-4	1,4-Dinitrobenzene
1634-04-4	Methyl tert-butyl ether	106-46-7	1,4-Dichlorobenzene
287-92-3	Cyclopentane	108-39-4	m-Cresol
4170-30-3	2-Butenaldehyde (2-Butenal)	108-90-7	Chlorobenzene
50-00-0	Formaldehyde	108-95-2	Phenol
540-84-1	2,2,4-Trimethylpentane	120-82-1	1,2,4-Trichlorobenzene
563-80-4	3-Methyl-2-butanone	120-83-2	2,4-Dichlorophenol
591-78-6	2-Hexanone	25551-13-	Trimethyl benzene
60-29-7	Ethyl ether	541-73-1	1,3-Dichlorobenzene
60-35-5	Acetamide	88-06-2	2,4,6-Trichlorophenol
624-83-9	Methyl isocyanate	88-75-5	2-Nitrophenol
64-17-5	Ethyl alcohol	95-48-7	o-Cresol
64-18-6	Formic acid	95-49-8	2-Chlorotoluene
64-19-7	Acetic acid	95-50-1	1,2-Dichlorobenzene
67-56-1	Methyl alcohol	95-57-8	2-Chlorophenol
67-63-0	2-Propyl alcohol	95-95-4	2,4,5-Trichlorophenol
67-64-1	2-Propanone (Acetone)	98-82-8	Cumene
71-23-8	n-Propyl alcohol	98-83-9	alpha-Methylstyrene
71-36-3	n-Butyl alcohol	98-95-3	Nitrobenzene
74-99-7	Methylacetylene	**Other Light Semivolatile Compounds**	
75-05-8	Acetonitrile	100-21-0	p-Phthalic acid
75-07-0	Acetaldehyde	101-84-8	Diphenyl ether
75-12-7	Formamide	106-93-4	Ethylene dibromide
75-15-0	Carbon disulfide	108-93-0	Cyclohexanol
75-21-8	Oxirane	109-99-9	Tetrahydrofuran
75-50-3	Trimethylamine	110-86-1	Pyridine
75-52-5	Nitromethane	111-76-2	Ethylene glycol monobutyl ether
75-55-8	2-Methylaziridine	121-69-7	Dimethylaniline
75-65-0	2-Methyl-2-propanol	122-39-4	N,N-Diphenylamine
78-83-1	2-Methylpropyl alcohol	144-62-7	Oxalic acid
78-92-2	1-Methylpropyl alcohol	542-88-1	Dichloromethyl ether
78-93-3	2-Butanone	57-14-7	1,1-Dimethylhydrazine
79-09-4	Propionic acid	59-89-2	N-Nitrosomorpholine
79-20-9	Methyl acetate	60-34-4	Methylhydrazine
96-22-0	3-Pentanone	621-64-7	Di-n-Propylnitrosamine
98-51-1	p-tert-Butyltoluene	627-13-4	Nitric acid, propyl ester
Non-aromatic Halogenated Hydrocarbons		62-75-9	N-Nitroso-N,N-dimethylamine
10061-01-	cis-1,3-Dichloropropene	79-10-7	2-Propenoic acid
10061-02-	trans-1,3-Dichloropropene	91-22-5	Quinoline
107-05-1	3-Chloropropene (Allyl chloride)	92-52-4	1,1'-Biphenyl
107-06-2	1,2-Dichloroethane	98-86-2	Acetophenone

Table E-1. List of Organic Constituents for Which There are Estimated Emission Rates and No Emission Rates

CAS Registry Number	Constituent of Potential Concern With Emission Rates	CAS Registry Number	Constituent of Potential Concern With Emission Rates
127-18-4	1,1,2,2-Tetrachloroethene	**Other Heavy Semivolatile Compounds**	
156-60-5	trans-1,2-Dichloroethylene	101-55-3	4-Bromophenylphenyl ether
27154-33-	Trichlorofluoroethane	107-66-4	Dibutylphosphate
540-59-0	1,2-Dichloroethylene	118-74-1	Hexachlorobenzene
56-23-5	Carbon tetrachloride	126-73-8	Tributyl phosphate
630-20-6	1,1,1,2-Tetrachloroethane	128-37-0	2,6-Bis(tert-butyl)-4-methylphenol
67-66-3	Chloroform	1582-09-8	Trifluralin
684-16-2	Hexafluoroacetone	1836-75-5	Nitrofen
71-55-6	1,1,1-Trichloroethane	2385-85-5	Mirex
74-83-9	Bromomethane	26140-60-	Terphenyls
74-87-3	Chloromethane	3825-26-1	Ammonium perfluorooctanoate
74-97-5	Bromochloromethane	603-34-9	Triphenylamine
75-00-3	Chloroethane	67-72-1	Hexachloroethane
75-01-4	1-Chloroethene	82-68-8	Pentachloronitrobenzene
75-09-2	Dichloromethane	87-68-3	Hexachlorobutadiene
75-27-4	Bromodichloromethane	87-86-5	Pentachlorophenol
75-34-3	1,1-Dichloroethane	88-85-7	2-sec-Butyl-4,6-dinitrophenol
75-35-4	1,1-Dichloroethene	88-89-1	Picric acid
75-43-4	Dichlorofluoromethane	96-69-5	Bis(3-tert-butyl-4-hydroxy-6-methyl-phenyl)sulf
75-45-6	Chlorodifluoromethane	**Herbicides and Organochlorinated Pesticides**	
75-61-6	Difluorodibromomethane	309-00-2	Aldrin
75-63-8	Trifluorobromomethane	319-84-6	alpha-BHC
75-69-4	Trichlorofluoromethane	319-85-7	beta-BHC
75-71-8	Dichlorodifluoromethane	319-86-8	Delta-BHC
75-99-0	2,2-Dichloropropionic acid	465-73-6	Isodrin
76-03-9	Trichloroacetic acid	50-29-3	4,4-DDT
76-11-9	1,1,1,2-Tetrachloro-2,2-difluoroethane	58-89-9	gamma-BHC (Lindane)
76-12-0	1,1,2,2-Tetrachloro-1,2-difluoroethane	60-57-1	Dieldrin
76-13-1	1,2,2-Trichloro-1,1,2-trifluoroethane	72-20-8	Endrin
76-14-2	1,2-Dichloro-1,1,2,2-tetrafluoroethane	72-43-5	Methoxychlor
76-15-3	Chloropentafluoroethane	72-54-8	4,4-DDD
78-87-5	1,2-Dichloropropane	72-55-9	4,4-DDE
79-00-5	1,1,2-Trichloroethane	76-44-8	Heptachlor
79-01-6	1,1,2-Trichloroethylene	8001-35-2	Toxaphene
79-34-5	1,1,2,2-Tetrachloroethane	93-72-1	Silvex (2,4,5-TP)
96-18-4	1,2,3-Trichloropropane	93-76-5	2,4,5-T
		94-75-7	2,4-D and esters

Table E-1. List of Organic Constituents for Which There are Estimated Emission Rates and No Emission Rates

Constituent of Potential Concern	Constituent of Potential Concern
Without Emission Rates	Without Emission Rates
Aromatic Halogenated Hydrocarbons	**Light Substituted Benzene Compounds (continued)**
Aromatic Nonhalogenated Hydrocarbons	n-Butyl benzene
Benzyl alcohol	2,4-Dimethylphenol
Benzaldehyde	4-Chlorotoluene
Non-aromatic Nonhalogenated Hydrocarbons	p-Cresol
Propargyl alcohol	p-Chloroaniline
Propylene gylcol monomethyl ether	p-Toluidine
2-Methoxyethanol	1,3,5-Trimethyl benzene
2-Ethoxyethanol	Bromobenzene
Cyanogen	2,4-Dinitrotoluene
Phosgene	sec-Butyl benzene
Glycidylaldehyde	2,4-Dinitrophenol
Methyl methacrylate	o-Dinitrobenzene
Ethyl methacrylate	4,6-Dinitro-o-cresol
p-Cymene	2,6-Dinitrotoluene
Non-aromatic Halogenated Hydrocarbons	Aniline
Chlorodibromomethane	Toluene-2,6-diamine
cis-1,2-Dichloroethene	1,2,3-Trichlorobenzene
Cyanogen bromide	o-Nitroaniline
Cyanogen chloride	o-Toluidine
1,3-Dichloropropene	1,2,4-Trimethyl benzene
Bromoethene	tert-Butyl benzene
Iodomethane	Benzotrichloride
Methylene bromide	1,3-Dinitrobenzene
Bromoform	**Other Light Semivolatile Compounds**
2-Chloropropane	4,4-Methylenedianiline
Pentachloroethane	Quinone
1,4-Dichloro-2-butene	Epichlorohydrin
1,2-Dibromo-3-chloropropane	Ethylene glycol
Dioxin and Furan Compounds [a]	Dichloroisopropyl ether
1,2,3,7,8,9-Hexachlorodibenzo(p)dioxin	Malononitrile
Octachlorodibenzo(p)dioxin	Ethylene glycol monoethyl ether acetate
1,2,3,4,6,7,8-Heptachlorodibenzo(p)dioxin	Bis(2-chloroethyl) ether
Octachlorodibenzofuran	Bis(2-chloroethoxy)methane
1,2,3,4,7,8-Hexachlorodibenzo(p)dioxin	1,3-Propane sultone
1,2,3,7,8-Pentachlorodibenzo(p)dioxin	1,2-Diphenylhydrazine
2,3,7,8-Tetrachlorodibenzofuran	Maleic hydrazide
1,2,3,4,7,8,9-Heptachlorodibenzofuran	Methyl styrene (mixed isomers)
2,3,4,7,8-Pentachlorodibenzofuran	Chlorocyclopentadiene
1,2,3,7,8-Pentachlorodibenzofuran	Ethyl Carbamate (urethane)
1,2,3,6,7,8-Hexachlorodibenzofuran	2-Chloroacetophenone
1,2,3,6,7,8-Hexachlorodibenzo(p)dioxin	1,2-Dimethylhydrazine
2,3,4,6,7,8-Hexachlorodibenzofuran	2,4-Toluene diisocyante
1,2,3,4,6,7,8-Heptachlorodibenzofuran	Ethyl methanesulfonate
1,2,3,4,7,8-Hexachlorodibenzofuran	Benzoic acid

Table E-1. List of Organic Constituents for Which There are Estimated Emission Rates and No Emission Rates

Constituent of Potential Concern	Constituent of Potential Concern
Without Emission Rates	**Without Emission Rates**
1,2,3,7,8,9-Hexachlorodibenzofuran	Dimethyl sulfate
PCBs [b]	Phthalic anhydride
2,3',4,4',5- Pentachlorobiphenyl	o-Anisidine
3,3',4,4'-Tetrachlorobiphenyl	N-Nitrosodi-n-butylamine
2,3,3',4,4'-Pentachlorobiphenyl	Safrole
3,3',4,4',5,5'- Hexachlorobiphenyl	Ethylene thiourea
2,2',3,4,4',5,5'- Heptachlorobiphenyl	Furfural
2,2',3,3',4,4',5- Heptachlorobiphenyl	Dichloropentadiene
2,3,3',4,4',5- Hexachlorobiphenyl	**Other Heavy Semivolatile Compounds**
2,3,3',4,4',5'- Hexachlorobiphenyl	Azobenzene
3,4,4',5-Tetrachlorobiphenyl	3,3'-Dimethoxybenzidine
2,3,4,4',5- Pentachlorobiphenyl	2-Cyclohexyl-4,6-dinitrophenol
2,3,3',4,4',5,5'- Heptachlorobiphenyl	Captan
2',3,4,4',5- Pentachlorobiphenyl	Pronamide
2,3',4,4',5,5'- Hexachlorobiphenyl	Chlorobenzilate
3,3',4,4',5- Pentachlorobiphenyl	Strychnine
Phthalates	Dimethyl aminoazobenzene
Dimethylphthalate	Pentachlorobenzene
Light Polycyclic Aromatic Hydrocarbons	Hexachlorophene
2-Methyl naphthalene	Hexachlorocyclopentadiene
Heavy Polycyclic Aromatic Hydrocarbons	Hexamethylene-1,5-diisocyanate
Benzo(e)pyrene	3,3-Dichlorobenzidine
Dibenzo(a,e)fluoranthene	1,2,4,5-Tetrachlorobenzene
Dibenzo(a,h)fluoranthene	1,3,5-Trinitrobenzene
Light Substituted Benzene Compounds	**Herbicides and Organochlorinated Pesticides**
4-Nitrophenol	Endothall
Benzyl chloride	Chlordane
n-Propyl benzene	

[a] 2,3,7,8-TCDD requirements will be used to represent emissions of other 2,3,7,8 substituted dioxin and furan compounds.

[b] All coplanar PCBs will be modeled together as 2,3,7,8-TCDD equivalents.

Table E-2. List of Inorganic Constituents for Which There are Estimated Emission Rates [a]

CAS Registry Number	Constituent	CAS Registry Number	Constituent
	Metals		**Non-metals and Anions**
7429-90-5	Aluminum	7664-41-7	Ammonia/Ammonium
7440-36-0	Antimony	24959-67-9	Bromide
7440-38-2	Arsenic	16887-00-6	Chloride
7440-39-3	Barium	57-12-5	Cyanide
7440-41-7	Beryllium	16984-48-8	Flouride
7440-69-9	Bismuth	14280-30-9	Hydroxide
7440-42-8	Boron	7553-56-2	Iodine (I2)
7440-43-9	Cadmium	14797-55-8	Nitrate
7440-70-2	Calcium	14797-65-0	Nitrite
18540-29-9	Chromium (total)	14265-44-2	Phosphate
7440-48-4	Cobalt	7723-14-0	Phosphorous
7440-50-8	Copper	14808-79-8	Sulfate
7439-89-6	Iron	63705-05-5	Total Sulfur
7439-92-1	Lead		**Criteria Pollutants**
7439-93-2	Lithium	124-38-9	Carbon dioxide
7439-95-4	Magnesium	10102-44-0	Nitrogen dioxide
7439-96-5	Manganese	10028-15-6	Ozone
7439-97-6	Mercury	no cas #	Particulate matter
7439-98-7	Molybdenum	7446-09-5	Sulfur dioxide
7440-02-0	Nickel		
7440-09-7	Potassium		
7440-16-6	Rhodium		
7782-49-2	Selenium		
7440-21-3	Silicon		
7440-22-4	Silver		
7440-23-5	Sodium		
7440-25-7	Tantalum		
7440-28-0	Thallium		
7440-31-5	Tin		
7440-33-7	Tungsten		
7440-61-1	Uranium		
7440-62-2	Vanadium		
7440-65-5	Yttrium		
7440-66-6	Zinc		
7440-67-7	Zirconium		

[a] Strontium does not have an emission rate.

Table E-3. List of Radionuclides for Which There are Estimated Emission Rates

CAS Registry Number	Radionuclide	CAS Registry Number	Radionuclide
With Emission Rates		**Without Emission Rates**	
13981-97-0	Barium-137	10045-97-3	Cesium-137
14762-75-5	Carbon-14	15510-73-3	Curium-242
13967-70-9	Cesium-134	15757-87-6	Curium-243
10198-40-0	Cobalt-60	13981-15-2	Curium-244
15585-10-1	Europium-154	14683-23-9	Europium-152
14391-16-3	Europium-155	14336-70-0	Nickel-59
15046-84-1	Iodine-129	13981-37-8	Nickel-63
13994-20-2	Neptunium-237	7440-03-1	Niobium-93
13981-16-3	Plutonium-238	13982-10-0	Plutonium-242
15117-48-3	Plutonium-239	14331-85-2	Protactinium-231
14119-33-6	Plutonium-240	13982-63-3	Radium-226
14119-32-5	Plutonium-241	15262-20-1	Radium-228
10098-97-2	Strontium-90	13967-48-1	Ruthenium-106
14133-76-7	Technetium-99	15715-94-3	Samarium-151
10028-17-8	Tritium	no cas #	Selenium-79
15117-96-1	Uranium-235	15594-54-4	Thorium-229
13982-70-2	Uranium-236	7440-29-1	Thorium-232
10098-91-6	Yttrium-90	15832-50-5	Tin-126
Without Emission Rates		14158-29-3	Uranium-232
14952-40-0	Actinium-227	13968-55-3	Uranium-233
14596-10-2	Americium-241	13966-29-5	Uranium-234
14993-75-0	Americium-243	7440-61-1	Uranium-238
14234-35-6	Antimony-125	15751-77-6	Zirconium-93
no cas #	Cadmium-113		