



WASHINGTON STATE
DEPARTMENT OF
E C O L O G Y

Results of a Screening Analysis for Pharmaceuticals in Wastewater Treatment Plant Effluents, Wells, and Creeks in the Sequim-Dungeness Area

November 2004

Publication No. 04-03-051

printed on recycled paper



This report is available on the Department of Ecology home page on the World Wide Web at <http://www.ecy.wa.gov/biblio/0403051.html>

For a printed copy of this report, contact:

Department of Ecology Publications Distributions Office

Address: PO Box 47600, Olympia WA 98504-7600

E-mail: ecypub@ecy.wa.gov


Phone: (360) 407-7472

Refer to Publication Number 04-03-051

Any use of product or firm names in this publication is for descriptive purposes only and does not imply endorsement by the author or the Department of Ecology.

The Department of Ecology is an equal-opportunity agency and does not discriminate on the basis of race, creed, color, disability, age, religion, national origin, sex, marital status, disabled veteran's status, Vietnam-era veteran's status, or sexual orientation.

If you have special accommodation needs or require this document in alternative format, please contact Joan LeTourneau at 360-407-6764 (voice) or 711 or 1-800-833-6388 (TTY).



WASHINGTON STATE
DEPARTMENT OF
E C O L O G Y

Results of a Screening Analysis for Pharmaceuticals in Wastewater Treatment Plant Effluents, Wells, and Creeks in the Sequim-Dungeness Area

by
Art Johnson, Barbara Carey, and Steve Golding

Environmental Assessment Program
Olympia, Washington 98504-7710

November 2004

Waterbody Numbers:
WA-18-1100, Bell Creek, and WA-18-1300, Cassalery Creek

Publication Number: 04-03-051

printed on recycled paper



This page is purposely left blank for duplex printing.

Table of Contents

	<u>Page</u>
List of Figures and Tables.....	ii
Abstract.....	iii
Acknowledgements.....	iv
Introduction.....	1
Study Area	2
Chemicals Analyzed	4
Sampling Design.....	7
Methods.....	9
Field Procedures.....	9
Laboratory Procedures	10
Data Quality	11
Results.....	13
Effluent and Creek Flows	13
Sunland WWTP and Related Samples.....	13
Sequim WWTP and Related Samples.....	16
Residential Area Well.....	18
Summary of Detection Frequencies.....	20
Significance of Findings	22
Recommendations.....	23
References.....	24
Appendices	
A. Latitude and Longitude of PPCP Sampling Sites	
B. Summary of PPCP Results for the Sequim-Dungeness Project	

List of Figures and Tables

	<u>Page</u>
Figure 1. Study Area.....	2
Table 1. PPCPs Analyzed	5
Table 2. PPCPs of Potential Interest but Not Analyzed.....	6
Table 3. Wells Sampled	8
Table 4. Number and Type of Samples Analyzed	8
Table 5. Field Procedures	9
Table 6. Laboratory Procedures.....	10
Table 7. Comparison of Results on Replicate Samples	12
Table 8. Effluent and Creek Flow Rates During PPCP Sample Collection	13
Table 9. Results for Sunland WWTP, Adjacent Wells, and Nearby Cassalery Creek	15
Table 10. Results for Sequim WWTP, Wells Adjacent to Water-Reuse Project, and Lower Bell Creek.....	17
Table 11. Results for Private Well Affected by Septic Systems.....	19
Table 12. Detection Frequency for PPCPs in Sequim-Dungeness Effluent, Well, and Creek Sample	21

Abstract

A screening analysis was conducted for 24 pharmaceuticals and personal care products (PPCPs) in tertiary wastewater treatment plant effluents and nearby wells and creeks in the Sequim-Dungeness area of northwest Washington. The objective was to investigate the potential for and status of PPCP contamination of area waters from application of treated wastewater via re-use programs and conventional land application. A well located in an area served by on-site septic systems also was sampled.

The following 16 compounds were detected in one or both effluents: Acetaminophen, Caffeine, Carbamazepine, Cimetidine, Codeine, Cotinine, Diltiazem, Hydrocodone, Ketoprofen, Metformin, Nicotine, Paraxanthine, Salbutamol, Sulfamethoxazole, Trimethoprim, and Estrone. Concentrations ranged from 0.26 ng/L (Estrone) to 200 ug/L (Paraxanthine). Only Caffeine, Nicotine, and the diabetes drug Metformin (tentatively identified) were consistently detected in the well and creek samples; concentrations were 25 ug/L or less. In at least some samples, Nicotine appeared to be an artifact of sampling handling procedures.

These limited results give no indication that PPCPs represent a significant concern in the wells or creeks sampled. Additional monitoring for PPCPs appears to be a low priority in connection with the Sequim and Sunland wastewater treatment plants.

Acknowledgements

The authors would like to thank the home owners and wastewater treatment plant operators in the Sequim-Dungeness area who gave permission to collect samples for this project.

The following people made important contributions to the study:

- Ann Soule, Clallam County, advised on the field work and reviewed the QA project plan and draft project report.
- Cynthia Nelson, Ecology Southwest Regional Office, provided helpful information for study planning and reviewed the draft project report.
- Dave Dougherty, Ecology Southwest Regional Office, provided helpful information on the Sequim and Sunland wastewater treatment plants.
- Karin Feddersen, Ecology Manchester Laboratory, arranged for contract laboratory services and performed the QA review of the PPCP data.
- Dale Norton, Ecology Environmental Assessment Program, supervised the work and reviewed the QA project plan and draft project report.
- Erika Wittmann, Ecology Environmental Assessment Program, did a literature search on PPCPs.
- Joan LeTourneau, Ecology Environmental Assessment Program, formatted the project report.

Introduction

Identifying emerging risks posed by previously unrecognized pollutants is one of the top five goals of the U.S. Environmental Protection Agency (EPA) Strategic Plan. Pharmaceuticals and certain personal care products (PPCPs) are a large and growing class of bioactive chemicals that, until recently, has received little attention.

Although some PPCPs are resistant to degradation, most have shorter environmental half-lives than conventional pollutants. PPCPs generally occur in surface and groundwater at ug/L to ng/L concentrations¹, far below therapeutic thresholds. There is growing concern about these compounds because many have been detected and little is known of their potential for adverse human or ecological effects. EPA has a website devoted to this issue (<http://www.epa.gov/nerlesd1/chemistry/pharma>).

In response to a request from Clallam County, the Washington State Department of Ecology (Ecology) conducted a screening analysis for PPCPs in the Sequim-Dungeness area of western Washington during November 2003. Sequim is a popular retirement center. Almost half the residents are over the age of 59, and more than 20 percent are 65 or older (2003). Pharmaceutical use is therefore likely higher than average for Washington communities. These facts, coupled with discharge of land applied reclaimed water and wastewater into or adjacent to surface waters, and a vulnerable surficial aquifer, made the Sequim-Dungeness a good candidate for assessing the potential for pharmaceutical contamination of state waters.

Samples for the screening study were collected of wastewater treatment plant (WWTP) effluents from the City of Sequim and the Sunland development in the Dungeness River valley, as well as from wells and creeks that could be impacted by these discharges. The objective was to investigate the potential for and status of pharmaceutical contamination of area waters from application of treated wastewater via re-use programs (Sequim) and conventional land application (Sunland). A well located in an area served by on-site septic systems also was sampled.

¹ parts-per-billion to parts-per-trillion

Study Area

The Sequim WWTP is a tertiary, high performing, reclaimed water plant that treats 0.5 - 0.6 million gallons per day (mgd). The influent is oxidized, coagulated, filtered, and disinfected. Final effluent meets Washington State Class A standards.

The treatment plant produces about 0.6 mgd of reclaimed water which goes to the city's Reuse Demonstration Site, constructed in 1999-2000 immediately north of Carrie-Blake Park (Figure 1). It is one of the first facilities of its kind in the Pacific Northwest. The reclaimed water is used for garden irrigation and wetland creation, and cooling, aeration, and flow augmentation of Bell Creek. For the past year, the effluent has been used to increase creek flow by 0.1 cfs (0.07 mgd). Unused water is currently discharged to a marine outfall outside of Sequim Bay. (David Dougherty, Ecology, personal communication, 2003; Pacific Groundwater Group, 2000).




Figure 1. Study Area

Sunland has a quasi-tertiary plant that has not been approved for reclaimed water. It is a land treatment facility that applies at or above agronomic rates. The sprayfield is adjacent to the plant and has been in use since 1979. Cassalery Creek flows along the north boundary of the sprayfield. There is normally no overland discharge to the creek, but effluent could enter the creek during runoff events or via groundwater. Sunland plans an upgrade to achieve Class A reclaimed water status and intends to use treated effluent on their golf course. This upgrade may be years away. (Dougherty, 2003).

Bell and Cassalery creeks are typically fed by groundwater discharge and irrigation tailwater. Irrigation diversions are from the Dungeness River and occur year-round, but are greatest during the growing season, mid-April to mid-September. The lowest flows are in September and October. Highest flows occur during winter rains and in the spring. (Pacific Groundwater Group, 2002).

The number and density of on-site sewage systems have increased in non-sewered portions of the Sequim-Dungeness area corresponding with the population increase in recent years. Thomas et al. (1999) estimated that 7,000 on-site systems existed here in 1996. The relatively shallow depth to groundwater and lack of a low-permeability layer in some areas makes the surficial aquifer vulnerable to contamination from above.

Nitrate is an indicator of groundwater contamination from various sources, including on-site sewage systems. A statistically significant increase in nitrate since 1980, though slight, was reported in this area by Thomas et al. (1999). The largest area of high nitrate concentrations is east of the Dungeness River and north of Bell Creek, where values up to 4.3 mg/L are found. The median nitrate concentration in residential areas (1.3 mg/L) is also higher than in agricultural (0.55 mg/L) or natural grassland or forest areas (0.12 mg/L). Nitrate values are highest in residential areas that have a high density of on-site systems compared to medium density systems, and lowest in low density areas.

The Sequim-Dungeness area is a rural region with a mild and relatively dry climate (15-20" or less annual rainfall). Approximately 4,440 of Clallam County's 65,900 residents live in the city (2003). 17,400 live in the unincorporated area around Sequim.

Chemicals Analyzed

Table 1 lists the PPCPs that were analyzed for the Sequim-Dungeness project and shows the basis for their selection. An initial target list was developed from recommendations in Daughton and Ternes (1999), a synthesis of the literature on the environmental occurrence, distribution, and effects of PPCPs. Other chemicals were then added that had been detected in a nationwide reconnaissance on the occurrence of PPCPs in surface waters, conducted by the U.S. Geological Survey (USGS) (Kolpin et al., 2002) or were under consideration for groundwater monitoring in the state of California (CDHS, 2003). The final list included several additional analytes that are routine target compounds for EPA-sponsored PPCP research being conducted by the State University of New York at Stony Brook (SUNYSB), the contractor selected for the present study.

Two steroid estrogens, Estrone and beta-Estradiol, were analyzed for this project. These are components or transformation products of drugs used in hormone replacement therapy. The birth control additive, Ethynyl Estradiol, was not analyzed as it is typically lost in the clean-up step for the method employed. It occurs in sewage effluents in lower concentrations than Estrone or beta-Estradiol, and their levels provide a relative indication of how much Ethynyl Estradiol could be present.

Table 2 lists other compounds that were considered for analysis in this project, but ultimately dropped due to lack of an adequate method, high cost, instrument problems at SUNYSB, or their being phased out of use (e.g., clofibrate).

Table 1. PPCPs Analyzed

Chemical	Use/Origin	Basis for Selection	Rank Among Top 200 U.S. Prescriptions 2002*	Rank Among Top 25 Washington State Prescriptions 2003**
Fenofibrate	Lipid regulator	1,4	129	
Carbamazepine	Antiepileptic	1,3,4		
Fluoxetine	Antidepressant	1,2,4	31	5
Codeine	Analgesic	2,4	1	3
Hydrocodone	Analgesic	2,4	1	3
Antipyrine	Analgesic	4		
Caffeine	Stimulant	2,3,4		
Paraxanthine	Caffeine metabolite	4		
Nicotine	Stimulant	4		
Cotinine	Nicotine metabolite	2,4		
Cimetidine	Ulcer drug	2,4		
Ranitidine	Ulcer drug	2,4	39	11
Diltiazem	Antianginal	2,4	92	
Nifedipine	Antianginal	4	194	
Salbutamol	Bronchial dilator	2,4	12	9
Sulfamethoxazole	Antibacterial	2,4		
Trimethoprim	Antibacterial	2,4		
Warfarin	Anticoagulant	2,4	57	23
Estrone	Hormone component	2,3,4		
beta-Estradiol	Hormone component	1,2,3,4		
Erythromycin	Antibiotic	2,4		
Acetaminophen	Anti-inflammatory	2,3,4	32	14
Ketoprofen	Anti-inflammatory	4		
Metformin	Antihyperglycemic	2,4	38	15

1 = Recommended for monitoring (Daughton and Ternes, 1999)

2 = Detected in USGS national study (Koplin et al., 2002)

3 = Under consideration for groundwater monitoring in California (CDHS 2003-draft)

4 = Routine target compound for Stony Brook University, NY

*<http://www.rxlist.com/top200a.htm>

**Group Health Cooperative (R. Johnson, May 29, 2003 email)

Table 2. PPCPs of Potential Interest but Not Analyzed

Chemical	Use/Origin
Gemfibrozil	Lipid regulator
Phenytoin	Antiepileptic
Primidone	Antiepileptic
17a-Ethynyl Estradiol	Oral contraceptive
Fluoroquinolone carboxylic acids	Antibiotics
Ibuprofen	Anti-inflammatory
Triclosan	Antiseptic
Clofibric acid	Lipid regulator (metabolite)
Fluvoxamine	Antidepressant
Paroxetine	Antidepressant
Ifosfamide	Antineoplastic
Cyclophosphamide	Antineoplastic
Diatrizoate (Na)	X-ray media
Iopamidol	X-ray media
Iopromide	X-ray media
Nitromusks	Fragrances
Amino musks	Fragrances
Sulfonamides	Antibiotics
Acetylsalicylic acid	Anti-inflammatory
Sildenafil citrate	Impotence drug
Methylbenzylidene camphor	Sunscreen agent
Diphenhydramine	Antihistamine

Sampling Design

Samples for PPCP screening were collected on November 17-18, 2003. A timeframe of mid-October had been initially selected for the field work as being worst-case for surface water, with the irrigation season ended and the creeks at low flow. However, the services of the contract laboratory could not be secured in time. The November data are more representative of general water quality conditions in these creeks, as opposed to extreme low flow.

The samples were analyzed for the chemicals listed in Table 1. Ancillary parameters included temperature, pH, conductivity, nitrate+nitrite-N, total suspended solids (TSS), and dissolved oxygen (groundwater only). Effluent flow rates were recorded from WWTP records and streamflow was gauged when the surface water samples were taken.

One sample each was collected of final effluents from the Sequim and Sunland WWTPs. The samples were composites of a morning and early afternoon grab. Effluent flow in WWTPs is highest in the morning, but can be more concentrated in the afternoon. A replicate effluent sample was prepared at Sunland. Sunland was selected for the replicate because of the lower level of treatment it provides.

A total of five wells were sampled: two near the Sunland sprayfield, two near the Sequim Reuse Demonstration Site, and one in an area of high nitrate concentrations (Figure 1, Table 3). The wells were selected in consultation with Ann Soule, hydrogeologist with Clallam County. The criteria for selecting wells included the following:

- The well is down-gradient of the site of interest.
- A driller's report (well log) is available for the well (if possible).
- The well is screened in as shallow an aquifer as possible – above any clay layer (if possible).
- The well is capable of producing samples representative of the groundwater.
- The well does not have a water treatment device (such as a water softener or iron treatment system) or a large storage tank that cannot be bypassed during well purging and sampling.
- The current well owner must grant access to the well.

One water sample each was collected from Bell Creek and Cassalery Creek. Bell was sampled at Schmuck Road, just before it flows into Sequim Bay (Figure 1). This site is approximately one mile below Carrie Blake Park. Cassalery was sampled at Jamestown Road, approximately ½ mile below the Sunland WWTP. A replicate sample was prepared for Bell Creek. Bell Creek was selected for the replicate because it receives effluent directly.

Table 4 shows the numbers and types of samples analyzed. This project was conducted following a Quality Assurance Project Plan prepared by Johnson and Carey (2004).

Table 3. Wells Sampled

Well ID	Location	Type of Well	Depth (ft)	Well Log?
Sunland WWTP Area				
5N2	Sunland land application site	monitoring	44	yes
Taylor Ranch Road	444 Taylor Ranch Rd.	private	49	yes
Sequim WWTP Area				
Bell Meadows Lane	154 Bell Meadows Lane	private	60?	no
Rhodefer Road	N. end N. Rhodefer Rd.	community	?	?
High Nitrate Area				
Beverage Street	885 N. Beverage St.	private	67	yes

Table 4. Number and Type of Samples Analyzed

Analysis	WWTP Effluent	Ground-water	Surface Water	Bottle Blank	Total Samples
Pharmaceuticals	3	5	3	1	12
Steroid Estrogens	3	3*	2*	0*	8
Nitrate+Nitrite-N	0	5	2	0	7
TSS	2	0	2	0	4

*Samples were lost in transport or not analyzed due to laboratory oversight

Methods

Field Procedures

Sample containers, preservation, and holding times for this project are shown in Table 5. The containers were obtained from the Ecology Manchester Environmental Laboratory, pre-cleaned to EPA QA/QC specifications (EPA, 1990).

Table 5. Field Procedures

Parameter	Minimum Sample Size	Container	Preservation	Holding Time
Pharmaceuticals	1 L	1L amber glass, teflon lid	Cool to 4°C	filter within 2 days
Steroid Estrogens	1 L	1L amber glass, teflon lid	10mL formalin, 4°C	*
Nitrate+Nitrite-N	125 mL	125 mL poly bottle	H ₂ SO ₄ to pH<2, 4°C	28 days
TSS	1 L	1 L poly bottle	Cool to 4°C	7 days

*holding time not established

The PPCP samples were shipped by FedEx and arrived at SUNYSB the morning after collection. Formalin preservative for the PPCP analysis was added within 24 hours of sample arrival. Samples for conventional parameters were returned to the Ecology Headquarters and picked up by the Manchester Laboratory courier the next day. All samples were accompanied by a chain-of-custody record and the coolers sealed with chain-of-custody tags or tape.

Effluents

Effluent composites from the Sequim and Sunland WWTPs were collected by filling the sample containers with equal amounts of water from morning and afternoon grabs. The grabs were taken with the same glass jars used for PPCPs. The samples were kept on ice and in the dark during the compositing period. Field personnel wore nitrile gloves while doing this work.

pH and conductivity were measured when the grabs are taken. pH was determined with an Orion Model 25A meter and conductivity with a Beckman Model RB-5 conductivity bridge. A Magellan 320 GPS was used to obtain the latitude and longitude of the effluent sampling sites (Appendix A).

Groundwater

Wells selected for sampling were field located on USGS 1:24,000 quad maps. Groundwater levels were measured at each of the study wells prior to sampling, where possible. Water level measurements were made using a calibrated electric well probe or steel tape in accordance with standard USGS methods (Stallman, 1983).

Wells were purged prior to sampling. A peristaltic pump was used for the Sunland monitoring well, which does not have a pump. Domestic wells were purged using the existing pump in the well. Roughly three well volumes were purged before samples are collected. Purge water was discharged to an enclosed flow cell where temperature, pH, conductivity, and dissolved oxygen were monitored and recorded every three minutes using a WTW P4 Multi-parameter field meter. Samples were collected after flow cell measurements stabilized.

Surface Water

Water samples from Bell and Cassalery creeks were collected from the center channel directly into the appropriate sample containers. The samples were put on ice immediately on collection. pH and conductivity were measured as described above for WWTP effluents. Temperature was recorded from a meter. Streamflow was gauged with a Swoffer Model 2100 meter and top-setting rod. A Magellan 320 GPS was used to determine the latitude and longitude of the sampling sites.

Laboratory Procedures

Table 6 shows the laboratory procedures used by SUNYSB in the project.

Table 6. Laboratory Procedures

Parameter	Sample Prep Method	Extraction Method	Analytical Method
Pharmaceuticals	Filter	SPE	HPLC-MS
Steroid Estrogens	Filter	SPE	HPLC-Electrospray-MS
Nitrate+Nitrite-N	na	na	EPA 353.2
TSS	na	an	EPA 2540D

na = not applicable

The pharmaceuticals method is outlined in Kolpin et al. (2002). The compounds are extracted from filtered, one-liter water samples using solid-phase extraction cartridges. The adsorbed compounds are then eluted with methanol. The extract is reduced to near dryness under nitrogen gas and brought to a final volume of 1 mL in acetonitrile. Compounds are separated and measured by High Pressure Liquid Chromatography-Mass Spectrometry (HPLC-MS) in positive ion mode.

The estrogen method is described in Ferguson et al. (2001). Filtered, one-liter water samples are extracted by solid-phase extraction and the resulting extract purified by passing over a selective immuno-affinity extraction column. The only significant change from Ferguson et al. is that time-of-flight MS was used instead of single quadrupole MS, providing additional sensitivity and confirmation based on accurate mass.

Data Quality

The PPCP data reported here is from a wide sweep analysis to get multi-class analyte detection. Unlike the estrogen analysis, where there is a labeled surrogate for each analyte, the PPCP data are corrected for suppression of only a single isotopically labeled standard; more isotope standards have become available recently.

SUNYSB has a high level of confidence in the estrogen data. They consider the PPCP data to be equal or better in quality to what USGS has reported in their national studies (e.g., Kolpin et al., 2002), the most important PPCP studies to date. (Brownawell, 2004).

Metformin appeared to be present in most of the samples for the present study. This compound has poor recovery and does not separate well from the solvent peak on the HPLC. Because of uncertainties associated with its detection, Metformin is reported here as being tentatively identified.

An appreciation for the total variability associated with the data generated for this project can be gained from Table 7 which has results on the replicate effluent and surface water samples. For most of the PPCPs analyzed, the results were similar. In 11 of the 17 instances where a compound was quantified in the replicates, the results agreed within a factor of 2 or better. Five compounds were either detected in one replicate only and/or were below the limit of quantification. For unknown reasons, there was a high degree of variability associated with the Estrone analysis.

A bottle blank, prepared at Manchester Laboratory, was analyzed to detect contamination arising from sample handling procedures. This sample was carried through the field work and treated the same as the other samples. Only Nicotine was detected in the blank; the concentration was 17 ug/L. There were no known sources of Nicotine during the field work. Due to an oversight by the laboratory, the field blank was not analyzed for estrogen compounds.

Table 7. Comparison of Results on Replicate Samples
(PPCPs reported in ug/L, except Estrone and beta-Estradiol in ng/L)

Compound	Sunland WWTP Effluent	Sunland WWTP Effluent – Replicate	RPD*	Bell Creek	Bell Creek - Replicate	RPD*
Acetaminophen	47	nd	--	nd	nd	--
Antipyrine	nd	nd	--	nd	nd	--
Caffeine	30	25	17	nd	nd	--
Carbamazepine	0.8	1.7	74	nd	nd	--
Cimetidine	nd	nd	--	nd	nd	--
Codeine	2.8	2.5	12	nd	nd	--
Cotinine	11	19	55	nd	nd	--
Diltiazem	<LOQ	<LOQ	--	nd	nd	--
Erythromycin	nd	nd	--	nd	nd	--
Fenofibrate	nd	nd	--	nd	nd	--
Fluoxetine	nd	nd	--	nd	nd	--
Hydrocodone	nd	nd	--	nd	nd	--
Ketoprofen	45	42	6	nd	nd	--
Metformin**	150	115	26	11	12	11
Nicotine	25	27	8	25	16	45
Nifedipine	nd	nd	--	nd	nd	--
Paraxanthine	218	194	11	nd	nd	--
Ranitidine	4.3	3.3	29	nd	nd	--
Salbutamol	nd	13	--	nd	nd	--
Sulfamethoxazole	<LOQ	10	--	nd	nd	--
Trimethoprim	14	19	32	nd	nd	--
Warfarin	nd	nd	--	nd	nd	--
Estrone	3.5	0.29	169	0.26	nd	--
beta-Estradiol	nd	nd	--	nd	nd	--

*Relative Percent Difference: range of values as percent of mean value

**tentatively identified

nd = not detected

<LOQ = below the limit of quantification

Results

Effluent and Creek Flows

Table 8 has flow data that pertain to the PPCP samples collected for the Sequim-Dungeness project.

Table 8. Effluent and Creek Flow Rates During PPCP Sample Collection (mgd)

Date (2003)	Sunland Effluent		Sequim Effluent		Cassalery Creek	Bell Creek
	Average	Irrigation Flow	Average	During Sampling		
17-Nov	0.132	0.152	0.588	0.75	1.4	2.9
18-Nov	0.117	0.181	1.06	1.5	--	--

Sunland effluent was collected on November 18, at which time the plant was experiencing a normal flow rate of 0.117 mgd. Effluent is stored in a pond and applied to the sprayfield only during weekdays. Irrigation rates during sampling for the PPCP study were 0.152 and 0.180 mgd.

A rainstorm occurred on November 18, becoming heavy between the morning and afternoon samples at the WWTPs. Although Sunland's flow did not appear to be affected, there was a large increase in the daily average flow for Sequim. Sequim typically has an effluent flow rate of 0.45 – 0.60 mgd compared to an average of 1.06 mgd on the 18th, one of the highest recorded for that facility. The first PPCP sample collected on the 18th was during a flow of approximately 0.75 mgd. The second sample was during a very high flow of approximately 1.5 mgd.

The creeks were sampled the day before the WWTPs. Cassalery Creek was flowing at 1.4 mgd. No other flow data were available for comparison.

Bell Creek was at 2.9 mgd during sample collection, which is close to the historical average of 3.0 mgd. Flow in Bell Creek ranges from 2.1 – 12 mgd (NPDES pre-permit application).

Sunland WWTP and Related Samples

The results for the Sunland WWTP effluent, a Sunland monitoring well, an adjacent private well, and nearby Cassalery Creek are summarized in Table 9. No data was obtained on estrogen compounds in the wells or creek, because the well 5N2 sample was broken in transport. The other two samples were not analyzed due to an oversight by the laboratory.

Two replicate samples of Sunland effluent were collected. Fourteen of the 24 compounds analyzed were detected, with 11 of these being identified in both samples. Concentrations were similar between the replicates, except for Acetaminophen, Sulfamethoxazole, and Estrone.

Almost half of PPCPs detected in Sunland effluent are common, non-prescription substances. Two compounds, Paraxanthine, a caffeine metabolite, and Metformin, used in the treatment of diabetes, were detected at concentrations ranging from approximately 100 – 200 ug/L. As noted previously, Metformin was tentatively identified. Seven compounds were detected at approximately 10 – 100 ug/L. In addition to Caffeine, Nicotine, and Cotinine (a nicotine metabolite), these included the anti-inflammatory Ketoprofen; Trimethoprim, used to treat a variety of bacterial infections; and the pain reliever Acetaminophen.

Five additional compounds - Codeine, Carbamazepine, Ranitidine, Sulfamethoxazole, and Estrone - were present at low concentrations of 10 ug/L or less. Carbamazepine is an antiepileptic, Ranitidine an ulcer drug, Sulfamethoxazole an antibacterial agent, and Estrone is a steroid hormone.

Only Nicotine, Caffeine, and Metformin were detected in Sunland monitoring well 5N2. Concentrations were 14 ug/L, 3.0 ug/L, and 1.0 ug/L, respectively. These and related compounds were among the most abundant in the effluent samples. Effluent concentrations were 2-to-20 times higher than the groundwater.

Except for Nicotine, no PPCPs were detected in the Taylor Ranch Road well. This well lies along the north edge of the WWTP sprayfield, approximately 400 yards from well 5N2.

As noted earlier, Nicotine was detected in the field blank for this project at 17 ug/L. The results reported here (Tables 8-10) have been corrected for the blank. For the Taylor Ranch Road well, the Nicotine concentration was 22 ug/L before blank correction, only slightly higher than the blank. Therefore the detection of Nicotine in this well can largely be attributed to contamination of the sample.

The same compounds identified in the Sunland monitoring well were also detected in Cassalery Creek, i.e., Nicotine, Caffeine, and Metformin. Concentrations were slightly higher than in the monitoring well and ranged from 1.9 – 25 ug/L. In this case Nicotine was substantially higher than the blank, so its presence may not be solely due to sample contamination.

There was some evidence of extremely low concentrations of Codeine, Cotinine, and Sulfamethoxazole in the wells and/or Cassalery Creek, but the concentrations were lower than the limit of quantification. Diltiazem, used in the treatment of angina, may have been present in Sunland effluent, but was also below the quantification limit.

